Running head: Listening difficulties in children

Listening Difficulties in Children with Normal Audiograms: Relation to Hearing and Cognition

Lauren Petley[1,4], Lisa L. Hunter[1], Lina Motlagh Zadeh[1], Nicholette T. Sloat[1], Audrey Perdew[1], Li Lin[1], and David R. Moore[1,2,3]

[1]Communication Sciences Research Center, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

[2]Department of Otolaryngology, College of Medicine, University of Cincinnati

[3]Manchester Centre for Audiology and Deafness, University of Manchester, Manchester, UK

[4]Department of Psychology, Clarkson University, Potsdam, NY, USA

Financial disclosures/conflicts of interest: The research was funded by NIH Grant 5R01DC014078 and Cincinnati Children's Hospital Research Foundation. There are no conflicts of interest, financial, or otherwise.

All correspondence should be addressed to:

David R. Moore, Communication Sciences Research Center, Cincinnati Children's Hospital Medical Center, S1.603 240 Albert Sabin Way, Cincinnati, OH 45229, USA. Email: david.moore2@cchmc.org.

Keywords (5-8): hearing loss, speech-in-noise, listening difficulties, cognition, pediatric

Vignette: *"There's a lot of difference between listening and hearing"* – G.K. Chesterton, Dr Hyde and the White Pillars Murder, 1925.

Abstract

Objectives: Children presenting at audiology services with listening difficulties often have normal audiograms, yet the appropriate approach for the further assessment and clinical management of these children is currently unclear. In this study we assessed listening ability in a large (n = 146) and heterogeneous sample of 6-13 year-old children with normal audiograms using a reliable and validated caregiver questionnaire. The results were related to a multifaceted laboratory assessment of the children's audiological, psychoacoustic, physiological and cognitive abilities. This report is an overview of the study and focuses on the children's behavioral performance. The overall goals of the study are to understand the auditory and other neural mechanisms underlying childhood listening difficulties and to translate that understanding into clinical assessment and, ultimately, intervention.

Design: Cross-sectional behavioral assessment of children recruited through relevant clinics of Cincinnati Children's Hospital, and advertisements placed in the hospital online information service. We aimed recruitment into two demographically-matched groups, a 'listening difficulties' group and a 'typically developing' control group. Caregivers completed the 'ECLiPS' questionnaire and the resulting 'total' standardized composite score formed the basis of further descriptive statistics, univariate and multivariate modeling of experimental data.

Results: All scores of the ECLiPS, the SCAN-3:C, a standardized clinical test suite for auditory processing, and the NIH Cognition Toolbox were significantly lower for children with listening difficulties than for their typically developing peers, using group comparisons via t-tests and

Wilcoxon Rank Sum tests. A similar effect was observed on the LiSN-S test for speech sentence-in-noise intelligibility, but only reached significance for the Low Cue and High Cue conditions, and the Talker Advantage derived score. Stepwise regression to examine factors contributing to the ECLiPS Total scaled score (pooled across groups) yielded a model that explained 42% of its variance based on the SCAN composite, LiSN-S Talker Advantage, and the NIH Toolbox Picture Vocabulary and Dimensional Change Card Sorting scores ($F_{4,95} = 17.35$, $p < 0.001$). High correlations were observed between many test scores including the ECLiPS, SCAN and NIH Toolbox composite measures. LiSN-S Advantage measures generally correlated weakly and non-significantly with non-LiSN-S measures. However, a significant interaction was found between extended high frequency threshold and LiSN-S Talker Advantage.

Conclusions: Children with listening difficulties but normal audiograms have problems with the cognitive processing of auditory and non-auditory stimuli that include both fluid and crystallized reasoning. Analysis of poor performance on the LiSN-S Talker Advantage measure identified subclinical hearing loss as a possible minor contributing factor to talker segregation. Beyond auditory tests, evaluation of children with complaints of listening difficulties should include standardized caregiver observations and consideration of broad cognitive abilities.

Introduction

Most people have listening difficulties (LiD) at one time or another, and some people have LiD continuously. While these difficulties are often accompanied by audiometric hearing loss, a substantial proportion of both children (Hind et al., 2011) and adults (Parthasarathy et al., 2020) attending audiology clinics are found to have normal audiometric sensitivity. Nationally within the US it has recently been estimated that about 80 million adults have LiD, of whom half have hearing loss, while the other half have clinically normal hearing (Edwards, 2020).

Individuals with LiD commonly have difficulty understanding speech in challenging situations such as noise, but also have other symptoms that can include difficulty following instructions and understanding rapid or degraded speech (Jerger & Musiek, 2000). Such tasks require not only auditory processing, but also speech decoding and the successful deployment of selective attention towards task-relevant stimuli (Moore et al., 2010). Both childhood and older age are marked by changes in attention, with the maturation of attention networks persisting through adolescence (Rueda & Posner, 2013), and losses that may be specific to dual set maintenance later in life (Verhaeghen & Cerella, 2002). A host of other top-down cognitive (Davis & Johnsrude, 2007) and bottom-up auditory (B. C. J. Moore, 2012) abilities underlie speech perception and, in almost all of these abilities that have been studied, children generally perform more poorly than adults (Moore et al., 2011; Rueda & Posner, 2013). Listening in individual children occurs along a continuum, as evidenced by caregiver questionnaires, including the Children's Communication Checklist (CCC-2; Bishop, 2006; Moore et al., 2010) and the Evaluation of Children's Listening and Processing Skills (ECLiPS; Barry & Moore, 2014; Barry et al., 2015; Roebuck & Barry, 2018). For some children, LiD are problematic because they impair communication to a degree that substantially delays their social, academic

and personal development relative to their peers.

Hearing and listening are often used to mean the same thing, but Merriam-Webster defines hearing as the "sense by which noises and tones are received as stimuli", while listening means "to pay attention to sound". Following these definitions, by 'hearing' we mean sound-evoked activation of the ear and central auditory nervous system (CANS), usually associated with, but sometimes independent of awareness. 'Listening' may be described as the conscious perception of sound, which is dependent on attention and is influenced by abilities including memory, executive function, emotion and, as above, linguistic processes. Crucially, it is possible to hear several speech sources simultaneously, but listen to only one (Cherry, 1953). Listening necessarily involves higher areas of brain function, beyond the CANS, for example the "posterior hot zone", a broad region of the posterior parietal, occipital and temporal cortex that appears to be the common locus of activation during different modes of conscious experience (Koch, 2018; Koch et al., 2016). The primary auditory cortex, in contrast, does "not directly contribute to the content of auditory . . . experience" (Koch, 2018). Despite the prevalence of LiD without hearing loss, listening is poorly understood clinically. For example, much of the controversy surrounding the concept of 'auditory processing disorder' (APD) centers on whether LiD in children is due to a 'hearing' or to a 'listening' problem (Moore, 2018).

There are various explanations for LiD in children. The most commonly-recognized cause is hearing loss, as typically indexed by pure-tone insensitivity. Recent large-scale studies have shown that children with even mild hearing loss (better ear pure tone average, PTA >20 dB HL) have reduced language skills, leading to impaired learning (Ching et al., 2018; Moeller & Tomblin, 2015; Moore et al., 2020). However, as above, there is also considerable evidence that children with normal PTA hearing can also have LiD (de Wit et al., 2016; Moore et al., 2010). In

the broader field of learning disabilities (e.g. developmental language disorder, reading disorder),

this is currently assumed to mean those children have 'normal hearing' and that their problems

lie outside the auditory system. Yet the current clinical definition of 'normal hearing' does not

guarantee the absence of peripheral pathology. Several forms of sub-clinical hearing loss have

been identified, including minimal or slight hearing loss (e.g. PTA = 15 – 20 dB HL) (Bess et al.,

1998; Moore et al., 2020), extended high frequency hearing loss (Hunter et al., 1996; Motlagh

Zadeh et al., 2019), auditory synaptopathy (Kujawa & Liberman, 2019), and reduced spectral or

temporal resolution of cochlear origin (Oxenham & Bacon, 2003).

Impaired function of the CANS is another possible source of LiD. For example,

reversibly impaired binaural interaction and spatial hearing has been found consequent on earlier

otitis media (Moore et al., 1991; Pillsbury et al., 1991; Whitton & Polley, 2011) and ear canal

atresia (Wilmington et al., 1994) in children. The impairment is considered to be 'central'

because it depends on binaural interaction, which only occurs in the brain, and because it can

remain after the peripheral pathology resolves, leaving the ears audiometrically and

tympanometrically normal. Otitis media earlier in life is also thought to underlie reduced spatial

unmasking, termed 'spatial processing disorder' (SPD; Cameron et al., 2014, 2015; Graydon et

al., 2017), as indexed by the Listening in Spatialized Noise – Sentences (LiSN-S) test. Spatial

unmasking is a well-documented phenomenon in which spatially separating a target sound from

a masking or competing sound renders the target more intelligible. The neural mechanisms of

SPD remain unclear, and studies reporting SPD have not ruled out sub-clinical hearing loss.

Reduced cognitive function is another possible mechanism of LiD, as introduced above.

For example, executive control is crucial for establishing sustained goal-directed attention and

dynamically switching attention based on task demands (Petersen & Posner, 2012). Working

memory supports speech comprehension, particularly under difficult listening conditions (Rudner & Rönnberg, 2008), and is associated with performance on even simple, non-speech hearing tests, such as frequency discrimination (Banai & Ahissar, 2004). Thus, reduced scores on those tasks in children may reflect either or both sensory and cognitive impairment (Moore et al., 2010).

The primary purpose of a large, US National Institutes of Health funded, longitudinal research project, of which this is one of the first experimental reports, is to advance the understanding of APD and to provide a basis for improvements in the diagnosis and management of children with LiD. Here, we present baseline behavioral data from the first full analysis of a large group of children with mixed listening abilities, as reported by their caregivers on the well-validated Evaluation of Children's Listening and Processing Skills (ECLiPS) questionnaire (Barry & Moore, 2014; Barry et al., 2015; Roebuck & Barry, 2018). The primary aim of this study was to examine the relationship between hearing, listening and cognitive performance through univariate group comparisons and multivariate statistical modeling using quantitative metrics from a battery of auditory and cognitive behavioral tests. Based on previous research, we predicted that these analyses would demonstrate broad differences between LiD and TD groups across demographic, cognitive, 'auditory processing' and speech-in-noise factors, and that all of these factors would be represented in the regression model predicting listening ability.

Materials and Methods

This study was approved by the Institutional Review Board of Cincinnati Children's Hospital (CCH) Research Foundation.

Participants

A total of 166 participants (74 with LiD, 92 typically developing - TD) was enrolled; 20 withdrew or otherwise exited the study (7 LiD, 13 TD). The remaining 146 participants ranged from 6.0 to 13.7 years of age at enrolment. Age is summarized along with other demographic variables (sex, race and maternal education) in Table 1. Due to attrition and updates to testing procedures, the sample size for each assessment was variable (see Results). All participants had normal hearing, bilaterally, defined as pure tone thresholds ≤ 20 dB at all octave frequencies between 0.25 – 8 kHz.

----------------------- Insert Table 1 here -----------------------

Procedure

Participants with LiD were recruited initially from a medical record review study of over 1,100 children assessed for Auditory Processing Disorder (APD) at Cincinnati Children's Hospital. Of those children, 179 were diagnosed by an audiologist as having an auditory processing "disorder" and 364 as having an auditory processing "weakness" (Moore et al., 2018). Caregivers of children in either of those categories who fit our inclusion criteria and responded positively to an invitation to participate in this study were sent a pack of three short questionnaires. Those who chose to complete and return the questionnaires were invited to bring their child into the lab for a study visit. Over time, recruitment efforts were expanded to include children both with and without LiD, and the use of IRB-approved materials, advertising, and messages via print, electronic, social and digital media at hospital locations and in the local and regional area. Flyers were posted in relevant hospital clinics (i.e., Audiology, Developmental and Behavioral Pediatrics, Speech/Language Pathology) and sent out via email to all hospital employees and families interested in research.

Interested caregivers completed eligibility screening procedures for their child by phone, or by paper or electronic questionnaires. Electronic questionnaires were delivered via a secure, unique link to REDCap, an electronic survey administration and database platform. Those who opted-in completed (i) a consent form, (ii) a medical and educational history 'Background' questionnaire, (iii) the ECLiPS questionnaire (Barry & Moore, 2014), and (iv) the Children's Communication Checklist, 2nd Edition (CCC-2; Bishop, 2006). If the child was deemed eligible by study staff, the caregiver was invited to schedule an initial visit.

Prior to completion of any other study-related procedures, caregivers reviewed the informed consent form with a study staff member and discussed the purpose, procedures, risks and benefits, duration, and expectations involved in the study. Children aged 11 and above were also assented using a child-friendly version of the consent document, per institutional policy.

About 70% (n=104) of the study sample scheduled individual visits. Most participants completed at least three separate visits, approximately 3 hours long each. The time between visits for the full sample varied an average of 18.6 days, but as some study procedures were added several months to a year after the start of the study, some participants returned to complete the series of visits several months apart over a 2-year period. Consequently, all age-related measures were adjusted for age at the time of testing (resolution = 0.1 years) rather than at enrolment. Hearing thresholds were re-checked if the participant returned more than six months following their most recent audiogram. To maintain engagement between visits, participants were sent handwritten birthday and holiday cards throughout their enrolment. Caregivers also received periodic email communications about updates on the study website and newsletters about study and lab activities. Caregivers were encouraged to access the website, which was built as a tool to

provide additional information about the study including interim results, staff members, and testing procedures.

As an alternative to scheduling individual visits, participants were offered the opportunity to attend a "summer camp" session, and approximately 30% of participants completed initial ('Wave 1') testing in this way (n = 44). Two-day camp sessions were offered throughout the summer over the 2-year period of data collection. Six children were invited to attend each camp session and six camps were held each year, for a total of 36 available camp spots per year. On the first day of camp, caregivers completed the informed consent process and then left for the day and returned to pick up their child in the afternoon. Over the two camp days, participants spent 8-10 hours completing study procedures and the remaining 5-7 hours playing board games, having lunch and snacks, watching movies, and doing arts and crafts with other campers and the study staff. Participants rotated between 5-6 testing stations (sound booths, quiet testing rooms) with study staff to complete all of the different procedures. Participants also received payment via ClinCard, a pre-paid debit card reloaded at the end of each appointment. Overall, participants reported that they enjoyed participating in the study via summer camp sessions and expressed interest in returning for future sessions.

Materials

Background questionnaire

This questionnaire was used to screen potential participants for study eligibility and gather background information. It asked about the participant's medical history, parental education, and demographic information. Key variables were date of birth, language, race and ethnicity, maternal education, history of ear and hearing problems (including pressure equalization, PE, tube placement), diagnosis of, or treatment for, learning problems (attention

disorders, developmental delays, speech-language disorders), neurological (e.g. history of head

trauma) or psychiatric conditions, school interventions, and birth history (prematurity, NICU

stay). Requirements for eligibility in the LiD group included age between 6 and 13 years old

upon enrolment, English native language, and no neurologic, psychiatric or intellectual (IQ < 80)

condition that would prevent or restrict the ability to complete testing procedures. Participants

were evaluated on a case-by-case basis to determine their ability to participate in the study. TD

participants were eligible based on the same criteria, with the addition that they could not have a

diagnosed developmental delay, attention or other learning disorder. Demographic variables

including maternal education, age, gender, and race, obtained from this questionnaire, are

summarized in Table 1 for the listening difficulty (LiD) group and their peers (TD).

Caregiver evaluation of children's listening (ECLiPS)

The ECLiPS (Barry & Moore, 2014) profiles children's listening and communication

abilities. It contains 38 simple statements (items) describing behaviors commonly observed in

children. Caregivers were asked to rate how much they agreed with each statement on a five-

point scale ranging from strongly disagree to strongly agree. The ratings were averaged to derive

scores, which were scaled by age, on five subscales (speech & auditory processing, SAP;

environmental & auditory sensitivity, EAS; language/ literacy/ laterality, LLL; memory &

attention, M&A, and pragmatic & social skills, PSS) each containing 6-9 distinct items. These

scales are further collected under Language, Listening, Social, and Total aggregate (composite)

scores. All scales and composites were standardized for a population mean of 10 (s.d. = 3) based

on British data (Barry et al., 2015). A secondary aim of this study was to provide initial US

normative data for the ECLiPS and to compare them with the British norms.

Caregiver evaluation of children's communication (CCC-2)

The CCC-2 (Bishop, 2006) asks caregivers to respond to 70 items relating to the child's communication skills. Subscales include speech, syntax, semantics, coherence, initiation, scripted language, context, nonverbal communication, social relations, and interests. The items are combined to create two composite scores, a General Communication Composite (GCC) and the Social Interaction Difference Index (SIDI). The GCC is a scaled composite of the scores on 8 of the 10 subscales, excluding subscales for social relations and interests.

Audiometry

All participants were screened for normal hearing at standard, octave intervals from 0.25 - 8 kHz bilaterally. All except 27 (5 LiD, 22 TD) participants additionally completed threshold audiometry at the extended high frequencies (EHFs) of 10, 12.5, 14 and 16 kHz using an Equinox audiometer (Interacoustics, Inc.) and the Hughson-Westlake adaptive method (American Speech-Language-Hearing Association, 2005). Early in data collection, insert earphones (E-A-R TONE GOLD 3A) were used for standard frequencies and Sennheiser HDA 300 circumaural headphones for EHFs. For most children (n = 114; Hunter et al., 2020), the Sennheiser headphones were used for all frequencies. Participants were tested alternately with each ear first. They were instructed to respond when they heard a tone by pressing a response button or raising one hand. Those with any elevated thresholds (> 20 dB HL) in the standard frequencies were excluded and were referred for further clinical care as appropriate. Participants were re-tested if they returned > 6 months after previous testing.

'Auditory Processing Disorder' (SCAN-3:C)

Participants were assessed in a private sound attenuated booth or quiet room using the SCAN-3:C (Keith, 2009), a standardized test of auditory processing for children aged 5-12 years

(Emanuel et al., 2011). Stimuli were presented via a laptop PC through Sennheiser HD 215 headphones. The four diagnostic subtests of the SCAN-3:C are low-pass filtered words (FW), auditory figure-ground (AFG+8), competing words – directed ear (CW-DE), and competing sentences (CS). For FW, participants are asked to repeat low-pass filtered, monosyllabic words in quiet. AFG+8 is a speech-in-noise test requiring repetition of unfiltered monosyllabic words against multi-talker speech (the 'noise') at a fixed +8 dB signal/noise ratio (SNR). In CW-DE, two different monosyllabic words are presented simultaneously, one in each ear (i.e. dichotically), and participants report them back either left or right ear first. Similarly, for CS, unrelated sentences are presented to the left and right ears but, in this case, participants repeat only the sentence heard in one, directed ear per trial. A complete SCAN battery results in age-scaled scores for each subtest as well as a standardized composite.

Speech hearing in noise (LiSN-S)

The Listening in Spatialized Noise – Sentences (LiSN-S) test (Brown et al., 2010; Cameron & Dillon, 2007; Phonak/NAL, 2011) measures the ability to listen and repeat simple, spoken sentences in the presence of informational masking, developed using the same criteria as the BKB sentences (Bench et al., 1979). The LiSN-S (US Edition; Brown et al., 2010) was administered using a commercial CD played on a laptop (Phonak/NAL, 2011), a task-specific soundcard, and Sennheiser HD 215 headphones. In the LiSN-S, participants are asked to repeat a series of target sentences ('T'), presented directly in front (0°; diotic), while ignoring two distracting speakers ('D1', 'D2'). There are four listening conditions, in which the distractors change voice (different or same as target) and/or position (0° and 90° virtual positions relative to the listener using generic head-related transfer functions; Humanski & Butler, 1988). The test is adaptive; the level of the target speaker decreases or increases in SNR relative to the distracting

speech if the listener responds correctly or incorrectly during up to 30 sentences in each

condition. The 50% correct SNR is either the Low cue speech reception threshold (SRT; same

voice, 0° relative to the listener) or the High cue SRT (different voice, 90° relative to the

listener). The three derived scores of the LiSN-S are the Talker Advantage, Spatial Advantage,

and Total Advantage, so-called because each is the difference between SRTs from two

conditions. This subtraction process should, to some extent, separate auditory from cognitive

influences (Moore & Dillon, 2018), as discussed below.

Cognition (NIH Toolbox)

Each participant's cognitive skills were assessed using the NIH Toolbox Cognition

Domain battery of tests (Wientraub, 2013). Participants completed testing online or via an iPad

app, in accordance with current Toolbox recommendations, in a private sound-attenuated booth

or quiet room. The battery contains up to eight standardized cognitive instruments measuring

different aspects of fluid or crystallized reasoning. The precise composition of the testing battery

is dependent on user choice and participant age.

All participants in this study completed the Picture Vocabulary test (PVT), Flanker

Inhibitory Control and Attention test (Flanker), Dimensional Change Card Sort test (DCCS), and

Picture Sequence Memory test (PSMT). Each test produced an age-corrected standardized score

and the scores of all four tests were combined to calculate a single Early Childhood Composite.

The PVT is an adaptive test in which the participant is presented with an audio recording of a

word and selects which of four pictures most closely matches the meaning of the word. In the

Flanker, which tests inhibition/attention, the participant reports the direction of a central visual

stimulus (left or right, fish or arrow) in a string of five similar, flanking stimuli that may be

congruent (same direction as target) or incongruent (opposite direction). The DCCS tests

cognitive flexibility (attention-switching). Target and test card stimuli vary along two dimensions, shape and color. Participants are asked to match test cards to the target card according to a specified dimension that varies for each trial. Both the PVT and DCCS score accuracy and reaction time. PSMT assesses episodic memory by presenting an increasing number of illustrated objects and activities, each with a corresponding audio-recorded descriptive phrase. Picture sequences vary in length from 6-18 pictures depending on age, and participants are scored on the cumulative number of adjacent pairs remembered correctly over two learning trials.

Additional NIH Toolbox Cognition tests were administered to all children 8 years of age and older. For the Fluid Composite measure, we used the list sorting working memory (LSWM) and the pattern comparison processing speed (PCPS) tests, in addition to the DCCS, Flanker, and PSMT, as above. LSWM assesses working memory by asking participants to arrange objects presented visually and auditorily (food and animals) in order of size. PCPS requires participants to respond as quickly as possible to whether two visually presented cards are the same or different. For the Crystallized Composite measure, participants completed the Reading Recognition (RR) test in addition to the PVT. The RR requires participants to read aloud words and letters accurately. A final Toolbox measure that was not part of a composite score, the Rey Auditory Verbal Learning (AVL) test measures verbal episodic memory as a supplement to the visual PSM test.

Analysis

The primary analysis was divided into three sequential parts. Part 1 examined subgroup differences between children with LiD who had a formal diagnosis of APD, including auditory processing 'weakness', (Dx subgroup; Moore et al., 2018) and children with listening difficulties

but no formal diagnosis (noDx subgroup). The purpose of this analysis was to determine whether these two subgroups could be treated as a homogeneous LiD group for subsequent analyses. These subgroups were tested for differences on the ECLiPS Total scaled score, SCAN-3:C composite scaled score, LiSN-S Talker Advantage, Spatial Advantage, and Low Cue scores, and the NIH Cognition Toolbox Fluid, Crystallized, and Early Childhood Composites, as well as the AVL test. A potential influence of maternal education was first explored via a separate two-way ANOVA for each of these variables with the factors diagnosis (Dx, noDx) and maternal education level (collapsed into two groups: graduated high school or less, and some college or more). In no case was there a main effect of, or an interaction with, maternal education ($p >$ 0.05). All group-level comparisons for the Dx versus noDx subgroups were therefore conducted using two-tailed, two-sample Student's t-tests, when the assumption of normality was met, and Wilcoxon Rank-Sum tests in all other cases.

Part 2 of the analysis examined group differences between TD children and children with LiD (combined Dx subgroups) on all of the same measures used for Part 1. As in Part 1, a potential influence of maternal education was first explored via separate two-way ANOVAs, this time with the factors listening difficulties (TD, LiD) and maternal education level (levels as above). In no case was there a main effect of, or an interaction with, maternal education ($p >$ 0.05). All group-level comparisons between LiD and TD groups were therefore assessed via two-tailed, two-sample Student's t-tests. To reduce the risk of alpha inflation through multiple comparisons, only descriptive statistics are provided for the remaining measures (e.g. subtests contributing to composite scores).

Part 3 of the analysis aimed to identify the functional domains that contribute to LiD across all participants by using stepwise multiple regression to predict ECLiPS Total scaled

scores. Candidate predictors for this analysis included only assessments that were available across the full age range of the study sample. As discussed above, some tests of the NIH Cognition Toolbox did not meet this criterion. Candidate predictors included maternal education level, race, SCAN-3:C Composite Score, four LiSN-S scores (Low Cue, High Cue, Talker Advantage, Spatial Advantage), and four NIH Cognition Toolbox tests (DCCS, Flanker, PST, PVMT). To explore a possible role of extended high frequency hearing thresholds, for which main effects were examined in a recent paper (Hunter et al., 2020), four interactions were included as candidate predictors. Three interactions with LiSN-S scores (Low Cue, Talker Advantage, Spatial Advantage) explored a possible contribution from extended high frequency thresholds to speech-in-noise performance in complex environments. In light of the known benefit that linguistic proficiency provides under difficult listening conditions (Kaandorp et al., 2016), an interaction with the NIH PVT was also explored to examine the possible role of language skills in mitigating the influence of extended high frequency hearing loss. In all four interactions, extended high frequency hearing thresholds were computed as a single average across both ears and all frequencies (10, 12.5, 14, and 16 kHz).

All candidate predictors for the stepwise multiple regression (Part 3) were first examined against the ECLiPS Total scaled score via univariate analyses. Continuous candidate predictors were examined using Spearman's Rho, while categorical predictors were examined via Student's t-tests. Variables with $p < 0.1$ in the univariate analysis were included in the stepwise regression. Collinearity among these variables was examined using the Variance Inflation Factor, which ranged from 1.31 to 3.01, indicating no collinearity. Statistical comparison of participants with complete and incomplete data across all demographic variables and candidate predictors yielded no significant differences between the two groups. It was therefore assumed that data were

missing at random. All data were included in the stepwise regression via the use of SAS v9.4

(The SAS Institute, Cary, NC) PROC MIXED with restricted maximum likelihood estimation.

Additional, secondary analyses examining correlations between factors other than the

ECLiPS were used to follow up on results revealed by the primary analysis. Multiple Spearman's

Rho correlations were used.

Results

Diagnosis of APD

For Part 1 of the analysis, among children with LiD, those with a diagnosis of APD (n =

14-20, depending on test) did not differ significantly ($p > 0.05$) from those without a diagnosis (n

= 25-44), with the exception of two variables. The exceptions were both cognitive verbal scores,

the NIH Crystallized Composite and the AVL. For the Crystallized Composite, a Wilcoxon

Rank-Sum test indicated that children with a diagnosis of APD had lower scores (mean = 88.0,

SD = 15.2) than children lacking a diagnosis (mean = 94.47, SD = 10.36; S = 347.5, $p = 0.026$).

Similarly, the AVL scores were significantly [$t(39) = -2.55$, $p = 0.015$] lower for children with an

APD diagnosis (mean = -2.47, SD = 2.14) than for those without (mean = -0.85, SD = 1.87).

Further *t*-tests found that both subgroups (Dx, noDx) performed significantly worse than the TD

group on both of these verbal measures ($p < 0.005$).

Effects of LiD

All measures except the LiSN-S Spatial and Total Advantage scores showed significantly

poorer performance for the LiD group than for the TD group (Table 2; Figs. 1-3). By design, the

mean ECLiPS Total scaled score was much lower [$t(141.03) = 22.67$, $p < 0.001$] for children in

the LiD group than for those in the TD group (Fig. 1). A repeated-measures ANOVA within the

TD group demonstrated that those children had similar mean standardized scores across

subscales and the Total score [$F(4, 312) = 1.18$, $p = 0.32$], as would be expected if British and

US samples were drawn from the same population. An identical ANOVA performed within the

LiD group, by contrast, yielded significant differences amongst the ECLiPS subscale scores,

which remained significant following Greenhouse-Geisser correction [$F(3.12, 206.2) = 15.94$, p

< 0.001]. Note that, among the ECLiPS subscales, the SAP scale reflects the profile statements

that are most closely aligned with listening. SAP ratings of children in the LiD group were all

within the lowest 30[th] percentile based on the test's standard (UK) normalization, while at least

one child in the LiD group scored in the top 25[th] percentile on each of the other subscales.

Descriptive statistics for the ECLiPS and CCC-2 are provided in Table 2 for the TD and LiD

groups. Outcomes for the GCC correlated highly with those for the ECLiPS total score [$r_s(143) =$

0.75, $p < 0.001$], while the SIDI did not [$r_s(143) = -0.060$, $p = 0.477$]. ECLiPS subscales and

their relation to other measures will be considered further in a separate report.

----------------------- Insert Figure 1 here -----------------------

        Scores on the SCAN-3:C composite scale were lower [$t(126) = 7.44$, $p < 0.001$] for the

LiD group than for the TD group. Details, including descriptive statistics on the SCAN-3:C

subscores are shown in Table 2 and Figure 2A. Further *t*-tests for group differences on each of

the SCAN-3:C subtests revealed significant differences between the LiD group and their TD

peers on all measures ($p < 0.005$ in all cases). Results obtained on the composite measure thus

reflect all functional domains tested by the SCAN-3:C and the subtests were not further explored

in the primary analysis.

----------------------- Insert Table 2 here -----------------------

For the LiSN-S, both the Low Cue [$t(138) = 3.32$, $p = 0.001$] and Talker Advantage [$t(138) = 3.22$, $p = 0.002$] were significantly lower for children with listening difficulties. The Spatial Advantage score demonstrated a nonsignificant trend in the same direction ($p = 0.10$). Scores on the LiSN-S are summarized by group in Figure 2B and Table 2. Note that Spatial Advantage had more variance, particularly on the lower scoring side, than the other measures. A follow-up analysis of the LiSN-S used a spatial 'Pattern Score', developed by Cameron and Dillon (2011) as a quantitative clinical measure of the benefit of adding virtual spatial cues to the information in the Low Cue condition of the LiSN-S (i.e., target and distracting stimuli presented diotically). Pattern Scores did not differ significantly between the LiD (mean = 7.25 dB) and TD (mean = 7.29 dB) groups. However, 7 children in the LiD group and 5 children in the TD group had Pattern Scores within the range to diagnose a 'spatial processing disorder' (Cameron & Dillon, 2008; see Discussion).

----------------------- Insert Figure 2 here -----------------------

As can be seen from the Fluid Composite [$t(102) = 7.64$, $p < 0.001$], Crystallized Composite [$t(115) = 7.69$, $p < 0.001$], Early Childhood Composite [$t(102) = 7.64$, $p < 0.001$], and Rey AVL test [$t(72.03) = 5.38$, $p < 0.001$], scores in the LiD group were considerably lower than those for the TD group on all domains of the NIH Cognition Toolbox (Table 2, Fig. 3). Group-level differences were also present for all of the individual tests contributing to the composite scores, including the Rey AVL test, not shown in Figure 3 ($p < 0.001$ in all cases).

----------------------- Insert Figure 3 here -----------------------

Four test scores predicted parent-reported listening skills

Spearman's Rho and Student's *t*-test results for each candidate predictor are summarized in Table 3 for all candidate predictors that demonstrated univariate association with ECLiPS

Total Score with $p < 0.1$. Notable among these was a specific, significant interaction between the

LiSN-S Talker Advantage and EHF threshold. One hundred participants had complete datasets

across all these predictors. Following stepwise regression, the final model explained 42% of the

variance in ECLiPS Total scaled score ($F_{4,95} = 17.35$, $p < 0.001$). This model had an intercept of

-14.15 (SE = 3.10) and included four predictors: the SCAN-3:C composite score, the LiSN-S

Talker Advantage score, and the NIH Picture Vocabulary and Dimensional Change Card Sorting

scores. Table 3 includes the regression coefficients, standard errors, t- and p-values for all of

these predictors. Figure 4 compares actual ECLiPS Total scaled scores to those predicted by this

model, where perfect predictions lie along the diagonal. As expected, the separation between

groups was preserved in the predicted scores, though predicted scores tended to be higher than

actual scores for the LiD group and lower for the TD group.

----------------------- Insert Table 3 here -----------------------

----------------------- Insert Figure 4 here -----------------------

Correlations between measures

In the secondary analysis, pairwise correlations were computed between all variables,

treating each as a continuous predictor across groups (i.e. not separated into LiD or TD). These

correlations are shown in Tables 4 and 5. The data in Table 4 show that many tests were highly

correlated with the ECLiPS, SCAN and NIH Toolbox composite measures. In contrast, the

LiSN-S Advantage measures correlated poorly with most other, non-LiSN-S measures. The

SCAN composite correlated to a highly significant level with the Low and High Cue measures of

the LiSN-S, but only weakly and non-significantly with the Spatial and Total Advantage

measures. Talker Advantage, however, did correlate significantly with the SCAN Composite.

Correlations between LiSN-S and SCAN sub-tests (Table 5), showed that the Filtered Words and

dichotic (Competing Words and Competing Sentences) tests, and the Low and High Cue

measures accounted for almost all of the relationship between SCAN and LiSN-S. Talker

Advantage was marginally related to Competing Sentences and Spatial Advantage to Competing

Words. However, when corrected for multiple comparisons (Bonferroni, $p < 0.01$), these

relatively weak relationships became non-significant.

----------------------- Insert Table 4 here -----------------------

----------------------- Insert Table 5 here -----------------------

Discussion

In this study, we found that children with LiD, identified primarily on the basis of the

ECLiPS caregiver questionnaire, had impaired performance on a broad range of auditory and

cognitive tasks relative to age-matched TD children. Tasks included a suite of tests normally

used for diagnosing APD, a test for repeating sentences presented against a background of

distracting sentences, and a battery of tests measuring fluid and crystallized cognition.

Given normal peripheral function (Hunter et al., 2020), the pattern of deficits observed

here with LiD could reflect central auditory (e.g., Cameron et al., 2014, 2015; Graydon et al.,

2017; Moore et al., 1991; Pillsbury et al., 1991) or general cognitive (e.g., Moore et al., 2010;

Moore, 2018) deficits, or a combination of both. For this reason, the protocol that was developed

for our research program cast a wide net, to provide a relatively detailed characterization of the

constellation of features that can accompany this complex clinical construct.

Diagnosis of APD

A crucial first step in the analysis was to identify any inhomogeneities in our sample of

children with caregiver-reported LiD (as reported on the ECLiPS), some of whom had received

clinical diagnoses of APD (Dx), and some of whom had not (noDx). Comparisons between these two subgroups yielded no differences on the ECLiPS Total scaled score, SCAN-3:C composite scaled score, LiSN-S scores (Talker Advantage, Spatial Advantage, and Low Cue scores), or the Fluid Composite and Early Childhood Composite scores of the NIH Cognition Toolbox. However, the Dx subgroup performed significantly more poorly on the NIH Crystallized Composite score and the Rey AVL test. These results suggest that children in the Dx subgroup had more severe problems than those in the noDx subgroup specifically in the cognitive domain.

Nature of listening difficulties

Comparisons between children with LiD and their TD peers yielded differences across nearly all tested domains of auditory and cognitive function, including the ECLiPS Total scaled score, SCAN-3:C composite scaled score, and all of the NIH Cognition Toolbox composite scores (Fluid, Crystallized, Early Childhood), as well as the Rey AVL test. Several of these tests may tap the same underlying deficit. To disentangle such overlapping contributions, a multivariate regression approach was used. The regression successfully explained 42% of the variance in ECLiPS Total scaled scores using four predictors: the SCAN-3:C composite score, the LiSN-S Talker Advantage score, and the NIH Picture Vocabulary and Dimensional Change Card Sorting scores. These findings suggest that both auditory and cognitive factors make unique, significant contributions to predicting listening skills as captured on the ECLiPS. However, we reason below that just one measure, the LiSN-S Talker Advantage, provided some limited evidence of impaired auditory processing.

ECLiPS

It is unusual in hearing research for a case-control study such as this to use a subjective assessment, the ECLiPS, as an independent variable. However, that is an accepted clinical

practice in the diagnosis of other neurobehavioral disorders, for example developmental language disorder (Bishop & McDonald, 2009) and attention deficit hyperactivity disorder (Pediatrics, 2011). We suggest that a parent or guardian is usually in the best position to judge their children's behavior, in this case, how their child responds to the challenges inherent in everyday communication. Several aspects of the ECLiPS results assured us that this was a reasonable decision. First, when caregivers responded to our advertisements, there was a high probability that their child's ECLiPS score would place them into the group identified by the parent (LiD or TD). Second, we were impressed that mean and variance standardized scores placed this mid-Western US TD sample closely alongside the English normalization data on which the standard scores were derived (Barry et al., 2015). Third, for the children with LiD, the distribution of ECLiPS scores between the various sub-scales seemed consistent with predominantly receptive speech and language deficits. Whether those deficits are of an auditory, cognitive or mixed origin is not directly addressed by the ECLiPS.

Because of the modeling applied to these data, it was possible to observe systematic disparities between caregiver reports and objective measures. To the extent that objective measures capture everyday listening skills, caregivers of children in both groups tended to express a more extreme view of their children's abilities than the objective data suggested. However, this does not imply that the objective data are more meaningful.

SCAN-3:C

For many years test batteries such as the SCAN (Keith, 2009) have been used to diagnose APD (American Academy of Audiology, 2010; American Speech-Language-Hearing Association, 1996). These batteries test aspects of listening purported to be mediated by central auditory processing, for example 'closure', the perceptual capacity to fill in missing or distorted

parts of an auditory stimulus, and 'binaural integration' (Moncrieff, 2006), the ability to identify different words simultaneously presented to each ear. According to the SCAN manual, closure is represented by the Filtered Words subtest and binaural integration by the Competing Words – Directed Ear subtest. In this study, SCAN composite scores differentiated between the two groups so successfully that the SCAN composite was one of the four measures left in the final multivariate regression that explained nearly half the total sample variance on the ECLiPS. It was thus a strong predictor of LiD.

LiSN-S

We chose the LiSN-S (Cameron & Dillon, 2007) in this study because it is a multi-faceted test of spatial and speech hearing, and was designed specifically for pediatric assessments. It also features 'derived' (also called 'difference' or 'subtraction') testing that we and others have argued can separate cognitive and sensory contributions to auditory perception when comparing performance within and between listeners (Dillon et al., 2014; Moore et al., 2010; Moore, 2012; Moore & Dillon, 2018). For the LiSN-S, performance on individual conditions, for example the Low Cue and High Cue SRT, reflects both the ability to hear the target sentence against the distracting sentences *and* the ability of the listener to attend selectively to the target, to decode the speech signal, to remember the words in the target sentence, and to repeat each sentence orally. In contrast, for the three derived Advantage measures (Talker, Spatial and Total), performance reflects only the ability to hear the target sentence, assuming that the attention, decoding, memory, linguistic, and reproduction aspects of the task are identical for each of the two contributing individual measures from which each Advantage measure is derived. This assumption appears to be only partly fulfilled for the LiSN-S. The Spatial Advantage relies only on the spatial separation of the distracting talkers, a

physical acoustic manipulation. However, the Talker Advantage employs different individual talkers as the distractors, introducing a decoding, a linguistic and, possibly, an attention difference, as well as an acoustic difference between the underlying test conditions.

While Low and High Cue SRTs were higher (i.e. poorer) in the LiD than in the TD group, among the Advantage scores, only the Talker Advantage was significantly poorer in the LiD group, while the Spatial Advantage showed a non-significant trend in that direction. These results are surprising in two respects. The poorer Talker Advantage represents the first and only example of which we are aware of a group difference in this measure in any study (e.g., Cameron & Dillon, 2008). Furthermore, despite the analysis in the previous paragraph, the finding of a significant deficit of the LiD group on a derived measure of hearing may still be direct evidence for impaired auditory system function. This possibility is considered further below in the context of correlations between LiSN-S thresholds, SCAN and EHF measures.

It has been suggested that deficits in the LiSN-S spatial measures are the basis for a specific and treatable disorder termed spatial processing disorder (SPD; Cameron et al., 2011, 2012, 2014; Cameron & Dillon, 2008) that is diagnosed using the LiSN-S Pattern Score. Here, we found evidence for SPD among both LiD and TD groups of children. However, neither the proportion of children with SPD nor the Pattern Score differed significantly between groups, suggesting that SPD is not predictive, and may not be representative, of the listening problems identified by the ECLiPS.

NIH Cognition Toolbox

One of the most consistent and largest deficits experienced by the children with LiD in this study was broadly-specified cognitive function, as also recognized in other recent studies (Moore et al., 2010; Seeto et al., in press; Tomlin et al., 2015). Scores on individual tests were

highly correlated and uniformly reduced relative to TD children, suggesting that cognitive

function is a major contributor to LiD. Both the Picture Vocabulary and Dimensional Change

Card Sorting tests contributed to the final regression model, so further discussion will focus on

them. PVT is an index of language and is highly associated with crystallized intelligence and

success in school and work. As above, DCCS indexes attention switching, an aspect of executive

function (Weintraub et al., 2013). Scores on the PVT and DCCS were highly significantly

correlated, despite their attribution to different subdomains of cognition and correspondingly

different neural pathways. Further studies in our laboratory are investigating cortical structure

and function in this same sample of children using MRI. However, the finding of language and

executive function deficits suggests a LiD profile that extends well beyond any notion of a

specific auditory processing disorder.

Correlations

Although the SCAN correlated highly with the ECLiPS, a key question remains whether

the SCAN tests are primarily markers of central auditory or of cognitive function (Moore et al.,

2018). Correlations between the SCAN and the cognitive and LiSN-S variables provide insight

into this question. The SCAN Total scaled score correlated significantly with all of the domain

composites of the NIH Cognition Toolbox and both the Low cue and High cue individual scores

of the LiSN-S. Critically, the SCAN correlated less well with the derived scores of the LiSN-S

(Spatial, Talker, Total Advantage). Similarly, the Low Cue score of the LiSN-S correlated well

with tests of cognition, as above, but the derived scores of the LiSN-S notably lacked such

correlations. Because the derived scores of the LiSN-S reflect a more sensory measure of

auditory perception, these data suggest overall that the SCAN is more sensitive to cognitive than

to auditory factors that contribute to perception. Although the LiSN-S Talker Advantage was

significantly correlated with the SCAN Composite, we suggest above that this could be due to remaining cognitive contributions to the Talker Advantage. The inclusion of the SCAN in the final regression model, and the strong correlation between the ECLiPS Total scaled score and the SCAN Composite suggest the possibility that one or more of the SCAN subtests may be valid predictors of a truly auditory processing disorder. However, as shown in Table 5, even the relation between SCAN dichotic subtests and LiSN-S Advantage measures is weak relative to that between SCAN dichotic subtests and LiSN-S Cue measures.

A final point concerning LiSN-S Talker Advantage is that it, alone among the LiSN-S measures, interacted significantly with EHF threshold. Recently, we have shown that EHF hearing plays a role in speech-in-noise perception in adults (Motlagh Zadeh et al., 2019). Recurrent otitis media in childhood, especially that treated with tympanostomy tubes, appears to contribute to reduced EHF thresholds, as shown in both groups of the sample examined here (Hunter et al., 2020) and in other studies (Hunter et al., 1996). The robust relationship between Talker Advantage and ECLiPS scores may thus receive a contribution from a truly auditory process, EHF hearing, the sensitivity of which is presumably determined by ear function. However, these influences are likely to be minor compared with the dominance of cognitive function in listening.

Overall, the results reported here demonstrate a major influence of impaired cognitive function on listening difficulties of children with normal audiograms, and provide some support for an additional auditory contribution of subclinical hearing loss. Evaluation of children with complaints of listening difficulties should include standardized caregiver observations, extended high frequency audiometry, and consideration of broad cognitive abilities, beyond auditory tests.

Acknowledgments

This research was supported by grant DC014078 from the National Institute of Deafness and other Communication Disorders, and by the Cincinnati Children's Research Foundation. DRM was supported in part by the NIHR Manchester Biomedical Research Centre.

Contributions of authors to research

L.P. analyzed and interpreted data, and co-wrote the paper; L.H. contributed to the design of the study, provided input to the analysis, and was involved in critical revisions; L.M.Z. collected, analyzed, and interpreted data, contributed to the writing, and was involved in critical revisions; N.T.S. collected, analyzed, and interpreted data, contributed to the writing, and was involved in critical revisions; A.P. collected, analyzed, and interpreted data, contributed to the writing, and was involved in critical revisions; L.L. analyzed and interpreted data and was involved in critical revisions; D.R.M. designed the study, analyzed and interpreted data, and co-wrote the paper.

References

American Academy of Audiology. (2010). Clinical practice guidelines: Diagnosis, treatment and management of children and adults with central auditory processing disorder. EUA: AAA.

American Speech-Language-Hearing Association. (1996). Central auditory processing: Current status of research and implications. American Journal of Audiology, 5(2), 41–52.

American Speech-Language-Hearing Association. (2005). Guidelines for manual pure-tone threshold audiometry [Guidelines]. https://doi.org/Available from www.asha.org/policy

Banai, K., & Ahissar, M. (2004). Poor frequency discrimination probes dyslexics with
        particularly impaired working memory. Audiology and Neurotology, 9(6), 328–340.

Barry, J.G., & Moore, D. R. (2014). Evaluation of Children's Listening and Processing Skills.
        MRC-T.

Barry, Johanna G., Tomlin, D., Moore, D. R., & Dillon, H. (2015). Use of questionnaire-based
        measures in the assessment of listening difficulties in school-aged children. Ear and
        Hearing, 36(6), e300–e313.

Bench, J., Kowal, Å., & Bamford, J. (1979). The BKB (Bamford-Kowal-Bench) stentence lists
        for partially-hearing children. British Journal of Audiology, 3, 108–112.

Bess, F. H., Dodd-Murphy, J., & Parker, R. A. (1998). Children with minimal sensorineural
        heaering loss: Prevalence, educational performance, and functional status. Ear and
        Hearing, 19(5), 339–354.

Bishop, D. V. M. (2006). Children's Communication Checklist-2: United States Edition. The
        Psychological Corporation.

Bishop, D. V. M., & McDonald, D. (2009). Identifying language impairment in children:
        Combining language test scores with parental report. International Journal of Language &
        Communication Disorders, 44(5), 600–615. https://doi.org/10.1080/13682820802259662

Brown, D. K., Cameron, S., Martin, J. S., Watson, C., & Dillon, H. (2010). The North American
        listening in spatialized noise—Sentences test (NA LiSN-S): Normative data and test-
        retest reliability studies for adolescents and young adults. Journal of the American
        Academy of Audiology, 21(10), 629–641.

Cameron, S., & Dillon, H. (2007). Development of the Listening in Spatialized Noise-Sentences
        Test (LISN-S). Ear and Hearing, 28(2), 196–211.

Cameron, S., & Dillon, H. (2008). Spatial hearing deficits as a major cause of auditory

      processing disorders: Diagnosis with the LiSN-S and management options. 235–241.

Cameron, S., Dillon, H., Glyde, H., Kanthan, S., & Kania, A. (2014). Prevalence and

      remediation of spatial processing disorder (SPD) in Indigenous children in regional

      Australia. International Journal of Audiology, 53(5), 326–335.

Cameron, S., Glyde, H., & Dillon, H. (2011). Listening in Spatialized Noise—Sentences Test

      (LiSN-S): Normative and retest reliability data for adolescents and adults up to 60 years

      of age. Journal of the American Academy of Audiology, 22(10), 697–709.

Cameron, S., Glyde, H., & Dillon, H. (2012). Efficacy of the LiSN & Learn auditory training

      software: Randomized blinded controlled study. Audiology Research, 2(1), e15.

Cameron, S., Glyde, H., Dillon, H., King, A., & Gillies, K. (2015). Results from a national

      central auditory processing disorder service: A real-world assessment of diagnostic

      practices and remediation for central auditory processing disorder. Seminars in Hearing,

      36(4), 216–236.

Cherry, E. C. (1953). Some experiments on the recognition of speech, with one and with two

      ears. The Journal of the Acoustical Society of America, 25(5), 975–979.

Ching, T. Y. C., Dillon, H., Leigh, G., & Cupples, L. (2018). Learning from the Longitudinal

      Outcomes of Children with Hearing Impairment (LOCHI) study: Summary of 5-year

      findings and implications. International Journal of Audiology, 57(sup2), S105–S111.

Davis, M. H., & Johnsrude, I. S. (2007). Hearing speech sounds: Top-down influences on the

      interface between audition and speech perception. Hearing Research, 229(1–2), 132–147.

de Wit, E., Visser-Bochane, M. I., Steenbergen, B., van Dijk, P., van der Schans, C. P., &

 Luinge, M. R. (2016). Characteristics of auditory processing disorders: A systematic

 review. Journal of Speech, Language, and Hearing Research, 59(2), 384–413.

Dillon, H., Cameron, S., Tomlin, D., & Glyde, H. (2014). Comments on "Factors influencing

 tests of auditory processing: A perspective on current issues and relevant concerns" by

 Tony Cacace and Dennis McFarland. Journal of the American Academy of Audiology,

 25(7), 699–703.

Edwards, B. (2020). Emerging technologies, market segments, and MarkeTrak 10 insights in

 hearing health technology. Seminars in Hearing, 41, 37–54.

Emanuel, D. C., Ficca, K. N., & Korczak, P. (2011). Survey of the diagnosis and management of

 auditory processing disorder. American Journal of Audiology, 20(1), 48–60.

Graydon, K., Rance, G., Dowell, R., & Van Dun, B. (2017). Consequences of early conductive

 hearing loss on long-term binaural processing. Ear and Hearing, 38(5), 621–627.

Hind, S. E., Haines-Bazrafshan, R., Benton, C. L., Brassington, W., Towle, B., & Moore, D. R.

 (2011). Prevalence of clinical referrals having hearing thresholds within normal limits.

 International Journal of Audiology, 50(10), 708–716.

Humanski, R. A., & Butler, R. A. (1988). The contribution of the near and far ear toward

 localization of sound in the sagittal plane. The Journal of the Acoustical Society of

 America, 83(6), 2300–2310.

Hunter, L. L., Blankenship, C. M., Lin, L., Sloat, N. T., Perdew, A., Stewart, H., & Moore, D. R.

 (2020). Peripheral auditory involvement in childhood listening difficulty. Ear and

 Hearing, Advance online publication. https://doi.org/doi:

 10.1097/AUD.0000000000000899

Hunter, L. L., Margolis, R. H., Rykken, J. R., Le, C. T., Daly, K. A., & Giebink, G. S. (1996). High frequency hearing loss associated with otitis media. Ear and Hearing, 17(1), 1–11.

Jerger, J., & Musiek, F. (2000). Report of the consensus conference on the diagnosis of auditory processing disorders in school-aged children. Journal of the American Academy of Audiology, 11(9), 8.

Kaandorp, M. W., De Groot, A. M. B., Festen, J. M., Smits, C., & Goverts, S. T. (2016). The influence of lexical-access ability and vocabulary knowledge on measures of speech recognition in noise. International Journal of Audiology, 55(3), 157–167.

Keith, R. W. (2009). SCAN-3 for Children: Tests for Auditory Processing Disorders. Pearson.

Koch, C. (2018). What is consciousness? Nature, 557(7704), S8–S12.

Koch, C., Massimini, M., Boly, M., & Tononi, G. (2016). Neural correlates of consciousness: Progress and problems. Nature Reviews Neuroscience, 17(5), 307–321.

Kujawa, S. G., & Liberman, M. C. (2019). Translating animal models to human therapeutics in noise-induced and age-related hearing loss. Hearing Research, 377, 44–52.

Moeller, M. P., & Tomblin, J. B. (2015). Epilogue: Conclusions and implications for research and practice. Ear and Hearing, 36, 92S-98S.

Moncrieff, D. W. (2006). Identification of binaural integration deficits in children with the Competing Words Subtest: Standard score versus interaural asymmetry: Identificación del déficit de integración binaural en niños, utilizando la prueba competitiva con palabras: calificación estándar vs asimetría interaural. International Journal of Audiology, 45(9), 545–558.

Moore, B. C. J. (2012). Aspects of auditory processing related to speech perception. In W. J.

    Hardcastle, J. Laver, & F. E. Gibbon (Eds.), The Handbook of Phonetic Sciences (1st ed.,

    pp. 454–488). Wiley.

Moore, D. R., Ferguson, M. A., Edmondson-Jones, A. M., Ratib, S., & Riley, A. (2010). Nature

    of auditory processing disorder in children. PEDIATRICS, 126(2), e382–e390.

Moore, D. R. (2012). Listening difficulties in children: Bottom-up and top-down contributions.

    Journal of Communication Disorders, 45(6), 411–418.

Moore, D. R. (2018). Auditory processing disorder (APD). Ear and Hearing, 39(4), 617–620.

Moore, D. R., Cowan, J. A., Riley, A., Edmondson-Jones, A. M., & Ferguson, M. A. (2011).

    Development of auditory processing in 6- to 11-yr-old children. Ear and Hearing, 32(3),

    269–285.

Moore, D. R., & Dillon, H. (2018). How should we detect and identify deficit specific auditory

    processing disorders. ENT & Audiology News, 27, 73–74.

Moore, D. R., Hutchings, M. E., & Meyer, S. E. (1991). Binaural masking level differences in

    children with a history of otitis media. Audiology, 30, 91–101.

Moore, D. R., Sieswerda, S. L., Grainger, M. M., Bowling, A., Smith, N., Perdew, A., Eichert,

    S., Alston, S., Hilbert, L. W., Summers, L., Lin, L., & Hunter, L. L. (2018). Referral and

    diagnosis of developmental auditory processing disorder in a large, United States

    hospital-based audiology service. Journal of the American Academy of Audiology,

    29(05), 364–377.

Moore, D. R., Zobay, O., & Ferguson, M. A. (2020). Minimal and mild hearing loss in children:

    Association with auditory perception, cognition, and communication problems. Ear and

    Hearing, 41(4), 720–732.

Motlagh Zadeh, L., Silbert, N. H., Sternasty, K., Swanepoel, D. W., Hunter, L. L., & Moore, D.

    R. (2019). Extended high-frequency hearing enhances speech perception in noise.

    Proceedings of the National Academy of Sciences, 116(47), 23753–23759.

Oxenham, A. J., & Bacon, S. P. (2003). Cochlear Compression: Perceptual Measures and

    Implications for Normal and Impaired Hearing: Ear and Hearing, 24(5), 352–366.

Parthasarathy, A., Hancock, K. E., Bennett, K., DeGruttola, V., & Polley, D. B. (2020). Bottom-

    up and top-down neural signatures of disordered multi-talker speech perception in adults

    with normal hearing. ELife, 9, e51419.

Pediatrics. (2011). ADHD: Clinical practice guideline for the diagnosis, evaluation, and

    treatment of attention-deficit/hyperactivity disorder in children and adolescents.

    PEDIATRICS, 128(5), 1007–1022.

Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years after.

    Annual Review of Neuroscience, 35(1), 73–89.

Pillsbury, H. C., Grose, J. H., & Hall, J. W. (1991). Otitis media with effusion in children:

    Binaural hearing before and after corrective surgery. Archives of Otolaryngology - Head

    & Neck Surgery, 117(7), 718–723.

Roebuck, H., & Barry, J. G. (2018). Parental perception of listening difficulties: An interaction

    between weaknesses in language processing and ability to sustain attention. Scientific

    Reports, 8(1), 6985.

Rudner, M., & Rönnberg, J. (2008). The role of the episodic buffer in working memory for

    language processing. Cognitive Processing, 9(1), 19–28.

Rueda, M. R., & Posner, M. I. (2013). Development of attention networks. In P. D. Zelazo (Ed.), The Oxford Handbook of Developmental Psychology, Vol. 1: Body and Mind (pp. 683–705). Oxford University Press.

Seeto, M., Tomlin, D., & Dillon, H. (in press). The relations between auditory processing scores and cognitive, listening and reading abilities. Ear and Hearing.

Tomlin, D., Dillon, H., Sharma, M., & Rance, G. (2015). The Impact of Auditory Processing and Cognitive Abilities in Children: Ear and Hearing, 36(5), 527–542. https://doi.org/10.1097/AUD.0000000000000172

Verhaeghen, P., & Cerella, J. (2002). Aging, executive control, and attention: A review of meta-analyses. Neuroscience & Biobehavioral Reviews, 26(7), 849–857.

Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J., Carlozzi, N. E., Slotkin, J., Blitz, D., Wallner-Allen, K., Fox, N. A., Beaumont, J. L., Mungas, D., Nowinski, C. J., Richler, J., Deocampo, J. A., Anderson, J. E., Manly, J. J., Borosh, B., … Gershon, R. C. (2013). Cognition assessment using the NIH Toolbox. Neurology, 80(Issue 11, Supplement 3), S54–S64.

Whitton, J. P., & Polley, D. B. (2011). Evaluating the perceptual and pathophysiological consequences of auditory deprivation in early postnatal life: A comparison of basic and clinical studies. Journal of the Association for Research in Otolaryngology, 12(5), 535–547.

Wilmington, D., Gray, L., & Jahrsdoerfer, R. (1994). Binaural processing after corrected congenital unilateral conductive hearing loss. Hearing Research, 74, 99–114.

Figure Captions

Figure 1: ECLiPS scores show better caregiver-assessed listening skills in the typically developing (TD) than in the listening difficulty (LiD) group of children. Violin plots with trimmed tails demonstrate the probability density of the data and are overlaid with boxplots indicating the median and interquartile range. Scores are shown for (A) all five ECLiPS subscales (speech & auditory processing, SAP; environmental & auditory sensitivity, EAS; language/ literacy/ laterality, LLL; memory & attention, M&A, and pragmatic & social skills, PSS) as well (B) as its four composite scores (Language, Listening, Social, and Total aggregate). Horizontal, dashed lines reflect expected standard score (here 10) for all scales in all figures.

Figure 2: TD children scored more highly than children with LiD on auditory tests. A – SCAN-3:C test for auditory processing disorder in children (Keith, 2009). Violin plots of scores on all subtests and the aggregate (Composite) score. B – LiSN-S listening in spatialized noise test (Cameron & Dillon, 2007). Cue and Advantage scores.

Figure 3: TD children scored more highly than children with LiD on cognitive tests. Scaled scores from the core NIH Cognition Toolbox (Weintraub et al., 2013) tests by group. A – individual tests of the Fluid Composite scale. B – individual tests of the Crystallized Composite scale. C – the Early Childhood Composite.

Figure 4: Actual and predicted ECLiPS Total scaled scores. Accuracy of the final equation yielded by stepwise multiple regression, which included four predictors: the SCAN-3:C Composite score, the LiSN-S Talker Advantage score, and the NIH Picture Vocabulary Test (PVT) and Dimensional Change Card Sort test (DCCS) scores. Distance from the diagonal reflects the accuracy of the prediction for each individual. Scores below the diagonal reflect instances in which the predicted scores were higher than actual scores, while those above the line were lower than observed.

Tables

Table 1: Participant numbers and demographic variables in the typically developing (TD) and listening difficulty (LiD) groups. Age is expressed as mean (SD). All other values are frequencies.

	TD	LiD
Total N	79	67
Age	9.42 (2.04)	9.79 (1.96)
Maternal Education Level		
High school and under	1	12
Some college and above	78	55
Gender		
Female	33	23
Male	46	44
Race		
Non-Caucasian	10	13
Caucasian	66	50
Mixed Race	3	4

Table 2: Group standardized means, standard deviations, and Wilcoxon Rank Sum test results for the ECLiPS, SCAN-3:C, LiSN-S, and NIH Cognition Toolbox.

	N LiD	N TD	LiD Mean	LiD 95% CI	TD Mean	TD 95% CI
ECLiPS Subtests						
SAP	67	79	3.00	2.58, 3.42	10.39	9.85, 10.93
EAS	67	79	4.84	4.21, 5.47	10.78	10.17, 11.40
L/L/L	67	79	3.90	3.33, 4.47	10.52	9.99, 11.05
M&A	67	79	4.15	3.66, 4.63	10.72	10.19, 11.25
PSS	67	79	5.12	4.70, 5.54	10.92	10.38, 11.47
ECLiPS Composite Scores						
Listening	67	79	3.37	2.98, 3.77	10.73	10.21, 11.26
Language	67	79	3.61	3.10, 4.12	10.58	10.07, 11.09
Social	67	79	4.46	3.94, 4.99	11.00	10.42, 11.58
Total	67	79	2.96	2.53, 3.38	10.81	10.28, 11.34
CCC-2 Composite Scores						
GCC	64	79	81.70	78.33, 85.07	111.85	109.94, 113.76
SIDI	64	79	2.27	-0.57, 5.10	-0.06	-1.29, 1.16
SCAN3:C						
FW	59	69	11.81	11.32, 12.30	13.00	12.54, 13.46
AFG	59	69	9.59	8.92, 10.27	11.41	10.86, 11.95
CW-DE	59	69	7.00	6.31, 7.69	10.13	9.53, 10.73
CS	59	69	8.71	7.96, 9.46	10.84	10.30, 11.38
Composite Score	59	69	94.85	91.73, 97.96	110.09	107.50, 112.67
LiSN-S Cue Scores						
Low Cue	64	76	-0.74	-1.03, -0.45	-0.13	-0.35, 0.09
High Cue	64	76	-0.40	-0.73, -0.08	0.28	0.05, 0.52
LiSN-S Advantage Scores						
Talker	64	76	-0.54	-0.79, -0.30	-0.05	-0.24, 0.13
Spatial	64	76	-0.65	-1.07, -0.24	-0.23	-0.53, 0.08
Total	64	76	-0.03	-0.33, 0.28	0.40	0.16, 0.64
NIH Cognition Toolbox Tests						
Flanker	57	57	89.40	85.59, 93.20	100.64	96.78, 104.49
DCCS	57	57	88.17	84.68, 91.66	104.11	100.25, 107.96
LSWM	56	67	92.80	89.11, 96.50	109.77	106.09, 113.46
PCPS	54	52	84.18	77.98, 90.39	103.29	97.27, 109.31
PSMT	52	73	94.51	89.10, 99.93	110.90	106.32, 115.48
PVT	59	72	97.36	94.04, 100.68	112.88	109.41, 116.35
RR	55	64	89.91	86.77, 93.05	106.85	103.76, 109.94
AVL	41	48	-1.48	-2.13, -0.84	0.66	0.22, 1.09
NIH Cognition Toolbox Composite Scores						
Fluid	45	51	84.93	80.10, 89.77	109.13	104.28, 113.97
Crystallized	54	63	92.43	89.14, 95.71	110.71	107.44, 113.97
Total	44	50	86.56	82.40, 90.72	112.26	108.13, 116.40
Early Childhood	48	56	88.97	84.91, 93.02	111.94	107.74, 116.15

Table 3: Correlations with the ECLiPS Total Score for all continuous predictors that were entered into the stepwise regression. Regression coefficients and their standard errors, *t* values, and *p* values are provided for predictors that were retained in the final model.

	Spearman Rho	Regression Coefficient	Standard Error	*t* value	*p* value
SCAN3:C					
Composite Score	0.51***	0.077	0.030	2.54	0.013
LiSN-S					
Low Cue	0.23*	-	-	-	-
High Cue	0.30**	-	-	-	-
Talker Advantage	0.28**	1.06	0.372	2.84	0.006
Spatial Advantage	0.27**	-	-	-	-
NIH Cognition Toolbox					
PVT	0.51***	0.085	0.026	3.28	0.002
PSMT	0.34***	-	-	-	-
Flanker	0.20	-	-	-	-
DCCS	0.37***	0.047	0.023	2.02	0.046
Interactions with Extended High Frequency Threshold (EHFT)					
EHFT * LiSN-S Low Cue	0.19	-	-	-	-
EHFT * LiSN-S Talker Adv.	0.27**	-	-	-	-
EHFT * LiSN-S Spatial Adv.	0.17	-	-	-	-

* $p < 0.05$
** $p < 0.01$
*** $p < 0.001$

Table 4: Correlation matrix (Spearman's Rho) comparing test scores as continuous variables across both LiD and TD children.

	ECLiPS Total Comp.	SCAN3:C Composite	LiSN-S Low Cue	LiSN-S High Cue	LiSN-S Talker Advantage	LiSN-S Spatial Advantage	LiSN-S Total Advantage	NIH Fluid Comp.	NIH Crystallized Comp.	NIH Total Comp.	NIH Early Child. Comp.
SCAN3:C											
Composite	0.461***										
LiSN-S											
Low Cue	0.232**	0.402***									
High Cue	0.245**	0.330***	0.384***								
Talker Advantage	0.323***	0.177*	0.051	0.420***							
Spatial Advantage	0.182*	0.111	-0.007	0.591***	0.384***						
Total Advantage	0.153	0.157	-0.022	0.897***	0.421***	0.662***					
NIH Cognition Toolbox											
Fluid Composite	0.508***	0.510***	0.452***	0.281**	0.228*	0.169	0.076				
Crystallized Comp.	0.529***	0.601***	0.35***	0.068	0.096	-0.014	-0.075	0.557***			
Total Composite	0.603***	0.612***	0.446***	0.249*	0.176	0.12	0.044	0.904***	0.845***		
Early Child. Comp.	0.528***	0.532***	0.387***	0.207*	0.106	0.136	0.039	0.900***	0.717***	0.921***	

** $p < 0.01$
*** $p < 0.001$

Table 5: Correlations between subtests of SCAN and LiSN-S for all children (total n = 124) who completed both tests.

	LiSN-S				
	Low Cue	*High Cue*	*Talker Advantage*	*Spatial Advantage*	*Total Advantage*
SCAN3:C					
Filtered Words	0.293**	0.187	0.040	-0.050	0.058
Auditory Figure Ground	0.097	0.128	0.114	0.069	0.108
Competing Words	0.343***	0.308***	0.172	0.212	0.159
Competing Sentences	0.360***	0.298***	0.193	0.101	0.128
Composite	0.406***	0.331***	0.177	0.111	0.156

** $p < 0.01$
*** $p < 0.001$

ECLiPS Scores by Group