

Title: When lockdown policies amplify social inequalities in COVID-19 infections. Evidence from a cross-sectional population-based survey in France.

Authors

Nathalie Bajos, PhD^{1*}, Prof Florence Jusot, PhD², Ariane Pailhé, PhD³, Alexis Spire, PhD¹,
Claude Martin, PhD⁴, Prof Laurence Meyer, PhD⁵, Nathalie Lydié, PhD⁶, Jeanna-Eve Franck,
PhD¹, Prof Marie Zins, PhD⁷, Prof Fabrice Carrat, PhD⁸; for the SAPRIS study group*

Affiliations

¹ IRIS, Inserm/EHESS/CNRS, Aubervilliers, France

² Université Paris Dauphine, Paris, France

³ INED, Aubervilliers, France

⁴ ARENES UMR 6051, CNRS, EHESP, Rennes

⁵ CESP Inserm UMR1018, Université Paris-Saclay, UVSQ, le Kremlin-Bicêtre, France

⁶ Santé publique France, Saint-Maurice, France

⁷ Département de Santé Publique, AHP. Sorbonne Université, Paris, France

⁸ Sorbonne Université, Inserm, Institut Pierre-Louis d'Epidémiologie et de Santé Publique, Paris, France

Email addresses:

nathalie.bajos@inserm.fr

florence.jusot@dauphine.psl.eu

ariane.pailhe@ined.fr

alexis.spire@gmail.com

claudio.martin@ehesp.fr

laurence.meyer@inserm.fr

nathalie.lydie@santepubliquefrance.fr

jeanna-eve.franck@inserm.fr

marie.zins@inserm.fr

fabrice.carrat@iplesp.upmc.fr

Correspondence to : Dr Nathalie Bajos, Orcid number [0000-0001-8073-9056](https://orcid.org/0000-0001-8073-9056)

Institut de Recherche Interdisciplinaire sur les enjeux Sociaux - Sciences sociales, politique, santé, IRIS (UMR 8156 CNRS - EHESS - U997 Inserm), 5 cours des humanités, 93322 Aubervilliers cedex, France.

Tel : +33 (0)6 66 32 30 00. nathalie.bajos@inserm.fr

Contributor and guarantor information

Dr Bajos, guarantor, had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis. She accepts full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish. She attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Study concept and design: Bajos, Carrat

Acquisition of data: Zins.

Analysis and interpretation of data: Bajos, Jusot, Pailhé, Spire, Martin, Meyer

Drafting of the manuscript: Bajos

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Franck

Obtained funding: Bajos, Carrat, Zins

Administrative, technical, or material support: Lydié

Study supervision: Bajos and Carrat

Copyright/license for publication

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, [a worldwide licence](#) to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above.

Competing interests declaration

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declared no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Key words

Social inequalities, lockdown, COVID-19, general population, risk factors

Word count

Abstract (300 words), text (2,631 words), references (27)

Abstract

Objectives

To assess social inequalities in the trends in COVID-19 infections following lockdown

Design

A cross-sectional survey conducted among the general population in France in April 2020, during COVID-19 lockdown.

Participants

10 401 participants aged 18-64, from a national cohort who lived in the three metropolitan French regions most affected by the first wave of COVID-19.

Main outcome

The main outcome was occurrence of possible COVID-19 symptoms, defined as the occurrence of sudden onset of cough, fever, dyspnea, ageusia and/or anosmia, that lasted more than three days in the 15 days before the survey. We used multinomial regression models to identify social and health factors related to possible COVID-19 before and during the lockdown.

Results

In all, 1,304 (13.0%; 95% CI: 12.0%-14.0%) reported cases of possible COVID-19. The effect of lockdown on the occurrence of possible COVID-19 was different across social hierarchies. The most privileged class individuals saw a significant decline in possible COVID-19 infections between the period prior to lockdown and during the lockdown (from 8.8% to 4.3%, $P=0.0001$) while the decline was less pronounced among working class individuals (6.9% before lockdown and 5.5% during lockdown, $P=0.03$). This differential effect of lockdown

remained significant after adjusting for other factors including history of chronic disease. The odds of being contaminated during lockdown as opposed to the prior period increased by 57% among working class individuals (OR=1.57; 95% CI: 1.0-2.48). The same was true for those engaged in in-person professional activities during lockdown (OR=1.53; 95% CI: 1.03-2.29).

Interpretation

Lockdown was associated with social inequalities in the decline in COVID-19 infections, calling for the adoption of preventive policies to account for living and working conditions. Such adoptions are critical to reduce social inequalities related to COVID-19, as working-class individuals also have the highest COVID-19 related mortality, due to higher prevalence of comorbidities.

Section 1: What is already known on this topic

Significant differences in COVID-19 incidence by gender, class and race/ethnicity are recorded in many countries in the world. Lockdown measures implemented throughout the globe have been effective in reducing transmission risks.

Section 2: What this study adds

Our study shows that lockdown's impact was socially differentiated and has benefited the working classes the least. Such results underline the need to design COVID-19 preventive policies that take into account living and working conditions, as working-class individuals also have the highest COVID-19 related mortality, due to higher prevalence of comorbidities.

Introduction

Given the pre-existing social inequalities in health within societies¹ and the significant differences in COVID-19 mortality by gender, class and origin recorded in countries such as France²⁻³, the United Kingdom⁴, the USA⁵ and other countries around the world⁶, several studies address issues of social inequalities related to COVID-19⁷⁻¹¹. However, to our knowledge, no study has investigated potential social inequalities in the effects lockdown policies, widely implemented around the globe.

Our hypothesis is that lockdown measures, shown to be effective in reducing the number of new cases¹², have not been effective in the same way for all, failing to protect the most vulnerable populations. While more privileged social classes may have had greater exposure to the virus prior to lockdown, due to more frequent social interactions in public spaces (e.g. bars, restaurants) and travelling, they may have better adapted to lockdown measures, through telework, while working classes may have benefited less from lockdown conditions, due to their professional obligations as essential workers and their living conditions in overcrowded housing.

Our objective was to study the differential effect of lockdown measures on possible COVID-19 infections according to social class in France, one of the most affected countries in Europe by the first wave of COVID-19.

Methods

Study design and participants

The SAPRIS (*SAnté, Pratiques, Relations et Inégalités Sociales en population générale pendant la crise COVID-19*) survey was set-up mid-March 2020, with the general aim of understanding the main epidemiological, social and behavioural challenges of the SARS-CoV2 epidemic in France. It relies on a *consortium* of four prospective cohort studies involving three general population-based adult cohorts and a population-based children cohort. The analysis presented here is based on data from one of the three adult cohorts, the *Constances* cohort, which is the only cohort to have accurate data on professional status and preventive measures in the workplace. *Constances* is a generalist cohort made up of a national sample of 215 000 adults aged 18 to 69 at inclusion and recruited from 2012 onwards¹⁴.

All cohort members of *Constances* who had regular access to the internet (n=66,848) were invited to complete the SAPRIS questionnaire online. 69.0% participated in the survey (46,107). To best highlight the impact of the lockdown on possible COVID-19 symptoms, we chose to center this analysis on individuals (18-64 years) who have already been employed, living in one of the three metropolitan French regions most affected by the first wave of COVID-19 *i.e.* Grand Est, Ile-de-France (Paris Region) and Hauts-de-France. 10,101 participants met this criteria and were included in the analysis.

Ethics and public involvement

The survey was approved by the National Institute for Health and Medical Research (Inserm) ethics evaluation committee (approval #20-672 dated March 30th, 2020).

Data collection

Data collected online from April 6th to May 5th, 2020 solicited information on socio-demographic characteristics, household size and composition, employment characteristics, daily life conditions, childcare arrangements, alcohol and tobacco use, sexual life, comorbidities, health care utilization and treatments. The questionnaire also addressed COVID-19 related topics including preventive behaviors (gel, mask, social distancing) for individuals and in the workplace, risk perceptions and COVID-19 related beliefs as well as a detailed description of COVID-19 symptoms over the last two weeks.

Symptoms were reported if they were unusual and occurred at least once in the past 15 days. The duration of symptoms were graded on a scale of one to five (less than 1 day, 1 to 3 days, 4 to 7 days, 8 to 14 days, >14 days). Finally, the total time (in days) between the onset of the first symptom and the date of the survey was reported.

Measures

Our main outcome was a three-category measure, distinguishing 1) No suspicion of Covid-19 contamination, 2) probable contamination before the lockdown and 3) probable contamination during the lockdown. We used the following criteria defined by the European Centre for Disease Prevention to identify “possible COVID-19 contamination: at least cough or fever or dyspnea or sudden onset of ageusia, dysgeusia or anosmia occurring during the at-risk period”¹⁵. We added an additional criterion of duration, including symptoms lasting more than 3 days to add additional specificity to our definition.

The likely period of contamination (LPC) was identified as a function of *i*) the duration between the onset of the first symptoms and the date of the survey (DFS), *ii*) the duration of incubation (DI: the 75th percentile duration between exposure and the onset of the first COVID-19 symptom) of 7 days¹⁶ and *iii*) the date of survey (DS). LPS was defined as follows:

$$LPC = DS - (DFS + DI)$$

Based on this information probable contamination before the lockdown included LPC before March 17th while contamination during the lockdown included LPC on or after March 17th.

Participants' social position was defined according to 3 criteria: current professional status (Inactive, retired or unemployed before the beginning of the pandemic/ employed but stopped working since the beginning of the pandemic, Full-time teleworking, Full-time or part-time in-person professional activities), socio-professional class and financial situation as perceived by respondents (comfortable/no problems/difficult). Socio-professional class was based on current or previous occupation, and distinguished health professions with specific exposure to the virus. The following 5 categories were constructed: Health professionals (doctors, nurses, caregivers), Upper class (senior managers), High middle class (intermediate professions), Low middle class (employees and skilled workers with a diploma of higher or equal to two years university degree), working class (unskilled employees and workers with a diploma lower than a two years university degree).

Statistical methods

We used inverse probability weighting to correct for selection and non-participation biases. Weights were estimated using logistic regression models, with selection or participation as the response variables, and socio-demographics characteristics as covariates: sex, age group, occupational status (active, inactive), social affiliation and department of residence.

Since the information on the number of rooms in the housing unit was only asked in a second survey in June 2020, this information was missing for the 22% of the sample who

didn't complete the second questionnaire. We imputed this data using predictions obtained by logistic regression.

We first conducted bivariate analysis to explore the association between sociodemographic characteristics area (size of the agglomeration and region), number of individuals living in the household *per* room, educational level, nationality (French or not French), professional status, smoking, body mass index, health status (chronic diseases), and COVID-19 related behaviors (individual and workplace preventive measures (gel, mask, social distancing) and possible COVID-19 contamination in three categories.

We then conducted a multinomial logistic regression to compare the risk of contamination before (reference category) and during the lockdown according to social class, with successive and additional adjustments for other socio-demographic and health factors. The final model presents the variables that allow us to test our hypotheses on the effect of living conditions: housing, social class and professional status. We performed a sensitivity analysis including those with symptoms lasting less than 3 days and found results similar in magnitude but some became statistically non-significant (not shown).

All analyses were performed using R software. A P-value <0.05 was considered statistically significant. All percentages are weighted to account for the complex sampling design and post stratification.

Multivariable analyses were performed on unweighted data.

Role of the funding source

The funders had no role in the design, analysis, interpretation or writing. All the authors had full access to all the data and NB and FC had final responsibility for the decision to submit for publication.

Results

Table 1 describes the sociodemographic and health characteristics distribution of study participants and the frequency of possible COVID-19, according to the probable date of contamination.

The sample was equally divided between men (47.6%) and women (52.4%) and the mean age was 43.50 years (95%CI: 43.17-43.83). More than a third (38.8%) of the sample lived in cities with more than 1,000,000 inhabitants while a minority (6.9%) lived in rural areas. About a third of the sample (32.7%) were considered upper class while 21.8% were working class. 15.3% of the sample had in person professional activities during the lockdown period. Altogether, 13.0% (95% CI: 12.0%-14.0%) of participants reported symptoms compatible with possible cases of COVID-19 (n=1335) in the two weeks preceding the survey.

Residents from the Paris region (Ile-de-France) (P=0.02), participants facing financial difficulties (P=0.046) and those who reported chronic conditions (asthma or respiratory pathologies specifically) (P<0.0001) were more likely to report possible COVID-19 while older participants (P=0.003), and those who did not work before lockdown (P=0.033) were less likely to report those symptoms. Reporting possible COVID-19 was unrelated to social class.

While the percentage of participants reporting possible COVID-19 infection during lockdown was lower than participants reporting possible COVID-19 infection before lockdown (5.0% versus 8.0%), this decrease was uneven across social groups. As shown in Figure 1, the decline was most pronounced among privileged classes (from 8.8% before lockdown to 4.3% during lockdown, $P=0.001$) while the decline was least pronounced among the working class (from 6.9% before lockdown to 5.5% during lockdown, $P=0.03$).

In addition, those living in housings with less than one room *per* person were slightly more likely to report a possible case of COVID-19 than others (16.3% versus 12.8%, $P=0.08$), with no difference between before and during lockdown.

The multivariable analyses presented in Table 2 indicated that the odds of no contamination relative to probable infection prior lockdown was unrelated to social class but depended on the region of residence, with increased odds among residents from the Hauts-de-France region relative to those residing in the Paris region (Ile de France) (OR= 1.39; 95% CI: 1.13-1.71).

Regarding the risk of infection during lockdown relative to the risk of infection before lockdown, it was higher among participants who had in-person professional activities compared to those who worked remotely (OR=1.53; 95% CI: 1.03-2.29). This risk was also increased among working class compared to upper class participants (OR=1.57; 95% CI: 1-2.48). It is worth noting that the odds-ratio for working class was 1.53 (95% CI: 0.96-2.42) when adjusting for smoking and it was 1.49 (95%CI: 0.93-2.4) when adjusting for history of chronic disease and obesity. Finally, this odds-ratio was reduced to 1.39 (95% CI: 0.87-2.21) when adjusting for perceived financial situation.

To study the stability of our results, we performed a sensitivity analysis check by repeating the multivariate analyses without excluding individuals who reported symptoms lasting less than 3 days. The odds ratio remained of the same magnitude but the degrees of significance were lower (not shown).

Discussion

To our knowledge, SAPRIS is the largest general population-based COVID-19 study in Europe that simultaneously collects detailed data on symptoms and social characteristics to investigate the impact of lockdown on possible COVID-19 infections.

Analyses by time period, corresponding to whether individuals may have been infected before or during lockdown, show differential trends by social class that were masked in an overall analysis. The issue of temporality is essential because confinement measures affected individuals differently according to their housing and working conditions^{5,17}. Individuals at the top of the social hierarchy saw a greater decline in COVID-19 symptoms after the lockdown than those from the working class. In fact, working-class individuals were more likely than those in the upper class to have been contaminated during lockdown rather than before.

Our results show that this overexposure during lockdown was partly a result of their health status (smoking and history of chronic disease and obesity). It was also partly an effect of their economic precariousness since the OR of the working class decreased when it was adjusted on this variable, a result consistent with economic work that has recently been

established at a macroeconomic level in France¹⁸. We also found that living in housing with less than one room per person tended to be linked to the risk of having been contaminated. Finally, our results do not reflect a lower propensity of the working class to adopt individual prevention measures.

One can think that the overexposure to the virus of the working class during lockdown may reflect, at least in part, the fact that more individuals belonging to this class live in neighborhoods with high population density. Such an effect is not completely captured by the size of the agglomeration. For example, the density in some neighborhoods in the Paris suburbs, where excess mortality by COVID-19 is particularly high, is higher than that observed in larger cities¹⁹. Residents of these dense cities could have faced a higher risk of being exposed to the virus by encountering contagious individuals in shops, in the streets, or in public transports.

In any case, the data suggest that working class individuals were less protected by the lockdown measures than the more privileged categories.

This analysis has several limitations. First, the sample is socially diverse but is not fully representative of the French population as it only represents three regions in France and respondents from the Constances cohort who have internet connectivity are not representative of all residents in France. In particular, the study fails to capture particularly vulnerable groups such as undocumented migrants and homeless people, who are particularly affected by the pandemic⁸.

While the study provides information on social status based on education and employment, it doesn't capture other forms of social disadvantage including race and ethnicity that are

shown to increase the risk of COVID-19 infection in many settings and the risk COVID-19 related mortality in France³ and other countries²⁰⁻²².

Additionally, it should be noted that our analyses are based on reported symptoms rather than on biologically tested cases, thus excluding asymptomatic individuals. However, the shortage of tests did not permit the use of testing in this study conducted in the early stages of the pandemic, especially before lockdown, as the use of RT-PCR testing was limited to patients with severe symptoms. Our symptom-based analysis is nevertheless consistent with epidemiological surveillance data by region¹⁹ and data on over-exposition of individuals with chronic respiratory diseases²³.

Another limitation relates to the fact that some people may have had COVID-19 symptoms prior to the 15 days of the survey and are not counted in our possible COVID-19 cases. Since the socio-demographic structure of the respondents is stable during the study period (not shown), it is reasonable to think that the *de facto* exclusion of these situations does not affect results on association of possible Covid19 with social class.

In addition, although symptom reporting may risk being socially differentiated, it is reasonable to assume that any social reporting bias does not vary during the month of the survey.

In any case, from a prevention perspective, it is important to characterise the most exposed social groups and to try to uncover the social logics that favour this exposure, particularly those referring to living conditions^{24,25}.

In conclusion, we showed that the effect of a lockdown policy designed and applied without taking into account social characteristics can contribute to increasing social inequalities in exposure to the risk of contracting the virus, as was rightly pointed out recently by Anderson

et al.²⁶ In this sense, the biomedical approach to prevention, which promotes preventive measures based on clinical knowledge without taking into account the socially differentiated effects of living conditions shows its limits, as was the case in the fight against previous epidemics^{9,27}. Our results call for the implementation of future preventive policies that tackle these social inequalities. Such implementation is critical to reduce social inequalities related to COVID-19, as working-class individuals also have the highest COVID-19 related mortality, due to higher prevalence of comorbidities.

References

- 1- Marmot M. and Bell R. (2016) 'Social inequalities in health: a proper concern of epidemiology', *Ann Epidemiol.*, 26(4), pp 238-40.
- 2- Salje H, Tran Kiem C, Lefrancq N, Courtejoie N, Bosetti P, Paireau J, Andronico A, Hoze N, Richet J, Dubost CL, Le Strat Y, Lessler J, Levy-Bruhl D, Fontanet A, Opatowski L, Boelle PY, Cauchemez S. Estimating the burden of SARS-CoV-2 in France. *Science*. 2020.
- 3- Papon S and Robert-Bobée I. Une hausse des décès deux fois plus forte pour les personnes nées à l'étranger que pour celles nées en France en mars-avril 2020. *Insee Focus* No 198 - juillet 2020.
- 4- Williamson, E. J. et al. OpenSAFELY: factors associated with COVID-19 death in 17 million patients. *Nature* <https://doi.org/10.1038/s41586-020-2521-4> (2020).
- 5- Baker MG, Peckham TK, Seixas NS (2020) Estimating the burden of United States workers exposed to infection or disease: A key factor in containing risk of COVID-19 infection. *PLoS ONE* 15(4): e0232452. <https://doi.org/10.1371/journal.pone.0232452>
- 6- Shadmi E , Chen Y, Dourado I, Faran-Perach I , Furler J, Hangoma P, Hanvoravongchai P , Obando C, Petrosyan V, Rao C, Ruano A, Shi L, Eugenio de Souza L, Spitzer-Shohat S, Sturgiss E, Suphanchaimat R, Villar Uribe M, Willems S. Health equity and COVID-19: global perspectives. *Int J Equity Health*. 2020 Jun 26;19(1):104. doi: 10.1186/s12939-020-01218-z.
- 7- Wang Z and Tang K. Combating COVID-19: health equity matters. *Nature Medicine | VOL 26 | April 2020. | 458–464.* <https://doi.org/10.1038/s41591-020-0823-6>
- 8- Okonkwo NE, Aguwa UT, Jang M, et al. COVID-19 and the US response: accelerating health inequities [published online ahead of print, 2020 Jun 3]. *BMJ Evid Based Med*. 2020;bmjebm-2020-111426. doi:10.1136/bmjebm-2020-111426
- 9- Bamba C, Riordan R, Ford J, Matthews F. The COVID-19 pandemic and health inequalities. June 2020. *Journal of Epidemiology & Community Health*. DOI: 10.1136/jech-2020-214401

10 Chung R, Dong D, Ming Li M. Socioeconomic gradient in health and the covid-19 outbreak. *BMJ* 2020; 369 doi: <https://doi.org/10.1136/bmj.m1329> (Published 01 April 2020)
Cite this as: *BMJ* 2020;369:m1329

11- Wright L, Steptoe A, Fancourt D. Are we all in this together? Longitudinal assessment of cumulative adversities by socioeconomic position in the first 3 weeks of lockdown in the UK [published online ahead of print, 2020 Jun 5]. *J Epidemiol Community Health*. 2020;jech-2020-214475. doi:10.1136/jech-2020-214475

12- Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, Whittaker C, Zhu H, Berah T, Eaton JW, Monod M, Imperial College C-RT, Ghani AC, Donnelly CA, Riley SM, Vollmer MAC, Ferguson NM, Okell LC, Bhatt S. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020.

13- Khalatbari-Soltani S, Cumming RC, Delpierre C, Kelly-Irving M. Importance of collecting data on socioeconomic determinants from the early stage of the COVID-19 outbreak onwards. *J Epidemiol Community Health*. 2020;74(8):620-623.
doi:10.1136/jech-2020-214297

14- Zins M, Goldberg M, team C. The French Constances population-based cohort: design, inclusion and follow-up. *Eur J Epidemiol*. 2015;30(12):1317-28.

15- European Center for Diseases Control (ECDC). Case definition for coronavirus disease 2019 (COVID-19), as of 29 May 2020 [Available from: <https://www.ecdc.europa.eu/en/covid-19/surveillance/case-definition>. Accessed July 15th, 2020.

16- Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med*. 2020;172(9):577-582. doi:10.7326/M20-0504

17- Chen T and Krieger N, Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county vs. ZIP code analyses. HCPDS Working Paper Volume 19, Number 1. April 21, 2020.

18- Brandily P, Brébion C, Briole S, Khoury L. A Poorly Understood Disease? The Unequal Distribution of Excess Mortality Due to COVID-19 Across French Municipalities. 2020. halshs-02895908.

19- Ministère de la santé, 2020.

<https://solidarites-sante.gouv.fr/soins-et-maladies/maladies/maladies-infectieuses/coronavirus/etat-des-lieux-et-actualites/article/indicateurs-de-l-activite-epidemie>

20- Pollán M, Pérez-Gómez B, Pastor-Barriuso R, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet*. 2020;396(10250):535-544. doi:10.1016/S0140-6736(20)31483-5

21- Bhala N, Curry G, Martineau A, Agyemang C, Bhopal R. Sharpening the global focus on ethnicity and race in the time of COVID-19. *The Lancet* Published Online. May 8, 2020.

- 22- Niedzwiedz, C.L., O'Donnell, C.A., Jani, B.D. et al. Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank. *BMC Med* 18, 160 (2020). <https://doi.org/10.1186/s12916-020-01640-8>
- 23- Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet Respir Med* 2020; 8(5): 436-8.
- 24- Napier AD. Rethinking vulnerability through Covid-19. *Anthropol Today*. 2020;36(3):1-2. doi:10.1111/1467-8322.12571
- 25- Lancet T. Redefining vulnerability in the era of COVID-19. *Lancet* 2020;395:1089.
- 26- Anderson G, Frank JW, Naylor CD, Wodchis W, Feng P. Using socioeconomics to counter health disparities arising from the covid-19 pandemic. *BMJ*. 2020;369:m2149. Published 2020 Jun 8. doi:10.1136/bmj.m2149
- 27- Peckham R. COVID-19 and the anti-lessons of history. *The Lancet*, Published Online March 2, 2020. [https://doi.org/10.1016/S0140-6736\(20\)30468-2](https://doi.org/10.1016/S0140-6736(20)30468-2)

Acknowledgments

The authors warmly thank all the volunteers of the Constances cohort.

We thank the staff of the Constances cohort that have worked with dedication and engagement to collect and manage the data used for this study and to ensure continuing communication with the cohort participants.

We thank A Sireyjol and L Kuhn for their help in statistical analysis and Pr J Bouyer for his advice in multivariable analysis.

We thank Rosalind Bell-Aldeghi for her help in editing the manuscript.

***The SAPRIS study group**

The SAPRIS study group: Nathalie Bajos (co-Principal investigator), Fabrice Carrat (co-Principal investigator), Pierre-Yves Ancel, Marie-Aline Charles, Florence Jusot, Claude Martin, Laurence Meyer, Alexandra Rouquette, Ariane Pailhé, Gianluca Severi, Alexis Spire, Mathilde Touvier, Marie Zins.

Data Statement: The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Funding

This study

ANR (Agence Nationale de la Recherche, #ANR-20-COVI-000,#ANR-10-COHO-06), Inserm (Institut National de la Santé et de la Recherche Médicale, #C20-26).

Cohort funding

The CONSTANCES Cohort Study is supported by the Caisse Nationale d'Assurance Maladie (CNAM), the French Ministry of Health, the Ministry of Research, the Institut national de la santé et de la recherche médicale. CONSTANCES benefits from a grant from the French National Research Agency [grant number ANR-11-INBS-0002] and is also partly funded by MSD, AstraZeneca, Lundbeck and L'Oreal.

Table 1: Participants characteristics and associated proportion of possible COVID-19 by period

	Weighted distribution	likely contamination	P-Value*	likely contamination prior the lockdown	likely contamination during the lockdown	P-Value**
Age			0.003			0.002
18-34	25.2 (1759)	13.8 (232)		8.3 (145)	5.5 (87)	
35-44	29.7 (3028)	14.8 (425)		8.3 (267)	6.6 (158)	
45-54	25.8 (2816)	12.6 (378)		8.7 (268)	3.9 (110)	
55-64	19.3 (2498)	9.7 (269)		6.2 (193)	3.5 (76)	
Sex			0.947			0.783
Female	52.4 (5164)	13 (663)		8.2 (442)	4.8 (221)	
Male	47.6 (4937)	13 (641)		7.8 (431)	5.2 (210)	

Nationality			0.437			0.709
French	96.1 (9703)	13.1 (1263)		8.1 (842)	5.1 (421)	
Not french	3.9 (348)	11.3 (37)		7.3 (28)	4 (9)	
Missing	50					
Social Class			0.965			0.234
Upper class	32.7 (5541)	13 (718)		8.8 (500)	4.3 (218)	
Upper middle class	20.9 (2397)	12.9 (303)		8.5 (205)	4.4 (98)	
Lower middle class	18.1 (894)	13.4 (114)		8 (72)	5.4 (42)	
Working class	21.8 (838)	12.4 (108)		6.9 (66)	5.5 (42)	
Health professional	6.5 (431)	13.9 (61)		6.2 (30)	7.7 (31)	
Professional status			0.018			0.002
Did not work before lockdown	24.5 (2178)	10.6 (241)		7.2 (177)	3.5 (64)	
Employed and stopped working since COVID	18.0 (1185)	15.7 (181)		10.3 (116)	5.4 (65)	
Switched to teleworking	38.8 (5469)	12.9 (719)		8.1 (492)	4.7 (227)	
Continued face-to-face working	18.7 (1269)	13.7 (163)		6.6 (88)	7.1 (75)	
Overcrowding			0.077			0.183
less or equal than one pers/room	94.4 (9557)	12.8 (1210)		7.9 (816)	4.9 (394)	
more than one pers/room	5.6 (544)	16.3 (94)		9.5 (57)	6.7 (37)	
Financial resources			0.046			0.077

At ease	29.5 (4082)	12.8 (493)		8.3 (346)	4.5 (147)	
No particular problem	46.5 (4369)	12.1 (563)		7.5 (382)	4.5 (181)	
Difficult	24.1 (1581)	15.2 (238)		8.7 (138)	6.6 (100)	
Missing	69					
Region			0.02			0.006
Ile-de-France	38.7 (5195)	14.4 (714)		8.9 (482)	5.5 (232)	
Grand Est	30.6 (2854)	13.4 (368)		9.1 (257)	4.3 (111)	
Hauts-de-France	30.7 (2052)	10.9 (222)		5.8 (134)	5.1 (88)	
Agglomeration size			0.21			0.268
Rural area	6.9 (488)	15.2 (67)		10.1 (47)	5.1 (20)	
< 50 000	4.3 (335)	12 (44)		7.3 (30)	4.7 (14)	
50-200 000	8.2 (536)	11.6 (67)		7.5 (49)	4.1 (18)	
200-500 000	17.1 (1972)	10.9 (229)		7.6 (157)	3.3 (72)	
500 000 -1 000 000	24.8 (1564)	12.2 (180)		6.5 (106)	5.6 (74)	
> 1 000 000	38.8 (5200)	14.4 (715)		8.9 (482)	5.5 (233)	
Missing	6					
Chronic disease			<0.00001			<0.00001
None	77.7 (7923)	12.9 (1001)		7.8 (664)	5 (337)	
Hypertension	4.7 (455)	10.2 (54)		4.7 (33)	5.5 (21)	
Asthma or other respiratory diseases	2.6 (246)	28.8 (61)		19.2 (44)	9.6 (17)	

Diabetes, cancer, heart disease, heart disease, immune diseases, liver, kidney, immunity,	2.4 (257)	12.6 (34)		10 (24)	2.6 (10)	
Others	12.7 (1220)	11.8 (154)		7.5 (108)	4.3 (46)	
Active smoking			0.807			0.856
Yes, daily	11.7 (819)	12.9 (93)		7.3 (57)	5.6 (36)	
Yes, sometimes (less than once a day)	4.6 (348)	11.3 (43)		6.2 (26)	5.1 (17)	
No	83.8 (8727)	13.1 (1136)		8.1 (770)	4.9 (366)	
Missing	207					
Obesity			0.811			0.448
BMI<30	85.9 (8889)	12.9 (1133)		7.8 (758)	5.1 (375)	
BMI>30	14.1 (868)	13.3 (124)		9.1 (84)	4.1 (40)	
Missing	344					
Individual preventive measures (mask, gel, social distancing) during outings in the last 7 days.			<0.00001			<0.00001
All 3	27 (2797)	16.2 (413)		9.7 (277)	6.5 (136)	
At least one	63 (6355)	11.3 (750)		7 (497)	4.4 (253)	
None	10 (949)	14.7 (141)		10 (99)	4.6 (42)	
Preventive measures at work (mask, gel, social distancing)			0.75			0.036
All 3	12.2 (851)	14.1 (113)		7.5 (65)	6.6 (48)	

At least one	9.4 (721)	13.0 (83)		5.8 (43)	7.2 (40)	
None	78.4 (8529)	12.8 (1108)		8.4 (765)	4.5 (343)	
All	100 (10,101)	13.0 (1304)		8.0 (873)	5.0 (431)	

* Chi2 test likely contamination/no contamination

** Chi2 test between no contamination/prior/during the lockdown

Table 2: Factors associated with possible COVID-19: adjusted OR (95% CI)

Multinomial regression results

Reference group: probable contamination prior to the lockdown

OR adjusted for all the variables presented in the table

	no symptoms/likely contaminated prior the lockdown	P-Value	likely contamination during the lockdown/likely contaminated prior the lockdown	P-Value
Age				
18-34	1	1	1	
35-44	0.95 (0.77-1.17)	0.622	1 (0.72-1.38)	0.99
45-54	0.87 (0.71-1.07)	0.198	0.69 (0.49-0.97)	0.035
55-64	1.09 (0.86-1.37)	0.475	0.73 (0.49-1.07)	0.101
Sex				
Female	1	1	1	
Male	1 (0.86-1.15)	0.95	1.03 (0.82-1.3)	0.809
Social Class				
Upper class	1	1	1	
Upper middle class	0.98 (0.82-1.18)	0.861	1.09 (0.8-1.48)	0.594
Lower middle class	1.15 (0.87-1.5)	0.322	1.28 (0.84-1.97)	0.251
Working class	1.11 (0.83-1.49)	0.467	1.57 (1-2.48)	0.051
Health professional	1.05 (0.69-1.61)	0.82	1.66 (0.91-3.04)	0.098
Professional status				
Switched to teleworking	1	1	1	
Did not work before lockdown	1.03 (0.85-1.25)	0.77	0.79 (0.56-1.12)	0.181
Employed and stopped working since COVID	0.84 (0.67-1.05)	0.126	1.11 (0.78-1.58)	0.568
Continued face-to-face working	1.25 (0.96-1.62)	0.104	1.53 (1.03-2.29)	0.037
Overcrowding housing	0.8 (0.6-1.06)	0.121	1.24 (0.8-1.9)	0.338
Region				
Ile-de-France	1	1	1	
Grand Est	0.99 (0.83-1.18)	0.924	0.84 (0.63-1.13)	0.258

Hauts-de-France	1.39 (1.13-1.71)	0.002	1.26 (0.91-1.74)	0.168
-----------------	------------------	-------	------------------	-------

* 111 (1%) participants excluded from the multivariate model due to missing values including 41 with possible COVID-19. Chronic disease, obesity, smoking and individual and work preventive measures are not presented in the final model since the odds ratio for the social class remained of the same magnitude (not shown).

Figure 1: Percentage of individuals likely to be contaminated before or during lockdown by social class.

