Title Page

Title: A Large-Scale Observational Study on the Temporal Trends and Risk Factors of Opioid Overdose: Real-World Evidence for Better Opioids

Running Heading: From Opioid Overdose Patterns to Better Opioids

Authors

Jianyuan Deng, M.Phil.[1], Wei Hou, Ph.D.[2], Xinyu Dong, M.Sc.[3], Janos Hajagos, Ph.D. [1], Mary Saltz, M.D. [1], Joel Saltz, M.D., Ph.D. [1], Fusheng Wang, Ph.D.[1,3]

[1] Department of Biomedical Informatics, Stony Brook University, Stony Brook, New York, United States

[2] Department of Family, Population and Preventive Medicine, Stony Brook University, Stony Brook, New York, United States

[3] Department of Computer Science, Stony Brook University, Stony Brook, New York, United States

Corresponding Author:

Fusheng Wang

Department of Biomedical Informatics

Department of Computer Science

Stony Brook University

2313D Computer Science

Stony Brook, New York 11794-8330

Tel: +1 (631)632-2594

Email: fusheng.wang@stonybrook.edu

Abstract

Background

The United States is in the midst of an opioid overdose epidemic. We evaluated the temporal trends and risk factors of inpatient opioid overdose. Based on the opioid overdose patterns, we further examined the innate properties underlying less overdose events.

Methods

We conducted a retrospective cross-sectional study based a large-scale inpatient electronic health records database, Cerner Health Facts®. We included patients admitted between January 1, 2009 and December 31, 2017. Opioid overdose prevalence by year, demographics and prescription opioid exposures.

Results

A total of 4,720,041 patients with 7,339,480 inpatient encounters were retrieved from Cerner Health Facts®. Among them, 30.2% patients were aged 65+, 57.0% female, 70.1% Caucasian, 42.3% single, 32.0% from South and 80.8% in urban area. From 2009 to 2017, annual opioid overdose prevalence per 1,000 patients significantly increased from 3.7 to 11.9 with an adjusted odds ratio (aOR): 1.16, 95% confidence interval (CI): [1.15-1.16]. Comparing to the major demographic counterparts above, being in 1) age group: 41-50 (overall aOR 1.36, 95% CI: [1.31-1.40]) or 51-64 (overall aOR 1.35, 95% CI: [1.32-1.39]), 2) marital status: divorced (overall aOR 1.19, 95% CI: [1.15-1.23]), 3) census region: West (overall aOR 1.32, 95% CI: [1.28-1.36]), were significantly associated with higher odds of opioid overdose. Prescription opioid exposures were also associated with increased odds of

opioid overdose, such as meperidine (overall aOR 1.09, 95% CI: [1.06-1.13]) and tramadol

(overall aOR 2.20. 95% CI: [2.14-2.27]).   Examination on the relationships between opioid

agonists' properties and their association strengths, aORs, in opioid overdose showed that

lower aORs values were significantly associated with 1) high molecular weight, 2) negative

interaction with multi-drug resistance protein 1 (MDR1) or positive interaction with

cytochrome P450 3A4 (CYP3A4) and 3) negative interaction with delta opioid receptor

(DOR) or kappa opioid receptor (KOR).

Conclusions

The significant increasing trends of opioid overdose at the inpatient care setting from 2009

to 2017 indicated an ongoing need of efforts to combat the opioid overdose epidemic in the

US. Risk factors associated with opioid overdose included patient demographics and

prescription opioid exposures. Different prescription opioids were associated with opioid

overdose to different extents, indicating a necessity to better differentiate them during

prescribing practice. Moreover, there are physicochemical, pharmacokinetic and

pharmacodynamic properties underlying less overdose events, which can be utilized to

develop better opioids.

Key Points

- There were significant increasing trends of opioid overdose at the US inpatient care setting from 2009 to 2017, showing an ongoing need for opioid overdose prevention.
- Different prescription opioids were associated with opioid overdose to different extents, indicating a necessity to differentiate prescription opioids during prescribing.
- The optimal properties underlying less overdose events mined from the large-scale, real-world electronic health records hold high potential to guide the development of

better opioids with reduced overdose effects.

Keywords

opioid overdose – temporal trends – risk factors – prescription opioids – opioid agonists
electronic health records – real-world data – real-world evidence

1 Introduction

Over the past two decades, drug overdose has been a leading cause of injury-related deaths in the United States (US), of which 70% involved illicit or prescription opioids [1]. Prescription opioids are among the most effective drugs to treat pain, which are ligands to the endogenous mu opioid receptor (MOR) and can exert agonistic, partially agonistic or antagonistic effect [2]. When activated by agonists, MOR can mediate analgesic effects as well as modulate respiratory responses [3]. However, lethal respiratory depression can happen in case of overdose [4]. In 2018, opioid overdose was attributed to 47,761 deaths, which imposed an enormous public health burden on the US [5, 6]. The epidemic of opioid overdose is dynamic and complex [7, 8]. As opined by Jalal et al, the current epidemic can be a recent manifestation of an ongoing longer-term process [8]. Close monitoring the temporal trends of opioid overdose is, therefore, crucial for developing and evaluating relevant policies to prevent and control the epidemic [8-12]. Furthermore, despite the complexity of the opioid overdose epidemic, there exist some patterns on patient demographics and opioid prescriptions, which can provide essential knowledge for the planning of effective prevention measures [12-15].

By now, various guidelines and programs have already been launched to curb opioid prescribing [16-18]. Notwithstanding the lethal respiratory depression effect, opioids are still among the most commonly prescribed medications, presumably due to that pain is the one of most common reasons for patients to visit their doctors [19, 20]. To combat the overdose epidemic, better opioid agonists with reduced overdose effects are indisputably

needed [21]. Novel treatments have been proposed based on optimizing their innate properties. For example, one selective MOR agonist was shown to be both effective for analgesia and devoid of respiratory depression [22]. Another MOR agonist was also reported to have low abuse potential due to a reduced rate of entry across the blood brain barrier (BBB) [23]. Nonetheless, whether these new chemical entities will eventually work in human subjects remains unclear yet [21]. In fact, new drug development is well known for its low success rate [24]. Thus, given the severity and urgency of the opioid overdose epidemic, early-stage optimization strategies for the lead compounds are the key to successfully bring better opioids into clinical use, which include physicochemical, pharmacokinetic and pharmacodynamic aspects [25].

Recently, real-world data (RWD), such as Electronic Health Records (EHRs), has received substantial attention for large-scale drug safety study [26]. Real-world evidence (RWE) generated from RWD holds high potential to guide drug discovery and subsequent drug development [27, 28]. In the case of opioid overdose, the fundamental question would be "What are the optimal properties underlying opioids with less overdose effects?". The properties associated with less overdose events in a large human population can serve as the valid target product profiles to clearly define the desired attributes during early drug discovery, thereby maximizing the success rate of better opioid agonists eventually getting approved [29].

Hence, in this large-scale observational study on opioid overdose, we firstly examined the temporal trends of opioid overdose and its risk factors, including patient demographics and

prescription opioid exposures based on a large-scale EHRs database. Secondly, we examined opioid agonists' relevant properties underlying lower association strengths in opioid overdose so as to generate RWEs on optimal properties for better opioids with less overdose events.

2 Methods

2.1 Data and Measurements

This retrospective cross-sectional study was based on Cerner Health Facts®, one of the largest EHRs databases in the United States. Health Facts® stores real-world, de-identified patient data, such as records on encounters, diagnoses and medications, from over 600 healthcare facilities. A recent study showed that Health Facts® and the Healthcare Cost and Utilization Project Nationwide Inpatient Sample have similar distribution across all data elements [30].

For the assembly of our study dataset, we retrieved patients at the inpatient care setting admitted between January 1, 2009 and December 31, 2017. To identify patients with opioid overdose, we used the International Classification of Diseases codes, Ninth Revision (ICD-9) and Tenth Revision (ICD-10), for poisoning or adverse effect by opium, heroin, methadone, other related narcotics, etc (Electronic Supplementary Material, Table S1) [31]. Patients with at least one relevant diagnosis code in all inpatient encounters within a year were added to the opioid overdose cohort in that year. Other independent variables are described as follows.

2.2 Patient Demographics

We measured patient demographics, including age group (<18, 18-30, 31-40, 41-50, 51-64,

65+), gender (female, male), race/ethnicity (African American, Asian, Caucasian, Hispanic, Native American, others), marital status (divorced, married, single, widowed, others), census region (Midwest, Northeast, South, West) and urbanicity (rural, urban). For patients with varied demographics, such as migration between rural and urban areas, records from their first encounter were used.

2.3 Prescription Opioids

For prescription opioids, we categorized them by their action types on MOR [2], namely, 1) opioid agonists: codeine, fentanyl and morphine, among others, 2) opioid partial agonists: buprenorphine, butorphanol and nalbuphine, and 3) opioid antagonist: naloxone. For each category, we collected the corresponding FDA-approved active ingredients from DrugBank 5.1.6, which were then mapped to medications containing the active ingredients in Health Facts® [32]. Illicit opioids and opioids withdrawn from market were excluded, such as diamorphine and propoxyphene. To ensure sufficient statistical significance, prescription opioids with annual prevalence less than 1‰ were also excluded from analyses, such as dihydrocodeine, naltrexone and pentazocine. Medications in all inpatient encounters within one year were aggregated and a patient was defined as exposed if medications containing the active ingredient were prescribed in that year, otherwise as non-exposed.

For prescription opioids' innate properties, they were also collected from DrugBank 5.1.6 [32], including, 1) physicochemical properties underlying BBB permeability: lipophilicity (logP), polar surface area (PSA) and molecular weight (MW) [33], 2) pharmacokinetic interaction with efflux transporters and metabolizing enzymes: multi-drug resistance protein 1 (MDR1), cytochromes P450 2D6 (CYP2D6) and 3A4 (CYP3A4) [34-36], and 3)

pharmacodynamic interaction with receptors: delta opioid receptor (DOR), kappa opioid receptor (KOR) and N-methyl-D-aspartate receptor (NMDAR) [22, 37].

2.2 Statistical Analyses

Demographic characteristics were determined using descriptive analyses. Proportions of opioid overdose, i.e., opioid overdose prevalence, among all patients and patient subgroups divided by demographics, were calculated on a yearly basis.

Inferential analyses were conducted to examine the temporal trends of opioid overdose and risk factors by calculating the adjusted odds ratios (aORs) and 95% confidence intervals (CIs). Firstly, for each patient subgroup stratified by demographics, multivariable logistic regression was used with 1) the inclusion of whether a patient was diagnosed as opioid overdose within a year as the dependent variable, and 2) year and a demographic factor as independent variables plus an interaction item between the two independent variables to examine the temporal trends, further adjusted for all demographics (e.g., age_group, gender, race/ethnicity, marital status, census region and rural/urban area). Secondly, for risk factors, including demographics and prescription opioid exposures, multivariable logistic regression adjusted for all demographics was also performed based on both data in each year for the yearly aORs and data across all years for the overall aORs. For demographics, age group in 65+, gender as female, race/ethnicity as Caucasian, marital status as single, census region in south and urban area were set as the reference. For prescription opioid exposures, the non-exposed was set as the reference.

To examine the relationship between MOR agonists' innate properties and their aORs (i.e.,

association strengths) in opioid overdose, power analysis was conducted to only exploit aORs with statistical power greater than 80% [38, 39]. For continuous physicochemical properties, we first discretized them into two groups, low or high, with median as the cutoff. Then, Wilcoxon test was used to examine the relationship between aORs of opioid agonists and their attributes in the physicochemical, pharmacokinetic and pharmacodynamic aspects. Dataset was processed in Python 3.7.3. Statistical analyses were conducted in R 3.6.1, packages: "stats", "epiR". All data analyses were performed between Feb 1, 2020 and Sept 1, 2020.

3 Results

3.1 Patient Characteristics

Between 2009 and 2017, a total of 7,339,480 inpatient encounters for 4,720,041 patients were retrieved from Health Facts®. Among them, 1,423,744 (30.2%) patients were in the 65+ age group, 2,689,898 (57.0%) patients were female, 3,310,471 (70.1%) patients were Caucasian, 1,995,380 patients (42.3%) were single, 1,511,746 patients (32.0%) were from South and 3,813,817 patients (80.8%) were in urban area. Encounter counts for patients were summarized in Electronic Supplementary Material, Figure S1.

3.2 Temporal Trends of Opioid Overdose

Annual opioid overdose prevalence from 2009 to 2017 stratified by demographics was detailed in Table 1. There were significant increasing trends of opioid overdose. From 2009 to 2017, the overall opioid overdose prevalence increased from 3.7‰ to 11.9‰ (3.2-fold increase; aOR: 1.16, 95% CI: [1.15-1.16]; p < 0.0001). The increasing trends in all patient

subgroups were also significant. Among them, patient subgroup from South had a 6.3-fold

increase (from 2.0‰ to 12.9‰; aOR: 1.22, 95% CI: [1.20-1.22]; p < 0.0001), followed by

Asian patients (5.2-fold increase: from 1.2‰ to 6.0‰; aOR: 1.20, 95% CI: [1.14-1.27]; p <

0.0001), African American patients (4.8-fold increase: from 2.2‰ to 10.8‰; aOR: 1.20, 95%

CI: [1.19-1.22]; p < 0.0001) and patients in rural area (4.2-fold increase: from 2.1‰ to

8.8‰; aOR: 1.14, 95% CI [1.13-1.16]; p < 0.0001). These prominent trends indicated that

prevention and control measures of opioid overdose at the US inpatient care setting had

been be inadequate over the years.

3.3 Demographics as Risk Factors for Opioid Overdose

Demographics as risk factors for opioid overdose were summarized in Table 2. Compared to

patients aged 65+, patients aged 41-50 (overall aOR: 1.36, 95% CI [1.31-1.40]; p < 0.0001)

or 51-64 (overall aOR: 1.35, 95% CI [1.32-1.39]; p < 0.0001) had higher odds of opioid

overdose whereas patients aged <18 (overall aOR: 0.18, 95% CI [0.17-0.19]; p < 0.0001) or

18-30 (overall aOR: 0.74, 95% CI [0.71-0.76]; p < 0.0001) had much lower odds. When

comparing to single patients, patients who were divorced (overall aOR: 1.19, 95% CI

[1.15-1.23]; p < 0.0001) had higher odds of opioid overdose whereas the odds were lower

for patients who were married (overall aOR: 0.73, 95% CI [0.71-0.75]; p < 0.0001) or

widowed (overall aOR: 0.89, 95% CI [0.86-0.93]; p < 0.0001). For patients in rural area

(overall aOR: 0.84, 95% CI [0.82-0.86]; p < 0.0001), their odds of opioid overdose were

lower than patients in urban area. Note that the yearly aORs were also varying. For example,

across all years, male patients (overall aOR: 0.98, 95% CI [0.96-1.00]; p = 0.05) had

comparable odds with female patients. However, male patients were more likely to have opioid overdose in 2009 whereas they became comparably or less likely to have it in subsequent years. Similarly, patients from Midwest, Northeast or West census regions also exhibited varying aORs of opioid overdose from 2009 to 2017, which partially reflected the changing dynamics of the opioid overdose epidemic [8].

3.4 Prescription Opioid Exposures as Risk Factors for Opioid Overdose

Prescription opioids included for evaluation were listed in Table 3, along with their relevant properties and the overall prevalence between 2009 and 2017. Prescription opioid exposures as risk factors for opioid overdose were detailed in Table 4. Among MOR agonists, morphine (342.1‰) was most frequently prescribed, followed by fentanyl (297.5‰), oxycodone (263.3‰), hydromorphone (242.1‰) and hydrocodone (182.4‰), the high usage rates of which manifested the prescription opioid abuse [40]. The MOR antagonist, naloxone (157.8‰), was also commonly prescribed and compared to the non-exposed patients, patients exposed to naloxone (overall aOR: 3.24, 95% CI [3.18-3.31]; $p < 0.0001$) had a higher odds of opioid overdose, presumably due to its therapeutic effect to reverse overdose [3].

For MOR partial agonists, buprenorphine (3.2‰) was less frequently prescribed than nalbuphine (45.7‰) and butorphanol (29.5‰). Besides, exposures to partial agonists, nalbuphine (overall aOR: 0.565, 95% CI [0.53-0.60]; $p < 0.0001$) and butorphanol (aOR: 0.09, 95% CI [0.07 - 0.11]; $p < 0.0001$), were associated with reduced odds of overdose whereas buprenorphine (overall aOR: 2.89, 95% CI [2.63-3.19]; $p < 0.0001$) was positively

associated with opioid overdose. This could be explained by that buprenorphine is for

medication assisted treatment (MAT), which is closely related to opioid overdose [41].

Exposures to opioid agonists were also positively associated with opioid overdose except

for codeine (overall aOR: 0.78, 95% CI [0.74-0.83]; $p < 0.0001$), consistent with a previous

finding that codeine had lowest risk for severe respiratory depression [13]. Moreover,

different opioid agonists posed unique aORs, reflecting varied association strengths in

opioid overdose.

3.5 Optimal Properties for Better Opioid Agonists

To explore the optimal properties underlying better opioid agonists with less overdose

effects, we examined the relationships between their relevant properties (Table 3) and the

yearly aORs of opioid agonists (Table 4), depicted in Fig. 1. Note that opioid agonists

targeting for peripheral MORs, namely loperamide and diphenoxylate, and methadone for

MAT purpose, were excluded [41, 42]. For the physicochemical properties, lower aORs, i.e.,

weakened association strengths, were observed for opioid agonists with high MW (median

decreased from 1.62 to 1.24; $p < 0.001$). The interpretation is that when an opioid agonist's

MW is high, its association with opioid overdose would be weakened, possibly due to

reduced BBB permeability [33]. For logP (median decreased from 1.49 to 1.40; $p = 0.2$) and

PSA (median increased from 1.25 to 1.60; $p = 0.45$), no significant associations were

observed.

From the pharmacokinetic perspective, a negative interaction with MDR1 was associated

with lower aORs (median decreased from 1.51 to 1.25; $p < 0.05$), which indicates that if an

opioid agonist does not interact with MDR1, its association strength with opioid overdose

would be lowered. This could be due to that an interaction between opioids and MDR1 at

the BBB can up-regulate the expression of MDR1, which leads to opioid tolerance, abuse,

dependence and eventually overdose [43, 44]. For the interaction with CYP3A4, a positive

interaction was associated with lower aORs (median decreased from 1.79 to 1.32; $p < 0.01$),

which suggests that CYP3A4 metabolism may be protective for opioid agonists from

overdose as a result of reduced bioavailability during first-pass metabolism [35]. From the

pharmacodynamic perspective, lower aORs were observed when opioid agonists do not

interact with DOR (median decreased from 1.52 to 1.13; $p < 0.01$) or KOR (median

decreased from 1.53 to 1.23; $p = 0.07$). The interpretation is that when opioid agonists are

selective towards MOR, overdose events would be reduced. For NMDAR (median

increased from 1.36 to 1.69; $p = 0.16$), no significant difference in aORs was observed.

4 Discussion

4.1 An Ongoing Need for Targeted Overdose Prevention

In this study, we found that opioid overdose increased significantly at the US inpatient care

setting from 2009 to 2017. As pointed out by Danovitch et al [45], inpatient opioid overdose

is a serious harm, yet preventable, and is likely to be underestimated in much of current

literature. The significant increasing trends observed from Health Facts®, a large-scale

EHRs database, indicated that prevention and control measures for opioid overdose had

been inadequate at the US inpatient setting over these years, especially for patient

subgroups with prominent fold increases, such as patients from South and patients in rural

area.

We also quantified how patient demographics were associated with inpatient opioid overdose and found that certain demographics increased the odds of opioid overdose. For instance, compared to the major demographic counterpart age 65+, patients aged 51-64 and 41-50 were more likely to have opioid overdose, which was partly consistent with a recent Mortality Disparities in American Communities study, where patients aged 40-59 were attributed to the most part of opioid overdose deaths from 2008 to 2015 [46]. Patients in urban area had higher odds of opioid overdose than those in rural area, which reflected the urbanicity of the opioid overdose epidemic [46]. The associated risk factors suggest the need for targeted measures in patient subgroups at higher risk.

4.2 A Necessity to Differentiate Prescription Opioids

In addition to patient demographics, prescription opioid exposures were also evaluated as risk factors. Our study showed that prescription opioids were commonly prescribed. For instance, 342.1‰ patients were prescribed with medications containing morphine, the high prevalence of which manifested inappropriate opioid prescribing practice and, eventually, prevalent opioid overdose [47]. Previous studies showed that opioid overdose was associated with opioid prescription patterns, such as dosage and duration [48-51]. Yet, measures to reduce prescription opioids supply alone may not be enough to combat the opioid crisis [52].

In current literature, prescription opioids were usually aggregated for the morphine milligram equivalents (MME) calculation, a widely adopted metric [16-18]. One potential

misconception arising from MME is that all opioids are interchangeable, which can falsely justify the transitioning of one opioid to other "equivalent" opioids [53, 54]. A previous study based on a cohort of 307 patients showed that some prescription opioids were associated with a much higher risk for severe respiratory depression than others [13]. For example, fentanyl had 20-fold higher relative risk compared to codeine, the lowest-risk opioid. In fact, a wide range of differences exist among prescription opioids. For instance, during distribution to the site of action, i.e., MOR in the central nervous system, they exhibit varied BBB permeability [55]. Besides, opioids also have disparate binding profiles to the opioid receptors [56, 57]. Our study revealed varied associations with opioid overdose for different prescription opioids, indicating a necessity to differentiate prescription opioids in opioid prescribing practice [13].

4.2 Real-World Evidence on Optimal Properties for Better Opioids

Given the foreseeable continuing trends of opioid overdose in the US and the high prevalence of prescription opioids for pain, there is an undisputed need to develop better opioids with less overdose effects. Our study generated RWEs on optimal properties for better opioid agonists. For instance, from the physicochemical perspective, opioid agonists with low MW had significantly higher association strengths with opioid overdose, possibly due to increased BBB permeability. From pharmacokinetic and pharmacodynamic perspectives, when opioid agonists interact with MDR1, DOR or KOR, the association strengths in opioid overdose were significantly increased. Thus, the interactions with the efflux transporter and other opioid receptors should be taken into consideration during early

drug discovery for better opioid agonists. Although some optimal properties have been proposed before, they were mostly based on preclinical studies and there still lacks a clear link to the safety profiles in human. With the emergence of artificial intelligence in drug discovery, molecule structure design biased towards pre-defined attributes has become a reality [58, 59]. The RWEs on the optimal properties for opioids with less overdose effects can function as actionable knowledge to develop better opioids [26].

4.3 Limitations

Our study also had some limitations. Firstly, we only used ICD-9 /ICD-10 codes to identify patients with opioid overdose, which may not be fully accurate [60, 61]. Besides, there was a transition of ICD-9 codes to ICD-10 codes in 2015, which can lead to discontinuities in the opioid overdose trends [62]. Secondly, we only focused on patient demographics and prescription opioid exposures as risk factors for opioid overdose. Other potential risk factors like comorbidities and surgery procedures were not examined yet [63, 64]. In addition, opioid prescription patterns including dosage and duration also need further assessment. Thirdly, because of the observational study design, where we did not strictly distinguish the temporal order of being prescribed with a medication and getting diagnosed as opioid overdose, the association strengths, thus, may not necessarily represent causality. Lastly, our study was based on a single database. Exploitation on other RWD sources would also be needed.

Nevertheless, despite the limitations, we conducted a comprehensive study on opioid overdose through investigating its patterns and generating RWEs on the optimal properties

for better opioids, which provided critical knowledge to combat the opioid overdose epidemic.

5 Conclusions

A retrospective observational study was conducted using a large-scale EHRs database to examine the opioid overdose patterns, including temporal trends and associated risk factors. The significant increasing trends of opioid overdose from 2009 to 2017 indicated that prevention and control measures for opioid overdose had been inadequate at the US inpatient care setting. Besides, patient demographics and prescription opioid exposures were associated with higher odds of opioid overdose, suggesting that targeted prevention and control measures are needed. Furthermore, there are optimal properties underlying opioid agonists with reduced overdose effects, which hold great potential for the development of better opioids.

References

1. Hedegaard H, Minino AM, Warner M. Drug Overdose Deaths in the United States, 1999-2018. NCHS Data Brief. 2020 Jan(356):1-8.

2. James A, Williams J. Basic Opioid Pharmacology - An Update. Br J Pain. 2020 May;14(2):115-21.

3. Boyer EW. Management of Opioid Analgesic Overdose. New Engl J Med. 2012 Oct 4;367(14):1372-3.

4. Boyer EW. Management of opioid analgesic overdose. N Engl J Med. 2012 Jul 12;367(2):146-55.

5. Wilson N, Kariisa M, Seth P, Smith Ht, Davis NL. Drug and Opioid-Involved Overdose Deaths - United States, 2017-2018. MMWR Morb Mortal Wkly Rep. 2020 Mar 20;69(11):290-7.

6. Gomes T, Tadrous M, Mamdani MM, Paterson JM, Juurlink DN. The Burden of Opioid-Related Mortality in the United States. JAMA Netw Open. 2018 Jun 1;1(2):e180217.

7. Jalal H, Buchanich JM, Sinclair DR, Roberts MS, Burke DS. Age and generational patterns of overdose death risk from opioids and other drugs. Nat Med. 2020 May;26(5):699-704.

8. Jalal H, Buchanich JM, Roberts MS, Balmert LC, Zhang K, Burke DS. Changing dynamics of the drug overdose epidemic in the United States from 1979 through 2016. Science. 2018 Sep 21;361(6408):1218-+.

9. Tedesco D, Asch SM, Curtin C, Hah J, McDonald KM, Fantini MP, et al. Opioid Abuse And Poisoning: Trends In Inpatient And Emergency Department Discharges. Health Aff (Millwood). 2017 Oct 1;36(10):1748-53.

10. Hasegawa K, Espinola JA, Brown DF, Camargo CA, Jr. Trends in U.S. emergency department visits for opioid overdose, 1993-2010. Pain Med. 2014 Oct;15(10):1765-70.

11. Gaither JR, Shabanova V, Leventhal JM. US National Trends in Pediatric Deaths From Prescription and Illicit Opioids, 1999-2016. JAMA Netw Open. 2018 Dec 7;1(8):e186558.

12. Burton BN, Lin TC, Said ET, Gabriel RA. National Trends and Factors Associated With Inpatient Mortality in Adult Patients With Opioid Overdose. Anesth Analg. 2019 Jan;128(1):152-60.

13. Fox LM, Hoffman RS, Vlahov D, Manini AF. Risk Factors for Severe Respiratory Depression from Prescription Opioid Overdose. Addiction. 2018 Jan;113(1):59-66.

14. Lo-Ciganic WH, Huang JL, Zhang HH, Weiss JC, Wu YH, Kwoh CK, et al. Evaluation of Machine-Learning Algorithms for Predicting Opioid Overdose Risk Among Medicare Beneficiaries With Opioid Prescriptions. JAMA Netw Open. 2019 Mar;2(3).

15. Nicol AL, Colquhoun DA, Brummett CM. The More You Know: Identifying Factors Associated With Inpatient Mortality Related to Opioid Overdose Can Drive Progress in the Opioid Health Crisis. Anesth Analg. 2019 Jan;128(1):16-8.

16. Dowell D, Zhang K, Noonan RK, Hockenberry JM. Mandatory Provider Review And Pain Clinic Laws Reduce The Amounts Of Opioids Prescribed And Overdose Death Rates.

Health Aff (Millwood). 2016 Oct 1;35(10):1876-83.

17. Rubin R. CMS to Improve Drug Programs and Opioid Overuse Oversight. JAMA. 2018 Mar 27;319(12):1189.

18. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain-United States, 2016. JAMA. 2016 Apr 19;315(15):1624-45.

19. Herzig SJ, Rothberg MB, Cheung M, Ngo LH, Marcantonio ER. Opioid Utilization and Opioid-Related Adverse Events in Nonsurgical Patients in US Hospitals. J Hosp Med. 2014 Feb;9(2):73-81.

20. Sauver JLS, Warner DO, Yawn BP, Jacobson DJ, McGree ME, Pankratz JJ, et al. Why Patients Visit Their Doctors: Assessing the Most Prevalent Conditions in a Defined American Population. Mayo Clin Proc; 2013: Elsevier; 2013. p. 56-67.

21. Machelska H, Celik MO. Advances in Achieving Opioid Analgesia Without Side Effects. Front Pharmacol. 2018 Nov 29;9.

22. Manglik A, Lin H, Aryal DK, McCorvy JD, Dengler D, Corder G, et al. Structure-Based Discovery of Opioid Analgesics with Reduced Side Effects. Nature. 2016 Sep 8;537(7619):185-+.

23. Miyazaki T, Choi IY, Rubas W, Anand NK, Ali C, Evans J, et al. NKTR-181: A Novel Mu-Opioid Analgesic with Inherently Low Abuse Potential. J Pharmacol Exp Ther. 2017 Oct;363(1):104-13.

24. Dowden H, Munro J. Trends in Clinical Success Rates and Therapeutic Focus. Nat Rev Drug Discov. 2019 Jul;18(7):495-6.

25. Hughes JP, Rees S, Kalindjian SB, Philpott KL. Principles of Early Drug Discovery. Br J Pharmacol. 2011 Mar;162(6):1239-49.

26. Singh G, Schulthess D, Hughes N, Vannieuwenhuyse B, Kalra D. Real World Big Data for Clinical Research and Drug Development. Drug Discov Today. 2018 Mar;23(3):652-60.

27. Wise J, Moller A, Christie D, Kalra D, Brodsky E, Georgieva E, et al. The Positive Impacts of Real-World Data on the Challenges Facing the Evolution of Biopharma. Drug Discov Today. 2018 Apr;23(4):788-801.

28. Cepeda MS, Kern DM, Seabrook GR, Lovestone S. Comprehensive Real-World Assessment of Marketed Medications to Guide Parkinson's Drug Discovery. Clin Drug Investig. 2019 Nov;39(11):1067-75.

29. Wyatt PG, Gilbert IH, Read KD, Fairlamb AH. Target Validation: Linking Target and Chemical Properties to Desired Product Profile. Curr Top Med Chem. 2011 May;11(10):1275-83.

30. DeShazo JP, Hoffman MA. A Comparison of a Multistate Inpatient EHR Database to the HCUP Nationwide Inpatient Sample. Bmc Health Serv Res. 2015 Sep 15;15.

31. Bramer GR. International statistical classification of diseases and related health problems. Tenth revision. World Health Stat Q. 1988;41(1):32-6.

32. Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a Major Update to the DrugBank Database for 2018. Nucleic Acids Res. 2018 Jan 4;46(D1):D1074-D82.

33. Rankovic Z. CNS Drug Design: Balancing Physicochemical Properties for Optimal

Brain Exposure. J Med Chem. 2015 Mar 26;58(6):2584-608.

34.   Yu CC, Yuan M, Yang HY, Zhuang XM, Li H. P-Glycoprotein on Blood-Brain Barrier Plays a Vital Role in Fentanyl Brain Exposure and Respiratory Toxicity in Rats. Toxicol Sci. 2018 Jul;164(1):353-62.

35.   Smith HS. Opioid Metabolism. Mayo Clin Proc. 2009 Jul;84(7):613-24.

36.   Feng XQ, Zhu LL, Zhou Q. Opioid Analgesics-Related Pharmacokinetic Drug Interactions: from the Perspectives of Evidence based on Randomized Controlled Trials and Clinical Risk Management. J Pain Res. 2017;10:1225-39.

37.   Hingne PM, Sluka KA. Blockade of NMDA Receptors Prevents Analgesic Tolerance to Repeated Transcutaneous Electrical Nerve Stimulation (TENS) in Rats. J Pain. 2008 Mar;9(3):217-25.

38.   Quartey G, Wang J, Kim J. A Review of Risk Measures in Pharmacoepidemiology with Tips for Statisticians in the Pharmaceutical Industry. Pharm Stat. 2011 Nov-Dec;10(6):548-53.

39.   Chen HN, Cohen P, Chen S. How Big is a Big Odds Ratio? Interpreting the Magnitudes of Odds Ratios in Epidemiological Studies. Commun Stat-Simul C. 2010;39(4):860-4.

40.   Edlund MJ, Martin BC, Russo JE, DeVries A, Braden JB, Sullivan MD. The Role of Opioid Prescription in Incident Opioid Abuse and Dependence Among Individuals With Chronic Noncancer Pain The Role of Opioid Prescription. Clin J Pain. 2014 Jul;30(7):557-64.

41.   Volkow ND, Frieden TR, Hyde PS, Cha SS. Medication-Assisted Therapies—Tackling the Opioid-Overdose Epidemic. N Engl J Med. 2014 May 29;370(22):2063-6.

42.   Camilleri M, Lembo A, Katzka DA. Opioids in Gastroenterology: Treating Adverse Effects and Creating Therapeutic Benefits. Clin Gastroenterol H. 2017 Sep;15(9):1338-49.

43.   Chaves C, Remiao F, Cisternino S, Decleves X. Opioids and the Blood-Brain Barrier: A Dynamic Interaction with Consequences on Drug Disposition in Brain. Curr Neuropharmacol. 2017 Nov 14;15(8):1156-73.

44.   Morgan MM, Christie MJ. Analysis of Opioid Efficacy, Tolerance, Addiction and Dependence from Cell Culture to Human. Br J Pharmacol. 2011 Oct;164(4):1322-34.

45.   Danovitch I, Vanle B, Van Groningen N, Ishak W, Nuckols T. Opioid Overdose in the Hospital Setting: A Systematic Review. J Addict Med. 2020 Jan/Feb;14(1):39-47.

46.   Altekruse SF, Cosgrove CM, Altekruse WC, Jenkins RA, Blanco C. Socioeconomic risk factors for fatal opioid overdoses in the United States: Findings from the Mortality Disparities in American Communities Study (MDAC). PLoS One. 2020;15(1):e0227966.

47.   West NA, Severtson SG, Green JL, Dart RC. Trends in abuse and misuse of prescription opioids among older adults. Drug Alcohol Depen. 2015 Apr 1;149:117-21.

48.   Dunn KM, Saunders KW, Rutter CM, Banta-Green CJ, Merrill JO, Sullivan MD, et al. Opioid prescriptions for chronic pain and overdose: a cohort study. Ann Intern Med. 2010 Jan 19;152(2):85-92.

49.   Dasgupta N, Funk MJ, Proescholdbell S, Hirsch A, Ribisl KM, Marshall S. Cohort Study of the Impact of High-Dose Opioid Analgesics on Overdose Mortality. Pain Med.

2016 Jan;17(1):85-98.

50. Volkow ND, McLellan AT. Opioid Abuse in Chronic Pain--Misconceptions and Mitigation Strategies. N Engl J Med. 2016 Mar 31;374(13):1253-63.

51. Chua KP, Brummett CM, Conti RM, Bohnert A. Association of Opioid Prescribing Patterns With Prescription Opioid Overdose in Adolescents and Young Adults. JAMA Pediatr. 2019 Dec 16.

52. Chen Q, Larochelle MR, Weaver DT, Lietz AP, Mueller PP, Mercaldo S, et al. Prevention of Prescription Opioid Misuse and Projected Overdose Deaths in the United States. JAMA Netw Open. 2019 Feb 1;2(2):e187621.

53. Schnoll SH. Misconceptions and Realities of the Prescription Opioid Epidemic. Clin Pharmacol Ther. 2018 Jun;103(6):963-5.

54. Fudin J, Pratt Cleary J, Schatman ME. The MEDD myth: the impact of pseudoscience on pain research and prescribing-guideline development. J Pain Res. 2016;9:153-6.

55. Schaefer CP, Tome ME, Davis TP. The Opioid Epidemic: a Central Role for the Blood Brain Barrier in Opioid Analgesia and Abuse. Fluids Barriers CNS. 2017 Nov 29;14(1):32.

56. Drewes AM, Jensen RD, Nielsen LM, Droney J, Christrup LL, Arendt-Nielsen L, et al. Differences between opioids: pharmacological, experimental, clinical and economical perspectives. Brit J Clin Pharmaco. 2013 Jan;75(1):60-78.

57. Pasternak GW, Pan YX. Mu Opioids and Their Receptors: Evolution of a Concept. Pharmacol Rev. 2013 Oct;65(4):1257-317.

58. Chen H, Engkvist O, Wang Y, Olivecrona M, Blaschke T. The Rise of Deep Learning in Drug Discovery. Drug Discov Today. 2018 Jun;23(6):1241-50.

59. Deng J, Yang Z, Li Y, Samaras D, Wang F. Towards Better Opioid Antagonists Using Deep Reinforcement Learning. arXiv preprint arXiv:200404768. 2020.

60. Green CA, Perrin NA, Janoff SL, Campbell CI, Chilcoat HD, Coplan PM. Assessing the accuracy of opioid overdose and poisoning codes in diagnostic information from electronic health records, claims data, and death records. Pharmacoepidemiol Drug Saf. 2017 May;26(5):509-17.

61. Wei WQ, Teixeira PL, Mo H, Cronin RM, Warner JL, Denny JC. Combining billing codes, clinical notes, and medications from electronic health records provides superior phenotyping performance. JAMIA. 2016 Apr;23(E1):E20-E7.

62. Heslin KC, Owens PL, Karaca Z, Barrett ML, Moore BJ, Elixhauser A. Trends in Opioid-related Inpatient Stays Shifted After the US Transitioned to ICD-10-CM Diagnosis Coding in 2015. Med Care. 2017 Nov;55(11):918-23.

63. Dilokthornsakul P, Moore G, Campbell JD, Lodge R, Traugott C, Zerzan J, et al. Risk Factors of Prescription Opioid Overdose Among Colorado Medicaid Beneficiaries. J Pain. 2016 Apr;17(4):436-43.

64. Hah JM, Bateman BT, Ratliff J, Curtin C, Sun E. Chronic Opioid Use After Surgery: Implications for Perioperative Management in the Face of the Opioid Epidemic. Anesth Analg. 2017 Nov;125(5):1733-40.

Figure and Tables

(A).Physicochemical Properties

(B).Pharmacokinetic Interaction

(C).Pharmacodynamic Interaction

Fig. 1 Relationships between Innate Properties of Opioid Agonists and the Adjusted Odds Ratios in Opioid Overdose

(**A**). Wilcoxon test on opioid agonists' aORs in opioid overdose and their discretized physicochemical properties: lipophilicity (logP), polar surface area (PSA) and molecular weight (MW). (**B**). Wilcoxon test on opioid agonists' aORs in opioid overdose and their interaction type with efflux transporter and metabolizing enzymes: multi-drug resistance protein 1 (MDR1), cytochromes P450 2D6 (CYP2D6) and 3A4 (CYP3A4). (**C**). Wilcoxon test on opioid agonists' aORs in opioid overdose and their interaction type with receptors: delta opioid receptor (DOR), kappa opioid receptor (KOR) and N-methyl-D-aspartate receptor (NMDAR).

Table 1. Temporal Trends of Opioid Overdose Stratified by Demographics from 2009 to 2017

Year	No. of Opioid Overdose Patients (Opioid Overdose Prevalence, ‰)									aORs [95% CI]
	2009	2010	2011	2012	2013	2014	2015	2016	2017	
Total	395,040	298,317	529,611	645,504	530,520	1,040,515	1,034,306	771,903	569,153	
All Patients	1,477	1,240	2,749	4,022	3,844	7,689	8,313	8,527	6,783	1.16 ****
	(3.7)	(4.2)	(5.2)	(6.2)	(7.2)	(7.4)	(8.0)	(11.0)	(11.9)	[1.15-1.16]
Age Groups										
<18	86	77	77	162	102	307	305	200	169	1.08 ****
	(1.2)	(1.4)	(0.9)	(1.5)	(1.5)	(1.6)	(1.6)	(1.7)	(2.3)	[1.06-1.10]
18-30	207	165	366	448	406	790	900	793	612	1.10 ****
	(4.2)	(4.5)	(5.4)	(5.1)	(5.5)	(5.6)	(6.5)	(8.0)	(9.2)	[1.09-1.12]
31-40	147	124	291	387	363	714	804	774	583	1.12 ****
	(4.2)	(4.8)	(5.7)	(6.1)	(6.8)	(7.0)	(7.8)	(9.9)	(10.3)	[1.10-1.14]
41-50	192	212	474	613	545	1,016	1,000	1,063	839	1.14 ****
	(5.0)	(7.5)	(8.7)	(9.6)	(10.8)	(11.3)	(11.4)	(15.8)	(16.9)	[1.13-1.15]
51-64	357	273	700	1,105	1,102	2,152	2,339	2,506	2,036	1.18 ****
	(5.0)	(5.0)	(6.6)	(8.9)	(10.5)	(11.2)	(12.3)	(16.5)	(17.3)	[1.17-1.19]
65+	488	389	841	1,307	1,326	2,710	2,965	3,191	2,544	1.18 ****
	(3.8)	(4.0)	(5.0)	(6.6)	(7.4)	(8.3)	(9.2)	(12.4)	(12.4)	[1.17-1.18]
Gender										
Female	841	692	1,560	2,314	2,235	4,448	4,771	4,873	3,836	1.16 ****
	(3.7)	(4.0)	(5.2)	(6.2)	(7.3)	(7.4)	(8.1)	(11.2)	(12.1)	[1.15-1.17]
Male	636	548	1,187	1,708	1,609	3,240	3,542	3,654	2,947	1.15 ****
	(3.9)	(4.3)	(5.2)	(6.2)	(7.2)	(7.3)	(8.0)	(11.0)	(11.7)	[1.14-1.16]
Race										
African American	143	120	371	517	389	821	975	1,060	1,051	1.20 ****

(2.2)	(2.4)	(3.9)	(4.6)	(4.6)	(5.6)	(6.6)	(9.0)	(10.8)	[1.19-1.22]
Asian									
6	2	17	29	37	65	90	98	71	1.20 ****
(1.2)	(0.5)	(2.2)	(2.5)	(3.4)	(3.1)	(4.2)	(5.2)	(6.0)	[1.14-1.27]
Caucasian									
1,240	1,054	2,198	3,216	3,130	6,207	6,604	6,717	5,068	1.15 ****
(4.3)	(4.9)	(5.9)	(7.3)	(8.5)	(8.4)	(9.1)	(12.5)	(13.4)	[1.14-1.16]
Hispanic									
23	16	39	41	19	44	29	25	17	1.14 ****
(2.4)	(2.2)	(2.9)	(3.7)	(4.0)	(4.3)	(3.5)	(6.9)	(6.1)	[1.08-1.20]
Native American									
3	5	14	25	51	123	137	135	92	1.20 ****
(3.0)	(7.8)	(3.6)	(3.9)	(7.3)	(7.7)	(8.7)	(15.2)	(11.6)	[1.14-1.26]
Others									
62	43	110	194	218	429	478	492	484	1.14 ****
(2.4)	(2.3)	(2.8)	(3.1)	(4.1)	(4.1)	(4.2)	(5.6)	(6.7)	[1.11-1.16]

Marital Status

Divorced									
153	117	364	502	491	936	1,045	1,082	850	1.16 ****
(6.6)	(6.6)	(9.0)	(10.6)	(12.1)	(13.0)	(14.9)	(19.4)	(19.7)	[1.14-1.17]
Married									
425	339	980	1,443	1,400	2,733	2,975	3,247	2,587	1.18 ****
(3.3)	(3.6)	(5.1)	(6.1)	(6.8)	(7.2)	(8.1)	(11.4)	(12.1)	[1.17-1.18]
Single									
403	328	855	1,338	1,242	2,583	2,799	2,723	2,230	1.13 ****
(3.0)	(3.2)	(4.3)	(5.1)	(6.2)	(5.9)	(6.3)	(8.4)	(9.8)	[1.12-1.14]
Widowed									
152	87	390	535	562	1,030	1,059	1,078	766	1.16 ****
(3.7)	(2.9)	(6.2)	(7.3)	(8.8)	(9.4)	(10.2)	(13.4)	(12.4)	[1.15-1.18]
Others									
344	369	160	204	149	407	435	397	350	1.16 ****
(5.0)	(6.7)	(4.5)	(7.1)	(8.4)	(9.2)	(9.5)	(14.5)	(15.1)	[1.14-1.17]

Census Region

Midwest									
328	272	564	656	625	1,409	1,703	1,934	1,070	1.18 ****
(3.6)	(4.2)	(4.7)	(5.9)	(6.6)	(6.9)	(7.4)	(13.3)	(11.9)	[1.17-1.19]
Northeast									
828	736	961	1,412	835	2,155	1,720	1,210	1,143	1.11 ****

	(4.7)	(5.3)	(5.7)	(6.5)	(8.2)	(7.9)	(7.8)	(9.2)	(10.9) [1.10-1.12]
South	201	185	902	987	1,020	1,990	2,519	3,109	3,296	1.22 ****
	(2.0)	(2.4)	(4.9)	(5.3)	(5.6)	(6.2)	(7.9)	(11.7)	(12.9) [1.20-1.22]
West	110	47	322	967	1,364	2,135	2,371	2,274	1,274	1.11 ****
	(4.2)	(2.6)	(5.7)	(7.3)	(8.9)	(8.8)	(8.9)	(9.9)	(10.7) [1.10-1.12]
Urbanicity									
Rural	112	117	435	777	528	1,404	1,494	1,172	967	1.14 ****
	(2.1)	(2.2)	(5.0)	(5.7)	(5.5)	(6.7)	(6.9)	(8.0)	(8.8) [1.13-1.16]
Urban	1,365	1,123	2,314	3,245	3,316	6,285	6,819	7,355	5,816	1.16 ****
	(4.0)	(4.6)	(5.2)	(6.4)	(7.6)	(7.6)	(8.3)	(11.8)	(12.7) [1.15-1.16]

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

Table 2. Demographics as Risk Factors for Opioid Overdose and Adjusted Odds Ratios from 2009 to 2017

Year	2009	2010	2011	2012	2013	2014	2015	2016	2017	Overall
Adjusted Odds Ratios [95% CI]										
Age Groups [a]										
<18	0.28 ****	0.32 ****	0.20 ****	0.22 ****	0.21 ****	0.18 ****	0.17 ****	0.14 ****	0.19 ****	0.18 ****
	[0.21-0.36]	[0.24-0.42]	[0.15-0.25]	[0.18-0.26]	[0.17-0.26]	[0.16-0.21]	[0.15-0.19]	[0.12-0.16]	[0.16-0.22]	[0.17-0.19]
18-30	1.10	1.07	1.24 **	0.81 ***	0.78 ****	0.67 ****	0.72 ****	0.66 ****	0.74 ****	0.74 ****
	[0.91-1.32]	[0.88-1.31]	[1.08-1.43]	[0.72-0.91]	[0.69-0.88]	[0.61-0.73]	[0.66-0.78]	[0.60-0.72]	[0.67-0.82]	[0.71-0.76]
31-40	1.19	1.22	1.37 ****	1.01	1.03	0.88 **	0.90 **	0.84 ****	0.86 **	0.93 ****
	[0.98-1.44]	[0.99-1.51]	[1.18-1.58]	[0.89-1.14]	[0.91-1.16]	[0.81-0.97]	[0.82-0.97]	[0.78-0.92]	[0.78-0.94]	[0.89-0.96]
41-50	1.31 **	1.82 ****	1.95 ****	1.50 ****	1.54 ****	1.37 ****	1.24 ****	1.31 ****	1.38 ****	1.36 ****
	[1.10-1.57]	[1.53-2.17]	[1.72-2.21]	[1.35-1.66]	[1.39-1.72]	[1.27-1.48]	[1.15-1.34]	[1.21-1.41]	[1.27-1.50]	[1.31-1.40]
51-64	1.33 ****	1.21 *	1.46 ****	1.39 ****	1.50 ****	1.36 ****	1.33 ****	1.34 ****	1.39 ****	1.35 ****
	[1.16-1.54]	[1.03-1.42]	[1.31-1.63]	[1.28-1.52]	[1.38-1.64]	[1.28-1.44]	[1.26-1.41]	[1.26-1.41]	[1.31-1.48]	[1.32-1.39]
Gender [b]										
Male	1.12 *	1.09	1.08	0.99	0.97	0.97	0.98	0.95 *	0.92 **	0.98 *
	[1.00-1.24]	[0.97-1.22]	[1.00-1.17]	[0.93-1.06]	[0.91-1.04]	[0.92-1.02]	[0.93-1.02]	[0.91-0.99]	[0.88-0.97]	[0.96-1.00]
Race [c]										
African American	0.66 ****	0.61 ****	0.62 ****	0.64 ****	0.58 ****	0.69 ****	0.73 ****	0.70 ****	0.78 ****	0.68 ****
	[0.55-0.79]	[0.50-0.75]	[0.56-0.70]	[0.58-0.70]	[0.52-0.65]	[0.64-0.75]	[0.68-0.78]	[0.65-0.74]	[0.73-0.84]	[0.66-0.70]
Asian	0.38 *	0.15 **	0.39 ***	0.37 ****	0.44 ****	0.41 ****	0.52 ****	0.49 ****	0.50 ****	0.47 ****
	[0.17-0.85]	[0.04-0.61]	[0.24-0.63]	[0.26-0.53]	[0.31-0.60]	[0.32-0.52]	[0.42-0.64]	[0.40-0.60]	[0.39-0.63]	[0.42-0.51]
Hispanic	0.74	0.57 *	0.54 ***	0.58 ***	0.54 **	0.74 *	0.56 **	0.68	0.54 *	0.53 ****
	[0.49-1.12]	[0.35-0.94]	[0.39-0.74]	[0.42-0.79]	[0.34-0.84]	[0.55-1.00]	[0.39-0.81]	[0.46-1.01]	[0.33-0.87]	[0.46-0.60]
Native American	0.65	1.58	0.56 *	0.47 ***	0.67 **	0.79 *	0.85	1.17	0.76 *	0.84 ****
	[0.21-2.01]	[0.65-3.83]	[0.33-0.95]	[0.32-0.70]	[0.50-0.88]	[0.66-0.95]	[0.71-1.01]	[0.98-1.39]	[0.61-0.94]	[0.77-0.91]

Others	0.57 ****	0.51 ****	0.54 ****	0.46 ****	0.56 ****	0.57 ****	0.56 ****	0.57 ****	0.56 ****	0.57 ****
	[0.44-0.74]	[0.37-0.69]	[0.44-0.65]	[0.40-0.53]	[0.49-0.65]	[0.51-0.63]	[0.51-0.61]	[0.52-0.62]	[0.51-0.62]	[0.55-0.60]
Marital Status [d]										
Divorced	1.23 *	1.19	1.26 ***	1.15 *	1.14 *	1.16 ***	1.28 ****	1.25 ****	1.22 ****	1.19 ****
	[1.00-1.50]	[0.95-1.49]	[1.11-1.44]	[1.03-1.28]	[1.02-1.27]	[1.07-1.25]	[1.19-1.38]	[1.16-1.35]	[1.12-1.32]	[1.15-1.23]
Married	0.65 ****	0.72 ****	0.74 ****	0.71 ****	0.69 ****	0.70 ****	0.75 ****	0.82 ****	0.82 ****	0.73 ****
	[0.56-0.75]	[0.61-0.85]	[0.67-0.82]	[0.65-0.77]	[0.64-0.75]	[0.66-0.74]	[0.71-0.80]	[0.77-0.86]	[0.77-0.87]	[0.71-0.75]
Widowed	0.82	0.66 **	1.12	0.90	0.97	0.94	0.96	0.96	0.85 ***	0.89 ****
	[0.66-1.03]	[0.50-0.86]	[0.96-1.29]	[0.80-1.01]	[0.86-1.09]	[0.86-1.02]	[0.89-1.05]	[0.88-1.04]	[0.77-0.93]	[0.86-0.93]
Others	0.97	1.20 *	0.79 **	1.03	0.93	1.10	1.12 *	1.20 ***	1.20 **	0.92 ****
	[0.83-1.14]	[1.02-1.42]	[0.66-0.94]	[0.89-1.20]	[0.78-1.11]	[0.99-1.22]	[1.01-1.24]	[1.08-1.34]	[1.07-1.35]	[0.88-0.96]
Census Region [e]										
Midwest	1.78 ****	1.60 ****	0.92	1.11 *	1.14 **	1.15 ****	0.97	1.19 ****	0.94	1.01
	[1.49-2.13]	[1.32-1.94]	[0.83-1.02]	[1.01-1.23]	[1.03-1.26]	[1.08-1.24]	[0.91-1.03]	[1.13-1.26]	[0.88-1.01]	[0.98-1.04]
Northeast	2.39 ****	2.12 ****	1.17 **	1.28 ****	1.49 ****	1.30 ****	1.02	0.84 ****	0.96	1.00
	[2.03-2.80]	[1.78-2.52]	[1.06-1.29]	[1.18-1.40]	[1.35-1.63]	[1.22-1.39]	[0.96-1.09]	[0.79-0.90]	[0.89-1.03]	[0.97-1.02]
West	2.59 ****	1.47 *	1.33 ****	1.52 ****	1.78 ****	1.55 ****	1.23 ****	1.03	1.13 **	1.32 ****
	[2.04-3.31]	[1.06-2.05]	[1.16-1.52]	[1.38-1.67]	[1.63-1.94]	[1.45-1.65]	[1.16-1.31]	[0.97-1.09]	[1.05-1.21]	[1.28-1.36]
Urbanicity [f]										
Rural	0.46 ****	0.51 ****	0.97	0.91 *	0.69 ****	0.90 ***	0.86 ****	0.83 ****	0.81 ****	0.84 ****
	[0.38-0.56]	[0.41-0.62]	[0.87-1.08]	[0.83-0.98]	[0.62-0.76]	[0.85-0.96]	[0.81-0.92]	[0.78-0.89]	[0.75-0.88]	[0.82-0.86]

a: Reference - patients in 65+ age group. **b**: Reference - female patients. **c**: Reference - Caucasian patients. **d**: Reference - patients in single marital status. **e**: Reference - patients from South census region. **f**: Reference - patients in urban area. * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001.

Table 3. Relevant Properties of Prescription Opioids and the Overall Prevalence (‰) between 2009 and 2017

Prescription Opioids [a]	Physicochemical Properties [b]			Pharmacokinetic Properties [c]			Pharmacodynamic Properties [d]			Overall Prevalence
	logP	PSA	MW	MDR1	CYP2D6	CYP3A4	DOR	KOR	NMDAR	
Agonist										
Morphine	1.0	52.9	285.3	+	-	+	+	+	-	342.1
Fentanyl	4.1	23.6	336.5	+	-	+	+	+	-	297.5
Oxycodone	1.0	59.0	315.4	-	+	+	+	+	-	263.3
Hydromorphone	1.7	49.8	285.3	-	-	-	+	+	-	242.1
Hydrocodone	2.1	38.8	299.4	-	+	+	+	-	-	182.4
Meperidine	2.9	29.5	247.3	-	+	+	-	+	+	82.3
Tramadol	2.7	32.7	263.4	+	+	+	+	+	+	55.7
Codeine	1.2	41.9	299.4	-	+	+	+	+	-	43.9
Loperamide	4.4	43.8	477.0	+	+	+	+	+	-	23.1
Diphenoxylate	5.7	53.3	452.6	-	-	-	+	-	-	8.4
Methadone	4.1	20.3	309.4	+	+	+	+	-	+	7.8
Partial Agonist										
Nalbuphine	2.0	73.2	357.4	-	-	-	+	+	-	45.7
Butorphanol	3.7	43.7	327.5	-	-	-	+	+	-	29.5
Buprenorphine	4.5	62.2	467.6	+	+	+	+	+	-	3.2
Antagonist										
Naloxone	1.5	70.0	327.4	-	-	+	+	+	-	157.8

a: Prescription opioids are categorized by their action on opioid mu receptor (MOR), ordered by their overall prevalence. **b**: Relevant physicochemical properties include lipophilicity (logP), polar surface area (PSA) and molecular weight (MW). **c**: Relevant pharmacokinetic properties focus on whether the opioid interacts with 1) drug efflux transporter, multi-drug resistance protein 1 (MDR1) 2) metabolizing enzymes, cytochromes P450 2D6 (CYP2D6) and 3A4 (CYP3A4). **d**: Relevant pharmacodynamic properties focus on whether the opioid interacts with 1) other opioid receptors, delta opioid receptor (DOR) and kappa opioid receptor (KOR) 2) non-opioid receptor in modulating opioid analgesia, N-methyl-D-aspartate receptor (NMDAR). N.A. = Not Applicable.

Table 4. Prescription Opioid Exposures as Risk Factors for Opioid Overdose and Adjusted Odds Ratios from 2009 to 2017

	Adjusted Odds Ratios [95% CI]									
Year	2009	2010	2011	2012	2013	2014	2015	2016	2017	Overall
Agonist										
Morphine	1.58 ****	1.31 ****	1.36 ****	1.36 ****	1.33 ****	1.49 ****	1.56 ****	1.53 ****	1.59 ****	1.44 ****
	[1.41-1.76]	[1.15-1.48]	[1.26-1.47]	[1.28-1.45]	[1.25-1.42]	[1.42-1.56]	[1.49-1.63]	[1.46-1.60]	[1.51-1.67]	[1.42-1.47]
Fentanyl	1.05	1.22 **	1.23 ****	1.24 ****	1.56 ****	1.27 ****	1.33 ****	1.47 ****	1.51 ****	1.22 ****
	[0.94-1.17]	[1.07-1.38]	[1.14-1.33]	[1.16-1.32]	[1.45-1.68]	[1.22-1.33]	[1.27-1.39]	[1.40-1.54]	[1.42-1.60]	[1.20-1.25]
Oxycodone	1.72 ****	1.68 ****	1.74 ****	1.60 ****	1.78 ****	1.73 ****	1.82 ****	1.89 ****	1.90 ****	1.70 ****
	[1.54-1.92]	[1.49-1.90]	[1.60-1.88]	[1.50-1.71]	[1.66-1.91]	[1.65-1.82]	[1.74-1.91]	[1.80-1.98]	[1.80-2.01]	[1.67-1.73]
Hydromorphone	1.97 ****	1.66 ****	1.88 ****	1.64 ****	1.71 ****	1.76 ****	1.79 ****	1.95 ****	1.89 ****	1.81 ****
	[1.74-2.23]	[1.45-1.91]	[1.73-2.04]	[1.54-1.75]	[1.59-1.83]	[1.68-1.84]	[1.71-1.87]	[1.86-2.04]	[1.80-1.99]	[1.77-1.84]
Hydrocodone	1.24 **	1.01 a	1.19 ***	1.31 ****	1.17 ****	1.26 ****	1.23 ****	1.25 ****	1.18 ****	1.12 ****
	[1.08-1.42]	[0.85-1.20]	[1.08-1.30]	[1.22-1.41]	[1.08-1.27]	[1.20-1.33]	[1.17-1.30]	[1.18-1.33]	[1.10-1.27]	[1.09-1.15]
Meperidine	1.23	0.98 a	1.09	1.13 *	1.15 *	1.07	1.08 *	1.14 **	1.01 a	1.09 ****
	[0.94-1.61]	[0.64-1.50]	[0.96-1.24]	[1.02-1.30]	[1.02-1.30]	[0.99-1.15]	[1.00-1.16]	[1.05-1.23]	[0.93-1.11]	[1.06-1.13]
Tramadol	3.70 ****	3.47 ****	3.01 ****	2.69 ****	2.11 ****	2.21 ****	2.16 ****	1.88 ****	1.50 ****	2.20 ****
	[3.10-4.41]	[2.86-4.20]	[2.68-3.38]	[2.45-2.95]	[1.91-2.34]	[2.08-2.36]	[2.03-2.30]	[1.76-2.01]	[1.36-1.65]	[2.14-2.27]
Codeine	0.96	0.81	0.96	0.95	0.71 ***	0.89	0.89 *	0.69 ****	0.73 **	0.78 ****
	[0.75-1.22]	[0.58-1.13]	[0.79-1.15]	[0.81-1.12]	[0.58-0.85]	[0.78-1.01]	[0.78-1.00]	[0.60-0.80]	[0.60-0.89]	[0.74-0.83]
Loperamide	2.08 ****	1.41	2.20 ****	2.16 ****	2.11 ****	2.12 ****	2.17 ****	1.96 ****	1.63 ****	2.03 ****
	[1.57-2.75]	[0.95-2.10]	[1.85-2.61]	[1.87-2.49]	[1.80-2.47]	[1.92-2.35]	[1.98-2.39]	[1.77-2.17]	[1.44-1.85]	[1.94-2.12]
Diphenoxylate	1.63 *	1.69	2.06 ****	1.88 ****	1.57 **	1.33 **	1.39 ***	1.69 ****	1.79 ****	1.48 ****
	[1.01-2.64]	[0.98-2.93]	[1.56-2.72]	[1.45-2.43]	[1.16-2.12]	[1.10-1.62]	[1.16-1.67]	[1.37-2.09]	[1.38-2.32]	[1.36-1.61]
Methadone	7.86 ****	6.36 ****	7.12 ****	7.07 ****	6.59 ****	6.78 ****	6.47 ****	6.53 ****	5.46 ****	6.51 ****
	[6.34-9.74]	[4.88-8.28]	[6.10-8.32]	[6.23-8.03]	[5.66-7.67]	[6.14-7.49]	[5.86-7.1]	[5.85-7.30]	[4.78-6.22]	[6.23-6.80]

Partial Agonist

Nalbuphine	0.38 ****	0.38 ****	0.46 ****	0.54 ****	0.63 ****	0.46 ****	0.57 ****	0.52 ****	0.91 ****	0.56 ****
	[0.26-0.57]	[0.25-0.60]	[0.34-0.61]	[0.43-0.66]	[0.50-0.79]	[0.39-0.54]	[0.50-0.66]	[0.45-0.60]	[0.78-1.07]	[0.53-0.60]
Butorphanol	0.19 ****	0.08 ***	0.10 ****	0.13 ****	0.06 ****	0.11 ****	0.05 ****	0.11 ****	0.08 ****	0.09 ****
	[0.09-0.44]	[0.02-0.34]	[0.05-0.19]	[0.06-0.29]	[0.03-0.14]	[0.08-0.17]	[0.03-0.08]	[0.06-0.18]	[0.04-0.16]	[0.07-0.11]
Buprenorphine	3.14 ****	1.90	2.33 ****	3.37 ****	5.14 ****	3.35 ****	2.77 ****	2.56 ****	1.74 ****	2.89 ****
	[1.80-5.46]	[0.78-4.61]	[1.58-3.44]	[2.48-4.60]	[3.73-7.10]	[2.73-4.11]	[2.19-3.49]	[2.05-3.21]	[1.29-2.35]	[2.63-3.19]

Antagonist

Naloxone	6.40 ****	5.70 ****	4.19 ****	3.93 ****	2.80 ****	3.50 ****	3.13 ****	2.49 ****	2.53 ****	3.24 ****
	[5.70-7.19]	[5.00-6.51]	[3.85-4.55]	[3.68-4.20]	[2.61-3.01]	[3.34-3.67]	[3.00-3.28]	[2.38-2.61]	[2.39-2.67]	[3.18-3.31]

a: Power less than 80%. * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. N.A. = Not Applicable.