

1 Global projections of lives saved from COVID-19 with universal 2 mask use

3 IHME COVID-19 Forecasting Team

4 Emmanuela Gakidou, Ph.D.,^{1,2} David M. Pigott, Ph.D.,^{1,2} Christopher E. Troeger, M.P.H.,¹ Erin N. Hulland,
5 M.P.H.,^{1,3} Robert C. Reiner Jr, Ph.D.,^{1,2} Christopher Adolph, Ph.D.,^{4,5} Aleksandr Y. Aravkin, Ph.D.,^{6,1,2}
6 Steven D. Bachmeier, M.S.,¹ Bree Bang-Jensen, M.A.,⁴ Ryan M. Barber, B.S.,¹ Catherine Bisignano,
7 M.P.H.,¹ Sabina Bloom, B.A.,¹ Austin Carter, M.P.H.,¹ Emma Castro, M.S.,¹ Suman Chakrabarti, M.A.,¹
8 Jhilik Chattopadhyay, M.S.,¹ Christine Chen, B.A.,¹ Rebecca M. Cogen, B.A.,¹ James K. Collins, B.S.,¹ Emily
9 Combs, B.A.,¹ Natalie Maria Cormier, M.P.S.A.,¹ Xiaochen Dai, Ph.D.,¹ William James Dangel, M.Ed.,¹
10 Amanda Deen, M.P.H.,¹ Samuel B. Ewald, M.S.,¹ Abraham D. Flaxman, Ph.D.,^{1,2} Nancy Fullman, M.P.H.,¹
11 Gaorui Guo, M.P.H.,¹ Kathryn Hackman, B.A.,¹ Jiawei He, M.Sc.,¹ Nathaniel J. Henry, B.S.,^{1,7} Casey
12 Johanns, B.S.,¹ Samantha Leigh Larson, B.S.,¹ Alice Lazzar-Atwood, B.S.,¹ Kate E. LeGrand, M.P.H.,¹ Haley
13 Lescinsky, B.A.,¹ Stephen S. Lim, Ph.D.,^{1,2} Emily Linebarger, B.A.,¹ Rafael Lozano, M.D.,^{1,2} Rui Ma, M.S.,^{1,2}
14 Beatrice Magistro, M.A.,⁴ Johan C. Månsson, M.S.,¹ Laurie B. Marczak, Ph.D.,¹ Molly K. Miller-Petrie,
15 M.Sc.,⁸ Ali H. Mokdad, Ph.D.,^{1,2} Meghan D. Mooney, B.S.,¹ Christopher M. Odell, M.P.P.,¹ James K.
16 O'Halloran, M.S.I.M.,¹ Samuel M. Ostroff, Ph.D.,^{1,9} Maja Pasovic, M.A.,¹ Disha J. Patel, M.P.H.,¹ Louise
17 Penberthy, M.S.,¹ Rebecca E. Ramshaw, M.P.H.,¹ Gregory A. Roth, M.D.,^{10,1,2} David H. Shaw, B.A.,¹
18 Brittney S. Sheena, B.S.,¹ Aleksei Sholokhov, B.S.,^{6,1} Reed J. D. Sorensen, M.P.H.,^{1,3} Gianna Sparks, B.A.,¹
19 Emma Elizabeth Spurlock, B.A.,¹ Michelle L. Subart, B.A.,¹ Ally Walker, M.A.,¹ Alexandria Watson, B.A.,¹
20 Catherine A. Welgan, B.S.,¹ Kirsten E. Wiens, Ph.D.,¹¹ Lauren Woyczynski, B.A.,¹ Liming Xu, M.E.C.,¹ Simon
21 I. Hay, F.Med.Sci.,^{1,2} Michael Brauer, D.Sc.,^{12,1} Theo Vos, Ph.D.,^{1,2} Christopher J. L. Murray, Ph.D.^{1,2,*}

22
23 Affiliations

24 ¹Institute for Health Metrics and Evaluation, University of Washington, Seattle, WA, United States

25 ²Department of Health Metrics Sciences, School of Medicine, University of Washington, Seattle, WA,
26 United States

27 ³Department of Global Health, University of Washington, Seattle, WA, United States

28 ⁴Department of Political Science, University of Washington, Seattle, WA, United States

29 ⁵Center for Statistics and the Social Sciences, University of Washington, Seattle, WA, United States

30 ⁶Department of Applied Mathematics, University of Washington, Seattle, WA, United States

31 ⁷Big Data Institute, University of Oxford, Oxford, United Kingdom

32 ⁸Department of Environmental and Occupational Health Sciences, University of Washington, Seattle,
33 WA, United States

34 ⁹Henry M Jackson School of International Studies, University of Washington, Seattle, WA, United States

35 ¹⁰Division of Cardiology, University of Washington, Seattle, WA, United States

36 ¹¹Department of Epidemiology, Johns Hopkins University, Baltimore, MD, United States

37 ¹²School of Population and Public Health, University of British Columbia, Vancouver, BC, Canada.

38 • Corresponding author Christopher J.L. Murray: cjlm@uw.edu

39

40 ABSTRACT

41 BACKGROUND. Social distancing mandates (SDM) have reduced health impacts from COVID-19 but also
42 resulted in economic downturns that have led many nations to relax SDM. Until deployment of an
43 efficacious and equitable vaccine, intervention options to reduce COVID-19 mortality and minimize
44 restrictive SDM are sought by society.

45 METHODS. A susceptible-exposed-infectious-recovered (SEIR) deterministic transmission model was
46 parameterized with data on reported deaths, cases, and select covariates to predict infections and
47 deaths from COVID-19 through March 01, 2021. We explore three scenarios: a “non-adaptive” scenario
48 where neither mask use or SDM adapt to changing conditions, a “reference” where current national
49 levels of mask use are maintained and SDM reintroduced when deaths rise, and an increase in mask use
50 to 95% coverage levels (“universal mask”). We reviewed published studies to set priors on the
51 magnitude of reduction in transmission through increasing mask use.

52 RESULTS. Mask use was estimated at 59.0% of people globally on October 19, 2020. Universal mask use
53 could avert 733,310 deaths (95% UI 385,981 to 1,107,759) between October 27, 2020 and March 01,
54 2021, the difference between the predicted 2.95 million deaths (95% UI 2.70 to 3.35) in the reference
55 scenario and 2.22 million deaths (95% UI 2.00 to 2.45) in the universal mask scenario over this time
56 period.

57 CONCLUSIONS. The cumulative toll of the COVID-19 pandemic could be substantially reduced by the
58 universal adoption of masks before the availability of a vaccine. This low-cost, low-barrier policy,
59 whether customary or mandated, has enormous health benefits with presumed marginal economic
60 costs.

61

62 COVID-19 has spread to all regions of the world and as of October 25, 2020 nearly 43 million cases and
63 1,195,034 deaths have been reported globally.¹ In March 2020, nearly all nations put in place various
64 social distancing mandates (SDM) (Figure 1) that have contributed to reducing the effects of COVID-19.²⁻
65 ⁵ The economic downturns associated with these measures subsequently led most nations to
66 progressively relax SDM. As COVID-19 transmission and deaths rise globally and rebound in many
67 countries,¹ policymakers are keen to identify policy options to reduce COVID-19 mortality without re-
68 imposing strict SDM before the widespread implantation of a vaccine.

69 Despite welcome news on vaccine development and efficacy of some candidates,^{6,7} safety, regulatory,
70 manufacturing and distribution hurdles all need to be overcome before there will be an impact at scale.⁸
71 One existing and attractive strategy is the wearing of masks in public spaces or when physical distancing
72 may not be possible. Several systematic reviews have suggested that cloth and other non-medical masks
73 worn by the general public can reduce transmission.^{9,10} Moreover, simple non-medical masks are
74 affordable and equitably accessible. Consequently, a growing number of national and local governments
75 have recommended mask use, with more than 150 countries now mandating mask use.¹¹ In nations
76 such as the United States (US) and Brazil, mask use has become politicized, with widely varying views on
77 the ethics, legality, and enforcement of mask use and mask use mandates.^{12,13} Evidence of the individual
78 benefits and population health impacts of increased mask use is still extremely pertinent to these
79 debates.

80 Masks reduce the transmission of respiratory pathogens *via* a physical barrier; the force of exhalation,
81 coughing, or sneezing leads to impaction and interception of viral droplets (> 5 μ m in diameter) and
82 aerosols (\leq 5 μ m in diameter) onto the fibers of the mask.¹⁴⁻¹⁶ Given greater impaction efficiency for
83 droplets, masks are most effective in blocking outward exhalation,¹⁷ although evidence from studies of
84 air pollution aerosol indicates some protection of non-medical masks in blocking a small fraction of

85 aerosol inhalation.¹⁸ Thus mask wearing is effective in intercepting particle emissions from the infected
86 and reducing large particle transmission to the uninfected.¹⁹

87 In this paper, we use online survey data to assess current levels and trends in use of masks at the
88 national level for 194 countries and quantify the benefits of universal mask use (here defined as 95% of
89 individuals always wearing a mask when outside their home) on COVID-19 mortality in these countries
90 using a COVID-19 transmission dynamics model. We then provide global and national estimates of lives
91 saved through universal mask use between October 27, 2020 and March 01, 2021.

92 **Methods**

93 Our analysis is comprised of two main components: an analysis of survey data on the levels and trends
94 in mask use; and a deterministic transmission dynamics model with categories for susceptible, exposed,
95 infected and recovered (SEIR) model of COVID-19 incidence and death. We explore three scenarios in
96 the SEIR model: 1) a “non-adaptive” where mask use and SDM remain constant 2) a plausible “reference
97 scenario” where SDMs are reinstated at a threshold level of daily deaths and current levels of mask use
98 maintained and 3) a “universal mask” scenario based on the adoption of universal mask use at 95%
99 coverage.

100 We used four main sources of data on self-reported use of masks to estimate past and current mask use
101 globally, and make forecasts for future use: Facebook Global Symptom Survey,²⁰ Facebook US Symptom
102 Survey²¹, PREMISE surveys,²² and YouGov COVID-19 Behaviour Tracker surveys.²³ Pooling estimates on
103 the prevalence of self-reported mask use from these survey series provides a time series on trends in
104 mask use in almost all countries. Between April 23, 2020 and October 19, 2020, Facebook has surveyed
105 30.86 million users from 198 countries and territories using an instrument with multiple items on
106 behaviors related to COVID-19, including mask use. The YouGov surveys cover 29 countries and
107 represent interviews with approximately 726,500 individuals since March 01, 2020 up until October 24,

108 2020. For the US, we used data collected through PREMISE and Facebook to inform the trend in mask
109 use. There were 273,424 total PREMISE responses representing all 50 states and the District of Columbia
110 with responses collected between April 21 and October 26. Facebook symptom surveys in the US started
111 asking about mask use in the beginning of September (data range from September 8 to October 20) and
112 included 1,813,000 respondents. Levels of mask use as reported by PREMISE are on average 30% lower
113 across the USA compared to data from Facebook and YouGov, but other data sources do not provide a
114 time trend in mask use. Therefore, we use the time trend implied by the PREMISE data and apply it to
115 the level of mask use reported in the Facebook data. Details on the specific questions and responses
116 used from each survey are identified in the Supplementary Appendix.

117 Mask use for each location was estimated using a spline-based smoothing process. For locations without
118 data on mask use, we used national level estimates or regional estimates (see Supplementary Appendix
119 for additional details).

120 COVID-19 SEIR model construction

121 The IHME COVID-19 prediction model has been described in detail elsewhere.²⁴ Briefly, we construct an
122 SEIR model for each location we model; the Supplementary Appendix shows the basic health states
123 included in the model and specific transition parameters. The critical driver of the epidemic is the rate at
124 which susceptible individuals become infected in each location which is represented as:

$$\frac{dS}{dt} = -\beta(t) \frac{S(I_1 + I_2)^\alpha}{N}$$

125 Where β_t is the transmission parameter for time period t , α represents a mixing coefficient to account
126 for imperfect mixing within each location, S is the fraction of each location's population that is
127 susceptible, and I_1 and I_2 is the fraction that are infectious. Effective R_t , the number of new infections
128 caused by each case is a simple monotonic transformation of β_t and the fraction of the population that is

129 susceptible. We use an efficient algorithm to directly estimate β_t in the past – see Supplementary
130 Appendix for details.

131 The following covariates are included in the model: population density measured as the share of the
132 population living in areas with more than 2,500 individuals per square kilometer, the fraction of the
133 population living below 100 meters above sea level, smoking prevalence, particulate matter air pollution
134 (PM2.5 population-weighted annual average concentration), mobility measured using cell phone apps,
135 mask use, COVID-19 testing *per capita*, and pneumonia seasonality.²⁴ To avoid over-estimating the
136 effects of pneumonia seasonality and mask use we used constraints on each in the regression – see
137 Supplementary Appendix for details. Specifically, for mask use, we did not let the regression estimate an
138 effect size larger than what was consistent with the mask use meta-analysis of the individual level effect
139 (Supplementary Appendix). To capture uncertainty in the input data, model parameters and regression
140 coefficients linking β_{it} to covariates, we generated 1,000 models for each location – see Supplementary
141 Appendix for details.

142 We evaluated out of sample predictive validity for this modeling approach by holding out the last twelve
143 weeks of data and compared predictions from the held-out data to what occurred; median absolute
144 percent error for cumulative deaths at six weeks was 11% and twelve weeks was 24%.²⁵ We also
145 compared this model to other COVID-19 prediction models that make their estimates publicly available;
146 overall, we find that our model is one of the best performing models at 12-weeks out-of-sample.²⁵

147 **Scenario construction**

148 We used the set of 1,000 SEIR models for each location to generate three scenarios: a plausible
149 reference scenario, a universal mask use scenario, and a scenario wherein SDM continued to be lifted. In
150 the plausible reference scenario key drivers such as mobility and testing *per capita* evolve according to
151 past trends – see Supplementary Appendix for details. In the universal mask use scenario, we

152 additionally assume that mask mandates and other campaigns lead to scale-up of mask use to 95%
153 within 7 days of enactment; this represents the highest reported level of mask use (in Victoria, Australia
154 Singapore) across the three survey platforms during the COVID-19 pandemic up to mid-October; other
155 locations including Chile, Peru, and Spain have also approached this 95% coverage level from previous
156 very low levels of coverage. The “universal mask” scenario assumes this level of mask use can be
157 achieved through the adoption of mask use around the world. Both the reference and universal mask
158 use scenarios additionally assume that SDM will be re-instated for 6 weeks if the daily death rate per
159 location reaches 8 per million people per day.²⁶ Finally, we estimate the trajectory of the epidemic in
160 each location if SDM are not reinstated in response to changing pandemic conditions (“non-adaptive
161 scenario”).

162 Our study follows the Guidelines for Accurate and Transparent Health Estimate Reporting. Data were
163 not obtained from subjects for this study; we used pre-existing datasets identified through online
164 searches, through outreach to institutions that hold relevant data such as ministries of health, or
165 through individual collaborator reference and identification. This study was approved by University of
166 Washington's Human Subjects Division Study ID: STUDY00009060. Author contributions, including the
167 identification of who analyzed data, are supplied in the Supplementary Appendix.

168 Results

169 Figure 2 shows the relative risk of viral respiratory infection among users of different types of
170 facemasks; additional details and results for this analysis are provided in the Supplementary Appendix.

171 We found that non-medical masks among the general population reduced risk of viral respiratory
172 infection by 40% (Relative risk 0.6, 95% CI 0.46 to 0.78).

173 Based on survey data collected through smartphone hosted questionnaires, Figure 3 shows a map of
174 mask use by location as of October 19, 2020, the last date of fully observed data in the model. Mask use

175 is high in several locations that were at the forefront of the COVID-19 peak in the spring of 2020
176 including Italy, Spain, and most parts of Latin America (*i.e.* Chile, and Panama). The highest national
177 mask use on October 19, 2020 was in Spain (93.9%), followed by Puerto Rico (92.6%), and Panama
178 (92.5%). The lowest rates are seen in Northern Europe (Sweden, Norway, and Denmark < 4%) and East
179 Africa (United Republic of Tanzania 2.1%). On October 19, women were more likely to report always
180 wearing a mask (64% women, 58% men), with pronounced sex differences in sub-Saharan Africa, Central
181 and Eastern Europe and Central Asia (See Supplementary Appendix). Mask use was highest among
182 people who live in cities and towns (about 60%) and lowest in rural communities (less than 50%; see
183 Supplementary Appendix). People over 65 years (56.4%) and young adults 18-24 (56.7%) were the least
184 likely age groups to report always wearing a mask (other adult age groups were over 60% at the global
185 level; see Supplementary Appendix).

186 Figure 4 shows the global and WHO regional trends in mask use since the beginning of the epidemic.
187 Mask use data start only in the beginning of April so the rapid expansion from likely a very low baseline
188 pre-COVID-19 outside of East Asia most probably occurred in March. Mask use was estimated at 59.0%
189 of people globally on October 19, 2020, ranging from 36.0% in the Eastern Mediterranean Region to
190 72.2% in the Region of the Americas (Figure 4). Mask use has increased in some countries where
191 mandates have been put into place such as Belgium, Ireland and the Ethiopia (SI Figure 7). Mask use has
192 declined in some settings in parallel with declining death rates such as Poland, Czechia, and Italy (SI
193 Figure 7).

194 Global projections of COVID-19 deaths with and without adoption of universal mask use are shown in
195 Figure 5. In our reference scenario, we expect daily COVID-19 deaths to increase in November through
196 December leading to as many as 2.95 million deaths (95% UI 2.70 million to 3.35 million) by March 01,
197 2021. Uncertainty in daily deaths and cumulative deaths increases over time such that there is
198 considerable uncertainty in global deaths by March 01, 2021. In contrast, the universal mask use

199 scenario leads to a mean estimate of 2.22 million deaths (95% UI 2.00 million to 2.45 million) by March
200 01, 2021. The difference in mortality between the reference and universal mask use scenario suggests
201 that 0.73 million lives (95% UI 0.39 million to 1.11 million) could be saved over this time period if 95% of
202 people were to always wear masks when outside their home. Table 1 provides estimates by location of
203 the expected deaths in the reference and universal mask use scenario along with the number of deaths
204 saved through universal mask use. The countries where universal mask use would have the largest effect
205 are populous nations where mask use is currently relatively low such as India (104,301 fewer deaths in
206 universal mask scenario, 95% UI 60,679 to 151,873 deaths), the Germany (90,005 fewer deaths; 95% UI
207 27,524 to 190,010 deaths), and United States of America (82,150 fewer deaths; 95% UI 46,079 to
208 116,120 deaths). The greatest magnitude difference in mortality rate occurred in the Czechia (12,419
209 fewer deaths, 117 deaths per 100,000), Germany (90,005 fewer deaths, 106 deaths per 100,000) and
210 Croatia (3,879 fewer deaths, 91 deaths per 100,000).

211 Figure 6 shows a map of the percent reduction in expected deaths from the reference scenario
212 compared to the mask use scenario in the deaths from October 27, 2020 to March 01, 2021. The largest
213 percent reduction in expected deaths occurred in Belarus (76.8% difference, 95% UI 66.9 to 82.3%),
214 Denmark (76.1% difference, 95% UI 61.1 to 87.3%), and Uganda (75.4% difference, 95% UI 38.4 to
215 97.0%). There are large variations in the number of lives saved globally, regionally, nationally, and
216 subnationally and these are detailed in full in Table 1.

217 Discussion

218 Given global mask use is currently at 59.0%, increasing mask use to 95% could decrease cumulative
219 COVID-19 deaths by March 01, 2021 by 0.73 million deaths (95% UI 0.39 to 1.11 million). This represents
220 a 24.8% reduction in the number of deaths expected from October 27, 2020 to March 01, 2021. Our
221 models also show that in many settings, increased mask use could delay the need re-imposition of SDM;

222 in addition to the lives saved, a delay in re-imposition of social mandates might also be accompanied by
223 substantial economic benefits, although this was not specifically modelled.

224 The published studies of SARS-CoV-2 included in our analysis of mask effectiveness (see Supplementary
225 Appendix for details) demonstrated reductions in relative risk of 30% - 100%, with the one study of non-
226 medical mask use amongst the general population, indicating a reduction of 42% for any mask use and
227 70% for consistent mask use.²⁷ Mask effectiveness is also supported by additional evidence from case
228 studies²⁸⁻³⁰ and from laboratory studies that report on the efficacy of masks in reducing exhalation of
229 both aerosols and droplets by those infected with SARS CoV-2.³¹ Based on all types of available
230 evidence, it seems wise to encourage mask use throughout the world as the benefits can be substantial
231 with low to zero contraindications. Guidance for specific materials, handling of face coverings and other
232 considerations is rapidly evolving as additional evidence emerges.³²

233 Given that the cost of masks is very low, mask mandates and/or the promotion of mask use seems
234 prudent, as the risk of adopting these policies is minimal and the potential benefits very large. In
235 recognition of this, over the last few months we have seen the number of countries and territories with
236 mask mandates in place increase substantially. Nevertheless, there remains reluctance to adopt mask
237 use and to impose mask mandates. In some settings, the current epidemiological context means that
238 mask wearing is not viewed as a necessary part of control, such as in Norway where the Norwegian
239 Institute of Public Health determined that, given their current low prevalence, “200,000 people would
240 need to wear facemasks to prevent one new infection per week”.³³ In other settings however, despite
241 increasing cases, public sentiment towards mask wearing hinders universal use. Past messages from
242 some governments have not encouraged mask use and may have actually discouraged mask use.³⁴⁻³⁶

243 Early on, WHO stated “*the wide use of masks by healthy people in the community setting is not*
244 *supported by current evidence and carries uncertainties and critical risks*”³⁷ and only changed their
245 official position on June 5th to encourage mask use.³⁸ For those decision-makers who are concerned with

246 the economic effects of SDM, mask use provides a low-cost strategy to reduce the risk of a further round
247 of SDM and the associated unemployment and economic downturn. Mask use scaled up rapidly in some
248 locations following mandated use. For example, at the start of July, mask use in Victoria state Australia
249 was less than 10% but following a mask mandate in that month, mask use exceeded 90% in early August.

250 While the effective R , the number of new infections created by a single infection under the auspices of
251 control, can potentially be reduced by one-third through universal mask use compared to a reference
252 scenario, mask use alone will likely be insufficient to control the epidemic in many locations. Even with
253 universal mask use, we expect the global death toll due to COVID-19 to reach as high as 1.71 million
254 deaths by the end of the year, and many more in 2021, assuming an efficacious vaccine is not widely
255 deployed in the interim. Despite encouraging news of candidates,^{6,7} safety procedures, regulatory
256 approval, and the problems of manufacturing and distribution at scale will likely delay the potential for
257 widespread impact.⁸ Countries will have to consider other policy strategies to reduce transmission,
258 including increasing testing, contact tracing, and isolation, along with “smart mandates”, that target
259 SDM or restrictions to particular subgroups of the population, such as specific age-groups or local
260 communities for short periods of time. A central policy challenge for many countries is understanding
261 which of these mandates targeting strategies makes the most sense in a given context and at what level
262 of COVID-19 transmission. This is the focus of our ongoing work.

263 The findings of this study should be interpreted while taking into account its limitations (further
264 elaborated in the Supplementary Appendix): (i) the number of studies on general mask use in the
265 population remains low; (ii) all the assumption of any type of SEIR model are also applicable here;²⁴
266 prediction accuracy is highly spatially variable and compound into the future;²⁵ (iv) death reporting
267 globally is subject to regionally and nationally specific errors and biases for which we cannot fully
268 control; (v) models is sensitive to death trend in last 7-14 days and (vi) self-reported mask use by surveys
269 will also have an set of errors and biases that are not completely known.

270 Universal mask use can save many lives and avoid or delay the need for re-imposition of SDM. This will
271 contribute to ameliorating the negative effects of COVID-19 on national economies. Until an affordable
272 and effective vaccine becomes universally available, we find encouraging or mandating mask use is the
273 least disruptive policy option available to countries currently experiences resurgences in infections and
274 deaths. Simple face coverings are cheap and effective; one of the few available interventions that is
275 widely available to everyone. Countries with currently low mask use will need to determine the optimal
276 balance between encouraging the use of masks through advocacy and information about their benefits,
277 and governance of a compulsory use associated with penalties for non-compliance.³⁹

278

279 [Code Availability Statement](#)

280 All code used for these analyses is publicly available online (<http://github.com/ihmeuw/>).

281 [Online content](#)

282 Results for each scenario are accessible through a visualization tool at <http://covid19.healthdata.org>.

283 The estimates presented in this tool will be iteratively updated as new data are incorporated and will
284 ultimately supersede the results in this paper.

285 [Competing interests](#)

286 This study was funded in part by the Bill & Melinda Gates Foundation and Bloomberg Philanthropies.

287 The funders of the study had no role in study design, data collection, data analysis, data interpretation,
288 writing of the final report, or decision to publish. CAdolph reports grants from Benificus Foundation,
289 grants from Center for Statistics and the Social Sciences, during the conduct of the study. ADFlaxman
290 reports personal fees from Kaiser Permanente and NORC, as well as other from Agathos, Ltd., outside
291 the submitted work. The corresponding author had full access to all of the data in the study and had
292 final responsibility for the decision to submit for publication.

293 **Acknowledgements**

294 We thank the various Departments of Health and frontline health professionals who are not only
295 responding to this epidemic daily, but also provide the necessary data to inform this work – IHME wishes
296 to warmly acknowledge the support of these and others who have made our COVID-19 estimation
297 efforts possible (<http://www.healthdata.org/covid/acknowledgements>). This work was supported in
298 part by the Bill & Melinda Gates Foundation, Bloomberg Philanthropies, as well as funding from the
299 state of Washington and the National Science Foundation (2031096). AD Flaxman was supported in part
300 by funding from the National Science Foundation (award DMS-1839116). We also extend a note of
301 particular thanks to John Stanton and Julie Nordstrom for their generous support. We are grateful to
302 Professor Wei Huang from Peking University for helping us extract information from scientific papers in
303 Chinese.

304

305

306 **List of Figures**

307 **Figure 1. Number of social distancing mandates by WHO region and selected countries on a timeline**
308 **starting February 01, 2020 through October 19, 2020.**

309
310 **Figure 2. Relative risk of viral respiratory infection among users of different types of facemasks.**

311 This analysis includes only variables addressing facemask type (non-medical versus medical) and
312 population type of mask user (general population versus healthcare setting). The size of the box is
313 proportional to the precision of the estimate, based on number of observations, with more precise
314 studies having larger boxes.

315
316 **Figure 3. Proportion of the population that self-report always wearing a facemask when outside the**
317 **home on October 19, 2020.**

318
319 **Figure 4. Proportion of the population that self-report always wearing a facemask when outside the**
320 **home by WHO region between February 1, 2020 and March 01, 2021. Values after the last**
321 **date of observed survey data are dashed lines and horizontal projections from the last**
322 **observed values.**

323
324 **Figure 5. Projected global deaths due to COVID-19 up to March 01, 2021 in the scenario for current**
325 **projections of mask use, the non-adaptive scenario, and the scenario with 95% coverage of**
326 **mask use. Projections are shown for the world and for the six WHO regions.**


327
328 **Figure 6. Percent reduction in cumulative deaths on March 01, 2021 in the universal mask use scenario**
329 **and the reference scenario.**

330 Areas have "NA" either because of no available data or because the population size, cases or deaths are
331 so small that the SEIR models do not run.

332

333


334 **Figure 1.**



335

336


337 **Figure 2.**



338

339


340 **Figure 3.**



341

342


343 **Figure 4.**



344

345


346 **Figure 5.**



347

348

349 **Figure 6.**



350

351

Table 1

The difference in cumulative deaths between the universal mask use scenario and the reference (current mask use) scenario globally and by country (and first administrative subnational units where applicable) within WHO regions, on March 01, 2021.

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Global	733,310 (385,981 to 1,107,759)	-24.6% (-34.6 to -14.0%)	2,954,098 (2,698,550 to 3,351,659)
African Region	31,730 (14,284 to 58,110)	-41.7% (-58.2 to -24.5%)	73,614 (54,785 to 101,214)
Algeria	10,282 (4,196 to 20,655)	-73.3% (-87.5 to -54.7%)	13,508 (7,358 to 24,041)
Angola	335 (82 to 1,130)	-25.2% (-45.7 to -11.6%)	1,171 (638 to 2,578)
Benin	140 (0 to 1,119)	-35.2% (-89.8 to -0.7%)	199 (42 to 1,274)
Botswana	14 (0 to 97)	-15.0% (-59.0 to -0.8%)	47 (23 to 196)
Burkina Faso	236 (1 to 1,567)	-44.0% (-94.0 to -2.1%)	321 (70 to 1,698)
Cabo Verde	144 (66 to 237)	-37.0% (-52.7 to -21.0%)	387 (248 to 488)
Cameroon	33 (2 to 135)	-6.6% (-23.8 to -0.6%)	463 (427 to 573)
Central African Republic	40 (1 to 271)	-23.7% (-77.0 to -2.0%)	107 (64 to 349)
Chad	160 (7 to 791)	-38.9% (-80.3 to -6.8%)	300 (109 to 1,047)
Comoros	0 (0 to 0)	-0.8% (-1.1 to -0.5%)	8 (8 to 8)
Congo	154 (9 to 1,044)	-34.5% (-85.8 to -7.2%)	284 (119 to 1,299)
Côte d'Ivoire	436 (3 to 3,393)	-42.1% (-95.1 to -2.0%)	577 (127 to 3,584)
Democratic Republic of the Congo	443 (80 to 2,037)	-38.4% (-78.1 to -15.6%)	918 (509 to 2,634)
Equatorial Guinea	3 (1 to 8)	-3.4% (-9.1 to -1.2%)	87 (85 to 93)
Eswatini	44 (2 to 217)	-20.1% (-60.5 to -1.7%)	174 (121 to 357)
Ethiopia	3,077 (1,214 to 7,835)	-52.2% (-75.0 to -34.6%)	5,587 (3,377 to 10,504)
Gabon	5 (0 to 39)	-5.2% (-35.5 to -0.1%)	63 (54 to 107)
Gambia	98 (4 to 425)	-31.2% (-72.6 to -3.1%)	236 (126 to 603)
Ghana	307 (75 to 1,195)	-33.3% (-67.0 to -14.4%)	810 (513 to 1,796)
Guinea	408 (13 to 2,148)	-60.5% (-93.3 to -14.8%)	519 (89 to 2,311)
Guinea-Bissau	177 (3 to 823)	-52.9% (-91.2 to -6.2%)	242 (49 to 911)
Kenya	9,914 (3,128 to 19,964)	-68.5% (-86.7 to -45.8%)	13,905 (5,902 to 24,512)
Lesotho	33 (0 to 265)	-19.5% (-80.7 to -0.2%)	81 (44 to 322)
Liberia	2 (0 to 10)	-2.7% (-11.0 to -0.4%)	85 (82 to 95)
Madagascar	5 (1 to 17)	-1.9% (-6.1 to -0.5%)	261 (249 to 290)
Malawi	4 (1 to 14)	-2.1% (-6.8 to -0.5%)	196 (188 to 212)
Mali	66 (5 to 342)	-23.5% (-67.8 to -3.3%)	210 (141 to 495)
Mauritania	54 (5 to 213)	-20.9% (-55.9 to -3.2%)	221 (170 to 383)
Mauritius	0 (0 to 0)	-0.3% (-0.5 to -0.2%)	11 (11 to 11)
Mozambique	55 (8 to 286)	-19.8% (-49.9 to -6.6%)	217 (124 to 589)
Namibia	18 (2 to 64)	-9.6% (-26.6 to -1.3%)	172 (142 to 245)
Niger	80 (2 to 759)	-25.8% (-89.6 to -3.1%)	154 (73 to 856)
Nigeria	266 (33 to 958)	-16.0% (-44.0 to -2.8%)	1,455 (1,191 to 2,207)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Rwanda	43 (0 to 316)	-17.6% (-55.7 to -0.7%)	126 (39 to 637)
Sao Tome and Principe	1 (0 to 5)	-6.4% (-23.7 to -1.0%)	17 (16 to 22)
Senegal	150 (69 to 337)	-27.2% (-46.6 to -15.4%)	532 (439 to 726)
Sierra Leone	44 (1 to 297)	-21.0% (-71.7 to -1.3%)	132 (76 to 413)
South Africa	1,457 (180 to 3,935)	-5.5% (-12.8 to -0.9%)	25,084 (20,625 to 32,461)
South Sudan	234 (0 to 1,977)	-39.0% (-92.8 to -0.3%)	316 (55 to 2,098)
Togo	193 (31 to 799)	-57.8% (-88.1 to -29.9%)	285 (101 to 921)
Uganda	1,992 (75 to 10,752)	-75.4% (-97.0 to -38.4%)	2,261 (193 to 11,088)
United Republic of Tanzania	477 (178 to 1,360)	-39.5% (-65.9 to -22.0%)	1,116 (780 to 2,062)
Zambia	9 (1 to 40)	-2.3% (-9.5 to -0.2%)	367 (344 to 434)
Zimbabwe	94 (7 to 527)	-16.5% (-54.7 to -2.7%)	402 (268 to 963)
Eastern Mediterranean Region	85,430 (37,549 to 151,164)	-33.8% (-51.8 to -16.6%)	248,219 (219,986 to 303,019)
Afghanistan	2,864 (599 to 10,488)	-52.2% (-82.9 to -26.2%)	4,735 (2,250 to 12,893)
Bahrain	382 (132 to 628)	-36.9% (-52.6 to -21.6%)	1,013 (575 to 1,271)
Djibouti	128 (0 to 778)	-28.6% (-87.9 to -0.2%)	215 (62 to 1,014)
Egypt	8,687 (3,941 to 22,233)	-51.3% (-75.3 to -34.4%)	16,057 (11,295 to 29,593)
Iran (Islamic Republic of)	10,794 (0 to 25,342)	-12.7% (-29.9 to 0.0%)	86,083 (81,032 to 91,487)
Iraq	14,273 (9,139 to 20,470)	-47.1% (-59.0 to -33.2%)	30,080 (25,712 to 35,048)
Jordan	468 (0 to 3,241)	-6.1% (-42.7 to 0.0%)	7,126 (6,072 to 8,383)
Kuwait	822 (280 to 1,399)	-30.2% (-47.6 to -9.9%)	2,719 (1,723 to 3,372)
Lebanon	1,540 (61 to 2,425)	-45.6% (-69.5 to -1.5%)	3,434 (1,841 to 4,080)
Libya	2,618 (1,318 to 3,400)	-55.2% (-71.0 to -26.5%)	4,765 (3,697 to 5,575)
Morocco	7,351 (1,887 to 13,297)	-30.5% (-55.6 to -7.7%)	24,183 (22,868 to 25,973)
Oman	450 (276 to 715)	-18.4% (-26.7 to -11.5%)	2,436 (2,074 to 2,840)
Pakistan	23,141 (5,635 to 56,754)	-64.7% (-86.0 to -38.9%)	32,756 (14,367 to 66,719)
Azad Jammu & Kashmir	1,187 (388 to 1,951)	-55.1% (-87.9 to -17.8%)	2,182 (1,795 to 2,753)
Balochistan	959 (43 to 4,380)	-65.1% (-94.6 to -21.0%)	1,181 (204 to 4,776)
Gilgit-Baltistan	338 (0 to 852)	-51.4% (-87.0 to -0.5%)	546 (91 to 1,186)
Islamabad Capital Territory	440 (41 to 761)	-39.8% (-67.3 to -3.4%)	1,117 (875 to 1,273)
Khyber Pakhtunkhwa	1,384 (172 to 5,646)	-41.4% (-80.1 to -11.8%)	2,752 (1,475 to 6,989)
Punjab	17,106 (2,067 to 47,426)	-76.1% (-94.0 to -44.9%)	20,339 (4,713 to 51,175)
Sindh	1,726 (375 to 5,214)	-33.2% (-63.4 to -12.0%)	4,639 (3,104 to 8,439)
Qatar	428 (138 to 882)	-41.0% (-62.9 to -19.9%)	1,022 (503 to 1,675)
Saudi Arabia	2,001 (1,060 to 3,395)	-20.2% (-30.6 to -12.2%)	9,759 (8,037 to 11,583)
Somalia	227 (4 to 1,886)	-34.1% (-90.4 to -3.7%)	357 (108 to 2,204)
Sudan	1,671 (2 to 18,812)	-23.6% (-94.6 to -0.2%)	2,539 (837 to 19,741)
Syrian Arab Republic	2,634 (1,015 to 5,748)	-71.9% (-88.8 to -50.9%)	3,536 (1,765 to 6,905)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Tunisia	4,212 (229 to 8,396)	-35.9% (-67.3 to -2.4%)	11,619 (7,962 to 17,998)
United Arab Emirates	735 (115 to 1,737)	-22.9% (-41.8 to -7.5%)	3,186 (1,007 to 5,329)
Yemen	6 (1 to 16)	-1.0% (-2.6 to -0.2%)	600 (593 to 613)
European Region	355,754 (175,846 to 583,089)	-31.1% (-44.2 to -17.6%)	1,126,561 (960,340 to 1,421,258)
Albania	1,238 (701 to 1,666)	-53.3% (-68.3 to -28.8%)	2,328 (2,074 to 2,719)
Andorra	7 (1 to 22)	-5.9% (-15.6 to -1.5%)	113 (91 to 150)
Armenia	1,539 (834 to 2,548)	-29.0% (-43.1 to -17.0%)	5,288 (4,120 to 7,104)
Austria	4,442 (1,472 to 8,797)	-44.1% (-72.9 to -18.9%)	10,005 (6,122 to 16,164)
Azerbaijan	3,320 (698 to 7,080)	-32.9% (-54.4 to -9.3%)	9,738 (6,631 to 15,271)
Belarus	4,835 (3,902 to 5,739)	-76.8% (-82.3 to -66.9%)	6,283 (5,552 to 7,070)
Belgium	6,473 (1,641 to 12,748)	-23.4% (-38.0 to -8.0%)	26,692 (19,781 to 34,565)
Bosnia and Herzegovina	982 (140 to 1,883)	-28.8% (-48.0 to -4.8%)	3,343 (2,669 to 4,733)
Bulgaria	4,847 (1,108 to 10,913)	-43.6% (-68.2 to -14.8%)	10,559 (7,051 to 16,914)
Croatia	3,879 (1,001 to 8,887)	-55.1% (-76.8 to -24.0%)	6,700 (3,853 to 12,411)
Cyprus	134 (1 to 727)	-42.9% (-86.8 to -3.7%)	203 (31 to 1,010)
Czechia	12,419 (6,969 to 20,681)	-49.4% (-67.0 to -30.5%)	25,242 (16,055 to 37,207)
Denmark	3,975 (1,998 to 7,339)	-76.1% (-87.3 to -61.1%)	5,142 (2,988 to 8,576)
Estonia	138 (3 to 548)	-37.7% (-73.8 to -3.2%)	279 (83 to 812)
Finland	501 (108 to 2,032)	-47.6% (-81.3 to -22.4%)	902 (478 to 2,592)
France	19,695 (9,480 to 39,305)	-20.1% (-31.6 to -11.5%)	95,640 (78,657 to 130,050)
Georgia	900 (0 to 2,099)	-29.3% (-61.8 to 0.0%)	2,991 (1,950 to 4,610)
Germany	90,005 (27,524 to 190,010)	-60.6% (-79.6 to -38.4%)	140,720 (69,488 to 257,049)
Baden-Württemberg	21,001 (5,002 to 55,334)	-61.6% (-82.6 to -35.4%)	31,229 (12,664 to 72,101)
Bavaria	12,983 (3,551 to 34,630)	-52.2% (-76.4 to -26.2%)	22,977 (11,657 to 47,058)
Berlin	2,914 (424 to 8,629)	-70.7% (-89.8 to -46.8%)	3,934 (798 to 10,620)
Brandenburg	1,831 (32 to 8,711)	-60.4% (-86.7 to -13.2%)	2,647 (249 to 11,638)
Bremen	145 (6 to 607)	-45.2% (-82.7 to -8.0%)	241 (75 to 765)
Hamburg	851 (81 to 2,629)	-55.8% (-81.8 to -20.1%)	1,340 (398 to 3,477)
Hesse	17,888 (3,016 to 41,332)	-61.7% (-82.3 to -37.2%)	27,695 (7,661 to 62,642)
Lower Saxony	4,273 (1,359 to 9,749)	-69.0% (-86.0 to -48.1%)	5,963 (2,632 to 11,982)
Mecklenburg-Vorpommern	81 (2 to 665)	-41.2% (-90.6 to -7.6%)	117 (28 to 709)
North Rhine-Westphalia	20,416 (5,689 to 59,418)	-58.8% (-81.3 to -31.4%)	32,632 (13,171 to 76,786)
Rhineland-Palatinate	1,521 (362 to 4,208)	-68.1% (-87.1 to -44.0%)	2,104 (796 to 5,067)
Saarland	149 (0 to 1,136)	-19.1% (-72.4 to -0.3%)	383 (179 to 1,673)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Saxony-Anhalt	614 (54 to 2,109)	-64.3% (-89.2 to -32.6%)	863 (165 to 2,819)
Saxony	4,009 (1,168 to 12,366)	-59.8% (-83.4 to -33.6%)	6,476 (2,043 to 17,251)
Schleswig-Holstein	327 (11 to 2,345)	-38.3% (-85.7 to -5.9%)	550 (180 to 2,870)
Thuringia	999 (91 to 3,715)	-58.3% (-82.5 to -25.9%)	1,569 (339 to 5,288)
Greece	1,011 (0 to 3,191)	-12.9% (-35.8 to 0.0%)	7,312 (6,175 to 9,524)
Hungary	2,624 (822 to 4,433)	-24.0% (-37.5 to -9.7%)	10,760 (8,287 to 13,921)
Iceland	55 (2 to 174)	-54.1% (-87.2 to -11.2%)	81 (16 to 232)
Ireland	3,278 (881 to 9,273)	-31.9% (-53.1 to -14.1%)	9,425 (5,793 to 17,793)
Israel	1,181 (413 to 2,446)	-14.5% (-32.3 to -4.6%)	8,489 (5,100 to 10,696)
Italy	21,491 (7,538 to 51,432)	-16.7% (-28.2 to -8.4%)	121,821 (87,133 to 202,584)
Piemonte	3,147 (839 to 8,351)	-21.9% (-37.8 to -9.7%)	13,292 (8,542 to 24,854)
Valle d'Aosta	25 (0 to 179)	-7.5% (-31.5 to 0.0%)	231 (154 to 548)
Liguria	568 (72 to 2,261)	-10.4% (-25.9 to -2.8%)	4,412 (2,493 to 10,417)
Lombardia	4,858 (1,126 to 14,130)	-12.1% (-24.6 to -4.4%)	36,233 (25,042 to 64,454)
Provincia autonoma di Bolzano	632 (98 to 1,300)	-18.9% (-33.0 to -7.7%)	3,535 (815 to 7,097)
Provincia autonoma di Trento	520 (60 to 1,272)	-21.4% (-36.0 to -6.8%)	2,190 (873 to 4,190)
Veneto	1,261 (279 to 3,881)	-15.8% (-32.3 to -5.0%)	7,282 (5,378 to 12,730)
Friuli-Venezia Giulia	882 (151 to 2,509)	-31.9% (-52.8 to -12.2%)	2,509 (1,251 to 4,920)
Emilia-Romagna	2,266 (387 to 7,448)	-16.3% (-33.4 to -5.1%)	12,118 (7,682 to 24,550)
Toscana	2,034 (432 to 6,234)	-20.4% (-35.9 to -9.4%)	9,031 (4,229 to 22,009)
Umbria	73 (0 to 243)	-15.4% (-39.1 to 0.0%)	503 (156 to 1,145)
Marche	590 (73 to 2,368)	-14.8% (-32.6 to -3.7%)	3,293 (1,893 to 8,644)
Lazio	1,295 (445 to 2,646)	-16.2% (-26.7 to -7.5%)	7,705 (5,695 to 10,650)
Abruzzo	230 (49 to 887)	-12.1% (-27.3 to -3.7%)	1,684 (1,243 to 3,438)
Molise	61 (0 to 341)	-23.6% (-67.7 to -0.2%)	157 (27 to 656)
Campania	1,192 (268 to 3,067)	-14.6% (-25.8 to -5.9%)	7,527 (4,324 to 13,315)
Puglia	414 (0 to 1,110)	-12.3% (-34.2 to 0.0%)	3,295 (2,731 to 4,438)
Basilicata	66 (0 to 219)	-20.4% (-49.9 to 0.0%)	305 (60 to 898)
Calabria	69 (0 to 463)	-14.9% (-51.2 to -0.4%)	263 (108 to 1,200)
Sicilia	1,133 (340 to 2,938)	-20.0% (-35.3 to -8.5%)	5,300 (3,748 to 8,716)
Sardegna	175 (0 to 425)	-19.8% (-43.8 to 0.0%)	953 (499 to 1,248)
Kazakhstan	2,232 (759 to 6,582)	-29.5% (-56.6 to -15.3%)	6,905 (4,790 to 12,017)
Kyrgyzstan	2,509 (868 to 4,459)	-41.6% (-67.7 to -16.4%)	6,094 (3,078 to 8,935)
Latvia	429 (27 to 1,435)	-61.3% (-88.9 to -24.0%)	600 (113 to 1,908)
Lithuania	682 (63 to 1,389)	-47.7% (-76.6 to -13.8%)	1,427 (301 to 2,654)
Luxembourg	115 (0 to 266)	-25.9% (-54.4 to 0.0%)	418 (176 to 600)
Malta	257 (90 to 677)	-45.9% (-68.5 to -23.6%)	555 (230 to 1,216)
Montenegro	156 (10 to 318)	-23.1% (-43.9 to -1.6%)	666 (519 to 793)
Netherlands	11,984 (7,206 to 18,654)	-47.6% (-62.0 to -29.4%)	25,101 (19,286 to 33,479)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
North Macedonia	498 (0 to 1,257)	-19.5% (-40.1 to 0.0%)	2,452 (2,034 to 3,410)
Norway	176 (15 to 776)	-29.4% (-71.8 to -5.0%)	469 (299 to 1,077)
Poland	8,040 (89 to 17,561)	-20.9% (-47.3 to -0.3%)	37,870 (30,568 to 49,559)
Portugal	3,303 (1,420 to 6,491)	-24.4% (-39.1 to -12.8%)	13,136 (10,891 to 17,051)
Republic of Moldova	464 (0 to 1,424)	-10.7% (-34.9 to 0.0%)	4,345 (3,965 to 4,861)
Romania	944 (0 to 3,070)	-4.5% (-14.4 to 0.0%)	20,120 (18,198 to 22,376)
Russian Federation	38,820 (8,738 to 66,611)	-32.1% (-56.1 to -7.1%)	121,841 (114,758 to 128,944)
San Marino	2 (1 to 8)	-4.9% (-15.7 to -1.4%)	46 (44 to 53)
Serbia	3,702 (1,484 to 6,084)	-70.3% (-83.7 to -50.3%)	5,192 (2,663 to 7,883)
Slovakia	1,046 (0 to 2,502)	-38.7% (-67.6 to 0.0%)	2,745 (968 to 4,544)
Slovenia	466 (0 to 1,367)	-22.6% (-53.9 to 0.0%)	1,900 (1,275 to 3,751)
Spain	401 (111 to 681)	-0.5% (-0.9 to -0.2%)	73,697 (69,330 to 78,890)
Andalucia	68 (0 to 185)	-1.4% (-3.4 to 0.0%)	4,728 (2,814 to 7,435)
Aragon	5 (0 to 37)	-0.2% (-1.5 to 0.0%)	2,268 (1,929 to 2,825)
Cantabria	2 (0 to 13)	-0.4% (-2.0 to 0.0%)	612 (554 to 757)
Castilla-La Mancha	7 (0 to 34)	-0.2% (-0.8 to 0.0%)	4,110 (3,480 to 4,899)
Community of Madrid	7 (0 to 87)	-0.1% (-0.5 to 0.0%)	15,531 (14,414 to 16,430)
Extremadura	3 (0 to 11)	-0.3% (-0.8 to 0.0%)	1,281 (1,117 to 1,529)
Balearic Islands	13 (0 to 33)	-1.3% (-3.5 to 0.0%)	1,030 (664 to 1,490)
Canary Islands	28 (6 to 59)	-2.1% (-4.3 to -0.4%)	1,438 (867 to 2,311)
Asturias	56 (14 to 109)	-1.6% (-3.2 to -0.6%)	3,646 (1,781 to 5,919)
Murcia	31 (0 to 78)	-1.7% (-3.7 to 0.0%)	1,830 (1,414 to 2,560)
Castile and León	25 (0 to 50)	-0.5% (-0.9 to 0.0%)	4,513 (3,708 to 6,148)
Catalonia	86 (0 to 198)	-0.4% (-0.9 to 0.0%)	19,392 (17,893 to 22,364)
Ceuta	1 (0 to 5)	-1.2% (-3.6 to 0.0%)	135 (23 to 743)
Navarre	8 (0 to 21)	-0.6% (-1.6 to 0.0%)	1,221 (1,030 to 1,451)
Valencian Community	23 (0 to 113)	-0.5% (-2.4 to 0.0%)	4,800 (4,169 to 5,908)
Galicia	17 (0 to 46)	-0.6% (-1.6 to 0.0%)	2,808 (2,360 to 3,575)
Melilla	3 (0 to 10)	-1.5% (-4.1 to -0.3%)	296 (16 to 648)
Basque Country	16 (0 to 57)	-0.5% (-1.6 to 0.0%)	3,421 (2,514 to 4,011)
La Rioja	0 (0 to 4)	0.0% (-0.6 to 0.0%)	637 (572 to 684)
Sweden	7,124 (4,052 to 12,595)	-51.4% (-66.3 to -38.5%)	13,563 (10,576 to 18,974)
Switzerland	5,195 (694 to 18,400)	-39.7% (-69.2 to -8.0%)	11,734 (7,465 to 26,782)
Tajikistan	108 (19 to 386)	-37.8% (-70.3 to -14.4%)	244 (125 to 553)
Turkey	9,793 (0 to 17,435)	-18.2% (-31.7 to 0.0%)	53,601 (47,235 to 57,687)
Ukraine	12,641 (4,410 to 22,207)	-31.0% (-56.9 to -10.7%)	41,226 (36,013 to 46,789)
United Kingdom	54,917 (25,955 to 100,855)	-31.7% (-47.1 to -18.2%)	168,540 (136,449 to 217,547)
Northern Ireland	1,059 (342 to 2,336)	-29.7% (-46.6 to -12.8%)	3,427 (2,431 to 5,443)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Scotland	4,625 (1,714 to 8,669)	-30.4% (-45.9 to -16.5%)	14,658 (10,126 to 19,836)
Wales	1,485 (573 to 2,944)	-22.5% (-36.3 to -10.3%)	6,440 (4,996 to 8,171)
England	47,749 (21,849 to 89,092)	-32.2% (-48.1 to -18.1%)	144,016 (116,422 to 190,937)
Uzbekistan	780 (313 to 1,852)	-37.2% (-57.1 to -22.3%)	2,018 (1,233 to 3,598)
Region of the Americas	123,358 (70,931 to 174,474)	-11.8% (-16.4 to -6.9%)	1,045,057 (987,186 to 1,098,209)
Argentina	647 (275 to 1,646)	-1.4% (-3.5 to -0.6%)	45,614 (41,519 to 51,590)
Bahamas	79 (30 to 165)	-24.3% (-40.0 to -12.0%)	313 (221 to 434)
Barbados	1 (1 to 2)	-14.3% (-20.7 to -9.0%)	10 (10 to 10)
Belize	69 (12 to 165)	-33.9% (-56.8 to -13.7%)	187 (85 to 316)
Bolivia (Plurinational State of)	90 (48 to 145)	-1.0% (-1.5 to -0.5%)	9,330 (9,092 to 9,633)
Brazil	9,707 (6,097 to 14,524)	-5.2% (-7.5 to -3.4%)	186,751 (178,214 to 195,088)
Acre	13 (5 to 25)	-1.7% (-3.2 to -0.7%)	731 (711 to 760)
Alagoas	175 (81 to 291)	-6.7% (-10.6 to -3.4%)	2,603 (2,436 to 2,762)
Amazonas	618 (0 to 1,758)	-8.9% (-23.4 to 0.0%)	6,636 (5,887 to 7,669)
Amapá	42 (7 to 125)	-4.8% (-12.8 to -1.0%)	840 (768 to 988)
Bahia	421 (213 to 799)	-4.7% (-8.1 to -2.5%)	8,929 (8,394 to 9,774)
Ceará	201 (40 to 575)	-2.0% (-5.6 to -0.4%)	9,753 (9,418 to 10,516)
Distrito Federal	177 (100 to 314)	-3.9% (-6.5 to -2.3%)	4,470 (4,200 to 4,813)
Espírito Santo	286 (131 to 546)	-6.2% (-11.0 to -3.1%)	4,570 (4,184 to 5,073)
Goiás	416 (79 to 955)	-5.6% (-11.1 to -1.3%)	7,208 (6,026 to 8,706)
Maranhão	414 (202 to 675)	-8.6% (-13.3 to -4.6%)	4,791 (4,412 to 5,167)
Minas Gerais	827 (267 to 1,741)	-7.2% (-13.6 to -2.7%)	11,146 (9,752 to 12,863)
Mato Grosso do Sul	87 (24 to 197)	-4.5% (-9.1 to -1.4%)	1,874 (1,673 to 2,175)
Mato Grosso	203 (90 to 403)	-4.6% (-8.4 to -2.2%)	4,383 (4,081 to 4,812)
Pará	119 (20 to 433)	-1.7% (-5.8 to -0.3%)	7,018 (6,796 to 7,587)
Paraíba	452 (197 to 1,013)	-10.9% (-20.7 to -5.4%)	4,049 (3,571 to 4,868)
Paraná	456 (174 to 1,096)	-6.6% (-14.2 to -2.9%)	6,683 (5,851 to 7,966)
Pernambuco	406 (59 to 1,125)	-4.1% (-10.4 to -0.7%)	9,532 (8,757 to 10,807)
Piauí	467 (212 to 975)	-13.5% (-24.3 to -7.1%)	3,378 (2,886 to 4,217)
Rio de Janeiro	1,218 (417 to 2,673)	-5.0% (-10.1 to -1.9%)	23,867 (21,613 to 26,942)
Rio Grande do Norte	297 (51 to 862)	-9.1% (-22.4 to -1.9%)	3,096 (2,686 to 3,910)
Rondônia	105 (40 to 236)	-6.0% (-12.6 to -2.5%)	1,690 (1,561 to 1,922)
Roraima	82 (38 to 147)	-9.0% (-14.8 to -4.7%)	906 (797 to 1,017)
Rio Grande do Sul	633 (150 to 1,844)	-7.6% (-18.2 to -2.4%)	7,821 (6,365 to 10,286)
Santa Catarina	135 (37 to 431)	-3.6% (-10.3 to -1.1%)	3,570 (3,230 to 4,236)
Sergipe	195 (98 to 376)	-7.0% (-12.6 to -3.7%)	2,742 (2,529 to 3,063)
São Paulo	1,018 (285 to 2,212)	-2.3% (-4.8 to -0.7%)	42,779 (40,303 to 46,440)
Tocantins	240 (96 to 468)	-13.9% (-23.5 to -6.7%)	1,683 (1,352 to 2,063)
Canada	11,350 (4,457 to 23,687)	-28.0% (-42.0 to -15.0%)	39,829 (22,778 to 65,957)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Alberta	2,730 (1,089 to 7,455)	-59.6% (-79.0 to -36.2%)	4,551 (2,039 to 11,060)
British Columbia	160 (35 to 440)	-30.8% (-55.1 to -11.0%)	473 (315 to 818)
Manitoba	1,098 (17 to 7,163)	-52.8% (-87.3 to -12.7%)	2,945 (68 to 18,524)
New Brunswick	430 (1 to 2,341)	-32.8% (-69.5 to -4.8%)	1,951 (11 to 10,535)
Nova Scotia	2 (0 to 10)	-2.6% (-11.1 to -0.3%)	70 (66 to 83)
Ontario	3,593 (1,185 to 9,776)	-23.9% (-39.7 to -10.9%)	14,395 (6,536 to 27,097)
Quebec	3,262 (1,054 to 8,067)	-20.3% (-36.2 to -8.3%)	15,333 (10,506 to 22,457)
Saskatchewan	76 (0 to 660)	-32.6% (-91.4 to -0.9%)	111 (25 to 783)
Chile	107 (46 to 216)	-0.6% (-1.2 to -0.3%)	16,635 (15,636 to 17,923)
Colombia	2,757 (1,480 to 4,992)	-5.9% (-9.7 to -3.4%)	46,438 (41,385 to 52,554)
Costa Rica	377 (162 to 698)	-11.1% (-17.8 to -5.9%)	3,361 (2,463 to 4,291)
Cuba	240 (14 to 1,751)	-21.5% (-53.3 to -5.8%)	721 (230 to 3,515)
Dominican Republic	40 (5 to 117)	-1.6% (-4.5 to -0.2%)	2,396 (2,244 to 2,648)
Ecuador	51 (23 to 111)	-0.2% (-0.5 to -0.1%)	20,713 (20,426 to 21,129)
El Salvador	213 (0 to 827)	-7.5% (-22.9 to 0.0%)	3,289 (1,250 to 4,763)
Guatemala	315 (84 to 949)	-4.6% (-9.9 to -1.8%)	6,388 (4,575 to 10,444)
Guyana	14 (2 to 35)	-5.0% (-10.5 to -1.3%)	247 (161 to 391)
Haiti	31 (1 to 152)	-9.8% (-38.0 to -0.6%)	268 (232 to 406)
Honduras	308 (64 to 781)	-7.5% (-15.8 to -2.0%)	3,867 (2,988 to 5,238)
Jamaica	44 (7 to 119)	-8.8% (-17.2 to -2.7%)	455 (273 to 795)
Mexico	13,148 (6,521 to 21,395)	-9.0% (-14.2 to -4.3%)	145,982 (131,662 to 156,820)
Aguascalientes	71 (25 to 164)	-5.3% (-11.4 to -1.7%)	1,373 (990 to 1,758)
Baja California	364 (0 to 698)	-6.2% (-11.4 to 0.0%)	5,829 (4,980 to 6,534)
Baja California Sur	24 (0 to 89)	-3.0% (-10.2 to 0.0%)	689 (574 to 928)
Campeche	6 (0 to 19)	-0.7% (-2.0 to 0.0%)	894 (859 to 951)
Coahuila	149 (0 to 543)	-3.4% (-12.6 to 0.0%)	4,143 (3,550 to 4,756)
Colima	77 (40 to 126)	-5.8% (-9.3 to -2.8%)	1,331 (1,144 to 1,577)
Chiapas	1,078 (1 to 3,232)	-27.5% (-56.3 to -0.1%)	3,114 (1,076 to 7,188)
Chihuahua	362 (0 to 657)	-7.2% (-14.0 to 0.0%)	5,037 (3,953 to 6,363)
Mexico City	699 (325 to 1,240)	-3.6% (-6.2 to -1.7%)	19,428 (18,260 to 20,727)
Durango	203 (40 to 472)	-14.1% (-27.2 to -4.2%)	1,362 (899 to 1,942)
Guanajuato	987 (162 to 2,063)	-16.4% (-29.3 to -4.3%)	5,744 (3,748 to 7,597)
Guerrero	618 (197 to 1,165)	-16.4% (-27.9 to -7.2%)	3,661 (2,677 to 4,534)
Hidalgo	439 (210 to 729)	-11.3% (-17.8 to -5.8%)	3,872 (3,225 to 4,439)
Jalisco	817 (0 to 2,301)	-10.1% (-24.4 to 0.0%)	8,210 (5,528 to 10,438)
México	1,762 (0 to 4,433)	-10.2% (-21.3 to 0.0%)	16,867 (12,214 to 22,609)
Michoacán de Ocampo	604 (0 to 1,269)	-12.6% (-24.7 to 0.0%)	4,890 (3,551 to 5,681)
Morelos	365 (121 to 675)	-17.4% (-29.3 to -7.2%)	2,051 (1,492 to 2,533)
Nayarit	163 (0 to 384)	-10.3% (-23.0 to 0.0%)	1,587 (1,200 to 1,841)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Nuevo León	344 (0 to 1,240)	-4.8% (-17.2 to 0.0%)	7,096 (6,168 to 8,036)
Oaxaca	638 (170 to 1,199)	-17.3% (-30.5 to -5.7%)	3,664 (2,192 to 4,718)
Puebla	983 (253 to 1,940)	-13.0% (-22.9 to -4.6%)	7,329 (5,469 to 9,045)
Querétaro	248 (0 to 552)	-10.5% (-22.1 to 0.0%)	2,374 (1,708 to 2,819)
Quintana Roo	68 (0 to 170)	-2.6% (-6.1 to 0.0%)	2,625 (2,339 to 2,996)
San Luis Potosí	245 (0 to 597)	-6.7% (-15.4 to 0.0%)	3,610 (3,099 to 4,137)
Sinaloa	294 (117 to 536)	-6.1% (-10.5 to -2.8%)	4,722 (4,082 to 5,397)
Sonora	315 (0 to 671)	-7.0% (-14.0 to 0.0%)	4,522 (3,485 to 5,169)
Tabasco	66 (18 to 145)	-2.0% (-4.1 to -0.6%)	3,298 (3,097 to 3,592)
Tamaulipas	117 (12 to 340)	-3.5% (-9.0 to -0.4%)	3,135 (2,676 to 4,023)
Tlaxcala	157 (0 to 339)	-8.4% (-17.6 to 0.0%)	1,886 (1,492 to 2,163)
Veracruz de Ignacio de la Llave	584 (148 to 1,448)	-8.4% (-18.2 to -2.7%)	6,583 (5,409 to 8,342)
Yucatán	84 (0 to 233)	-2.6% (-7.2 to 0.0%)	3,211 (2,794 to 3,789)
Zacatecas	216 (0 to 429)	-11.6% (-21.8 to 0.0%)	1,848 (1,458 to 2,162)
Nicaragua	725 (116 to 2,418)	-53.2% (-79.2 to -28.2%)	1,205 (386 to 3,431)
Panama	119 (58 to 213)	-2.5% (-4.4 to -1.3%)	4,692 (4,046 to 5,403)
Paraguay	195 (93 to 383)	-8.3% (-13.8 to -4.6%)	2,313 (1,946 to 2,880)
Peru	75 (12 to 241)	-0.2% (-0.5 to 0.0%)	44,830 (43,857 to 46,523)
Suriname	51 (3 to 191)	-19.7% (-48.4 to -2.5%)	212 (120 to 463)
Trinidad and Tobago	35 (11 to 88)	-14.8% (-28.2 to -6.6%)	221 (161 to 325)
United States of America	82,150 (46,079 to 116,120)	-18.0% (-25.1 to -10.1%)	456,375 (424,449 to 489,680)
Alabama	1,641 (702 to 3,007)	-22.5% (-38.9 to -10.3%)	7,327 (4,609 to 9,522)
Alaska	93 (9 to 321)	-36.6% (-69.0 to -9.8%)	211 (95 to 500)
Arizona	1,572 (0 to 3,036)	-15.6% (-28.7 to 0.0%)	10,123 (7,786 to 13,035)
Arkansas	432 (0 to 1,267)	-10.6% (-31.7 to 0.0%)	4,075 (3,476 to 4,659)
California	10,847 (4,969 to 17,611)	-26.1% (-39.3 to -12.8%)	41,492 (28,290 to 51,411)
Colorado	1,693 (432 to 3,131)	-30.6% (-49.4 to -12.7%)	5,444 (2,936 to 7,494)
Connecticut	1,653 (280 to 4,913)	-16.5% (-36.0 to -3.8%)	9,144 (7,209 to 14,005)
Delaware	161 (17 to 318)	-13.2% (-25.9 to -1.2%)	1,243 (954 to 1,445)
District of Columbia	20 (0 to 98)	-2.7% (-12.1 to 0.0%)	687 (643 to 841)
Florida	5,402 (0 to 9,274)	-18.6% (-30.8 to 0.0%)	29,050 (22,259 to 31,933)
Georgia	3,221 (33 to 5,226)	-20.9% (-34.0 to -0.2%)	15,497 (13,905 to 17,171)
Hawaii	251 (40 to 591)	-29.2% (-49.3 to -10.8%)	787 (355 to 1,350)
Idaho	900 (67 to 2,399)	-34.7% (-57.4 to -3.4%)	2,472 (1,624 to 4,497)
Illinois	1,924 (0 to 4,846)	-9.1% (-22.1 to 0.0%)	20,807 (19,124 to 23,122)
Indiana	2,125 (719 to 3,871)	-19.5% (-33.1 to -7.4%)	10,772 (9,489 to 12,811)
Iowa	1,161 (163 to 2,245)	-24.8% (-44.6 to -3.9%)	4,653 (3,753 to 5,818)
Kansas	1,156 (296 to 2,380)	-27.7% (-47.2 to -8.4%)	4,110 (2,946 to 5,782)
Kentucky	721 (0 to 1,978)	-13.2% (-33.2 to 0.0%)	5,110 (4,158 to 7,037)
Louisiana	1,054 (0 to 2,111)	-11.1% (-22.9 to 0.0%)	9,502 (8,901 to 10,445)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Maine	95 (4 to 455)	-25.6% (-64.5 to -2.8%)	282 (156 to 722)
Maryland	1,492 (656 to 2,362)	-18.4% (-29.9 to -7.9%)	8,142 (6,899 to 9,614)
Massachusetts	830 (0 to 2,556)	-5.2% (-14.9 to 0.0%)	15,397 (14,247 to 17,332)
Michigan	1,370 (0 to 3,559)	-8.7% (-20.6 to 0.0%)	15,329 (13,874 to 17,904)
Minnesota	693 (0 to 1,794)	-10.1% (-27.1 to 0.0%)	6,771 (6,163 to 7,876)
Mississippi	714 (0 to 1,402)	-13.2% (-26.3 to 0.0%)	5,418 (4,972 to 5,807)
Missouri	1,725 (0 to 2,988)	-25.1% (-43.0 to 0.0%)	6,938 (5,043 to 9,405)
Montana	854 (158 to 2,189)	-36.6% (-59.2 to -12.6%)	2,215 (1,078 to 4,738)
Nebraska	785 (53 to 1,883)	-29.0% (-51.0 to -2.5%)	2,586 (1,927 to 4,174)
Nevada	805 (351 to 1,352)	-21.7% (-34.7 to -8.2%)	3,752 (2,655 to 4,826)
New Hampshire	693 (42 to 2,253)	-36.3% (-61.2 to -7.5%)	1,645 (547 to 4,040)
New Jersey	2,797 (0 to 6,941)	-11.5% (-22.4 to 0.0%)	23,654 (19,210 to 30,875)
New Mexico	935 (288 to 2,209)	-22.5% (-38.8 to -9.4%)	3,919 (2,913 to 5,939)
New York	2,001 (248 to 5,895)	-5.1% (-13.9 to -0.7%)	37,152 (34,014 to 42,900)
North Carolina	2,917 (0 to 7,239)	-19.5% (-41.8 to 0.0%)	14,354 (11,699 to 20,333)
North Dakota	168 (48 to 346)	-17.5% (-32.4 to -6.5%)	920 (687 to 1,236)
Ohio	3,354 (0 to 9,408)	-23.0% (-41.3 to 0.0%)	14,065 (7,702 to 23,896)
Oklahoma	822 (0 to 1,788)	-20.0% (-45.7 to 0.0%)	4,118 (3,528 to 5,099)
Oregon	1,111 (311 to 2,044)	-35.6% (-57.2 to -17.4%)	3,138 (1,280 to 4,764)
Pennsylvania	4,686 (1,664 to 9,809)	-21.3% (-37.4 to -8.4%)	21,574 (17,997 to 27,933)
Rhode Island	267 (0 to 774)	-10.7% (-25.9 to 0.0%)	2,276 (1,885 to 3,114)
South Carolina	840 (0 to 2,267)	-11.1% (-30.3 to 0.0%)	7,491 (6,669 to 8,838)
South Dakota	244 (0 to 489)	-22.9% (-41.0 to 0.0%)	1,038 (766 to 1,449)
Tennessee	1,741 (0 to 3,224)	-22.7% (-40.9 to 0.0%)	7,782 (4,942 to 10,050)
Texas	6,967 (453 to 11,797)	-19.5% (-33.5 to -1.2%)	36,092 (32,097 to 37,875)
Utah	897 (254 to 1,552)	-34.4% (-59.8 to -11.9%)	2,689 (1,086 to 3,713)
Vermont	4 (1 to 12)	-5.6% (-14.4 to -1.8%)	69 (65 to 79)
Virginia	1,831 (9 to 3,684)	-20.8% (-40.3 to -0.1%)	8,992 (5,102 to 11,915)
Washington	1,895 (933 to 3,085)	-29.0% (-43.8 to -15.5%)	6,501 (4,972 to 7,792)
West Virginia	564 (85 to 977)	-35.2% (-53.7 to -5.1%)	1,623 (951 to 2,516)
Wisconsin	1,849 (269 to 3,439)	-23.9% (-41.7 to -4.1%)	7,566 (6,106 to 9,363)
Wyoming	176 (14 to 611)	-41.7% (-69.8 to -13.9%)	377 (100 to 1,137)
Uruguay	6 (1 to 20)	-7.0% (-19.9 to -1.3%)	72 (58 to 104)
Venezuela (Bolivarian Republic of)	414 (183 to 846)	-17.3% (-28.9 to -9.0%)	2,342 (1,859 to 3,101)
South-East Asia Region	121,638 (72,475 to 176,995)	-30.5% (-41.6 to -19.6%)	397,289 (345,640 to 470,913)
Bangladesh	4,423 (2,249 to 8,156)	-36.6% (-53.0 to -24.0%)	11,781 (8,953 to 16,187)
India	104,301 (60,679 to 151,873)	-31.2% (-42.3 to -19.9%)	332,258 (285,297 to 396,931)
Andhra Pradesh	2,252 (1,192 to 4,009)	-20.6% (-32.1 to -12.6%)	10,764 (8,870 to 13,079)
Arunachal Pradesh	375 (70 to 732)	-52.2% (-84.7 to -20.0%)	769 (128 to 1,221)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Assam	1,267 (187 to 3,910)	-39.8% (-65.4 to -14.2%)	2,841 (1,283 to 6,191)
Bihar	9,069 (1,081 to 42,104)	-66.8% (-92.6 to -39.9%)	11,687 (2,663 to 45,700)
Chhattisgarh	5,160 (2,652 to 9,277)	-48.8% (-66.5 to -30.3%)	10,467 (6,633 to 15,253)
Delhi	2,888 (1,541 to 4,581)	-20.3% (-30.9 to -11.2%)	14,187 (12,621 to 16,034)
Goa	174 (80 to 310)	-15.2% (-24.3 to -8.3%)	1,130 (864 to 1,381)
Gujarat	2,757 (1,036 to 6,313)	-28.2% (-47.2 to -14.2%)	9,329 (6,559 to 13,680)
Haryana	1,516 (139 to 4,338)	-27.8% (-52.0 to -6.4%)	4,691 (2,138 to 8,873)
Himachal Pradesh	1,114 (320 to 2,443)	-46.3% (-67.1 to -27.7%)	2,326 (973 to 4,217)
Jammu & Kashmir and Ladakh	1,103 (457 to 2,290)	-31.3% (-47.9 to -18.7%)	3,400 (2,345 to 5,008)
Jharkhand	1,338 (401 to 3,242)	-43.4% (-65.8 to -25.2%)	2,897 (1,555 to 5,266)
Karnataka	3,937 (1,279 to 7,934)	-18.6% (-30.9 to -8.6%)	20,426 (14,613 to 27,609)
Kerala	4,013 (1,446 to 7,893)	-28.4% (-45.1 to -15.3%)	13,783 (7,303 to 20,098)
Madhya Pradesh	4,728 (1,433 to 12,022)	-42.0% (-65.9 to -23.2%)	10,447 (5,617 to 20,250)
Maharashtra	9,055 (4,453 to 16,138)	-12.2% (-19.5 to -7.0%)	73,657 (60,033 to 87,612)
Manipur	612 (165 to 1,056)	-22.6% (-38.0 to -7.4%)	2,666 (2,045 to 3,286)
Meghalaya	384 (20 to 879)	-22.3% (-59.2 to -1.2%)	1,785 (1,270 to 2,299)
Nagaland	561 (206 to 926)	-60.3% (-86.1 to -25.5%)	967 (343 to 1,558)
Odisha	8,536 (2,962 to 16,109)	-55.8% (-76.7 to -35.0%)	14,773 (7,170 to 23,726)
Punjab	1,197 (343 to 3,245)	-17.5% (-36.1 to -6.9%)	6,362 (4,996 to 9,327)
Rajasthan	8,632 (3,132 to 19,753)	-52.5% (-74.0 to -31.6%)	15,556 (8,841 to 28,199)
Sikkim	101 (44 to 170)	-17.2% (-41.5 to -8.6%)	601 (376 to 778)
Tamil Nadu	4,778 (1,321 to 10,649)	-23.5% (-41.4 to -9.5%)	19,249 (13,975 to 26,437)
Telangana	1,076 (379 to 2,393)	-31.0% (-50.4 to -16.5%)	3,298 (2,177 to 5,038)
Tripura	410 (186 to 816)	-40.9% (-59.7 to -25.8%)	972 (684 to 1,420)
Uttar Pradesh	4,598 (2,482 to 9,857)	-31.2% (-48.4 to -21.1%)	14,325 (10,929 to 21,680)
Uttarakhand	1,662 (276 to 3,515)	-36.3% (-58.4 to -15.7%)	4,316 (1,654 to 6,787)
West Bengal	21,010 (11,694 to 31,455)	-38.6% (-58.0 to -20.1%)	54,583 (44,352 to 60,015)
Indonesia	2,518 (1,335 to 4,881)	-11.1% (-18.7 to -6.4%)	22,434 (19,590 to 26,666)
Maldives	20 (1 to 92)	-15.7% (-41.7 to -2.2%)	91 (45 to 271)
Myanmar	6,451 (2,609 to 13,658)	-44.6% (-64.6 to -26.3%)	14,178 (7,621 to 23,573)
Nepal	3,623 (0 to 9,217)	-23.7% (-67.8 to 0.0%)	16,041 (11,297 to 19,520)
Sri Lanka	268 (0 to 3,487)	-29.9% (-81.0 to -1.2%)	405 (16 to 4,377)
Thailand	34 (6 to 168)	-27.0% (-68.3 to -8.5%)	101 (68 to 247)
Western Pacific Region	14,031 (3,207 to 35,638)	-21.8% (-43.3 to -9.0%)	58,890 (33,758 to 100,123)

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Location	Lives saved (95% UI)	Percent difference in deaths (95% UI)	Cumulative deaths in reference scenario (95% UI)
Australia	72 (32 to 148)	-5.4% (-10.3 to -2.6%)	1,317 (1,231 to 1,457)
China	1,467 (41 to 18,772)	-10.7% (-72.2 to -0.8%)	6,680 (4,819 to 25,720)
Hong Kong Special Administrative Region of China	166 (1 to 1,355)	-16.1% (-52.7 to -0.5%)	542 (109 to 3,707)
Beijing	32 (1 to 211)	-42.0% (-86.7 to -6.2%)	49 (11 to 243)
Chongqing	4 (1 to 16)	-27.2% (-65.2 to -10.1%)	12 (9 to 26)
Guangdong	29 (3 to 127)	-53.8% (-87.6 to -20.8%)	45 (15 to 151)
Hainan	7 (1 to 54)	-28.5% (-80.7 to -11.9%)	16 (9 to 70)
Hebei	8 (2 to 27)	-43.0% (-72.9 to -21.8%)	17 (11 to 37)
Heilongjiang	1 (0 to 2)	-5.7% (-11.5 to -2.2%)	15 (15 to 17)
Hubei	1,175 (3 to 18,652)	-6.5% (-72.7 to -0.1%)	5,915 (4,565 to 25,385)
Shandong	35 (4 to 157)	-58.8% (-89.4 to -27.4%)	49 (15 to 178)
Shanghai	10 (2 to 35)	-43.2% (-76.3 to -19.6%)	20 (12 to 47)
Japan	6,373 (1,226 to 19,648)	-22.9% (-39.7 to -10.6%)	24,607 (10,311 to 61,165)
Malaysia	3,232 (318 to 9,074)	-47.3% (-71.2 to -25.3%)	6,227 (1,108 to 16,112)
New Zealand	1 (0 to 6)	-2.5% (-19.4 to -0.1%)	27 (26 to 33)
Papua New Guinea	0 (0 to 1)	-1.8% (-9.5 to -0.1%)	9 (8 to 11)
Philippines	2,233 (285 to 6,228)	-12.0% (-25.1 to -3.1%)	16,708 (9,036 to 29,294)
Republic of Korea	415 (73 to 1,664)	-12.1% (-23.7 to -5.1%)	2,969 (1,240 to 8,514)
Singapore	6 (0 to 40)	-7.6% (-27.4 to -0.5%)	52 (29 to 156)
Viet Nam	231 (5 to 1,757)	-52.7% (-94.3 to -10.3%)	294 (47 to 1,903)

References

1. Center for Systems Science and Engineering (CSSE). Coronavirus COVID-19 Global Cases [Internet]. Johns Hopkins Coronavirus Resource Center. [cited 2020 Oct 28]; Available from: <https://coronavirus.jhu.edu/map.html>
2. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *The Lancet* 2020;395(10228):931–934.
3. Flaxman S, Mishra S, Gandy A, et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* 2020;584(7820):257–261.
4. Lyu W, Wehby GL. Community Use Of Face Masks And COVID-19: Evidence From A Natural Experiment Of State Mandates In The US: Study examines impact on COVID-19 growth rates associated with state government mandates requiring face mask use in public. *Health Affairs* 2020;39(8):1419–1425.
5. Lai S, Ruktanonchai NW, Zhou L, et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. 2020;
6. Pfizer Inc. Pfizer and BioNTech Announce Vaccine Candidate Against COVID-19 Achieved Success in First Interim Analysis from Phase 3 Study | Pfizer [Internet]. [cited 2020 Nov 13]; Available from: <https://www.pfizer.com/news/press-release/press-release-detail/pfizer-and-biontech-announce-vaccine-candidate-against>
7. Bloom BR, Nowak GJ, Orenstein W. “When Will We Have a Vaccine?”—Understanding Questions and Answers about Covid-19 Vaccination. *New England Journal of Medicine* 2020;
8. Robbins R, Gelles D. How Pfizer Plans to Distribute Its Vaccine (It’s Complicated) [Internet]. *The New York Times*. 2020 [cited 2020 Nov 13]; Available from: <https://www.nytimes.com/2020/11/12/business/pfizer-covid-vaccine-coronavirus.html>
9. Liang M, Gao L, Cheng C, et al. Efficacy of face mask in preventing respiratory virus transmission: A systematic review and meta-analysis. *Travel Med Infect Dis* [Internet] 2020 [cited 2020 Jul 16]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7253999/>
10. Chu DK, Akl EA, Duda S, et al. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet* 2020;395(10242):1973–87.
11. Community Initiatives. What Countries Require Masks in Public or Recommend Masks? [Internet]. #Masks4All. [cited 2020 Jul 16]; Available from: <https://masks4all.co/what-countries-require-masks-in-public/>
12. The Editors. Dying in a Leadership Vacuum. *New England Journal of Medicine* 2020;383(15):1479–80.

13. Aratani L. How did face masks become a political issue in America? [Internet]. the Guardian. 2020 [cited 2020 Jul 16]; Available from: <http://www.theguardian.com/world/2020/jun/29/face-masks-us-politics-coronavirus>
14. Tcharkhtchi A, Abbasnezhad N, Seydani MZ, Zirak N, Farzaneh S, Shirinbayan M. An overview of filtration efficiency through the masks: Mechanisms of the aerosols penetration. *Bioactive materials* 2020;6(1):106–122.
15. Wang C, Otani Y. Removal of Nanoparticles from Gas Streams by Fibrous Filters: A Review [Internet]. 2012 [cited 2020 Jul 17]; Available from: <https://pubs.acs.org/doi/pdf/10.1021/ie300574m>
16. World Health Organization. Transmission of SARS-CoV-2: implications for infection prevention precautions [Internet]. Geneva, Switzerland: 2020. Available from: <https://apps.who.int/iris/rest/bitstreams/1286634/retrieve>
17. Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *The Lancet Respiratory Medicine* [Internet] 2020 [cited 2020 Aug 2];0(0). Available from: [https://www.thelancet.com/journals/lanres/article/PIIS2213-2600\(20\)30323-4/abstract](https://www.thelancet.com/journals/lanres/article/PIIS2213-2600(20)30323-4/abstract)
18. Rajagopalan S, Huang S, Brook RD. Flattening the curve in COVID-19 using personalised protective equipment: lessons from air pollution. *Heart* 2020;heartjnl-2020-317104.
19. MacIntyre CR, Chughtai AA. A rapid systematic review of the efficacy of face masks and respirators against coronaviruses and other respiratory transmissible viruses for the community, healthcare workers and sick patients. *International Journal of Nursing Studies* 2020;108:103629.
20. Fan J, Li Y, Stewart K, et al. COVID-19 World Survey Data API [Internet]. 2020 [cited 2020 Jul 19]; Available from: <https://covidmap.umd.edu/api.html>
21. COVID-19 Symptom Survey - Request for Data Access [Internet]. Facebook Data for Good. [cited 2020 Nov 9]; Available from: <https://dataforgood.fb.com/docs/covid-19-symptom-survey-request-for-data-access/>
22. Huprikar S, Ludolph T. Are Americans Wearing Face Masks? [Internet]. Premise. 2020 [cited 2020 Jul 16]; Available from: <https://www.premise.com/are-americans-wearing-face-masks/>
23. Jones SP, Imperial College London Big Data Analytical Unit, YouGov Plc. YouGov Covid Data Hub, v1.0 [Internet]. YouGov Plc.; 2020 [cited 2020 Jul 19]. Available from: <https://github.com/YouGov-Data/covid-19-tracker>
24. Reiner RC, Barber RM, Collins JK, et al. Modeling COVID-19 scenarios for the United States. *Nature Medicine* 2020;1–12.
25. Friedman J, Liu P, Gakidou E, Team IC-19 MC. Predictive performance of international COVID-19 mortality forecasting models. v.4. medRxiv [Internet] 2020 [cited 2020 Jul 23]; Available from: <https://www.medrxiv.org/content/10.1101/2020.07.13.20151233v4>

26. Friedman J, Liu P, Gakidou E, IHME COVID-19 Model Comparison Team. Predictive performance of international COVID-19 mortality forecasting models [Internet]. *Epidemiology*; 2020 [cited 2020 Jul 16]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.07.13.20151233>
27. Wang Y, Tian H, Zhang L, et al. Reduction of secondary transmission of SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing, China. *BMJ Glob Health* 2020;5(5):e002794.
28. Bielecki M, Züst R, Siegrist D, et al. Social distancing alters the clinical course of COVID-19 in young adults: A comparative cohort study. *Clinical Infectious Diseases* 2020;ciaa889.
29. Hendrix MJ. Absence of Apparent Transmission of SARS-CoV-2 from Two Stylists After Exposure at a Hair Salon with a Universal Face Covering Policy — Springfield, Missouri, May 2020. *MMWR Morb Mortal Wkly Rep* [Internet] 2020 [cited 2020 Jul 17];69. Available from: <https://www.cdc.gov/mmwr/volumes/69/wr/mm6928e2.htm>
30. Wang X, Ferro EG, Zhou G, Hashimoto D, Bhatt DL. Association Between Universal Masking in a Health Care System and SARS-CoV-2 Positivity Among Health Care Workers. *JAMA* [Internet] 2020 [cited 2020 Aug 2]; Available from: <https://jamanetwork.com/journals/jama/fullarticle/2768533>
31. Leung NHL, Chu DKW, Shiu EYC, et al. Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine* 2020;26(5):676–80.
32. CDC. Coronavirus Disease 2019 (COVID-19): Use of masks to help slow the spread of COVID-19 [Internet]. Centers for Disease Control and Prevention. 2020 [cited 2020 Aug 2]; Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/diy-cloth-face-coverings.html>
33. Iversen BG, Vestrheim D, Flottorp S, Denison E, Oxman A, Norwegian Institute of Public Health. Should individuals in the community without respiratory symptoms wear facemasks to reduce the spread of COVID-19? *Norwegian Institute of Public Health* 2020;45.
34. Zaken M van A. Avoid busy places and stay 1.5 metres away from others [Internet]. 2020 [cited 2020 Jul 16]; Available from: <https://www.government.nl/latest/news/2020/05/06/avoid-busy-places-and-stay-1.5-metres-away-from-others>
35. The Public Health Agency of Sweden. FAQ about COVID-19 [Internet]. [cited 2020 Jul 16]; Available from: <http://www.folkhalsomyndigheten.se/the-public-health-agency-of-sweden/communicable-disease-control/covid-19/>
36. Wong T. Coronavirus: Why some countries wear face masks and others don't - BBC News [Internet]. [cited 2020 Jul 16]; Available from: <https://www.bbc.com/news/world-52015486>
37. World Health Organization. Advice on the use of masks in the context of COVID-19: Interim guidance [Internet]. Geneva, Switzerland: 2020. Available from: https://apps.who.int/iris/bitstream/handle/10665/331693/WHO-2019-nCov-IPC_Masks-2020.3-eng.pdf?sequence=1&isAllowed=y
38. World Health Organization. Advice on the use of masks in the context of COVID-19: Interim guidance [Internet]. Geneva, Switzerland: 2020. Available from:

https://apps.who.int/iris/bitstream/handle/10665/332293/WHO-2019-nCov-IPC_Masks-2020.4-eng.pdf?sequence=1&isAllowed=y

39. Mills M, Rahal C, Akimova E, et al. Face masks and coverings for the general public: behavioural knowledge, effectiveness of cloth coverings and public messaging [Internet]. The Royal Society & The British Academy; 2020. Available from: <https://royalsociety.org/-/media/policy/projects/set-c/set-c-facemasks.pdf>