1 **Functional connectivity directionality between large-scale resting-state networks in**
2 **children and adolescence from the Healthy Brain Network sample**
3

4 Martina J. Lund[1*], Dag Alnæs[1,2], Jaroslav Rokicki[1,3], Simon Schwab[4,5], Ole A. Andreassen[1,6],
5 Lars T. Westlye[1,3,6], Tobias Kaufmann[1,7*].
6

[1] Norwegian Centre for Mental Disorders Research (NORMENT), Division of Mental Health
and Addiction, Oslo University Hospital, and Institute of Clinical Medicine, University of
Oslo, Norway
[2] Bjørknes College, Oslo, Norway
[3] Department of Psychology, University of Oslo, Oslo, Norway
[4] Center for Reproducible Science (CRS) & Epidemiology, Biostatistics and Prevention
Institute (EBPI), University of Zürich, Zurich, Switzerland
[5] Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, Nuffield
Department of Population Health, University of Oxford, Oxford, UK
[6] KG Jebsen Centre for neurodevelopmental disorders, University of Oslo, Oslo, Norway
[7] Department of Psychiatry and Psychotherapy, University of Tübingen, Tübingen, Germany
18

19 [*] Corresponding authors:
20   Martina J. Lund and Tobias Kaufmann, PhD
21   Email: m.j.lund@medisin.uio.no, tobias.kaufmann@medisin.uio.no
22   Postal address: OUS, PoBox 4956 Nydalen, 0424 Oslo, Norway
23   Telephone: +47 23 02 73 50, Fax: +47 23 02 73 33

24
25
26 Counts:
27   Abstract: 203 words
28   Main text body: 4817 words
29   Figures: 5
30
Keywords:
Directed functional connectivity
Brain Development
Dynamic graphical models
Resting-state fMRI
The Healthy Brain Network

Abstract

Objective: Mental disorders often emerge during adolescence and have been associated with age-related differences in connection strengths of brain networks (static connectivity), manifesting in non-typical developmental trajectories of brain maturation. However, little is known about the directionality of information flow (directed connectivity) in this period of brain development.

Methods: We employed dynamic graphical models (DGM) to estimate directed functional connectivity from resting state functional magnetic resonance imaging data on 1143 participants, aged 6 to 17 years from the healthy brain network (HBN) sample. We tested for effects of age, sex, cognitive abilities and psychopathology on directionality.

Results: We show robust bi-directionality in information flow between visual-medial and visual-lateral networks, in line with prior studies investigating this in adults. Furthermore, we found that age in this developmental sample was associated with directionality of information flow for sensorimotor, auditory, and control networks, and that cognition was linked to directionality between the visual occipital and the default mode network. However, we did not find associations with psychopathology.

Discussion: Our results revealed that directionality in information flow of large-scale brain networks is sensitive to age and cognition during adolescence, warranting further studies that may explore trajectories of development in more fine-grained network parcellations and in different populations.

Introduction

The brain undergoes tremendous changes throughout life, where childhood and adolescence is a particularly sensitive developmental period for brain maturation processes (Blakemore, 2012). A vital part of the brain's maturation happens in the functional networks that show pronounced reorganization in order to facilitate neural efficiency and integration of information, as indicated by connectome studies in healthy children and young individuals as well as in clinical populations in the same age range (Insel, 2010; Keshavan, Giedd, Lau, Lewis, & Paus, 2014; Kolskar et al., 2018; Rausch et al., 2016; Schweinsburg, Nagel, & Tapert, 2005). This supports that this time period is a sensitive phase, where there is large alterations in the functional interconnections between brain regions as a whole that link to when different mental disorders emerge (Insel, 2010).

The primary method to study functional brain interconnections so far has been to estimate static functional connectivity between brain regions, where research has shown that the functional connectome in healthy children and adolescents becomes more specialized (Li, Deng, He, Zhai, & Jia, 2019). Resting-state networks (RSNs) also become more coherent and stable during this time-period (Hoff, Van den Heuvel, Benders, Kersbergen, & De Vries, 2013), and evidence has indicated aberrant connectivity in individuals with pre-clinical psychiatric symptoms (Kaufmann et al., 2017), which is in line with the brain dysconnectivity hypothesis in mental disorders (Connolly et al., 2013; Di Martino et al., 2011; K. J. Friston & Frith, 1995; Hamm et al., 2014). Only recently has directed functional connectivity received interest as a field that could yield new knowledge about the connections between brain regions by estimating directionality in neural information flow. Riley et al. (2018) examined effective connectivity, the causal influence that one node exerts over another (Bielczyk et al., 2019; Karl J. Friston, 2011), in a neurodevelopmental sample and observed that there are differences between sexes in relation to how memories are encoded in the hippocampus. In

another neurodevelopmental study by Hwang, Velanova, and Luna (2010), the authors observed that improvements in inhibitory control were linked to strengthening of top-down connectivity for regions implicated in cognitive control networks, while similar findings were found in relation to top-down processing for a language network (Bitan et al., 2006). Further, it has been shown that there are differences in directionality in adolescent boys with externalizing behavior disorders compared to a control group (Shannon, Sauder, Beauchaine, & Gatzke-Kopp, 2009) and for children with Attention Deficit Hyperactivity Disorder (ADHD) (Zhao, Zheng, Yang, & Tian, 2017). However, the literature on connection directionality is scarce, calling for validation of earlier findings and for new insights using novel approaches. In particular, we do not currently know if and how the direction flow in the communication between brain networks is implicated in mental traits and disorders, including cognitive abilities. Given that cognitive deficits are seen across mental disorders, connecting these behavioral and clinical characteristics is crucial for gaining a better understanding of how higher-order processing is characterized in children and adolescence. Such integrative understanding is also important given that no consensus has been reached when it comes to the disorder-specific connectivity alterations that characterize particular mental disorders and the known overlap of symptoms (Craddock & Owen, 2010; Jollans & Whelan, 2018) and the pleiotropy between disorders (Demontis et al., 2019; Grove et al., 2019; Hibar et al., 2015; Schizophrenia Working Group of the Psychiatric Genomics, 2014; Talkowski et al., 2012).

Dynamic graphical models (DGM) is a Bayesian approach for examining directed functional connectivity using dynamic linear models (DLM) (Schwab et al., 2018). As such, it implements a state space model that is linear and Gaussian in form. This includes statistically stationary properties as it uses a hidden Markov modelling approach, although it incorporates time varying coefficients and as a result can provide information about directionality in the form of a binary view of coupling between two brain regions for each connectivity direction.

Here, we used DGM and publicly available resting-state functional magnetic

resonance imaging (rsfMRI) data from the Healthy Brain Network (HBN) project (Alexander

et al., 2017) to study brain network information flow in children and adolescents and its

associations with age, sex, cognitive abilities and psychopathology. We hypothesized that

information flow would be strengthened with increasing age (Bitan et al., 2006; Hwang et al.,

2010), that there would be differences between females and males in the maturation of brain

networks (Riley et al., 2018), as well as alterations associated with information flow for

control networks as these nodes are central to a range of disorders (Cortese, Kelly, & Di

Martino, 2012; Eisenberg & Berman, 2010; Francx et al., 2015; Geiger et al., 2016; P. Li et

al., 2017; Q. Zhao, Swati, Metmer, Sang, & Lu, 2019).

Methods

Study samples

The HBN is a project organized by the Child Mind Institute (Alexander et al., 2017) and is a

resource targeting novel insight into the critical time period when psychiatric and mental

disorders emerge. The HBN consortium aims to include 10,000 individuals in the age range

of 5-21 years from the New York area, where participants are included by use of

announcements that are distributed to community members, educators, local care providers

with the addition of sending information via email lists and events to parents, encouraging

participation of children with clinical concerns to this study (Alexander et al., 2017). Data on

these   individuals   include   a   package   consisting   of   MRI   scanning,   genetics,

electroencephalography   (EEG),   eye-tracking,   as   well   as   biological   testing   and   a

neuropsychological battery consisting of cognitive, lifestyle indices, behavioral and

psychiatric domains in addition to actigraphy and voice and video interviews (Alexander et

al., 2017). Exclusion criteria include serious neurological disorders, neurodegenerative

disorders, acute encephalopathy, hearing or visual impairment, lifetime substance abuse that

necessitated chemical replacement therapy/acute intoxication at time of study, recent

diagnosis of a severe mental disorder or manic/psychotic episode within the last 6 months

without ongoing treatment, in addition to the onset of suicidality/homicidality where there is

no current, ongoing treatment (Alexander et al., 2017). All participants over the age of 18

years provided signed informed consent, while legal guardians signed informed consent for

participants under the age of 18, in addition to participants giving a written assent (Alexander

et al., 2017). The Chesapeake Institutional Review Board approved the study (https://www.

chesapeakeirb.com/).

MRI acquisition and preprocessing

MR data was collected by the study team of HBN, where we included MRI data from the

following sites; Rutgers University Brain Imaging Center (RUBIC), Citigroup Biomedical

Imaging Center (CBIC) and a mobile scanner located in Staten Island. MRI data was

collected using one scanner at each site, giving a total of 3 scanners comprising our sample.

Rutgers applied a Siemens 3T Tim Trio scanner, while CBIC utilized a Siemens 3T Prisma,

and both sites applied the same MRI parameters, where resting- state blood-oxygen-level-

dependent (BOLD) fMRI data was collected for each subject using a T2*-weighted BOLD

echo-planar imaging (EPI) sequence with a repetition time (TR) of 800ms, echo time (TE) of

30ms, multiband acceleration factor = 6, 60 number of slices with the rsfMRI session

consisting of 375 volumes and with voxel size= 2.4×2.4×2.4 mm. The third mobile scanner

located in Staten Island used a 1.5T Siemens Avanto system equipped with 45 mT/m

gradients (Alexander et al., 2017), where the following parameters were implemented; TR=

1.45s, TE=40ms, multiband acceleration factor = 3, number of volumes=420, slices= 54,

resolution in mm= 2.5×2.5×2.5 mm (for more information about the MRI parameters, see

http://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/MRI%20Protocol.html).

Raw imaging data was downloaded from the HBN database

(http://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/sharing_neuro.html#Dir

ect%20Down), and analyzed on the secure data storage and computing facilities (TSD,

https://www.uio.no/tjenester/it/forskning/sensitiv) at University of Oslo. Image processing

tools, based largely on Smith et al. (2013), were used for the functional data while T1-

weighted data, which was applied as an intermediate in the registration, was processed using

FreeSurfer 5.3 (http://freesurfer.net), including removal of non-brain tissue.

As part of the HBN MRI protocol, multiple T1weighted sequences were acquired for

each subject, and we used MRIQC version 0.14.2 (Esteban et al., 2017) for automated quality

assessment (N=2427, CUNY scanning site was included for the MRIQC stage, yet this site

was dropped at a later stage due the low sample size, N=22). From this, T1 weighted images

with the best image quality metrics, based on the classifier ratings, were used as input for

registration, while structural scans that were flagged as low quality (classifier rating score

>0.5, N=160) were manually checked and excluded (N=117). In addition, 22 subjects had

errors when they were run through FreeSurfer and after manually inspecting these datasets,

these were also omitted due to motion artefacts and/or having MRI findings disrupting the

segmentation pipeline in FreeSurfer. We also made a global mask of the T1 data where we

manually checked subjects that had low coverage, excluding another 95 subjects leaving a

total of 2171 datasets that had a usable T1w sequence for registration.

Two of the MRI sites, RUBIC and CBIC, had two resting-state scans acquired as part

of the same MRI session. As the DGM method benefits from more time points, we merged

the time series for subjects with two resting state scans together to leverage all available

resting-state data. This was done prior to implementing feat as a means of optimizing the data

by improving the spatial alignment between the sessions, and to have the full set of volumes

to inform FIX.

Furthermore, the first five volumes for the fMRI dataset were discarded. We

preprocessed fMRI data using FSL 6.0.3 (http://fsl.fmrib.ox.ac.uk), including motion

correction and brain extraction. FSL's FEAT (Woolrich, Ripley, Brady, & Smith, 2001)

included spatial smoothing with a Gaussian kernel FWHM of 6 mm and a high-pass filter

cutoff of 100. FMRIB's Nonlinear Image Registration tool (FNIRT) was used to register

fMRI volumes to standard space (MNI-152) with the T1w volumes as intermediates.

We also implemented a cleaning step for fMRI data, where we made a global mask for

the fMRI datasets (which indicated data coverage across participants), and from this we

excluded 128 of 1813 datasets with poor coverage. To reduce the influence of noise in the

data and increase the tSNR (Kaufmann et al., 2017), we removed artefacts by use of ICA-

AROMA, a classifier that identifies and removes motion specific noise in fMRI data (Pruim,

Mennes, Buitelaar, & Beckmann, 2015; Pruim, Mennes, van Rooij, et al., 2015). Afterwards,

ICA was rerun and FSL's FIX (FMRIB's ICA-based X-noisiefier (Griffanti et al., 2014;

Salimi-Khorshidi et al., 2014)), was used with the recommended threshold of 20 to remove

remaining motion confounds and other artefacts in the data. Further, data was temporally

demeaned and variance normalized (Beckmann & Smith, 2004), and the quality controlled

fMRI dataset (N=1685) was submitted to a group ICA, utilizing FSL's Multivariate

Exploratory Linear Optimized Decomposition into Independent Components (MELODIC)

tool (Beckmann & Smith, 2004; Hyvärinen, 1999), where 25 components were extracted from

the ICA and used for further analysis. Dual regression was applied to estimate individual

spatial maps and corresponding time series from the group ICA (Beckmann & Smith, 2004;

Filippini et al., 2009), which were used as input for the DGM analysis.

 Mental and cognitive measures

From the HBN data release, N=1685 participants were included after quality assessment. Out

of these, information on age at MR was missing for N=45, sex for N=9, and cognitive/clinical

information for 496 individuals. The participants were 5-22 years (mean: 11.5, years, sd: 3.51

years) and 37.9% were females. This data was used to study group-level patterns of dFC

(average connectivity matrix). For the subsequent associations with age, sex, cognition and

mental health, we restricted the analysis to a subset based on data availability. Thus, the final

sample for the association analyses comprised N=1143, individuals aged 6-17 years (mean:

10.7 years, sd: 2.62 years, 37.4% females, where N=83 where from the Staten Island site,

N=503 from CBIC and N=557 from RUBIC scanning site (see SI SFig.2 for age distributions

within scanner sites). A large proportion of the sample had a diagnosis (N=1012), while

N=105 did not have a diagnosis, N=23 dropped out before a diagnosis could be determined,

and N=3 was missing information for clinical consensus diagnosis data and had as such not

received a diagnosis. Broadly, the majority of patients included in our analysis had disorders

in the following categories where comorbidities are included. Accordingly, there is a higher N

than the total number of subjects for each diagnosis category given here: Neurodevelopmental

disorders (N=1526), anxiety disorders (N=557), disruptive, impulsive control and conduct

disorders (N=171), elimination disorders (N=111), and depressive disorders (N=115), see SI;

SFig. 1 for further details.

We used the full-scale intelligence quotient (FSIQ) from the Wechsler Intelligence

Scale for Children (WISC-V) taken for participants aged 6-17 years as a proxy for cognitive

ability. This composite score includes the following domains; visual spatial, verbal

comprehension, fluid reasoning, working memory, and processing speed (Wechsler, 2003).

Mental health was measured on a continuum including both healthy subjects and patients as it

is difficult to uncover robust findings for psychiatric diagnosis that constitutes heterogeneous

disorders, showing a wide range in symptoms, severity, duration and prognosis, and as

patients often have more than one diagnosis. Such heterogeneity is also reflected in the brain,

making the search for biological markers a complex task. As such, for mental health, we

performed a principal component analysis (PCA) on The Extended Strengths and Weaknesses

Assessment of Normal Behavior (E-SWAN), which has been shown to be a valid

psychometric assessment tool for investigating behavior underlying DSM disorders

(Alexander, Salum, Swanson, & Milham, 2020). E-SWAN domains include depression,

social anxiety, disruptive mood dysregulation disorder (DMDD), and panic disorder. We

excluded 3 of the items relating to panic disorder from the questionnaire that had a high

degree (90%) of missing values, giving a total of 62 items for the PCA analysis. The

remaining items had available data for 2626 participants with no missing values. We

performed PCA using the "prcomp" function in R, where the first PC, often denoted as the p-

factor or pF (Caspi et al., 2013), explained 43.6% of the variance (Figure 1). From the

loadings from the PCA, this component was associated with items related to self-control and

depression/anxiety (Figure 1). In accordance with other studies showing more than one factor

being of importance (Alnaes et al., 2018; Mallard et al., 2019), we also included the second

principal component referred to as pF_2 , which was associated with items relating to mood

dysregulation. This component explained 11.3% of the variance.

Figure 1: Principal component analysis performed on the ESWAN questionnaire taken as part of the HBN protocol. For visualization purposes, only the components included in our analysis are shown. We used the first two principal components as proxies of psychopathology, referred to as "pF" and "pF₂".

Network analysis

We chose ten resting-state networks from the model order of 25 for inclusion in the dFC analysis. These networks were chosen based on spatial correlation and manually ensuring

overlapping ICs with the ten RSNs reported by Smith et al. (2009). The ten networks

comprised the default mode (DMN), cerebellar (Cer), visual occipital (VO), visual medial

(VM), visual lateral (VL), right frontoparietal (FPR), left frontoparietal (FPL), sensorimotor

(SM), auditory (Au), and the executive control (Ex) network (Figure 2). These networks were

also used in previous DGM studies (Lund et al., 2020; Schwab et al., 2018). Including the

same nodes allowed us to compare results with prior findings and to integrate the current

results obtained from a childhood and adolescent sample with previous adult studies. The

included timeseries for each node were mean centered before estimating dFC from individual

level RSN time series using the DGM package v1.7.2 in R. DGM is a set of regressions with

time-varying coefficients (DLMs) for every receiving node. The receiver node refers to the

network that receives information from a sending node that transmits information between a

node pair. For each DLM, all possible models for the sender nodes are tested and the model

with the highest model evidence is selected. Consequently, a model is stated by the sending

networks of a node, giving a binary outcome. The outcome indicates if there is an influence or

not between two networks. DGM has uncovered meaningful trajectories in rsfMRI mice data

where DGM revealed a link between areas in the hippocampus that fed information to the

cingulate cortex, in line with previous studies using viral tracers to delineate directed

anatomical connectivity in mice (Schwab et al., 2018). Additionally, even with systemic

hemodynamic lag confounds being introduced in the data, network simulations demonstrated

a sensitivity of 72%–77% for the DGM approach (Schwab et al., 2018).

Figure 2: group-level maps for the 10 selected independent components included in the analysis (where the z-score maps threshold is set at 4).

<u>Statistical analysis</u>

We included all available rsfMRI scans for the group ICA, as a higher number of subjects is beneficial for yielding more robust ICs, while restricting the association analyses to the subset that had all covariates available. We performed the same analysis as previously reported in an adult sample (Lund et al., 2020), examining dFC on the edge- and node-level. Edge-level analysis deployed logistic regression for every connection of the directed network using directed connectivity as the response variable and testing for associations with age, age-orthogonalized age squared (age², using the poly function in R), sex, cognitive abilities, mental health, tSNR, motion and scanning site (as data was acquired at multiple scanners). These covariates were included into one model (see SI; SFig.5, for additional analyses examining potential multicollinearity for covariates included in the model). P-values were Bonferroni corrected for a number of 90 analyses on the edge-level (alpha level of 0.05). In addition, we performed node-level analysis to examine which networks overall send and receive information to each other (Lund et al., 2020). We calculated the number of output connections or outgoing edges (referred to as out-degree) and the number of input

connections or incoming edges (referred to as in-degree) for a given node. We performed

linear regression using in-degree and out-degree as dependent variables and the same

independent variables as used in the logistic regression on edge-level. P-values were

Bonferroni corrected for a number of 10 analysis on the node-level (alpha level of 0.05).

Results

Figure 3 depicts the average directed functional connectivity matrix across all individuals.

Several connections indicated bi-directionality in information flow, especially directionality

estimates for the visual lateral and visual medial networks where information flow from VM

to VL was present in 94.4% of individuals, while connection from VL to VM was present in

94.2%. In addition, there was overall a high number of participants that showed a reciprocal

relationship between the VO to VM and VL (VO to VL: 89.1%, VO to VM: 80.2%) and

opposite (VL to VO: 91.5%, VM to VO: 82.7%). These bi-directional relationships are in

alignment with patterns of information flow found in adults from the UK Biobank sample

(Lund et al., 2020). However, we did not observe the bi-directional information flow found

previously for adults between the DMN and FPR network. Further, when observing receiver

networks, the FPL showed the same as found for adults, that it did not receive information to

a large extent (Lund et al., 2020), coherent with our findings in youths. Additionally, the edge

most present across all individuals, aside from the visual networks, was Cer which receives

input from the Au (85.6%). This connection was also relatively strongly expressed in the

other direction (68.6%). Moreover, the FPL sent information to Cer, and this edge was

present in 77.7% of individuals. Yet, in this neurodevelopmental sample we did not observe

that Cer and Au overall were mostly receivers, as found for adults (Lund et al., 2020).

Figure 3: Average directed connectivity matrix across HBN subjects showing the significant

proportions of edges between the 10 RSNs included in the analysis; VO, visual occipital;

DMN, default mode; VM, visual medial; VL, visual lateral; FPR, frontoparietal right; FPL,

frontoparietal left; SM, sensorimotor; Cer, cerebellum; Au, auditory; Ex, executive control

network. The y-axis indicates the sender node, and the x-axis indicates the receiver nodes.

Significance was assessed using a binomial test as implemented in the binom.nettest function

in R, with a predefined FDR threshold of 5% and the hypothesized probability p0 = .56.

<u>Effects of age and cognition on dFC measures</u>

Edge-level analysis of dFC showed significant effects of age, in line with our hypothesis of

age being associated with the functional networks' maturation processes occurring in

childhood and adolescence (Figure 4; SI Tables 1-6 show z-scores and p- values and SFig.3-4

show effects of scanner). Specifically, we observed a positive association with age and dFC

from the Cer to the Au network. Correspondingly, the Au sends more information with higher

age to the SM node. For cognitive test performance and dFC, we observed a significant

negative association between VO and DMN, with the DMN receiving less information from

the VO network with higher cognitive test performance.

On the node-level, we assessed similar phenotypic associations for the out-degree and

in-degree of the networks. Confirming edge-level results, age was significantly associated

with the in-degree (number of input connections) of SM (fig.4b), indicating that the SM

network overall receive more information with increasing age. Additionally, for out-degree

(number of output connections), there was a significant positive association for age and FPR,

signifying that the FPR network overall sends more information with increasing age. As

expected, there was significant effects for the control variables scanning site (SFig.3-4), tSNR

and motion on edge and node-level.

Based on prior studies, we expected that mental health would be associated with

information flow for control networks, however, this was not observed. We also expected

sex-related differences in the maturation of brain networks, but this was not found.

Figure 4: Matrices showing the effects of age (a), age² (b), sex (c), intellectual abilities (FSIQ) (d), mental health (pF and pF₂) (e,f), motion (g), and tSNR (h) on directed connectivity. The analysis was performed in HBN data that had no missing values (N = 1143,

6–17 years, df = 1132). Significant edges following Bonferroni correction are marked as X.

The legend shows the 10 RSNs included in the analysis; VO, visual occipital; DMN, default

mode; VM, visual medial; VL, visual lateral; FPR, frontoparietal right; FPL, frontoparietal

left; SM, sensorimotor; Cer, cerebellum; Au, auditory; Ex, executive control network. The y-

axis indicates the sender node, while the x-axis refers to the receiving node. The colors reflect

the z-value for the corresponding effects where red indicates a positive association and blue a

negative association.

Figure 5: Associations on the node level (N=1143, 6-17 years, df= 1132). a) Out-degree matrix with corresponding effects of covariates age, age², sex, cognition (FSIQ), mental health (pF and pF₂), tSNR, and motion in HBN data. B) In-degree matrix with corresponding effects for the same covariates as in panel a). The colors reflect the t-value for the corresponding effect where numbers inside the boxes indicate t-statistic and p-value, and significant effects are marked with a red border following Bonferroni correction (p < .0.05).

Discussion

Our analysis revealed marked dFC patterns for the visual networks. Specifically, we observed a reciprocal information flow between the VM and VL in the investigated sample of children and adolescence. This is in alignment with previous findings in adults (Lund et al., 2020; Schwab et al., 2018), and coherent with primary sensory and motor cortical regions maturing first, and being present early on in children (Casey, Tottenham, Liston, & Durston, 2005). In contrast, we did not find that the cerebellum or auditory networks are mostly receivers of information as observed in adults (Lund et al., 2020; Schwab et al., 2018). However, apart from the visual circuits, the Cer received most information overall from the other transmitter networks, potentially indicating its role as a receiver, as found for adults.

We observed significant associations between age and directed connectivity both at edge and node-level. At the edge-level and with increasing age, the Cer had higher information flow to the Au network, while the Au sent more information to the SM network. Of note, these connections demonstrating age-dependent dFC have been implicated in neurodevelopmental disorders such as autism, which is characterized by symptoms related to hyper- and hypo sensitivity in relation to sensory modalities (Marco, Hinkley, Hill, & Nagarajan, 2011). Moreover, it has been reported that the cerebellar network is mostly a receiver of information from other attention and memory related regions in children with

ADHD when estimating directed connectivity in rsfMRI using a seed-based approach (Zhao et al., 2017). The average connectivity across participants indicated that the FPL sent information to Cer in 77.7% of the participants, which is in line with the aforementioned findings. Hence, this could be an interesting avenue to investigate in relation to potential biomarkers in future studies.

Likewise, on the node-level, the SM received more information with increasing age, while the FPR sent more information with higher age to the other networks overall. These results indicate that directionality estimates in both bottom-up and top-down processing are involved in the transitional phase from child to adulthood across patients and healthy individuals. This is in line with prior findings implicating these networks in normal functional development (Muetzel et al., 2016). Likewise, alterations in the right frontoparietal network has been associated with ADHD, where children and adolescent ADHD combined type patients show a reduction in FPR regions, when probing spatial working memory by use of task-fMRI (Silk et al., 2005; Vance et al., 2007).

For cognitive abilities and dFC, we observed a significant negative association between VO and DMN, with the DMN receiving less information from the VO network with higher cognitive test performance. This could indicate that individuals that were more focused on the task at hand benefited from less information flowing from VO to DMN, potentially facilitating effective whole-brain connectivity which has been linked to cognitive efficiency (van den Heuvel, Stam, Kahn, & Hulshoff Pol, 2009). However, it is challenging to delineate information flow as this could be disrupted for many connections and give rise to different alterations in connectivity. Nonetheless we did expect to observe associations for control networks or the DMN in relation to cognitive abilities as this has been reported by others (de Lacy, Doherty, King, Rachakonda, & Calhoun, 2017). Cognitive flexibility in healthy adults has been associated with strong within-system connectivity in higher order systems and from

these networks to primary-sensory motor systems (Chen et al., 2019). Here we estimated

information flow between networks with different functions rather than within-system

connectivity (with the exception of the visual networks). In addition, we used a normed

cognitive composite test score as a proxy for cognition. However, this may not be an optimal

approach as test norms could have limitations when it comes to representativeness and

number of subjects included in the normative sample, making it less sensitive for clinical

populations and for the lowest scoring percentile.

We utilized two general p factors for measuring mental health in this sample. These

psychopathology loadings, were related to self-control and depression/anxiety (pF) and items

pertaining to mood dysregulation (pF$_2$). Top-down control functions for cortico-limbic

regions have been linked to modifications for symptoms relating to mood dysregulation in

development. For instance, it has been shown that increased centromedial amygdala-rostral

anterior cingulate cortex functional connectivity is related to a higher level of anxiety and

depression symptoms during early adulthood, while increased structural connectivity in

centromedial amygdala-anterior ventromedial prefrontal cortex white matter was linked to

augmented symptoms of anxiety and depression during late childhood (Jalbrzikowski et al.,

2017). We did not observe significant associations between dFC and symptom burden for any

of the networks. Our dimensional approach of estimating PCA on symptom data represents

both healthy subjects and individuals with a psychiatric disorder on a symptom continuum

rather than as cases and control. Such a dimensional approach may better capture intermediate

phenotypes which map to brain biology and the neuronal mechanisms underlying the

symptoms, compared to diagnostic categories (Hengartner & Lehmann, 2017; Krueger &

Bezdjian, 2009). While the lack of significant associations in our study indicates that

between-network dFC in large-scale brain networks is not sensitive to mental health in this

sample, future research will need to investigate dFC in distinct subnetworks and by different

symptom measures, and also consider structural connectivity.

We expected to find differences between sexes in relation to directed connectivity as

this has been found previously in adolescence (Riley et al., 2018) using task-fMRI, and when

examining dFC in adults (Lund et al., 2020). However, we did not find any significant sex

differences in dFC on edge- or node-level in rsfMRI. This may partly relate to sample

characteristics such as the uneven sex distribution and the inclusion of patients in our sample.

<u>Limitations</u>

As with any MRI study, participant motion during scanning may confound the results,

especially as it is known that both young individuals and those with a psychiatric diagnosis on

average move more (Satterthwaite et al., 2012). We therefore implemented a stringent

correction pipeline, where we implemented several steps of ICA and machine learning based

cleaning (FIX and AROMA). We also included the mean relative motion measure from FSL

as a covariate in our analysis step. Another relevant confound effect that could potentially

bias our results and the comparison to previous work is variability in the ICA decompositions.

We performed ICA in the study sample and selected components based on similarity to

components reported by Smith et al. (2009). While this approach yielded overall good

comparison to previous work, subtle differences between decompositions may retain and

might impact the derived dFC patterns. For instance, sensorimotor nodes were divided into

three independent components where the node with the best spatial overlap with the SM

component in Smith et al. (2009) was chosen. Most likely including another sensorimotor

related node would change the results of our analysis. Hence, our results call for follow-up

analysis in more fine-grained networks. In this framework, different ways to parcellate the

brain could be tested.

The majority of the individuals in the sample were patients. This was beneficial when

examining associations of psychopathology on directionality estimates as it yielded a high

number of patients in each group, but makes it difficult to delineate if average patterns of

information flow are for typical or non-typical functional development. In addition,

medication may have had an effect on our results, as participants were asked to discontinue

stimulant medication but were still enrolled if they chose not to discontinue due to personal

reasons or if recommended by their physician.

On a methodological note, DGM reflects instantaneous relationships and does not only

look at lagged relationships, however sensitivity of DGM drops depending on the total offset

between a node pair and can as such influence the estimation of these interactions. Further,

DGM estimates binary connections, which may have reduced the sensitivity of the association

analyses.

Lastly, whether differences in findings compared to previous studies in adults are of

methodological origin, such as differences in preprocessing pipelines or network definitions,

or can be explained by differences in dynamic connectivity between children and adults

remains to be further investigated.

Conclusions

To conclude, using a sample of children and adolescents, we showed that the direction of the

information flow was age dependent for auditory, sensorimotor and cerebellar connections. In

addition, the FPR had a higher degree of input connections with higher age. These findings

contribute to the existing knowledge in the brain development field, and warrant further

studies for replication in healthy samples as well as other clinical populations.

Funding

The authors were funded by the Research Council of Norway #276082 (LifespanHealth), #223273 (NORMENT), #249795, #298646, #300767, #283798; H2020 European Research Council #802998 (BRAINMINT); The South-East Norway Regional Health Authority #2019101, #2019107, #2020086; Swiss National Science Foundation #171598

Financial disclosures

OAA is a consultant for HealthLytix.

Data and code availability

The data incorporated in this work were gathered from the open access Healthy Brain

Network resource. Software needed to estimate directed connectivity is available at

https://github.com/schw4b/DGM.

Acknowledgements

This manuscript was prepared using a limited access dataset obtained from the Child Mind Institute Biobank, the HBN resource (http://www.healthybrainnetwork.org). Its initiatives are supported by philanthropic contributions from the following individuals, foundations and organizations: Margaret Bilotti; Brooklyn Nets; Agapi and Bruce Burkard; James Chang; Phyllis Green and Randolph Cōwen; Grieve Family Fund; Susan Miller and Byron Grote; Sarah and Geoff Gund; George Hall; Jonathan M. Harris Family Foundation; Joseph P. Healey; The Hearst Foundations; Eve and Ross Jaffe; Howard & Irene Levine Family Foundation; Rachael and Marshall Levine; George and Nitzia Logothetis; Christine and Richard Mack; Julie Minskoff; Valerie Mnuchin; Morgan Stanley Foundation; Amy and John Phelan; Roberts Family Foundation; Jim and Linda Robinson Foundation, Inc.; The Schaps Family; Zibby Schwarzman; Abigail Pogrebin and David Shapiro; Stavros Niarchos

Foundation; Preethi Krishna and Ram Sundaram; Amy and John Weinberg; Donors to the

2013 Child Advocacy Award Dinner Auction; Donors to the 2012 Brant Art Auction.

This manuscript reflects the views of the authors and does not necessarily reflect the

opinions or views of the Child Mind Institute.

This work was performed on the TSD (Tjeneste for Sensitive Data) facilities, owned

by the University of Oslo, operated and developed by the TSD service group at the University

of Oslo, IT-Department (USIT) (tsd-drift@usit.uio.no).

<u>References</u>

Alexander, L. M., Escalera, J., Ai, L., Andreotti, C., Febre, K., Mangone, A., . . . Milham, M.
P. (2017). An open resource for transdiagnostic research in pediatric mental health and
learning disorders. *Scientific Data, 4*(1). doi:10.1038/sdata.2017.181
Alexander, L. M., Salum, G. A., Swanson, J. M., & Milham, M. P. (2020). Measuring
strengths and weaknesses in dimensional psychiatry. *J Child Psychol Psychiatry,*
61(1), 40-50. doi:10.1111/jcpp.13104
Alnaes, D., Kaufmann, T., Doan, N. T., Cordova-Palomera, A., Wang, Y., Bettella, F., . . .
Westlye, L. T. (2018). Association of Heritable Cognitive Ability and
Psychopathology With White Matter Properties in Children and Adolescents. *JAMA*
Psychiatry, 75(3), 287-295. doi:10.1001/jamapsychiatry.2017.4277
Beckmann, C. F., & Smith, S. M. (2004). Probabilistic Independent Component Analysis for
Functional Magnetic Resonance Imaging. *IEEE Transactions on Medical Imaging,*
23(2), 137-152. doi:10.1109/tmi.2003.822821
Bielczyk, N. Z., Uithol, S., van Mourik, T., Anderson, P., Glennon, J. C., & Buitelaar, J. K.
(2019). Disentangling causal webs in the brain using functional magnetic resonance
imaging: A review of current approaches. *Netw Neurosci, 3*(2), 237-273.
doi:10.1162/netn_a_00062
Bitan, T., Burman, D. D., Lu, D., Cone, N. E., Gitelman, D. R., Mesulam, M. M., & Booth, J.
R. (2006). Weaker top-down modulation from the left inferior frontal gyrus in
children. *NeuroImage, 33*(3), 991-998. doi:10.1016/j.neuroimage.2006.07.007
Blakemore, S.-J. (2012). Imaging brain development: The adolescent brain. *NeuroImage,*
61(2), 397-406. doi:10.1016/j.neuroimage.2011.11.080
Casey, B. J., Tottenham, N., Liston, C., & Durston, S. (2005). Imaging the developing brain:
what have we learned about cognitive development? *Trends Cogn Sci, 9*(3), 104-110.
doi:10.1016/j.tics.2005.01.011
Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., . . .
Moffitt, T. E. (2013). The p Factor. *Clinical Psychological Science, 2*(2), 119-137.
doi:10.1177/2167702613497473

Chen, O. Y., Cao, H., Reinen, J. M., Qian, T., Gou, J., Phan, H., . . . Cannon, T. D. (2019). Resting-state brain information flow predicts cognitive flexibility in humans. *Sci Rep, 9*(1), 3879. doi:10.1038/s41598-019-40345-8

Connolly, C. G., Wu, J., Ho, T. C., Hoeft, F., Wolkowitz, O., Eisendrath, S., . . . Yang, T. T. (2013). Resting-state functional connectivity of subgenual anterior cingulate cortex in depressed adolescents. *Biol Psychiatry, 74*(12), 898-907. doi:10.1016/j.biopsych.2013.05.036

Cortese, S., Kelly, C., Chabernaud, C., Proal, E., , & Di Martino, A., Milham, M. P., Castellanos, F. X. . (2012). Toward systems neuroscience of ADHD: a meta-analysis of 55 fMRI studies. . *American Journal of Psychiatry, 169*(10), 1038-1055.

Craddock, N., & Owen, M. J. (2010). The Kraepelinian dichotomy - going, going... but still not gone. *Br J Psychiatry, 196*(2), 92-95. doi:10.1192/bjp.bp.109.073429

de Lacy, N., Doherty, D., King, B. H., Rachakonda, S., & Calhoun, V. D. (2017). Disruption to control network function correlates with altered dynamic connectivity in the wider autism spectrum. *Neuroimage Clin, 15*, 513-524. doi:10.1016/j.nicl.2017.05.024

Demontis, D., Walters, R. K., Martin, J., Mattheisen, M., Als, T. D., Agerbo, E., . . . Neale, B. M. (2019). Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet, 51*(1), 63-75. doi:10.1038/s41588-018-0269-7

Di Martino, A., Kelly, C., Grzadzinski, R., Zuo, X. N., Mennes, M., Mairena, M. A., . . . Milham, M. P. (2011). Aberrant striatal functional connectivity in children with autism. *Biol Psychiatry, 69*(9), 847-856. doi:10.1016/j.biopsych.2010.10.029

Eisenberg, D. P., & Berman, K. F. (2010). Executive function, neural circuitry, and genetic mechanisms in schizophrenia. *Neuropsychopharmacology, 35*(1), 258-277. doi:10.1038/npp.2009.111

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J. (2017). MRIQC: Advancing the automatic prediction of image quality in MRI from unseen sites. *PLoS One, 12*(9), e0184661. doi:10.1371/journal.pone.0184661

Filippini, N., MacIntosh, B. J., Hough, M. G., Goodwin, G. M., Frisoni, G. B., Smith, S. M., . . . Mackay, C. E. (2009). Distinct patterns of brain activity in young carriers of the APOE- 4 allele. *Proceedings of the National Academy of Sciences, 106*(17), 7209-7214. doi:10.1073/pnas.0811879106

Francx, W., Oldehinkel, M., Oosterlaan, J., Heslenfeld, D., Hartman, C. A., Hoekstra, P. J., . . . Mennes, M. (2015). The executive control network and symptomatic improvement in attention-deficit/hyperactivity disorder. *Cortex, 73*, 62-72. doi:10.1016/j.cortex.2015.08.012

Friston, K. J. (2011). Functional and Effective Connectivity: A Review. *Brain Connectivity, 1*(1), 13-36. doi:10.1089/brain.2011.0008

Friston, K. J., & Frith, C. D. (1995). Schizophrenia: a disconnection syndrome. *Clin Neurosci, 3*(2), 89-97.

Geiger, M. J., Domschke, K., Ipser, J., Hattingh, C., Baldwin, D. S., Lochner, C., & Stein, D. J. (2016). Altered executive control network resting-state connectivity in social anxiety disorder. *World J Biol Psychiatry, 17*(1), 47-57. doi:10.3109/15622975.2015.1083613

Griffanti, L., Salimi-Khorshidi, G., Beckmann, C. F., Auerbach, E. J., Douaud, G., Sexton, C. E., . . . Smith, S. M. (2014). ICA-based artefact removal and accelerated fMRI acquisition for improved resting state network imaging. *NeuroImage, 95*, 232-247. doi:10.1016/j.neuroimage.2014.03.034

Grove, J., Ripke, S., Als, T. D., Mattheisen, M., Walters, R. K., Won, H., . . . Borglum, A. D. (2019). Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet, 51*(3), 431-444. doi:10.1038/s41588-019-0344-8

Hamm, L. L., Jacobs, R. H., Johnson, M. W., Fitzgerald, D. A., Fitzgerald, K. D., Langenecker, S. A., . . . Phan K. L. (2014). Aberrant amygdala functional connectivity at rest in pediatric anxiety disorders. *Biology of mood & anxiety disorders, 4(1)*(15).

Hengartner, M. P., & Lehmann, S. N. (2017). Why Psychiatric Research Must Abandon Traditional Diagnostic Classification and Adopt a Fully Dimensional Scope: Two Solutions to a Persistent Problem. *Front Psychiatry, 8*, 101. doi:10.3389/fpsyt.2017.00101

Hibar, D. P., Stein, J. L., Renteria, M. E., Arias-Vasquez, A., Desrivieres, S., Jahanshad, N., . . . Medland, S. E. (2015). Common genetic variants influence human subcortical brain structures. *Nature, 520*(7546), 224-229. doi:10.1038/nature14101

Hoff, G. E., Van den Heuvel, M. P., Benders, M. J., Kersbergen, K. J., & De Vries, L. S. (2013). On development of functional brain connectivity in the young brain. *Front Hum Neurosci, 7*, 650. doi:10.3389/fnhum.2013.00650

Hwang, K., Velanova, K., & Luna, B. (2010). Strengthening of top-down frontal cognitive control networks underlying the development of inhibitory control: a functional magnetic resonance imaging effective connectivity study. *J Neurosci, 30*(46), 15535-15545. doi:10.1523/JNEUROSCI.2825-10.2010

Hyvärinen, A. (1999). Fast and robust fixed-point algorithms for independent component analysis. *IEEE Transactions on Neural Networks, 10*(3), 626–634.

Insel, T. R. (2010). Rethinking schizophrenia. *Nature, 468*(7321), 187-193. doi:10.1038/nature09552

Jalbrzikowski, M., Larsen, B., Hallquist, M. N., Foran, W., Calabro, F., & Luna, B. (2017). Development of White Matter Microstructure and Intrinsic Functional Connectivity Between the Amygdala and Ventromedial Prefrontal Cortex: Associations With Anxiety and Depression. *Biol Psychiatry, 82*(7), 511-521. doi:10.1016/j.biopsych.2017.01.008

Jollans, L., & Whelan, R. (2018). Neuromarkers for Mental Disorders: Harnessing Population Neuroscience. *Front Psychiatry, 9*, 242. doi:10.3389/fpsyt.2018.00242

Kaufmann, T., Alnaes, D., Doan, N. T., Brandt, C. L., Andreassen, O. A., & Westlye, L. T. (2017). Delayed stabilization and individualization in connectome development are related to psychiatric disorders. *Nat Neurosci, 20*(4), 513-515. doi:10.1038/nn.4511

Keshavan, M. S., Giedd, J., Lau, J. Y. F., Lewis, D. A., & Paus, T. (2014). Changes in the adolescent brain and the pathophysiology of psychotic disorders. *The Lancet Psychiatry, 1*(7), 549-558. doi:10.1016/s2215-0366(14)00081-9

Kolskar, K. K., Alnaes, D., Kaufmann, T., Richard, G., Sanders, A. M., Ulrichsen, K. M., . . . Westlye, L. T. (2018). Key Brain Network Nodes Show Differential Cognitive Relevance and Developmental Trajectories during Childhood and Adolescence. *eNeuro, 5*(4). doi:10.1523/ENEURO.0092-18.2018

Krueger, R. F., & Bezdjian, S. (2009). Enhancing research and treatment of mental disorders with dimensional concepts: toward DSM-V and ICD-11. . *World Psychiatry, 8(1)*(3).

Li, C. L., Deng, Y. J., He, Y. H., Zhai, H. C., & Jia, F. C. (2019). The development of brain functional connectivity networks revealed by resting-state functional magnetic resonance imaging. *Neural Regen Res, 14*(8), 1419-1429. doi:10.4103/1673-5374.253526

Li, P., Fan, T. T., Zhao, R. J., Han, Y., Shi, L., Sun, H. Q., . . . Lu, L. (2017). Altered Brain Network Connectivity as a Potential Endophenotype of Schizophrenia. *Sci Rep, 7*(1), 5483. doi:10.1038/s41598-017-05774-3

Lund, M. J., Alnaes, D., Schwab, S., van der Meer, D., Andreassen, O. A., Westlye, L. T., &
Kaufmann, T. (2020). Differences in directed functional brain connectivity related to
age, sex and mental health. *Hum Brain Mapp.* doi:10.1002/hbm.25116
Mallard, T. T., Linnér, R. K., Okbay, A., Grotzinger, A. D., de Vlaming, R., Meddens, S. F.
W., . . . Harden, K. P. (2019). Not just one p: Multivariate GWAS of psychiatric
disorders and their cardinal symptoms reveal two dimensions of cross-cutting genetic
liabilities. . *bioRxiv, 603134.* doi:10.1101/603134
Marco, E. J., Hinkley, L. B., Hill, S. S., & Nagarajan, S. S. (2011). Sensory processing in
autism: a review of neurophysiologic findings. *Pediatr Res, 69*(5 Pt 2), 48R-54R.
doi:10.1203/PDR.0b013e3182130c54
Muetzel, R. L., Blanken, L. M., Thijssen, S., van der Lugt, A., Jaddoe, V. W., Verhulst, F. C.,
. . . White, T. (2016). Resting-state networks in 6-to-10 year old children. *Hum Brain
Mapp, 37*(12), 4286-4300. doi:10.1002/hbm.23309
Pruim, R. H. R., Mennes, M., Buitelaar, J. K., & Beckmann, C. F. (2015). Evaluation of ICA-
AROMA and alternative strategies for motion artifact removal in resting state fMRI.
NeuroImage, 112, 278-287. doi:10.1016/j.neuroimage.2015.02.063
Pruim, R. H. R., Mennes, M., van Rooij, D., Llera, A., Buitelaar, J. K., & Beckmann, C. F.
(2015). ICA-AROMA: A robust ICA-based strategy for removing motion artifacts
from fMRI data. *NeuroImage, 112*, 267-277. doi:10.1016/j.neuroimage.2015.02.064
Rausch, A., Zhang, W., Haak, K. V., Mennes, M., Hermans, E. J., van Oort, E., . . . Groen, W.
B. (2016). Altered functional connectivity of the amygdaloid input nuclei in
adolescents and young adults with autism spectrum disorder: a resting state fMRI
study. *Mol Autism, 7*, 13. doi:10.1186/s13229-015-0060-x
Riley, J. D., Chen, E. E., Winsell, J., Davis, E. P., Glynn, L. M., Baram, T. Z., . . . Solodkin,
A. (2018). Network specialization during adolescence: Hippocampal effective
connectivity in boys and girls. *NeuroImage, 175*, 402-412.
doi:10.1016/j.neuroimage.2018.04.013
Salimi-Khorshidi, G., Douaud, G., Beckmann, C. F., Glasser, M. F., Griffanti, L., & Smith, S.
M. (2014). Automatic denoising of functional MRI data: combining independent
component analysis and hierarchical fusion of classifiers. *NeuroImage, 90*, 449-468.
doi:10.1016/j.neuroimage.2013.11.046
Satterthwaite, T. D., Wolf, D. H., Loughead, J., Ruparel, K., Elliott, M. A., Hakonarson, H., .
. . Gur, R. E. (2012). Impact of in-scanner head motion on multiple measures of
functional connectivity: relevance for studies of neurodevelopment in youth.
NeuroImage, 60(1), 623-632. doi:10.1016/j.neuroimage.2011.12.063
Schizophrenia Working Group of the Psychiatric Genomics, C. (2014). Biological insights
from 108 schizophrenia-associated genetic loci. *Nature, 511*(7510), 421-427.
doi:10.1038/nature13595
Schwab, S., Harbord, R., Zerbi, V., Elliott, L., Afyouni, S., Smith, J. Q., . . . Nichols, T. E.
(2018). Directed functional connectivity using dynamic graphical models.
NeuroImage, 175, 340-353. doi:10.1016/j.neuroimage.2018.03.074
Schweinsburg, A. D., Nagel, B. J., & Tapert, S. F. (2005). fMRI reveals alteration of spatial
working memory networks across adolescence. *J Int Neuropsychol Soc, 11*(5), 631-
644. doi:10.1017/S1355617705050757
Shannon, K. E., Sauder, C., Beauchaine, T. P., & Gatzke-Kopp, L. M. (2009). Disrupted
Effective Connectivity Between the Medial Frontal Cortex and the Caudate in
Adolescent Boys With Externalizing Behavior Disorders. *Criminal Justice and
Behavior, 36*(11), 1141-1157. doi:10.1177/0093854809342856
Silk, T., Vance, A., Rinehart, N., Egan, G., O'boyle, M., Bradshaw, J. L., & Cunnington, R.
(2005). Fronto-parietal activation in attention-deficit hyperactivity disorder, combined

type: functional magnetic resonance imaging study. *The British Journal of Psychiatry,*
187(3), 282-283.
Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., . . .
Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during
activation and rest. *Proc Natl Acad Sci U S A, 106*(31), 13040-13045.
doi:10.1073/pnas.0905267106
Talkowski, M. E., Rosenfeld, J. A., Blumenthal, I., Pillalamarri, V., Chiang, C., Heilbut, A., .
. . Gusella, J. F. (2012). Sequencing chromosomal abnormalities reveals
neurodevelopmental loci that confer risk across diagnostic boundaries. *Cell, 149*(3),
525-537. doi:10.1016/j.cell.2012.03.028
van den Heuvel, M. P., Stam, C. J., Kahn, R. S., & Hulshoff Pol, H. E. (2009). Efficiency of
functional brain networks and intellectual performance. *J Neurosci, 29*(23), 7619-
7624. doi:10.1523/JNEUROSCI.1443-09.2009
Vance, A., Silk, T. J., Casey, M., Rinehart, N. J., Bradshaw, J. L., Bellgrove, M. A., &
Cunnington, R. (2007). Right parietal dysfunction in children with attention deficit
hyperactivity disorder, combined type: a functional MRI study. *Mol Psychiatry, 12*(9),
826-832, 793. doi:10.1038/sj.mp.4001999
Wechsler, D. (2003). *Wechsler intelligence scale for children--Fourth Edition (WISC-IV)*
Woolrich, M. W., Ripley, B. D., Brady, M., & Smith, S. M. (2001). Temporal autocorrelation
in univariate linear modeling of FMRI data. *NeuroImage, 14*(6), 1370-1386.
doi:10.1006/nimg.2001.0931
Zhao, D., Zheng, S., Yang, L., & Tian, Y. (2017). Causal connectivity abnormalities of
regional homogeneity in children with attention deficit hyperactivity disorder: a rest-
state fMRI study. *ADMET and DMPK, 5*(4), 242-252. doi:10.5599/admet.5.4.485
Zhao, Q., Swati, Z. N. K., Metmer, H., Sang, X., & Lu, J. (2019). Investigating executive
control network and default mode network dysfunction in major depressive disorder.
Neurosci Lett, 701, 154-161. doi:10.1016/j.neulet.2019.02.045
736