

1 The role of masks, testing and contact tracing in preventing
2 COVID-19 resurgences: a case study from New South Wales,
3 Australia

4
5 RM Stuart^{1,2†}, Romesh G. Abeysuriya², Cliff C. Kerr^{3,4}, Dina Mistry³, Daniel J. Klein³, Richard Gray⁵,
6 Margaret Hellard^{2,6,7,8,9}, Nick Scott^{2,6}

7
8 1. Department of Mathematical Sciences, University of Copenhagen, Copenhagen, Denmark

9 2. Disease Elimination Program, Burnet Institute, Melbourne, Victoria, Australia

10 3. Institute for Disease Modeling, Global Health Division, Bill & Melinda Gates Foundation, Seattle,
11 USA

12 4. School of Physics, University of Sydney, Sydney, New South Wales, Australia

13 5. The Kirby Institute, UNSW Sydney, Sydney, New South Wales, Australia

14 6. School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria,
15 Australia

16 7. University of Melbourne, Parkville, Victoria, Australia

17 8. Department of Infectious Diseases, The Alfred Hospital and Monash University, Melbourne,
18 Victoria, Australia

19 9. The School of Population and Global Health and the Peter Doherty Institute for Infection and
20 Immunity, Melbourne, Victoria, Australia

21 † Corresponding author. E-mail: robyn@math.ku.dk

22

23

24

25 **Key words:** COVID-19, coronavirus, SARS-CoV-2, modelling, masks, testing, contact tracing,
26 Australia

27

28

1 Abstract

2 **Objectives:** The early stages of the COVID-19 pandemic illustrated that SARS-CoV-2, the virus that
3 causes the disease, has the potential to spread exponentially. Therefore, as long as a substantial
4 proportion of the population remains susceptible to infection, the potential for new epidemic waves
5 persists even in settings with low numbers of active COVID-19 infections, unless sufficient
6 countermeasures are in place. We aim to quantify vulnerability to resurgences in COVID-19
7 transmission under variations in the levels of testing, tracing, and mask usage.

8
9 **Setting:** The Australian state of New South Wales, a setting with prolonged low transmission, high
10 mobility, non-universal mask usage, and a well-functioning test-and-trace system.

11
12 **Participants:** None (simulation study)

13
14 **Results:** We find that the relative impact of masks is greatest when testing and tracing rates are
15 lower (and vice versa). Scenarios with very high testing rates (90% of people with symptoms, plus
16 90% of people with a known history of contact with a confirmed case) were estimated to lead to a
17 robustly controlled epidemic, with a median of ~180 infections in total over October 1 – December
18 31 under high mask uptake scenarios, or 260–1,200 without masks, depending on the efficacy of
19 community contact tracing. However, across comparable levels of mask uptake and contact tracing,
20 the number of infections over this period were projected to be 2-3 times higher if the testing rate was
21 80% instead of 90%, 8-12 times higher if the testing rate was 65%, or 30-50 times higher with a 50%
22 testing rate. In reality, NSW diagnosed 254 locally-acquired cases over this period, an outcome that
23 had a low probability in the model (4-7%) under the best-case scenarios of extremely high testing
24 (90%), near-perfect community contact tracing (75-100%), and high mask usage (50-75%), but a far
25 higher probability if any of these were at lower levels.

26
27 **Conclusions:** Our work suggests that testing, tracing and masks can all be effective means of
28 controlling transmission. A multifaceted strategy that combines all three, alongside continued
29 hygiene and distancing protocols, is likely to be the most robust means of controlling transmission
30 of SARS-CoV-2.

31 **Strengths and limitations of this study**

- A key methodological strength of this study is the level of detail in the model that we use, which allows us to capture many of the finer details of the extent to which controlling COVID-19 transmission relies on the balance between testing, contact tracing, and mask usage.
- Another key strength is that our model is stochastic, so we are able to quantify the probability of different epidemiological outcomes under different policy settings.
- A key limitation is the shortage of publicly-available data on the efficacy of contact tracing programs, including data on how many people were contacted for each confirmed index case of COVID-19.

1 Introduction

2 Across the world, governmental responses to the outbreak of the COVID-19 pandemic in the first
3 half of 2020 profoundly curtailed the spread of SARS-CoV-2, the virus that causes the disease (1–4).
4 By midway through the year, an increasing number of countries had moved from an initial crisis-
5 management phase into a new phase centred around minimising transmission risk while allowing
6 societal and economic activities to resume (5). However, by the end of 2020, many countries around
7 the world had experienced epidemic resurgences necessitating further shutdowns (6). This was true
8 even in settings that had come close to eliminating the virus, such as Vietnam (7), New Zealand (8),
9 and Australia (9). In low- or zero-transmission contexts, new outbreaks can emerge if community
10 transmission has not been eliminated, or if infected people arrive from abroad or interstate and
11 interact with the local community (10). It is therefore essential to be able to quantify the risk of
12 epidemic resurgence under different policy settings.

13
14 Given the complexities of COVID-19 transmission, including the duration of pre-symptomatic
15 infection (11,12), the proportion of infections that are asymptomatic (13), and the possibility of
16 transmission via surface contact (14), maintaining control of COVID-19 has proven challenging in
17 many jurisdictions (15–19). The often-cited success stories of Taiwan, Vietnam, Thailand, and South
18 Korea included high mask usage, high rates of testing, and fast, effective contact tracing (14–16). The
19 benefits of a multi-pronged approach have also been illustrated in the literature; in the UK, for
20 example, a recent study (23) found that mandating masks in secondary schools could achieve
21 approximately the same reduction in resurgence risk as having an 8-11% increase in symptomatic
22 testing.

23
24 In this work, we use an agent-based model to estimate the combination of testing, community-based
25 contact tracing, and mask usage required to maintain epidemic control in a low-transmission, high-
26 mobility setting. These three non-pharmaceutical interventions (NPIs) were key components in
27 reducing the probability of epidemic resurgences prior to the availability of a vaccine, and are likely
28 to remain so for some time even after vaccination coverage increases (24). When used in combination
29 with physical distancing and hand-washing/hygiene measures, all three strategies allow relatively
30 high mobility: testing and contact tracing means that only those at greatest risk of transmitting the
31 virus need to stay home and have been shown to be effective in numerous settings (15,25–29), while
32 masks mean that people with undiagnosed infections present less of a risk to others (30–34).

33
34 The context for our study is the Australian state of New South Wales (NSW), with a population of
35 7.5 million and a cumulative total of just over 4700 diagnosed cases as of December 31, 2020. After
36 an initial wave of COVID-19 infections in March and subsequent lockdown in April, New South
37 Wales began relaxing physical lockdown measures over May and was experiencing near-zero case
38 counts by the start of June, with students back at school, businesses reopening and social/community
39 activities resuming. In late June several clusters of new infections were detected, which subsequently
40 led to a 4-month long period of low but steady case counts, with a mean of ~5 cases per day over July
41 to October (excluding cases in quarantined travellers). However, NSW subsequently went on to
42 record ~180 cases in the last two weeks of 2020, a result of a localised outbreak whose containment
43 necessitated the introduction of stringent new restrictions on travel and gatherings that affected
44 many over the holiday period (35).

45
46 During the prolonged period of low transmission that NSW experienced in the second half of 2020
47 prior to December 15, 2020, mobility remained high (36) and transmission was controlled via NPIs.

1 Masks were recommended by the government for the general public and made mandatory for staff
2 in various businesses including supermarkets, but were not universally adopted. In a survey
3 undertaken in mid-September, 78% of the NSW population reported wearing a mask at some point
4 in the previous week (37), although CCTV footage from August registered ~30% of passengers on
5 urban public transport wearing masks (38). At the same time, high levels of testing were in place,
6 with ~20,000 people tested per day over June–September (~2.7/day per 1,000 people), resulting in an
7 average testing yield of 0.05%, one of the world’s lowest (39). The state also had a strong focus on
8 contact tracing, with all cases interviewed within one day of case notification (40). Over the four
9 months from June 1 – September 30, 2020, ~900 new cases were identified, but only 45 (5%) of were
10 classified as “source unknown”, meaning that they were neither acquired overseas/interstate nor
11 linked to known clusters (41).

12
13 Relative to many other contact tracing programs across the world, the NSW contact tracing program
14 was differentiated by its extensive efforts to identify a person’s community contacts in addition to
15 their household, social, school, and workplace contacts (42). The state’s health department (NSW
16 Health) required all businesses to have a COVID-19 Safety Plan, and for the majority of public-facing
17 businesses this included a requirement for customers to register their details upon entry. Upon
18 identifying a new case, NSW Health’s contact tracers would then (a) conduct an extended interview
19 to determine all possible venues in which transmission may have occurred, (b) place details of those
20 venues on their website, on social media, and in newspapers, urging people who had been at the
21 venue to self-isolate for 14 days, (c) attempt to contact all people who were registered to have been
22 at venues within a given window of the time that the diagnosed case was known to have been there
23 and instruct them to self-isolate for 14 days (42).

24
25 The dynamics of COVID-19 transmission are complex, and in low-transmission settings the
26 probability of maintaining epidemic control depends on numerous factors outside of policy control,
27 including the characteristics of people who get infected: the size of their households, the type of work
28 that they do, and a number of other socio-economic factors that may influence their contact
29 networks, access to testing and capacity to self-isolate. Several studies have pointed to the role of
30 superspreading events and overdispersion of infections in COVID-19 transmission (25,26,43,44). As a
31 result, even with physical distancing, high levels of testing, and rapid contact tracing, there is still a
32 non-zero probability that a sustained outbreak could occur depending on who gets infected and
33 where. In this study, we consider a range of testing and contact tracing levels, and assess the roles
34 of masks, testing and contact tracing as a means of controlling community-based transmission.

35

36 Methods

37 Transmission model


38 We used an open-source agent-based model, Covasim (45), developed by the Institute for Disease
39 Modeling with source code and documentation available at <https://covasim.org>, and previously
40 adapted by our group to model the Victorian epidemic (10,46). Covasim contains detailed
41 descriptions of age-dependent disease acquisition and progression probabilities, duration of disease
42 by acuity, and the effects of interventions including symptomatic and asymptomatic testing,
43 isolation, contact tracing, and quarantine, as well as other NPIs such as physical distancing, hygiene
44 measures, and protective equipment such as masks. Importantly, it also captures individual
45 variability, with viral loads varying both between individuals and over time.

46

1 We began by simulating a population representative of New South Wales by taking data on the age
 2 and sex composition of the population from the 2016 census (the latest available), and using it to
 3 create a model population of agents with similar characteristics. The simulations consist of 100,000
 4 individual agents, who are dynamically scaled based on prevalence to represent the total New South
 5 Wales population of 7.5 million. The dynamical scaling means that whenever the proportion of
 6 susceptible agents falls below a threshold of 5%, the number of agents in the model is increased;
 7 further implementation details can be found in Section 2.3.6 of Kerr et al (45).

8
 9 Next, we created contact networks for these agents. The governmental response to COVID-19 in New
 10 South Wales consisted of a set of highly context-specific policies covering individuals, businesses,
 11 schools, and other types of organisations. To model these policies, we allow agents in the model to
 12 interact over five types of contact network: households, schools, workplaces, and static and dynamic
 13 community networks. The static community network consists of interactions with friends,
 14 colleagues, or other known associates who come together on a regular and predictable basis, and
 15 contains four sub-networks: professional sports, community sports/fitness/leisure clubs, places of
 16 worship, and socialising with friends. The dynamic community network consists of interactions in
 17 which people interact with strangers or random groups of people, and contains seven separate sub-
 18 networks, representing: (1) arts venues such as museums, galleries, theatres, and cinemas, (2) large
 19 events such as concerts, festivals, sports games, (3) pubs and bars, (4) cafes and restaurants, (5) public
 20 parks and other outdoor settings, (6) public transport, (7) all other community settings. The method
 21 for constructing these networks is described in our previous study of the Victorian epidemic (46) and
 22 is based on the methodology of the SynthPops Python package (47).

23



24
 25 **Figure 1.** Relative changes in network structure and transmission risk across different settings in
 26 New South Wales over March–August. The absolute transmission risk varies by setting and is highest
 27 in household and lowest in outdoor settings (see (46) for details).

28
 29

1 Finally, to simulate the epidemic and policy environment in New South Wales over March 1 –
2 September 30, 2020, we include parameter changes that capture the testing, tracing, isolation,
3 quarantine, and lockdown policies that were enacted over this time. Figure 1 presents a summary of
4 how contact networks and the relative risk of COVID-19 transmission in different settings changed
5 as policies evolved. Some of these changes in transmission risk are derived from available data (42),
6 while others are taken from a similar modelling exercise conducted in Victoria, in which a panel of
7 Australia-based experts reviewed the likely effect of policies on transmission risks (46). Further
8 details of all policies and how we model their effects on transmission risk are contained in
9 Supplementary Table 1. We also model the introduction of interstate cases in late June prior to the
10 closure of the state border with neighbouring Victoria.
11

12 Data and model fitting

13 We calibrate the model by adjusting the per-contact transmission risk to minimise the absolute
14 differences between the model projections and data on the daily number of cases diagnosed,
15 excluding cases acquired overseas or interstate from NSW Health (41). Specifically, we drew 500
16 samples from a Gaussian prior distribution of values for the per-contact transmission risk
17 ($\sim N(0.025, 0.002)$). For each sample, we run $N=20$ simulations for each to produce 10,000 trial
18 simulations, and then retain the 1% of these with the minimum absolute differences between the
19 model projections and the data.
20

21 Model analysis and timeframe

22 As described above, we fit the model using data up until September 30, 2020. We then run the model
23 for three months (from October 1, 2020, until December 31, 2020), under a set of assumptions about
24 future testing rates, the efficacy of contact tracing, and mask uptake. For each scenario (described
25 below), we quantify the probability of the epidemic exceeding certain thresholds. We then compare
26 these projections to the observed epidemic outcomes over the period from October 1 to December
27 31, 2020.
28

29 For testing rates, we distinguish between symptom-based testing and testing of asymptomatic
30 contacts. NSW Health guidelines advise anyone with symptoms to get tested, as well as anyone
31 identified as a contact regardless of symptoms (42). To reflect this, we use a baseline assumption that
32 the testing rate for asymptomatic contacts is the same as for people with symptoms, and run a
33 counterfactual set of scenarios in which it is assumed to be half the rate for those with symptoms
34 (Table 1). We modelled scenarios in which 50%, 65%, 80%, or 90% of those with symptoms get tested.
35 Test results are assumed to be communicated within 24 hours (50).
36

37 We also considered variations on mask uptake and effectiveness. On uptake, we modelled scenarios
38 in which 0%, 25%, 50%, or 75% of the population will wear masks in dynamic community settings
39 (i.e., settings in which people interact with strangers or random groups of people), which in the model
40 includes arts venues, large events such as concerts, festivals, sports games, public parks and other
41 outdoor settings, public transport, and all other community settings. We do not consider scenarios
42 in which 100% of the population wear masks across these settings, due to the infeasibility of wearing
43 masks whilst eating or drinking. On efficacy, we note that although the body of evidence supporting
44 the effectiveness of masks for protecting against transmission between individuals is now
45 considerable, the size of the effect is difficult to determine, with estimates in the range of 20–80%
46 and varying depending on whether one or both parties wear masks (31,49,51), or whether spillover
47 behavioural changes on people's attention to other NPIs are captured (52). To capture the uncertainty

1 regarding the effectiveness of masks, we assume that masks will reduce the per-contact probability
2 of transmission by 30%, in line with estimates from (53), but also consider 15% and 45% in a sensitivity
3 analysis presented in the supplementary materials.

4
5 To model the efficacy of contact tracing, we use publicly-available data from NSW Health (42). We
6 note that, although the program reports the proportion of known contacts that were reached within
7 defined timeframes, we would ideally like to know the proportion of all contacts that were reached,
8 which will be lower than the reported values since it will also include contacts that the case did not
9 recall or disclose. Thus, although NSW Health’s published reports (42) indicate that 100% of contacts
10 are notified within 48 hours, we use slightly more conservative values, namely that 100% of household
11 contacts will be traced and notified on the same day that test results are communicated, and that
12 95% of school contacts and 90% of workplace contacts will be notified on the following day. We then
13 consider scenarios in which 0%, 25%, 50%, 75%, or 100% of all other contacts (which we refer to as
14 community contacts) can be traced within a week of a case notification. For each scenario, the time
15 to trace each contact is drawn from a scaled Poisson distribution with a mean of 1 day (Figure S1).
16 We run 100 simulations for each permutation of testing rates, mask usage and contact tracing
17 efficacy.

18
19 Finally, we assume that people who have been contact-traced will quarantine with 90% compliance
20 from their workplace, school, and community contacts. Whilst this assumption may be optimistic in
21 other global contexts, the lower case counts in NSW mean that contact tracers have far greater
22 capacity for rigorous ongoing follow-up of contacts, and breaches of isolation are escalated with local
23 authorities, as stipulated in national guideline documents (50).

Core scenarios:	
4x combinations of symptomatic testing	50%, 65%, 80% 90% tested over the course of their symptoms
Asymptomatic contact testing	Asymptomatic contact testing rate equal to symptomatic testing rate
5x combinations of contact tracing for community contacts	0%, 25%, 50%, 100% of contacts traced within 1 week (Poisson distribution with a mean of 1 day)
4x combinations of mask uptake	0%, 25%, 50%, 75%
Mask efficacy	30%
Sensitivity analyses:	
Sensitivity to asymptomatic contact testing rate assumption	We run the same 80 (4x5x4) scenarios described above, but with the asymptomatic contact testing rate half of the symptomatic testing rate
Sensitivity to the mask efficacy assumption	We run the same 80 (4x5x4) scenarios described above, but with individual mask efficacy assumed to be 15% or 45% instead of 30%

25 **Table 1:** overview of the scenarios analysed over Oct 1 – Dec 31, 2020.

1 Role of the funding source

2 The funders had no role in the design or execution of this study.

3


4 Patient and public involvement

5 This is a modelling study; no personal data was used so patient/public involvement was not required.

6

7 Results

8 According to our modelled estimates, 4,550 (95% projected interval: 3,700–6,660) people had acquired
9 COVID-19 in NSW by the end of September 2020 (excluding cases acquired overseas or interstate),
10 of which 39% (27–48%) had been diagnosed (Figure 2). The majority of undiagnosed infections
11 occurred during the March-April wave; since the beginning of June, we estimate that 65% of all
12 locally-acquired infections have been diagnosed, with the remaining 35% being primarily comprised
13 of asymptomatic infections (65%). We further estimate that 79,330 (60,100–129,020) people had been
14 required to self-isolate at some point by September 30, 2020 as a result of having potentially been in
15 contact with a confirmed case.



16
17 **Figure 2.** Calibration of the model to the NSW epidemic. Solid lines indicate the median model
18 projections over 100 model runs; shaded areas indicate 95% projected intervals over different
19 initialisations; blue diamonds indicate data on confirmed locally-acquired cases.


20

21 Estimates of daily new infections under each combination of testing, community contact tracing, and
22 mask usage are presented in Figure 3. A key finding highlighted by Figure 3 is how effective high
23 levels of testing are in maintaining epidemic control: all strategies in which there is at least some
24 contact tracing in place and testing rates are very high (90% of people with symptoms and 90% of
25 asymptomatic contacts of confirmed cases) lead to a robustly controlled epidemic, with a median of
26 ~180 infections in total estimated over October 1 – December 31 under high mask uptake scenarios,
27 or 260–1,200 without masks, depending on the efficacy of community contact tracing (Figure 3,
28 bottom row). However, holding mask uptake and contact tracing constant, we estimate that the
29 number of infections over October 1 – December 31, 2020, would be 2-3 times higher if the testing
30 rate was 80% instead of 90%, 8-12 times higher if the testing rate was 65%, or 30-50 times higher
31 with a 50% testing rate (Figure 3, third row).

32

33 A second key finding is that the lower the testing rate, the greater the impact of masks and contact
34 tracing (and vice versa). With medium-lower testing rates of 50% or 65%, the marginal impact of both
35 masks and contact tracing are considerable (Figure 3, top two rows). Under these scenarios, the most
36 robust strategies consist of a combination of masks and community contact tracing. Assuming a 50%
37 testing rate, a scenario with near-perfect tracing and no masks is approximately equivalent to a
38 scenario with no tracing and high mask uptake. However, without community contact tracing, a

1 reduction in mask usage from 75% to 50% would lead to a near-doubling in estimated infections (from
 2 7,900 to 17,000), whereas even moderate levels of community contact tracing (25% traceable within
 3 a week) would increase the robustness of the response to lower mask usage (with 6,400 infections
 4 estimated under 75% mask usage and 9,700 under 50% mask usage). Similarly, masks strengthen the
 5 resilience of the epidemic outcome to decreases in contact tracing efficacy: without masks, a
 6 reduction in community contact tracing from 75% to 50% would lead to a 70% increase in total
 7 infections, whereas it would have no impact if 75% of the population were wearing masks.
 8




9
 10 **Figure 3.** Trailing 14-day average of daily locally-acquired infections under different assumptions
 11 about the testing rate (rows), proportion of community contacts that can be traced within one week
 12 (columns), and mask uptake (line colours). Projections represent the median of 100 simulations. Text
 13 boxes in each panel displays the cumulative number of infections over October 1 – December 31,
 14 2020.

15
 16 Figure 4 captures the risks associated with different strategies, quantified in terms of the probability
 17 of the 14-day average of daily diagnoses exceeding a given number by December 31, 2020. With no
 18 community tracing and a 50% testing rate, the probability of exceeding 100 diagnoses/day by the end
 19 of 2020 was calculated to be 98% without masks, and to remain greater than 50% even under the
 20 most optimistic mask uptake scenario (Figure 4, top left panel). However, higher levels of community
 21 tracing reduce these risks considerably: even without any mask usage, the probability of exceeding
 22 100 cases/day is estimated to be half as high with perfect community tracing (Figure 4, top right
 23 panel) compared to the same scenario without community tracing (48% vs 98%), and the addition of
 24 mask usage in community settings is estimated to reduce this even further, to ~10%.

26 **Comparison with actual outcomes**

27 The text boxes on Figure 4 display the probability of observing the outcome that was seen in NSW,
 28 in which the two-week average of daily diagnoses exceeded 12 by December 31. We calculate that

1 this outcome would have been assigned a low probability (4-7%) under the best-case scenarios of
 2 extremely high testing (90%), near-perfect community contact tracing (75-100%), and high mask
 3 usage (50-75%), but that it would have been considered a far more likely outcome if any of these were
 4 assumed to be at lower levels.
 5




6
 7 **Figure 4.** Quantifying the likelihood of epidemic resurgence in New South Wales: the probability of
 8 the trailing 14-day average of locally-acquired cases exceeding a given number by December 31, 2020
 9 under different assumptions about the testing rate (rows), proportion of community contacts that
 10 can be traced within one week (columns), and mask uptake (line colours).
 11

12 Sensitivity analyses

13 If asymptomatic contacts only test at half the rate of people with symptoms, we estimate a more
 14 severe epidemic over the last quarter of 2020, but the key results regarding the roles of community
 15 contact tracing and masks do not change (Figure S2). Since our core analyses already assume high
 16 compliance with recommended self-isolation policies for known contacts of confirmed cases, the
 17 marginal benefit of high asymptomatic contact tracing is primarily to further bolster the efficacy of
 18 contact tracing, since it allows for the identification of chains of transmission even in the absence of
 19 symptoms. To highlight the role of testing asymptomatic contacts, in Figure 5 we highlight a
 20 particular set of scenarios corresponding to the most optimistic contact tracing assumptions, in
 21 which all community contacts are assumed to be traced within one week, with a mean time to trace
 22 of one day (the full set of scenarios are summarised in Figure S2). Within this set of scenarios, the
 23 total number of infections is estimated to be around 50% higher if asymptomatic contacts test at a
 24 lower rate than people with symptoms (averaged across all levels of mask usage).
 25

26 We also find that, if masks are assumed to be more or less effective, this would change the risks
 27 associated with diagnosing more than a given number of cases per day under the different mask
 28 scenarios, but would not qualitatively change the results (Figure S3-S4).



1
2 **Figure 5.** Estimated total infections over October 1 – December 31, 2020 under different assumptions
3 about testing rates and mask uptake, assuming all community contacts can be traced within a week
4 with a mean time to trace of 1 day. Projections represent the median of 100 simulations.
5

6 Discussion

7 In this work we presented estimates illustrating the extent to which masks, community testing, and
8 community contact tracing could reduce the spread of SARS-CoV-2 in New South Wales, Australia
9 in the absence of widespread vaccination. We found that the relative impact of masks is greatest
10 when testing and tracing rates are lower (and vice versa). With very high testing rates (90% of people
11 with symptoms, plus 90% of people with a known history of contact with a confirmed case), we
12 estimate that epidemic control would be possible even without widespread mask uptake, provided
13 that fast and effective contact tracing is in place. However, under any scenario where testing rates
14 are lower, we estimate that mask use can play an important role in reducing the potential for
15 epidemic resurgence.
16

17 Our findings have relevance for assessing the outbreak that occurred in the state towards the end of
18 December 2020, which saw an average of ~12 diagnoses/day over two weeks. According to our
19 estimates, this outcome would have been assigned a relatively low probability (4-7%) if it was
20 assumed that compliance with the state's testing guidelines, venue registration protocols, and non-
21 mandatory mask recommendations would be at very high levels, but that it would have been
22 evaluated as a far more likely outcome if compliance with any of these NPIs was assumed to be
23 lower. As a result of the December 2020 outbreak, the NSW government made masks mandatory in
24 all indoor community settings from January 4, 2021 (54), a decision which will likely increase the
25 robustness of the overall response to decreases in testing or contact tracing rates, but will still require
26 ongoing commitment to ensure these remain at high levels, at least until vaccination coverage is
27 sufficiently high. According to Australia's vaccination schedule, half the population may be
28 vaccinated by mid-2021, and NPIs will feature as part of the COVID-19 control strategy until well
29 beyond this point (55).
30

31 Of the interventions considered, this study suggests that maintaining high levels of symptomatic
32 testing, contact tracing, and testing of contacts are all important. We also quantified the importance
33 of testing asymptomatic contacts as a strategy that further strengthens the power of contact tracing
34 programs, since it allows the tracing and quarantining of their contacts in turn. Longer term,
35 'pandemic fatigue' may bring about challenges in maintaining high levels of testing, mask use, and
36 contact tracing (56). This study suggests that having high levels of any two of mask use, testing and
37 contact tracing can largely mitigate the need for the third; however practical challenges mean that
38 this is unlikely to occur, and a more multifaceted approach of aiming for high coverage of all three

1 and ending up with moderate coverage of all three may be an effective and more robust strategy.
2 Various studies have shown that the roll-out of vaccination plans is unlikely to mitigate the need for
3 NPIs such as these for some time (24,57–59), so these findings are likely to remain relevant for a
4 considerable portion of 2021.

5
6 Although various efforts have been made to synthesise the ever-expanding body of research
7 regarding the efficacy of different interventions (1–4,61), each country's epidemic has distinct
8 characteristics and there are very few standardised, globally-applicable guidelines on what
9 constitutes a best-practice public health strategy (62,63). As a result, jurisdictions have taken diverse
10 approaches in terms of which interventions to prioritise, from a list that includes physical distancing,
11 travel restrictions, wearing of masks or face coverings, isolation and/or testing of those with COVID-
12 like symptoms, isolation of those who test positive, and tracing the contacts of confirmed cases for
13 testing and/or quarantining. Although it may not always be articulated as such, numerous trade-offs
14 are being made between different policy options in an attempt to allow the highest degree of societal
15 activity commensurate with epidemic control. In New South Wales, mask use was encouraged in
16 particular settings since July 2020, but not mandated until January 2021; at the same time, there was
17 a strong focus on contact tracing. The results from this work suggest that the prioritisation of contact
18 tracing may mitigate the relative importance of masks to some extent, but that this relies on
19 continued high levels of community testing.

20
21 There are several limitations to this study. Firstly, the mathematical model that we use requires data
22 on various aspects of SARS-CoV-2 transmission and prevention that are still not known exactly,
23 including the effects of masks on preventing individual transmission and the proportion of infections
24 that are asymptomatic. Whilst we have used the best available data and sampled from appropriate
25 distributions where possible, this represents a source of uncertainty in all mathematical models of
26 COVID-19. Additional uncertainties are introduced by the evolution of new strains of SARS-CoV-2
27 with increased transmissibility and/or severity. Secondly, we have constructed sets of scenarios that
28 examine various combinations of parameters on mask uptake, contact tracing, testing of people with
29 symptoms, and asymptomatic contact testing, but there are many more parameters that determine
30 the dynamics of transmission, including the stringency of border control measures, people's
31 adherence to quarantine and isolation policies, and the effect of ongoing distancing policies. Changes
32 to any of these policies would affect the results presented here in ways that are not straightforward
33 to predict or extrapolate. Third, we have not considered any outbreak risk associated with newly
34 seeded cases in the community that may arise from international or interstate arrivals. Another
35 limitation is the relatively simplistic way that we have modelled mask usage, whereby we have not
36 included variation in (a) adherence to mask usage across different types of venue, or (b) individual
37 compliance. Assuming that mask-wearing reduces everyone's transmission risk by a certain
38 percentage disregards the individual behavioural changes that may adjust individual-level
39 transmission risk by varying amounts. Furthermore, it is very likely that mask usage would be higher
40 in certain settings (e.g., public transport) than others, especially restaurants which are suspected to
41 be important for transmission. Finally, the model used here does not contain a geospatial component,
42 and we have not considered heterogeneities in incidence, behaviour, or contact patterns across
43 different parts of the state in these analyses. This could be relevant for questions around mask
44 uptake, as uptake of masks is generally higher in more densely populated areas, so using a state-
45 wide average for the proportion of the population wearing masks may underestimate their impact,
46 especially since >90% of infections in NSW to date have occurred in the Sydney metropolitan area.

47

1 Conclusions

2 Our work suggests that testing, tracing and masks can be effective means of controlling transmission
3 in dynamic community settings, and higher compliance with one can offset lower compliance with
4 the other to some extent. However, pursuing a strategy that combines aggressive testing, high mask
5 usage, and effective contact tracing, alongside continued hygiene and distancing protocols, is likely
6 to be the most robust means of controlling community-based transmission of SARS-CoV-2.
7

8 List of supplementary materials

- 9 ● Figure S1: Assumed distribution of times required to find community contacts.
- 10 ● Table S1: Parameter values and effects of policies on transmission risk in New South Wales.
- 11 ● Figure S2: Sensitivity to asymptomatic testing assumption: the probability of the trailing 14-
12 day average of locally-acquired cases exceeding a given number by December 31, 2020 with
13 individual-level mask efficacy set to 15%, under different assumptions about the testing rate
14 (rows), proportion of community contacts that can be traced within one week (columns), and
15 mask uptake (line colours).
- 16 ● Figure S3: Sensitivity to mask efficacy assumption: the probability of the trailing 14-day
17 average of locally-acquired cases exceeding a given number by December 31, 2020 with
18 individual-level mask efficacy set to 15%, under different assumptions about the testing rate
19 (rows), proportion of community contacts that can be traced within one week (columns), and
20 mask uptake (line colours).
- 21 ● Figure S4: Sensitivity to mask efficacy assumption: the probability of the trailing 14-day
22 average of locally-acquired cases exceeding a given number by December 31, 2020 with
23 individual-level mask efficacy set to 45%, under different assumptions about the testing rate
24 (rows), proportion of community contacts that can be traced within one week (columns), and
25 mask uptake (line colours).

26
27

1 Declarations

2
3 **Competing interests statement:** The authors declare that they have no competing interests.

4
5 **Data sharing statement:** All code and data analysed during this study are available in
6 https://github.com/optimamodel/covid_nsw.

7
8 **Contributorship:** RMS wrote the manuscript and produced the results. The model parameters and
9 calibration were agreed upon by RMS, RA, NS, and RG. The scenarios and analyses were agreed upon
10 by NS, MH, and RMS. CK, RMS, RA, DM, and DJK led development of the model (along with
11 numerous other contributors listed below).

12
13 **Acknowledgements:** We thank all contributors to the Covasim model, including at the Institute for
14 Disease Modeling, including Brittany Hagedorn, Katherine Rosenfeld, Prashanth Selvaraj, Rafael
15 Nunez, Gregory Hart, Carrie Bennette, Marita Zimmermann, Assaf Oron, Dennis Chao, Michael
16 Famulare, Lauren George; at GitHub, including Michał Jastrzębski, Will Fitzgerald, Cory Gwin, Julian
17 Nadeau, Hamel Husain, Rasmus Wriedt Larsen, Aditya Sharad, and Oege de Moor; at Microsoft,
18 including William Chen, Scott Ayers, and Rolf Harms; and at the Burnet Institute, including Anna
19 Palmer, Dominic Delport, and Sherrie Kelly. This work benefited from a useful discussion with James
20 Wood, Deborah Cromer, and John Kaldor of the University of NSW, Australia.

23 References

- 24
25 1. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, et al. Estimating the
26 effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature*. 2020
27 Aug;584(7820):257–61.
- 28 2. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, et al. Ranking the
29 effectiveness of worldwide COVID-19 government interventions. *Nature Human Behaviour*.
30 2020 Dec;4(12):1303–12.
- 31 3. Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions |
32 *Science* [Internet]. [cited 2021 Jan 25]. Available from:
33 <https://science.sciencemag.org/content/369/6500/eabb9789>
- 34 4. Hsiang S, Allen D, Annan-Phan S, Bell K, Bolliger I, Chong T, et al. The effect of large-scale
35 anti-contagion policies on the COVID-19 pandemic. *Nature*. 2020 Aug;584(7820):262–7.
- 36 5. Han E, Tan MMJ, Turk E, Sridhar D, Leung GM, Shibuya K, et al. Lessons learnt from easing
37 COVID-19 restrictions: an analysis of countries and regions in Asia Pacific and Europe. *The*
38 *Lancet* [Internet]. 2020 Sep 24 [cited 2020 Sep 28];0(0). Available from:
39 [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)32007-9/abstract](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)32007-9/abstract)
- 40 6. Which countries are in lockdown? [Internet]. *The Independent*. 2020 [cited 2021 Jan 25].
41 Available from: [https://www.independent.co.uk/news/world/which-countries-currently-in-](https://www.independent.co.uk/news/world/which-countries-currently-in-lockdown-b1777806.html)
42 [lockdown-b1777806.html](https://www.independent.co.uk/news/world/which-countries-currently-in-lockdown-b1777806.html)
- 43 7. Nong VM, Le Thi Nguyen Q, Doan TT, Van Do T, Nguyen TQ, Dao CX, et al. The second wave
44 of COVID-19 in a tourist hotspot in Vietnam. *Journal of Travel Medicine* [Internet]. 2020 Sep
45 18 [cited 2020 Dec 16];(taaa174). Available from: <https://doi.org/10.1093/jtm/taaa174>
- 46 8. “Here we go again”: Auckland fears a long lockdown as coronavirus returns [Internet]. *the*
47 *Guardian*. 2020 [cited 2021 Jan 25]. Available from:
48 [http://www.theguardian.com/world/2020/aug/13/here-we-go-again-auckland-fears-a-long-](http://www.theguardian.com/world/2020/aug/13/here-we-go-again-auckland-fears-a-long-lockdown-as-coronavirus-returns)
49 [lockdown-as-coronavirus-returns](http://www.theguardian.com/world/2020/aug/13/here-we-go-again-auckland-fears-a-long-lockdown-as-coronavirus-returns)
- 50 9. Coronavirus: Victoria declares state of disaster after spike in cases. *BBC News* [Internet]. 2020

- 1 Aug 2 [cited 2021 Jan 25]; Available from: <https://www.bbc.com/news/world-australia-53627038>
- 2
- 3 10. Abeysuriya R, Delpont D, Stuart R, Sacks-Davis R, Kerr C, Mistry D, et al. Preventing a cluster
- 4 from becoming a new wave in settings with zero community COVID-19 cases. Pre-print.
- 5 11. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and
- 6 transmissibility of COVID-19. *Nature Medicine*. 2020 May;26(5):672–5.
- 7 12. Slifka MK, Gao L. Is presymptomatic spread a major contributor to COVID-19 transmission?
- 8 *Nature Medicine*. 2020 Aug 17;1–3.
- 9 13. He J, Guo Y, Mao R, Zhang J. Proportion of asymptomatic coronavirus disease 2019: A
- 10 systematic review and meta-analysis. *Journal of Medical Virology* [Internet]. [cited 2020 Sep
- 11 22];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jmv.26326>
- 12 14. Chin AWH, Chu JTS, Perera MRA, Hui KPY, Yen H-L, Chan MCW, et al. Stability of SARS-
- 13 CoV-2 in different environmental conditions. *The Lancet Microbe*. 2020 May 1;1(1):e10.
- 14 15. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of controlling
- 15 COVID-19 outbreaks by isolation of cases and contacts. *The Lancet Global Health*. 2020 Apr
- 16 1;8(4):e488–96.
- 17 16. Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease
- 18 outbreak controllable. *PNAS*. 2004 Apr 20;101(16):6146–51.
- 19 17. Keeling MJ, Hollingsworth TD, Read JM. Efficacy of contact tracing for the containment of
- 20 the 2019 novel coronavirus (COVID-19). *J Epidemiol Community Health*. 2020 Oct
- 21 1;74(10):861–6.
- 22 18. Iacobucci G. Covid-19: Is local contact tracing the answer? *BMJ* [Internet]. 2020 Aug 17 [cited
- 23 2020 Sep 22];370. Available from: <https://www.bmj.com/content/370/bmj.m3248>
- 24 19. Ferretti L, Ledda A, Wymant C, Zhao L, Ledda V, Dorner LA-, et al. The timing of COVID-19
- 25 transmission. medRxiv. 2020 Sep 16;2020.09.04.20188516.
- 26 20. Wang CJ, Ng CY, Brook RH. Response to COVID-19 in Taiwan: Big Data Analytics, New
- 27 Technology, and Proactive Testing. *JAMA*. 2020 Apr 14;323(14):1341–2.
- 28 21. Duong DM, Le VT, Ha BTT. Controlling the COVID-19 Pandemic in Vietnam: Lessons From a
- 29 Limited Resource Country: [Internet]. *Asia Pacific Journal of Public Health*. SAGE
- 30 PublicationsSage CA: Los Angeles, CA; 2020 [cited 2020 Sep 15]. Available from:
- 31 <https://journals.sagepub.com/doi/10.1177/1010539520927290>
- 32 22. You J. Lessons From South Korea’s Covid-19 Policy Response: The American Review of Public
- 33 Administration [Internet]. 2020 Jul 17 [cited 2020 Sep 22]; Available from:
- 34 <https://journals.sagepub.com/doi/10.1177/0275074020943708>
- 35 23. Panovska-Griffiths J, Kerr CC, Waites W, Stuart RM, Mistry D, Foster D, et al. The potential
- 36 contribution of face coverings to the control of SARS-CoV-2 transmission in schools and
- 37 broader society in the UK: a modelling study. medRxiv. 2020 Sep 29;2020.09.28.20202937.
- 38 24. Conlen M, Lu D, Glanz J. Why Vaccines Alone Will Not End the Pandemic. *The New York*
- 39 *Times* [Internet]. 2021 Jan 24 [cited 2021 Jan 25]; Available from:
- 40 <https://www.nytimes.com/interactive/2021/01/24/us/covid-vaccine-rollout.html>
- 41 25. Sun K, Wang W, Gao L, Wang Y, Luo K, Ren L, et al. Transmission heterogeneities, kinetics,
- 42 and controllability of SARS-CoV-2. *Science* [Internet]. 2021 Jan 15 [cited 2021 Jan
- 43 25];371(6526). Available from: <https://science.sciencemag.org/content/371/6526/eabe2424>
- 44 26. Kerr CC, Mistry D, Stuart RM, Rosenfeld K, Hart G, Nunez RC, et al. Controlling COVID-19
- 45 via test-trace-quarantine [Internet]. 2020 [cited 2020 Aug 19]. Available from:
- 46 <https://www.medrxiv.org/content/10.1101/2020.07.15.20154765v3>
- 47 27. Kretzschmar ME, Rozhnova G, Bootsma MCJ, Boven M van, Wiggert JHHM van de, Bonten
- 48 MJM. Impact of delays on effectiveness of contact tracing strategies for COVID-19: a
- 49 modelling study. *The Lancet Public Health*. 2020 Aug 1;5(8):e452–9.
- 50 28. Kucharski AJ, Klepac P, Conlan A, Kissler SM, Tang M, Fry H, et al. Effectiveness of isolation,
- 51 testing, contact tracing and physical distancing on reducing transmission of SARS-CoV-2 in
- 52 different settings [Internet]. *Epidemiology*; 2020 Apr [cited 2020 May 27]. Available from:
- 53 <http://medrxiv.org/lookup/doi/10.1101/2020.04.23.20077024>
- 54 29. Ferretti L, Wymant C, Kendall M, Zhao L, Nurtay A, Abeler-Dörner L, et al. Quantifying
- 55 SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*

- 1 [Internet]. 2020 May 8 [cited 2021 Jan 25];368(6491). Available from:
2 <https://science.sciencemag.org/content/368/6491/eabb6936>
- 3 30. Eikenberry SE, Mancuso M, Iboi E, Phan T, Eikenberry K, Kuang Y, et al. To mask or not to
4 mask: Modeling the potential for face mask use by the general public to curtail the COVID-19
5 pandemic. *Infectious Disease Modelling*. 2020 Jan 1;5:293–308.
- 6 31. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, et al. Physical distancing, face
7 masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and
8 COVID-19: a systematic review and meta-analysis. *The Lancet*. 2020 Jun 27;395(10242):1973–
9 87.
- 10 32. Catching A, Capponi S, Yeh MT, Bianco S, Andino R. Examining face-mask usage as an
11 effective strategy to control COVID-19 spread [Internet]. *Epidemiology*; 2020 Aug [cited 2020
12 Aug 20]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.08.12.20173047>
- 13 33. Javid B, Balaban NQ. Impact of Population Mask Wearing on Covid-19 Post Lockdown.
14 *Infectious Microbes & Diseases* [Internet]. 2020 Jun 5 [cited 2020 Aug 20]; Available from:
15 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7299124/>
- 16 34. Mitze T, Kosfeld R, Rode J, Wälde K. Face masks considerably reduce COVID-19 cases in
17 Germany. *PNAS* [Internet]. 2020 Dec 3 [cited 2020 Dec 4]; Available from:
18 <https://www.pnas.org/content/early/2020/12/02/2015954117>
- 19 35. Covid in Sydney: New restrictions announced as outbreak grows - BBC News [Internet].
20 [cited 2021 Jan 25]. Available from: <https://www.bbc.com/news/world-australia-55378180>
- 21 36. COVID-19 Community Mobility Reports [Internet]. [cited 2021 Jan 25]. Available from:
22 <https://www.google.com/covid19/mobility/>
- 23 37. Household Impacts of COVID-19 Survey, August 2020 | Australian Bureau of Statistics
24 [Internet]. 2020 [cited 2020 Sep 28]. Available from:
25 [https://www.abs.gov.au/statistics/people/people-and-communities/household-impacts-covid-
26 19-survey/latest-release](https://www.abs.gov.au/statistics/people/people-and-communities/household-impacts-covid-19-survey/latest-release)
- 27 38. Clun TR Kate Aubusson, Rachel. CCTV monitors how many are wearing masks on trains
28 [Internet]. *The Sydney Morning Herald*. 2020 [cited 2020 Sep 2]. Available from:
29 [https://www.smh.com.au/national/nsw/cctv-monitors-how-many-are-wearing-masks-on-
30 trains-20200813-p551h4.html](https://www.smh.com.au/national/nsw/cctv-monitors-how-many-are-wearing-masks-on-trains-20200813-p551h4.html)
- 31 39. Roser M, Ritchie H, Ortiz-Ospina E, Hasell J. Coronavirus Pandemic (COVID-19). *Our World
32 in Data* [Internet]. 2020 Mar 4 [cited 2021 Jan 25]; Available from:
33 <https://ourworldindata.org/coronavirus>
- 34 40. COVID-19 weekly surveillance reports - COVID-19 (Coronavirus) [Internet]. [cited 2020 Aug
35 28]. Available from: [https://www.health.nsw.gov.au/Infectious/covid-19/Pages/weekly-
36 reports.aspx](https://www.health.nsw.gov.au/Infectious/covid-19/Pages/weekly-reports.aspx)
- 37 41. NSW COVID-19 data | Data.NSW [Internet]. [cited 2020 Sep 28]. Available from:
38 <https://data.nsw.gov.au/nsw-covid-19-data>
- 39 42. COVID-19: Updated advice on testing, 24 September 2020 - COVID-19 (Coronavirus)
40 [Internet]. [cited 2020 Sep 28]. Available from:
41 <https://www.health.nsw.gov.au/Infectious/covid-19/Pages/case-definition.aspx>
- 42 43. Endo A, Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working
43 Group, Abbott S, Kucharski AJ, Funk S. Estimating the overdispersion in COVID-19
44 transmission using outbreak sizes outside China. *Wellcome Open Res*. 2020 Jul 10;5:67.
- 45 44. Althouse BM, Wenger EA, Miller JC, Scarpino SV, Allard A, Hébert-Dufresne L, et al.
46 Stochasticity and heterogeneity in the transmission dynamics of SARS-CoV-2.
47 *arXiv:200513689 [physics, q-bio]* [Internet]. 2020 May 27 [cited 2020 Aug 29]; Available from:
48 <http://arxiv.org/abs/2005.13689>
- 49 45. Kerr CC, Stuart RM, Mistry D, Abey Suriya RG, Hart G, Rosenfeld K, et al. Covasim: an agent-
50 based model of COVID-19 dynamics and interventions. 2020 [cited 2020 Aug 19]; Available
51 from: <https://www.medrxiv.org/content/10.1101/2020.05.10.20097469v1>
- 52 46. Scott N, Palmer A, Delpont D, Abey Suriya RG, Stuart RM, Kerr CC, et al. Modelling the
53 impact of reducing control measures on the COVID-19 pandemic in a low transmission
54 setting. 2020 [cited 2020 Aug 19]; Available from:
55 <https://www.medrxiv.org/content/10.1101/2020.06.11.20127027v1>

- 1 47. InstituteforDiseaseModeling/synthpops [Internet]. Institute for Disease Modeling; 2020 [cited
2 2020 Aug 21]. Available from: <https://github.com/InstituteforDiseaseModeling/synthpops>
- 3 48. ABC News. Recovering from coronavirus: Data shows how Australia is rebuilding as COVID-
4 19 restrictions ease. 2020 [cited 2020 Aug 20]; Available from:
5 [https://www.abc.net.au/news/2020-05-21/road-to-recovery-from-the-covid-19-coronavirus-](https://www.abc.net.au/news/2020-05-21/road-to-recovery-from-the-covid-19-coronavirus-pandemic/12229514?nw=0)
6 [pandemic/12229514?nw=0](https://www.abc.net.au/news/2020-05-21/road-to-recovery-from-the-covid-19-coronavirus-pandemic/12229514?nw=0)
- 7 49. Liang M, Gao L, Cheng C, Zhou Q, Uy JP, Heiner K, et al. Efficacy of face mask in preventing
8 respiratory virus transmission: A systematic review and meta-analysis. *Travel Med Infect Dis*.
9 2020 May 28;101751.
- 10 50. Health AGD of. Coronavirus (COVID-19) common operating picture [Internet]. Australian
11 Government Department of Health. Australian Government Department of Health; 2020
12 [cited 2020 Sep 24]. Available from:
13 [https://www.health.gov.au/resources/publications/coronavirus-covid-19-common-operating-](https://www.health.gov.au/resources/publications/coronavirus-covid-19-common-operating-picture)
14 [picture](https://www.health.gov.au/resources/publications/coronavirus-covid-19-common-operating-picture)
- 15 51. Leung NHL, Chu DKW, Shiu EYC, Chan K-H, McDevitt JJ, Hau BJP, et al. Respiratory virus
16 shedding in exhaled breath and efficacy of face masks. *Nature Medicine*. 2020 May;26(5):676-
17 80.
- 18 52. Doung-ngern P, Suphanchaimat R, Panjangampatthana A, Janekrongtham C, Ruampoom D,
19 Daochaeng N, et al. Case-Control Study of Use of Personal Protective Measures and Risk for
20 SARS-CoV 2 Infection, Thailand - Volume 26, Number 11—November 2020 - Emerging
21 Infectious Diseases journal - CDC. [cited 2020 Oct 21]; Available from:
22 https://wwwnc.cdc.gov/eid/article/26/11/20-3003_article
- 23 53. IHME. COVID-19: What's New for June 25, 2020.
- 24 54. NSW makes face masks mandatory as state records seven new Covid cases [Internet]. the
25 Guardian. 2021 [cited 2021 Jan 25]. Available from: [http://www.theguardian.com/australia-](http://www.theguardian.com/australia-news/2021/jan/02/nsw-makes-face-masks-mandatory-as-state-records-seven-new-covid-cases)
26 [news/2021/jan/02/nsw-makes-face-masks-mandatory-as-state-records-seven-new-covid-cases](http://www.theguardian.com/australia-news/2021/jan/02/nsw-makes-face-masks-mandatory-as-state-records-seven-new-covid-cases)
- 27 55. Aubusson KC Kate. Half of Australian population could get COVID vaccine by middle of the
28 year [Internet]. The Sydney Morning Herald. 2021 [cited 2021 Jan 25]. Available from:
29 [https://www.smh.com.au/politics/federal/half-of-australian-population-could-get-covid-](https://www.smh.com.au/politics/federal/half-of-australian-population-could-get-covid-vaccine-by-mid-year-20210107-p56sg8.html)
30 [vaccine-by-mid-year-20210107-p56sg8.html](https://www.smh.com.au/politics/federal/half-of-australian-population-could-get-covid-vaccine-by-mid-year-20210107-p56sg8.html)
- 31 56. WHO. Pandemic fatigue: Reinvigorating the public to prevent COVID-19 [Internet]. Available
32 from: [https://apps.who.int/iris/bitstream/handle/10665/335820/WHO-EURO-2020-1160-40906-](https://apps.who.int/iris/bitstream/handle/10665/335820/WHO-EURO-2020-1160-40906-55390-eng.pdf)
33 [55390-eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/335820/WHO-EURO-2020-1160-40906-55390-eng.pdf)
- 34 57. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission
35 dynamics of SARS-CoV-2 through the postpandemic period. *Science*. 2020 May
36 22;368(6493):860–8.
- 37 58. So AD, Woo J. Reserving coronavirus disease 2019 vaccines for global access: cross sectional
38 analysis. *BMJ*. 2020 Dec 15;371:m4750.
- 39 59. Microbe TL. COVID-19 vaccines: the pandemic will not end overnight. *The Lancet Microbe*.
40 2021 Jan 1;2(1):e1.
- 41 60. Robert A, Funk S, Kucharski AJ. The measles crisis in Europe—the need for a joined-up
42 approach. *The Lancet*. 2019 May 18;393(10185):2033.
- 43 61. Maier BF, Brockmann D. Effective containment explains subexponential growth in recent
44 confirmed COVID-19 cases in China. *Science*. 2020 May 15;368(6492):742–6.
- 45 62. Should countries aim for elimination in the covid-19 pandemic? | The BMJ [Internet]. [cited
46 2021 Jan 25]. Available from: <https://www.bmj.com/content/370/bmj.m3410>
- 47 63. Considerations for implementing and adjusting public health and social measures in the
48 context of COVID-19 [Internet]. [cited 2021 Jan 25]. Available from:
49 [https://www.who.int/publications-detail-redirect/considerations-in-adjusting-public-health-](https://www.who.int/publications-detail-redirect/considerations-in-adjusting-public-health-and-social-measures-in-the-context-of-covid-19-interim-guidance)
50 [and-social-measures-in-the-context-of-covid-19-interim-guidance](https://www.who.int/publications-detail-redirect/considerations-in-adjusting-public-health-and-social-measures-in-the-context-of-covid-19-interim-guidance)

1 Supplementary materials: The role of masks, testing and
2 contact tracing in preventing COVID-19 resurgences: a case
3 study from New South Wales, Australia

4
5 RM Stuart^{1,2†}, Romesh G. Abeysuriya², Cliff C. Kerr^{3,4}, Dina Mistry³, Daniel J. Klein³, Richard Gray⁵,
6 Margaret Hellard^{2,6,7,8,9}, Nick Scott^{2,6}

7
8 1. Department of Mathematical Sciences, University of Copenhagen, Copenhagen, Denmark

9 2. Disease Elimination Program, Burnet Institute, Melbourne, Victoria, Australia

10 3. Institute for Disease Modeling, Global Health Division, Bill & Melinda Gates Foundation, Seattle,
11 USA

12 4. School of Physics, University of Sydney, Sydney, New South Wales, Australia

13 5. The Kirby Institute, UNSW Sydney, Sydney, New South Wales, Australia

14 6. School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria,
15 Australia

16 7. University of Melbourne, Parkville, Victoria, Australia

17 8. Department of Infectious Diseases, The Alfred Hospital and Monash University, Melbourne,
18 Victoria, Australia

19 9. The School of Population and Global Health and the Peter Doherty Institute for Infection and
20 Immunity, Melbourne, Victoria, Australia

21 † Corresponding author. E-mail: robyn@math.ku.dk

22

23


24

25 **Key words:** COVID-19, coronavirus, SARS-CoV-2, modelling, masks, contact tracing, Australia

26

27

1



2
3
4
5


Figure S1. Assumed distribution of times required to find venue-based contacts.

Setting	Description of policy changes and their effects
Schools	School attendance rates in NSW had already dropped by 25% by 15 March 2020 (1), and on 23 March 2020 the NSW Premier advised that although schools remained open, parents were encouraged to keep their children at home for online learning (2). School attendance rates subsequently dropped to 5% of their pre-COVID levels (3). However, attendance quickly returned to pre-COVID levels shortly after schools reopened in mid-May (4). To model this, we removed 95% of contacts between school children and then restored them again as schools opened, but with the relative transmission risk set to 80% of its pre-March levels to account for additional safety measures in place for school activities (5).
Workplaces	According to survey data from the Australian Bureau of Statistics, almost half of working Australians were working from home in late April/early May (6), which is roughly consistent with Google movement data indicating that 40% fewer people were at work over that period compared to baseline. Workplace-based activities increased as COVID restrictions eased, but remained 15% lower over June-July compared to baseline. In the model, we removed 50% of workplace contacts and then restored them so that the workplace network was back to 85% of its pre-COVID size by the beginning of July (Figure 1). As with schools, we set the relative transmission risk set to 80% of its pre-March levels to account for the presence of NPIs.


Static community	We assume that almost no contacts occurred over these networks from March 23 to May 1 with the exception of the limited contacts arising from permitted single-person visits., These networks were gradually restored over the period from May 1 to July 9 as restrictions eased.
Dynamic community networks over May-July (negligible mask usage)	Within New South Wales, arts venues such as museums, galleries, theatres, and cinemas, large events such as concerts, festivals, sports games, and pubs/bars were all closed over the period from March 23 to May 15, after which the networks were gradually restored. Cafes and restaurants, public parks and other outdoor settings, public transport, and all other community settings including essential retail remained open in some capacity throughout the year but with decreased demand and operational restrictions to reduce the likelihood of transmission (e.g., takeaway service only, closure of playgrounds, capacity limits on transport, and physical distancing/hygiene).
Dynamic community networks over August (increased mask usage)	From August 3, 2020, the use of masks was mandated in Victoria, which led to a marked increase in mask usage across the country. Only 13% of Australians wore a mask at least once over the month of June, but 58% reported wearing one at least once over August 7–17, 2020 (99% of Victorian residents compared to 44% of residents of other states) (7). Within New South Wales, media sources reported that 30% of people were wearing face masks on public transport in central Sydney in mid-August (8). To reflect the gradual increase in mask uptake in the model, we adjust the relative transmission risk assuming that the proportion of adults who wore masks while at work and in dynamic community settings increased over August to reach 30% by the end of the month.
Testing, tracing, and isolation	<p>We assume that the daily testing probability for those with symptoms increased from 5% in April to 20% by the beginning of June. Assuming a symptomatic period of roughly 10 days, this implies that the proportion of symptomatic people who get tested increased from 40% to 90%. In addition to symptom-based testing, we also assume that 90% of people who are not symptomatic but have been told to quarantine as a result of having been in contact with a confirmed case will get tested. Our assumptions around test sensitivity coupled with our modelling of viral load kinetics are specified such that the probability of identifying a true positive increases and then decreases over the course of an infection, following a similar profile to that reported in the literature (9–10).</p> <p>To model the efficacy of contact tracing over the period from June 1 – September 30, 2020, we assume that all household contacts were traced and notified on the same day that test results were communicated, that 95% of school contacts and 90% of workplace contacts were notified on the following day, and that 50% of all other contacts (which we refer to as community contacts) were traced within a week of a case notification, with a mean time to trace of one day (Figure S1).</p> <p>We assume that confirmed cases will isolate with near-perfect effectiveness, meaning that the probability of them transmitting to school or workplace contacts is zero, and the probability of them transmitting to their household contacts is reduced by 80%. Similarly, we assume high levels of adherence to isolation policies</p>

	imposed on those who have been notified that they were in contact with a confirmed case, with a 90% reduction in their probability of transmitting to school or workplace contacts.
Quarantine, and border control measures	From early on in the epidemic, Australia imposed strict border control measures requiring all arrivals to quarantine for two weeks in a hotel. These policies are assumed to be maintained throughout the duration of the model simulations.


1 **Table S1.** Effects of policies on transmission risk in New South Wales
2
3



4 **Figure S2.** Sensitivity to asymptomatic testing assumption: the probability of the trailing 14-day
5 average of locally-acquired cases exceeding a given number by December 31, 2020 with individual-
6 level mask efficacy set to 15%, under different assumptions about the testing rate (rows), proportion
7 of community contacts that can be traced within one week (columns), and mask uptake (line colours).
8
9



1
2 **Figure S3.** Sensitivity to mask efficacy assumption: the probability of the trailing 14-day average of
3 locally-acquired cases exceeding a given number by December 31, 2020 with individual-level mask
4 efficacy set to 15%, under different assumptions about the testing rate (rows), proportion of
5 community contacts that can be traced within one week (columns), and mask uptake (line colours).
6



1 **Figure S4.** Sensitivity to mask efficacy assumption: the probability of the trailing 14-day average of
2 locally-acquired cases exceeding a given number by December 31, 2020 with individual-level mask
3 efficacy set to 45%, under different assumptions about the testing rate (rows), proportion of
4 community contacts that can be traced within one week (columns), and mask uptake (line colours).
5
6

7 Supplementary references

- 8 1. Baker J. Coronavirus: NSW school attendance falls 25 per cent as soap, toilet paper shortage
9 hit. The Sydney Morning Herald [Internet]. 2020 [cited 2020 Aug 20]; Available from:
10 [https://www.smh.com.au/national/school-attendance-falls-25-per-cent-as-soap-toilet-paper-](https://www.smh.com.au/national/school-attendance-falls-25-per-cent-as-soap-toilet-paper-shortage-hits-20200316-p54aok.html)
11 [shortage-hits-20200316-p54aok.html](https://www.smh.com.au/national/school-attendance-falls-25-per-cent-as-soap-toilet-paper-shortage-hits-20200316-p54aok.html)
- 12 2. National Centre for Immunisation Research and Surveillance,. COVID-19 in schools – the
13 experience in NSW. 2020.
- 14 3. Elsworthy E. Students are being sent back to school in NSW soon – here’s what you need to
15 know. ABC News [Internet]. 2020 Apr 21 [cited 2020 Aug 20]; Available from:
16 [https://www.abc.net.au/news/2020-04-21/coronavirus-return-for-nsw-schools-](https://www.abc.net.au/news/2020-04-21/coronavirus-return-for-nsw-schools-explained/12167826)
17 [explained/12167826](https://www.abc.net.au/news/2020-04-21/coronavirus-return-for-nsw-schools-explained/12167826)
- 18 4. Chrysanthos N. “There were bubbles and balloons”: High attendance on first day back at
19 school. The Sydney Morning Herald [Internet]. 2020 May 25 [cited 2020 Aug 20]; Available
20 from: [https://www.smh.com.au/national/nsw/there-were-bubbles-and-balloons-high-](https://www.smh.com.au/national/nsw/there-were-bubbles-and-balloons-high-attendance-on-first-day-back-at-school-20200525-p54w7o.html)
21 [attendance-on-first-day-back-at-school-20200525-p54w7o.html](https://www.smh.com.au/national/nsw/there-were-bubbles-and-balloons-high-attendance-on-first-day-back-at-school-20200525-p54w7o.html)
- 22 5. A guide to NSW school students for Term 3 [Internet]. [cited 2020 Aug 29]. Available from:
23 <https://education.nsw.gov.au/covid-19/advice-for-families.html>
- 24 6. Australian Bureau of Statistics. Media Release - Loneliness most common stressor during
25 COVID-19 (Media Release) [Internet]. c=AU; o=Commonwealth of Australia; ou=Australian
26 Bureau of Statistics; 2020 [cited 2020 Aug 20]. Available from:
27 [https://www.abs.gov.au/ausstats/abs%40.nsf/mediareleasesbyCatalogue/DB259787916733E4CA](https://www.abs.gov.au/ausstats/abs%40.nsf/mediareleasesbyCatalogue/DB259787916733E4CA25855B0003B21C?OpenDocument)
28 [25855B0003B21C?OpenDocument](https://www.abs.gov.au/ausstats/abs%40.nsf/mediareleasesbyCatalogue/DB259787916733E4CA25855B0003B21C?OpenDocument)
- 29 7. Statistics c=AU; o=Commonwealth of A ou=Australian B of. Media Release - COVID-19
30 anxiety in Victoria felt Australia-wide (Media Release) [Internet]. c=AU; o=Commonwealth of
31 Australia; ou=Australian Bureau of Statistics; 2020 [cited 2020 Sep 2]. Available from:
32 [https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4940.0Media%20Release1August](https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4940.0Media%20Release1August%202020?opendocument&tabname=Summary&prodno=4940.0&issue=August%202020&num=&view=)
33 [%202020?opendocument&tabname=Summary&prodno=4940.0&issue=August%202020&num=&](https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4940.0Media%20Release1August%202020?opendocument&tabname=Summary&prodno=4940.0&issue=August%202020&num=&view=)
34 [view=](https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4940.0Media%20Release1August%202020?opendocument&tabname=Summary&prodno=4940.0&issue=August%202020&num=&view=)
- 35 8. Clun TR Kate Aubusson, Rachel. CCTV monitors how many are wearing masks on trains
36 [Internet]. The Sydney Morning Herald. 2020 [cited 2020 Sep 2]. Available from:
37 [https://www.smh.com.au/national/nsw/cctv-monitors-how-many-are-wearing-masks-on-](https://www.smh.com.au/national/nsw/cctv-monitors-how-many-are-wearing-masks-on-trains-20200813-p55lh4.html)
38 [trains-20200813-p55lh4.html](https://www.smh.com.au/national/nsw/cctv-monitors-how-many-are-wearing-masks-on-trains-20200813-p55lh4.html)
- 39 9. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of controlling
40 COVID-19 outbreaks by isolation of cases and contacts. The Lancet Global Health. 2020 Apr
41 1;8(4):e488–96.
- 42 10. Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-Negative Rate of
43 Reverse Transcriptase Polymerase Chain Reaction–Based SARS-CoV-2 Tests by Time Since
44 Exposure. Ann Intern Med [Internet]. 2020 May 13 [cited 2021 Feb 1]; Available from:
45 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7240870/>
46