Predicting Olfactory Loss In Chronic Rhinosinusitis Using Machine Learning

Vijay R. Ramakrishnan, MD[1], Jaron Arbet, PhD[2], Jess C. Mace, MPH, CCRP[3], Zachary M. Soler, MD, MSc[4], Timothy L. Smith, MD, MPH[3]

Department of Otolaryngology-Head and Neck Surgery, University of Colorado, Aurora, CO.[1]

Department of Biostatistics, University of Colorado, Aurora, CO.[2]

Department of Otolaryngology-Head and Neck Surgery, Oregon Health Sciences University, Portland, OR.[3]

Department of Otolaryngology-Head and Neck Surgery, Medical University of South Carolina, Charleston, SC.[4]

Corresponding author:
Vijay R. Ramakrishnan, MD
Department of Otolaryngology, University of Colorado
12631 E 17th Ave, B205
Aurora, CO. 80045
Phone: 303-724-1965
Fax: 303-724-1961
E-mail: Vijay.Ramakrishnan@ucdenver.edu

Running Title: Machine Learning and CRS Olfactory Loss

Financial Disclosures: V.R.R., J.C.M., Z.M.S., and T.L.S. are supported by grants for this investigation from the National Institute on Deafness and Other Communication Disorders (NIDCD), one of the National Institutes of Health, Bethesda, MD., USA [R01 DC005805 (T.L.S.) and K23 DC014747 (V.R.R.)]. Public clinical trial registration (www.clinicaltrials.gov) ID# NCT01332136. JA is supported by NIH/NCATS Colorado CTSA Grant Number UL1 TR002535. Contents are the authors' sole responsibility and do not necessarily represent official NIH views.

Conflicts of Interest: None related to this study

38 **ABSTRACT**

39 **Objective**
40 Compare machine learning (ML) based predictive analytics methods compared to traditional
41 logistic regression in classification of olfactory dysfunction in chronic rhinosinusitis (CRS-OD),
42 and identify predictors within a large multi-institutional cohort of refractory CRS patients.
43
44 **Methods**
45 Adult CRS patients enrolled in a prospective, multi-institutional, observational cohort study were
46 assessed for baseline CRS-OD using a smell identification test (SIT) or brief SIT (bSIT). Four
47 different ML methods were compared to traditional logistic regression for classification of CRS
48 normosmics versus CRS-OD.
49
Results
Data were collected for 611 study participants who met inclusion criteria between April 2011
and July 2015. 34% of enrolled patients demonstrated olfactory loss on objective testing.
Differences between CRS normosmics and those with smell loss included objective disease
measures (CT and endoscopy scores), age, sex, prior surgeries, socioeconomic status, steroid use,
polyp presence, asthma, and aspirin sensitivity. Most ML methods outperformed traditional
logistic regression in terms of predictive ability. Top predictors include known factors reported
in the literature, as well as several socioeconomic factors.
58
Conclusion
Olfactory dysfunction is a variable phenomenon within a large multicenter cohort of refractory
CRS patients. ML methods outperform traditional logistic regression in classification of
normosmia versus smell loss in CRS, and are able to include numerous risk factors into
prediction models. These results carry implications for basic science and clinical research in
hyposmia secondary to sinonasal disease, the most common cause of persistent olfactory loss in
the general population.
66

INTRODUCTION

Olfactory loss is common, affecting up to 25% of the adult population, with chronic rhinosinusitis (CRS) being a major cause of persistent olfactory loss [1]. Smell loss is considered a hallmark of CRS, particularly in those with nasal polyps, and is one of the cardinal diagnostic symptoms for CRS [2-4]. Curiously, this symptom does not affect all CRS patients. Just as CRS is a heterogeneous disease [5], previous literature has estimated a 20-80% prevalence of olfactory loss in CRS cohorts, based on the sensitivity of olfactory assessment used and the proportion of enrolled subjects with nasal polyps (CRSwNP). A recent study using well-defined CRS diagnostic criteria and objective testing using Sniffin' Sticks assessment showed a 44% prevalence of smell loss in CRSsNP subjects and 58% in CRSwNP [6]. Risk factors associated with CRS-OD identified in this study included disease severity score on computed tomography (CT) the presence of comorbid conditions such as diabetes, allergy, and asthma. Adding a healthy control group for comparison, Schlosser et al. noted CRS-OD to affect CRSwNP patients more than CRSsNP, and identified age as an additional risk factor [7]. Given that most existing studies consist of single-institution cohorts, with subjects undergoing variable treatment regimens, and different assessments of smell function, a systematic review and meta-analysis was performed by Kohli et al [8] to clarify the presence of OD in CRS and identify consistent risk factors. The authors included 47 studies in the analysis, and identified age, CT score, and presence of nasal polyps as the three most consistent risk factors for CRS-OD in the published literature. Similarly, olfactory response to CRS treatment is highly variable, with resolution of smell disturbance after standard therapies occurring in only approximately 40% of refractory CRS patients [9].

Olfactory dysfunction in CRS is potentially complex and the interplay among possible mechanisms is poorly understood. The simple model of conductive loss due to impaired odorant transport to the olfactory cleft in CRS may explain a portion of the disease for some patients. Additional contributions of local and systemic inflammation likely contribute to a varying degree in some individuals, with underlying factors such as age and presence of comorbid diabetes influencing the capacity for olfactory sensory neuron regeneration. Given the complex and multifactorial nature of CRS-OD, compounded by inherent challenges in human cohort studies, it is no surprise that putative predictors and their relative importance have been inconsistently identified across studies. Beyond the risk of sampling bias found in most single institution studies, existing publications attempt to identify predictor variables for CRS-OD generally through univariate regression models. But, these analyses cannot account for all possible predictor variables or interactions among variables, and therefore, may offer limited conclusions. Additionally, attempting multivariable regressions that were not developed for predictive performance or assessed as such, cannot be relied upon [10].

Traditional statistical approaches can model how system variables relate to one another and generate corresponding metrics of statistical significance. However, novel data analytics approaches including machine learning (ML) methods have shown improved classification accuracy and can be utilized to predict future outcomes, rather than attempting to interpret individual components within a system. ML approaches are particularly useful when there is a role for including complex interactions that may otherwise be ignored or dismissed as noise when using traditional statistics [11], such as in the case of CRS-OD.

Understanding the heterogeneity of CRS-OD and grasping multi-dimensional risk factors for classification prediction is foundational for future clinical care and research. The objective of this study was to test different ML models used for predictive analytics against traditional regression

112 modeling for classification accuracy of OD in CRS patients. A secondary goal was to explore top
113 predictors in high-performing models to identify common predictor variables of high importance
114 that may help explain the heterogeneity of OD found among CRS patients.
115
MATERIALS and METHODS
Study Population
Adult patients (≥18 years old) were recruited into a prospective, multi-center,
observational cohort study developed to evaluate treatment outcomes of endoscopic sinus
surgery (ESS), for which multiple reports have been previously published [12-15]. All study
patients were diagnosed with medically refractory CRS as defined by both the American
Academy of Otolaryngology-Head and Neck Surgery 2007 guidelines and the European Position
Paper on Rhinosinusitis and Nasal Polyps 2012 [3,4]. The Institutional Review Board (IRB) at
each performance site governed all study protocols, informed consent documentation, and data
safety monitoring. Performance sites consisted of tertiary care sinus surgery centers located
within academic hospital systems in North America including: Oregon Health & Science
University (OHSU; Portland, OR; eIRB#7198), Stanford University (Palo Alto, CA; IRB#4947),
the Medical University of South Carolina (Charleston, SC; IRB#12409), and the University of
Calgary (Calgary, Alberta, Canada; IRB#E-24208), with central study coordination at OHSU.
At original study enrollment, participants were screened for demographics, social and
medical history including factors listed in **Table 1**. The diagnoses of asthma, allergy, and aspirin
sensitivity were made by presence of classic symptoms and self-report of prior testing. Subjects
with comorbid cystic fibrosis, ciliary dyskinesia, or autoimmune disease were not excluded from
final analysis. Participants were followed postoperatively through the standard of care for up to
18-months and asked to complete postoperative, study evaluations at regular 6-month intervals,
either during physician-directed appointments or follow-up study mailings utilizing the postal
service and self-addressed return envelopes.
138
Objective Measures of Disease Severity
Diagnostic measures of disease severity were collected as standard of care, and were also
used for investigational purposes. These well-established objective measures have been
recommended for clinical study [16], and include Lund-Kennedy nasal endoscopy scoring [17]
and high-resolution computed tomography (CT) radiographic imaging graded using the Lund-
Mackay scoring system [18]. Olfactory function was using the brief Smell Identification Test
(bSIT) in the years 2011-2013, and the Smell Identification Tests (SIT) in the years 2013-2015
(Sensonics Inc., Haddon Heights, NJ). Higher SIT total scores reflect a better sense of smell, and
categorical ratings of "smell loss" (anosmia + hyposmia) and "normosmia" were assigned based
on established scales [19-21].
149

Patient Reported Outcome Measures
Other patient reported outcome measures (PROMs) were documented but were not used
in the predictive analytics given the potential overlap between particular symptom ratings and
objective smell loss. Study PROMs included the 22-item SinoNasal Outcome Test (SNOT-22;
©2006, Washington University, St. Louis, MO) [22,23], the Rhinosinusitis Disability Index
(RSDI) [24], and the questionnaire on olfactory dysfunction (QOD-NS) [25].
157

Data Management and Statistical Analyses

Protected health information was previously removed and study data was safeguarded
using unique study identification number assignment for each participant. Study data was
securely transferred from a HIPAA compliant, relational database (Access, Microsoft Corp,
Redmond, WA.) to the University of Colorado Department of Otolaryngology-Head and Neck
Surgery password protected research server, according to specifications of a Data Use
Agreement between OHSU and the University of Colorado.
Study data were evaluated descriptively while continuous and ordinal variables were
evaluated for assumptions of normality. Comparing variables between normosmics and those
with smell loss, Fisher's exact test was used for categorical variables and Welch's two sample t-
test was used for continuous variables. Statistical comparisons were reported using a type-I error
probability at the 0.050 level of significance. All statistical and data analyses were completed
using the R software program [26].
171
Machine Learning Approaches to Classification
Several machine learning predictive analytics models were applied, in order to compare
classification accuracy to tradition *logistic regression*, which was performed here with
backwards step-wise AIC variable selection. *Random forests* was chosen as it allows for non-
linear and interaction effects; conditional inference trees were used with permutation-based
variable importance scores for unbiased variable selection [27]. Similar to traditional logistic
regression, least absolute shrinkage and selection operator (*LASSO*), assumes only linear and
additive affects (without interaction) where regression coefficients are shrunk toward zero to
perform variable selection. In this model, unimportant variables are given coefficients equal to
zero, effectively removing them from the model. Along these lines, multivariate adaptive
regression splines (*MARS*) was also applied in a similar fashion to traditional logistic regression,
but it uses stepwise methods for variable selection and allows for nonlinear and interaction
effects. Lastly, the *Support Vector Machine* approach was applied with a radial basis kernel.
To evaluate classification accuracy, repeated 10-fold cross validation (5 repeats) was
used to tune and evaluate each model area. To deal with the fact that the two classes are
unbalanced (smell loss vs normosmia), down sampling was used within the cross-validation
procedure, as implemented in the caret R package. Up sampling was also applied, but produced
equal classification accuracy in terms of AUC. The method of "surrogate splits" [28] was used to
handle missing values in Random Forest. The missForest R package [29] was used to impute
missing values for all other models.
To identify the important predictor variables included in each classification algorithm,
permutation-based variable importance scores were used to rank variables from most to least
important in Random Forests, while the absolute value of the standardized regression coefficient
was used for LASSO variable rankinds. The IML R package [30] was used to calculate variable
importance scores for SVM using a permutation method. All variable importance scores were
scaled from 0 to 100, where the most important variable for the model has a score of 100. Of
note, for a given categorical variable, RF and LASSO calculate separate variable importance
scores for each category, whereas the approach used for SVM only calculates an overall score.
200
RESULTS
Final Cohort Characteristics
A total of 611 study participants met all inclusion criteria and were prospectively enrolled
between April 2011 and July 2015. Demographics and comorbid disease characteristics,

examined by olfactory classification, are described in **Table 1**. The overall mean age for the final
cohort was 51.3 years (38.3, 61.7) years with slight female preponderance (53% female).
Approximately one-third of the cohort exhibited nasal polyps (35%).
Study participants were stratified by SIT category. Statistically significant differences
were observed between the normosmic and smell loss cohorts in terms of objective disease
measures (CT and endoscopy scores), age, sex, previous surgery, socioeconomic status, steroid
use, asthma/COPD, and aspirin sensitivity. Local conditions including nasal polyps, allergic
fungal rhinosinusitis, and deviated septum were also significant factors.
213
Smell loss vs Normosmia Classification Accuracy
A traditional logistic regression model for classification of baseline olfactory dysfunction
demonstrated on these criteria demonstrated acceptable accuracy in the context of CRS-OD
(AUC 0.707). Three of the four machine learning models outperformed the traditional statistical
approach (**Table 2**). The SVM approach was the most accurate, with a relatively good
classification accuracy (AUC 0.754) and much higher sensitivity than other the methods. LASSO
and Random Forest also outperformed logistic regression, whereas MARS performed worse than
all of the approaches.
222
Machine Learning Predictions: Top variables
ML algorithms are adept are recognizing patterns within complex data, but the lack of *a*
priori framework results in the inability to understand *why* the algorithm made its decisions. As
we see in the variable importance graphs (**Figs 1, 2, Suppl Fig 1**), the ML models each arrived at
somewhat different relative importances of a given predictor variable. SVM, the top predictor
model, included the most number of variables in its algorithm with nearly all inputs having
significant importance (**Fig 1**). Relative importance of predictor variables in Random Forests and
LASSO are shown in **Fig 2** (**and Suppl Fig 1**) and in these models, fewer variables were
important and the most weight was placed on the presence of nasal polyps and the nasal
endoscopy disease severity score.
Top predictors across the approaches contain many factors associated with olfactory
function in previous studies, including steroid use, prior surgery, nasal polyps, aspirin
exacerbated respiratory disease, age, sex, and smoking history. Inflammatory disease severity, as
measured by CT and nasal endoscopy scores, are not surprisingly very important predictors of
CRS-OD.
Interestingly, socioeconomic factors such as insurance status, household income, race,
were among top predictors across the three models displayed. Comorbid medical conditions
including asthma, diabetes, septal deviation, alcohol use, and sleep apnea, are also of moderate
importance. Notably, these factors can be influenced by unrelated medical care, suggesting that
other approaches to improve general health and well-being may be considered as part of the
holistic approach to CRS-OD management. Although the presence of nasal polyps, asthma, and
AERD, are significantly associated with CRS-OD here and in prior literature, the presence of
allergic rhinitis by history or by skin testing appeared to carry relatively minimal weight in the
CRS-OD classifications.
247
DISCUSSION
CRS is a heterogeneous process in terms of presentation, subtypes, natural history and
responses to treatment [2]. In terms of CRS-OD, current data from a multi-institutional cohort

studying olfactory loss in CRS with a sensitive olfactory test (Sniffin' Sticks) demonstrates that
nearly 70% of the CRS population has significant olfactory loss, with 20% exhibiting complete
anosmia [7]. The pathophysiology of CRS-OD is ascribed to both sensorineural and conductive
effects of local tissue inflammation, on top of the myriad factors that can affect olfaction in the
general population [1,31]. In previous study, factors associated with worse baseline olfactory
function in CRS patients included nasal polyposis, asthma, age, smoking, and eosinophilia [32].
Other works have shown CRS-OD response treatment in ~40% of patients, with risk factors such
as olfactory dysfunction severity, nasal polyps, female gender, high socio-economic status and
non-smoking associating with better quality of life results. In multivariate regression, only nasal
polyps and degree of baseline olfactory dysfunction maintained statistical significance,
highlighting the importance of sample size and analytical methods to glean useful information
from a prospectively collected human dataset [10].
Here, we utilized ML-based data science approaches to interrogate a large multicenter
CRS cohort to test whether novel predictive analytics approaches could outperform the gold-
standard traditional statistical method of logistic regression. ML is defined as the use of
algorithms to break down data, learn from it, and then make a determination or prediction about
some aspect. In doing so, ML provides a novel approach to uncover previously unrecognized
patterns among CRS-OD patients and thereby offers numerous advantages over regression
analyses, which have been traditionally employed in studies of CRS disease severity and
outcomes [33]. Whereas traditional statistical methods focus on modeling how system variables
relate to one another and what statistical inference (e.g. significance in p-values) can be made,
the goal of machine learning (ML) is not interpretation of individual components but prediction
of future outcomesTop predictorsIn this study, different supervised ML algorithms were used
map features of interest to the outcome or "label" of olfactory loss. We observed improved
prediction accuracy with most ML approaches, measured by AUC and sensitivity/specificity.
This advantage may be attributed to the ability of many ML models to handle high-dimensional
data in which more features exist than observations, whereas traditional statistical modeling with
a large number of potential features requires some form of dimension reduction or variable
selection. As a result, we believe that machine learning and artificial intelligence data analytics
are well-suited to prime research and eventually for precision medicine in olfactory disorders,
given the idiosyncrasies, nuances, and numerous possible predictor variables.
There are some limitations of the current study that should be addressed, the first is the
potential for sampling bias. Two major biases in healthcare ML are gender and race [34]. Details
of this cohort have been extensively published and include even distributions by age and sex.
Racial minorities and low socioeconomic status groups are underrepresented as a function of the
geography and practice referral patterns, the general lack of accessible and affordable healthcare
and racial disparities in the U.S. healthcare system [35]. A related concern is that models created
from our cohort data may "overfit" if new or unseen clinical data are applied. Certainly, these
approaches should be further validated but nonetheless demonstrate the important utility of novel
data science approaches to uncover unrecognized patterns and predictors from "noisy" human
data in a noisy disease process. There are likely potentially unrecognized but important data
inputs (i.e., molecular endotyping data) that are not yet clinically available. Serum markers, such
as peripheral eosinophilia and serum IgE, have been proposed for CRS in the past but have
generally been disappointing and do not appear to correlate with tissue levels [36,37] Some our
clinical data input fields may associate phenotypes with likely molecular endotypes, but such

overlap is not universal [5]. It is an exciting direction for the near future to include molecular
based biomarkers to the input data, and test additional accuracy gained.
298
CONCLUSION
Here we demonstrate the importance of novel predictive analytics approaches in study of
clinical olfactory disorders. Although founded in data science principles, and capable of
recognizing patterns within noisy data, it is challenging to understand *why* the models make
predictions with comparable accuracy despite use of some different predictor variable.
Nonetheless, consistently observed predictors include some weighted interest in SES and other
potentially modifiable conditions such as asthma/AERD. Such approaches may have value for
both clinical counseling and guidance for future basic science and translational research.
307

308 **ACKNOWLEDGMENTS**
309 The authors would like to thank Drs. Miranda Kroehl, PhD, John Rice, PhD, and Krithika
310 Suresh, PhD, for biostatistical expertise in machine learning and prediction models, as well as
311 data management considerations. We would also like to thank Drs. Peter Hwang, MD, and Luke
312 Rudmik, MD, for their contributions in original study enrollment and ongoing collaboration.

REFERENCES

1. Hummel T, Whitcroft KL, Andrews P et al. Position Paper on Olfactory Dysfunction. Rhinology 2016 31;56(1):1-30.

2. Orlandi RR, Kingdom TT, Hwang PH, et al. International Consensus Statement on Allergy and Rhinology: Rhinosinusitis. Int Forum Allergy Rhinol. 2016;6 Suppl 1:S22-209. Section X.B.

3. Fokkens WJ, Lund VJ, Mullol J, et al. European position paper on rhinosinusitis and nasal polyps 2012. Rhinol. Suppl 2012;23: 1-298.

4. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. Otolaryngol Head Neck Surg. 2015;152(2 Suppl):S1-39.

5. Akdis CA, Bachert C, Cingi C, et al. Endotypes and phenotypes of chronic rhinosinusitis: a PRACTALL document of the European Academy of Allergy and Clinical Immunology and the American Academy of Allergy, Asthma & Immunology. J Allergy Clin Immunol 2013;131(6):1479-90.

6. Soler ZM, Kohli P, Storck KA, Schlosser RJ. Olfactory Impairment in Chronic Rhinosinusitis Using Threshold, Discrimination, and Identification Scores. Chem Senses 2016;41(9):713-719.

7. Schlosser RJ, Smith TL, Mace JC, et al. Factors driving olfactory loss in patients with chronic rhinosinusitis: a case control study. Int Forum Allergy Rhinol 2020;10(1):7-14.

8. Kohli P, Naik AN, Harruff EE, et al. The prevalence of olfactory dysfunction in chronic rhinosinusitis. Laryngoscope 2017; 127(2):309-320.

9. DeConde AS, Mace JC, Alt JA, Schlosser RJ, Smith TL, Soler ZM. Comparative effectiveness of medical and surgical therapy on olfaction in chronic rhinosinusitis: a prospective, multi-institutional study. Int Forum Allergy Rhinol. 2014;4(9):725-33.

10. Katotomichelakis M, Simopolous E, Tripsianis G, et al. Predictors of quality of life outcomes in chronic rhinosinusitis after sinus surgery. Eur Arch Otorhinolaryngol 2014; 271(4):733-41.

11. Bzdok D, Altman N, Krzywinski M. Statistics versus machine learning. Nat Methods 2018;15(4):233-234.

12. Smith TL, Kern R, Palmer JN, et al. Medical therapy vs surgery for chronic rhinosinusitis: a prospective, multi-institutional study with 1-year follow-up. Int Forum Allergy Rhinol 2013;3(1):4-9.

13. DeConde AS, Mace JC, Alt JA, et al. Longitudinal improvement and stability of the SNOT-22 survey in the evaluation of surgical management for chronic rhinosinusitis. Int Forum Allergy Rhinol. 2015; 5(3):233-239.

14. Soler ZM, Smith TL, Alt JA, et al. Olfactory-specific quality of life outcomes after endoscopic sinus surgery. Int Forum Allergy Rhinol. 2016; 6(4): 407-413.

15. Ramakrishnan VR, Mace JC, Soler ZM, Smith TL. Examination of high-antibiotic users in a multi-institutional cohort of chronic rhinosinusitis patients. Int Forum Allergy Rhinol. 2017;7(4):343-51.

16. Hopkins C, Hettige R, Soni-Jaiswal A, et al. Chronic rhinosinusitis outcome measures
(CHROME), developing a core outcome set for trials of interventions in chronic rhinosinusitis.
Rhinology 2018;56(1):22-32.

17. Lund VJ and Kennedy DW. Quantification for staging sinusitis. The Staging and Therapy
Group. Ann Otol Rhinol Laryngol Suppl. 1995;167:17-21.

18. Lund VJ and Mackay IS. Staging in rhinosinusitis. Rhinology. 1992;31(4):183-4.

19. Doty RL. The Smell Identification Test Administration Manual. 3. Haddon Heights, NJ:
Sensonics, Inc; 1995.

20. Doty RL. The Brief Smell Identification Test Administration Manual. Haddon Heights, NJ:
Sensonics, Inc; 2001.

21. El Rassi E, Mace JC, Steele TO, et al. Sensitivity analysis and diagnostic accuracy of the
Brief Smell Identification Test in patients with rhinosinusitis. Int Forum Allergy Rhinol.
2016;6(3):287-92.

22. Piccirillo JF, Merritt MG, Jr, Richards ML. Psychometric and clinimetric validity of the 20-
Item Sino-Nasal Outcome Test (SNOT-20) Otolaryngol Head Neck Surg. 2002;126(1):41–47.

23. Hopkins C, Gillett S, Slack R, et al. Psychometric validity of the 22-item Sinonasal Outcome
Test. Clin Otolaryngol. 2009;34(5):447–454.

24. Senior BA, Glaze C, Benninger MS. Use of the rhinosinusitis disability index (RSDI) in
rhinologic disease. Am J Rhinol. 2001; 15(1):15-20

25. Simopoulos E, Katotomichelakis M, Gouveris H, Tripsianis G, Livaditis M, Danielides V.
Olfaction-associated quality of life in chronic rhinosinusitis: adaptation and validation of an
olfaction-specific questionnaire. Laryngoscope 2012;122:1450–1454.

26. R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing.

27. Strobl C, Boulesteix AL, Zeileis A, Hothorn T. Bias in random forest variable importance
measures: Illustrations, sources and a solution. BMC bioinformatics 2007;8(1):25.

28. Strobl C, Malley J, Tutz G. An introduction to recursive partitioning: Rationale, application,
and characteristics of classification and regression trees, bagging, and random forests.
Psychological methods 2009; 14:323.

29. Stekhoven DJ and Bühlmann, P. MissForest—non-parametric missing value imputation for
mixed-type data. Bioinformatics 2012;28(1):112-118.

30. Molnar C, Casalicchio G, Bischl B. iml: An R package for interpretable machine learning. J
Open Source Software 2018;3(26), 786.

31. Rombaux P, Huart C, Levie P, Cingi C, Hummel T. Olfaction in Chronic Rhinosinusitis.
Curr Allergy Asthma Rep 2016;16(5):41.

32. Litvack JR, Fong K, Mace J, et al. Predictors of olfactory dysfunction in patients with
chronic rhinosinusitis. Laryngoscope 2008;118:2225–30.

33. Michie D, Spiegelhalter D, Taylor C. Machine Learning, Neural, and Statistical
Classification. New York, NY: Ellis Horwood; 1994.

34. Rajkomar A, Dean J, Kohane I. Machine learning in medicine. N Engl J Med
2019;380(14):1347-1358.

35. 2018 National Healthcare Quality and Disparities Report. Content last reviewed April 2020.
Agency for Healthcare Research and Quality, Rockville, MD.
https://www.ahrq.gov/research/findings/nhqrdr/nhqdr18/index.html

36. Gitomer SA, Fountain CR, Kingdom TT, et al. Clinical Examination of Tissue Eosinophilia
in Patients with Chronic Rhinosinusitis and Nasal Polyposis. Otolaryngol Head Neck Surg
2016;155(1):173-8.

37. Settipane GA. Nasal polyps and immunoglobulin E (IgE). Allergy Asthma Proc
1996;17(5):269-73.

Table 1. Study population. Collected demographics and clinical metadata used as inputs for classification models are shown here for the overall cohort, and those with and without olfactory dysfunction. Data sorted from smallest to largest q-value (multiple testing adjusted p-value).

Characteristic	Overall, N = 611	N	Yes, N = 203[1]	No, N = 408[1]	p-value[2]	q-value[3]
CRS with POLYPS		611			**<0.001**	**<0.001**
No	399 (65%)		86 (42%)	313 (77%)		
Yes	212 (35%)		117 (58%)	95 (23%)		
TOTAL ENDOSCOPY SCORE	6.0 (4.0, 8.5)	607	8.0 (4.0, 11.0)	4.0 (2.0, 7.0)	**<0.001**	**<0.001**
TOTAL CT SCORE	12.0 (7.0, 16.0)	582	15.0 (10.0, 19.0)	11.0 (6.0, 14.0)	**<0.001**	**<0.001**
AERD		611			**<0.001**	**<0.001**
No	567 (93%)		173 (85%)	394 (97%)		
Yes	44 (7.2%)		30 (15%)	14 (3.4%)		
AGE	51.0 (38.3, 61.7)	611	56.6 (44.6, 65.4)	49.1 (36.9, 59.9)	**<0.001**	**<0.001**
PRIOR SURGERY		610			**<0.001**	**<0.001**
0	278 (46%)		66 (33%)	212 (52%)		
1	153 (25%)		56 (28%)	97 (24%)		
2	90 (15%)		34 (17%)	56 (14%)		
3	40 (6.6%)		24 (12%)	16 (3.9%)		
4+	49 (8.0%)		23 (11%)	26 (6.4%)		
INSURANCE		610			**<0.001**	**<0.001**
Employer provided	373 (61%)		112 (55%)	261 (64%)		
Medicaid	24 (3.9%)		4 (2.0%)	20 (4.9%)		
Medicare	129 (21%)		63 (31%)	66 (16%)		
None	6 (1.0%)		5 (2.5%)	1 (0.2%)		
Private	62 (10%)		13 (6.4%)	49 (12%)		
State Assisted	8 (1.3%)		4 (2.0%)	4 (1.0%)		
VA Benefits	3 (0.5%)		0 (0%)	3 (0.7%)		
VA Benefits / Tricare	5 (0.8%)		2 (1.0%)	3 (0.7%)		
ASTHMA		611			**<0.001**	**<0.001**
No	397 (65%)		110 (54%)	287 (70%)		
Yes	214 (35%)		93 (46%)	121 (30%)		

Characteristic	Overall, N = 611	N	Yes, N = 203[1]	No, N = 408[1]	p-value[2]	q-value[3]
SEPTAL DEVIATION		611			**<0.001**	**0.001**
No	408 (67%)		155 (76%)	253 (62%)		
Yes	203 (33%)		48 (24%)	155 (38%)		
RECURRENT ACUTE SINUSITIS		611			**<0.001**	**0.003**
No	571 (93%)		199 (98%)	372 (91%)		
Yes	40 (6.5%)		4 (2.0%)	36 (8.8%)		
STEROID USE		611			**0.003**	**0.009**
No	555 (91%)		174 (86%)	381 (93%)		
Yes	56 (9.2%)		29 (14%)	27 (6.6%)		
SEX		611			**0.013**	**0.037**
Female	324 (53%)		93 (46%)	231 (57%)		
Male	287 (47%)		110 (54%)	177 (43%)		
AFRS		611			**0.019**	**0.049**
No	593 (97%)		192 (95%)	401 (98%)		
Yes	18 (2.9%)		11 (5.4%)	7 (1.7%)		
COPD		611			**0.020**	**0.049**
No	579 (95%)		186 (92%)	393 (96%)		
Yes	32 (5.2%)		17 (8.4%)	15 (3.7%)		
HOUSEHOLD INCOME		595			**0.027**	0.064
0-25000	96 (16%)		34 (17%)	62 (16%)		
26000-50000	102 (17%)		47 (24%)	55 (14%)		
51000-75000	120 (20%)		37 (19%)	83 (21%)		
76000-100000	107 (18%)		28 (14%)	79 (20%)		
100000+	170 (29%)		51 (26%)	119 (30%)		
Inferior Turbinate Hypertrophy		611			0.091	0.200
No	520 (85%)		180 (89%)	340 (83%)		
Yes	91 (15%)		23 (11%)	68 (17%)		
CF or ciliary dysfunction		611			0.117	0.241
No	588 (96%)		199 (98%)	389 (95%)		
Yes	23 (3.8%)		4 (2.0%)	19 (4.7%)		
ALLERGY by HISTORY		611			0.143	0.277
No	480 (79%)		152 (75%)	328 (80%)		
Yes	131 (21%)		51 (25%)	80 (20%)		

Characteristic	Overall, N = 611	N	Yes, N = 203[1]	No, N = 408[1]	p-value[2]	q-value[3]
SITE		611			0.189	0.349
#1	222 (36%)		84 (41%)	138 (34%)		
#2	253 (41%)		77 (38%)	176 (43%)		
#3	136 (22%)		42 (21%)	94 (23%)		
OSA by HISTORY		611			0.215	0.376
No	584 (96%)		191 (94%)	393 (96%)		
Yes	27 (4.4%)		12 (5.9%)	15 (3.7%)		
EDUCATION (Yrs)	16.0 (13.0, 17.0)	608	15.0 (13.0, 16.0)	16.0 (13.0, 17.0)	0.235	0.392
RACE		610			0.247	0.393
African American	49 (8.0%)		18 (8.9%)	31 (7.6%)		
American Indian/Alaska Native	5 (0.8%)		2 (1.0%)	3 (0.7%)		
Asian	24 (3.9%)		10 (4.9%)	14 (3.4%)		
Native Hawaiian/Pacific Islander	1 (0.2%)		1 (0.5%)	0 (0%)		
Other	25 (4.1%)		12 (5.9%)	13 (3.2%)		
White	506 (83%)		160 (79%)	346 (85%)		
GERD		611			0.303	0.460
No	570 (93%)		186 (92%)	384 (94%)		
Yes	41 (6.7%)		17 (8.4%)	24 (5.9%)		
ETHNICITY		611			0.352	0.493
Hispanic / Latino	21 (3.4%)		9 (4.4%)	12 (2.9%)		
Non-Hispanic / Latino	590 (97%)		194 (96%)	396 (97%)		
ALCOHOL Use		608			0.344	0.493
No	313 (51%)		109 (54%)	204 (50%)		
Yes	295 (49%)		92 (46%)	203 (50%)		
ALLERGY_TESTING		611			0.382	0.510
No	362 (59%)		115 (57%)	247 (61%)		
Yes	249 (41%)		88 (43%)	161 (39%)		
DEPRESSION		611			0.403	0.510
No	518 (85%)		176 (87%)	342 (84%)		
Yes	93 (15%)		27 (13%)	66 (16%)		
FIBROMYALGIA		611			0.408	0.510
No	584 (96%)		192 (95%)	392 (96%)		

Characteristic	Overall, N = 611	N	Yes, N = 203[1]	No, N = 408[1]	p-value[2]	q-value[3]
Yes	27 (4.4%)		11 (5.4%)	16 (3.9%)		
OSA BY TESTING		611			0.450	0.543
No	557 (91%)		188 (93%)	369 (90%)		
Yes	54 (8.8%)		15 (7.4%)	39 (9.6%)		
ALCOHOL USE (#)	0.0 (0.0, 36.0)	608	0.0 (0.0, 24.0)	0.0 (0.0, 36.0)	0.473	0.552
DIABETES		609			0.536	0.605
No	558 (92%)		183 (91%)	375 (92%)		
Yes	51 (8.4%)		19 (9.4%)	32 (7.9%)		
AUTOIMMUNE DISEASE		610			0.589	0.645
No	573 (94%)		188 (93%)	385 (94%)		
Yes	37 (6.1%)		14 (6.9%)	23 (5.6%)		
IMMUNODEFICIENCY		611			0.626	0.664
No	592 (97%)		198 (98%)	394 (97%)		
Yes	19 (3.1%)		5 (2.5%)	14 (3.4%)		
SMOKER (current or former)		607			1.000	1.000
No	574 (95%)		190 (95%)	384 (95%)		
Yes	33 (5.4%)		11 (5.5%)	22 (5.4%)		

[1]Statistics presented: n (%); median (IQR)
[2]Statistical tests performed: Fisher's exact test; Wilcoxon rank-sum test
[3]False discovery rate correction for multiple testing

407 **Table 2**. **Comparing classification accuracy.** Three ML approaches outperform traditional
408 logistic regression. *AUC = area under receiver operating characteric curve; SVM-radial =*
409 *support vector machine with a radial basis kernel; Log Reg-Step = stepwise logistic regression.*
410
411
412

Model	AUC	Sensitivity	Specificity
SVM-Radial	0.754	0.726	0.665
Random Forest	0.744	0.649	0.696
LASSO	0.737	0.659	0.689
Log Reg-Step	0.707	0.635	0.676
MARS	0.682	0.604	0.667

413
414
415
416

416 **Figure 1. Variable importance display for most accurate classification model (Support**
417 **Vector Machine with a radial basis kernel).** Note the inclusion of many predictor variables –
418 32 are included with > 10% variable importance, suggesting significant interaction between
419 predictors.
420

421
422
423
424

Figure 2. **Top 10 Predictors and Variable importance for Random Forest and LASSO models.** *AERD = aspirin exacerbated respiratory disease; CF = cystic fibrosis.*

428 **Supplementary Figure 1**. **Complete list of predictors for Random Forest and LASSO**
429 **models.** Notably, the LASSO model incorporates significantly more features of interest than the
430 Random Forest algorithm (~25 features with >10% importance in the prediction algorithm).
431
432