# Chronic Obstructive Pulmonary Disease in adults exposed to fine particles from a coal mine fire

Shivonne Prasad[1,2], Caroline Gao[1,3], Brigitte Borg[1,2], Jonathan Broder[1], David Brown[1], Jillian Ikin[1], Annie Makar[2], Tom McCrabb[4], Ryan Hoy[1,2], Bruce Thompson[5], Michael J. Abramson[1]

[1]School of Public Health & Preventive Medicine, Monash University, Melbourne, AU.
[2]Respiratory Medicine, The Alfred Hospital, Melbourne, AU.
[3]Centre for Youth Mental Health (Orygen), University of Melbourne
[4]School of Population & Global Health, University of Melbourne, Parkville, AU
[5]School of Health Sciences, Swinburne University of Technology, Hawthorn, AU.

**Correspondence:**
Professor Michael J. Abramson
School of Public Health and Preventive Medicine
Monash University
Level 2, 553 St Kilda Road, Melbourne, Vic. 3004, Australia.
E-mail: michael.abramson@monash.edu
Tel: +61 3 9903 0573

## Abstract

In 2014 the Hazelwood open cut coal mine burned for six weeks, exposing nearby residents to fine particulate matter ($PM_{2.5}$). The long-term health consequences are being evaluated as part of the Hazelwood Health Study (HHS). These analyses explore the association between $PM_{2.5}$ and chronic obstructive pulmonary disease (COPD).

A sample of 346 exposed, and 173 unexposed, adults participated in the longitudinal Respiratory Stream of the HHS. Participants underwent spirometry and gas transfer measurements and answered validated respiratory questionnaires 3.5-4 years after the fire. Individual level mine fire-related $PM_{2.5}$ exposure was modelled. Multivariate linear regression and logistic models were fitted to analyse associations between mean and peak $PM_{2.5}$ exposure and clinical outcomes, stratified by smoking status.

A 10 μg/m$^3$ increase in mean $PM_{2.5}$ exposure was associated with a 69% (95%CI: 11% to 158%) increase in odds of spirometry consistent with COPD amongst non-smokers and increased odds of chest tightness (odds ratio [OR] 1.30, 95%CI 1.03 to 1.64) and chronic cough (OR 1.24, 95%CI 1.02 to 1.51) in the previous 12 months in all participants. For current smokers, increased mean $PM_{2.5}$ exposure was associated with higher odds of chronic cough in the preceding 12 months (OR 2.18, 95%CI 1.28 to 3.71) relative to other participants.

More than three years after a six-week period of coal fire $PM_{2.5}$ exposure, we identified a dose-response association between exposure and COPD in non-smokers. With climate change a likely contributor to increased risk of landscape fires, the findings will inform policy decisions during future sustained smoke events.

**Introduction**

The industrial era, largely fuelled by coal, has seen an increase in fires occurring within coal seams and at coal mines in association with human activity and changes in climate patterns [1]. Prominent examples of coal mine fires occurring in close proximity to human settlements have raised concerns about risks to human health around the world [2, 3]. Fires at the Jharia Coalfield in India and under the town of Centralia in Pennsylvania, illustrate instances where coal mine fires have led to mass relocation of residents arising from health and environmental effect concerns [1, 4].

The Centralia coal mine fire started in 1962 after local officials set fire to refuse in an abandoned coal pit. The fire then spread along an underground seam to tunnels beneath Centralia [5]. Emission of noxious gases and surface destabilization eventually made the town uninhabitable, with most residents evacuating between 1985 and 1991 [1, 5]. A health study comparing local residents with those of a nearby town suggested an increase in reported respiratory diseases in men and residents aged 40-79 years relative to their comparators [6]. Two years later a lower rate of diagnosed respiratory disease in relocated residents suggesting some attenuation of effects after exposure to the fire ceased [7].

Whilst coal mine fires near communities are widely reported, there remains limited peer reviewed literature on human health effects [1-3]. Analogous emissions from wildfires provide some insights into likely effects of exposure to smoke from coal mine fires [1]. Many of these studies assessed exposure to fine particulate matter with an aerodynamic diameter of less than 2.5 µm ($PM_{2.5}$)[8]. A review of the health impacts of wildfire smoke in 2016 identified consistent evidence of smoke related respiratory morbidity, particularly asthma and chronic obstructive pulmonary disease (COPD) [8]. Wildfire smoke exposure was associated with COPD related physician visits [9, 10], emergency department presentations [11, 12], hospitalisations [13, 14] as well as medication dispensing [9, 15]. However, almost all of the existing studies used administrative health records to evaluate short term smoke exposure effects. Clinical testing, including lung function measurements, to ascertain effects of bushfires or coal mine fire exposure was lacking.

In 2014, prolonged air pollution was generated from a fire at the Hazelwood open-cut brown coal mine in the Latrobe Valley in south-eastern Australia. Concerns were raised regarding the health of local residents, particularly in the nearby town of Morwell where smoke was visible for 6 weeks. The Hazelwood Health Study (HHS) was established to investigate potential longitudinal health outcomes in people who were exposed to smoke from the Hazelwood mine fire (www.hazelwoodhealthstudy.org.au). The HHS Hazelinks Stream which utilises administrative health datasets, has previously reported more COPD related emergency department presentations [16], medications dispensed for respiratory conditions [17] as well as visits to specialist respiratory services [18] during the mine fire period. The HHS Adult Survey, carried out 2.5 years after the mine fire, also identified higher risks of self-reported respiratory symptoms associated with individual-level mine fire $PM_{2.5}$ exposure [19]. The aim of this analysis was to evaluate clinical respiratory outcomes more than 3.5 years after the fire, particularly the risk of COPD and related respiratory symptoms, and their association with individual-level coalmine fire $PM_{2.5}$ exposure.

**Methods**

*Study design and setting*

This analysis examined cross-sectional data from the Hazelwood Health Study's Respiratory Stream, a longitudinal follow-up sub-study of the Adult Survey (see the methodology outlined elsewhere; [20]. Clinical testing was conducted in Morwell (exposed) between August and December 2017, and in Sale (unexposed) between January and March 2018. Study data were collected and managed using REDCap (Research Electronic Data Capture) [21] electronic data capture tools hosted at Monash University (Victoria, Australia).

*Participants*

The Respiratory Stream participants were drawn from the Adult Survey cohort which comprised residents of Morwell or Sale who were aged at least 18 years at the time of the mine fire [20]. Adult Survey cohort members were excluded from eligibility to participate in the Respiratory Stream if they had specified no further contact, had unknown age, sex or were aged over 90 years. Potential participants were also excluded if they had a contraindication to spirometry, such as recent surgery, myocardial infarction, pneumothorax, pulmonary embolus, open pulmonary tuberculosis or known aneurysms [22]. A target sample size of 339 from Morwell and 170 from Sale was derived based on the ability to detect a 5ml/year or greater $FEV_1$ decline in exposed compared with non-exposed residents using a two-sample t-test with a two-sided p-value of 0.05 and 80% power. A weighted random sample (to correct for lower response rate in some subgroups of participants, such as young people) of 1,346 Adult Survey cohort members were selected for invitation into the Respiratory Stream, with those who had reported an asthma attack or taking asthma medication oversampled (40%) for further evaluation in an asthmatic sample (see Figure 1). Invitation to participate was by mail, email and/or SMS, and recruitment continued until the target sample size was achieved.

*Participant characteristics*

For each participant, demographic characteristics, such as age, sex, ethnicity, employment status, educational background and occupational exposure (working in dusty or polluted environments, such as coal mines, farms or driving diesel trucks) were drawn from the previously completed Adult Survey. Physical characteristics, including height and weight, were measured by trained staff during clinical testing, and body mass index (BMI) was calculated [23]. Smoking history was taken and participants were classified as current smokers, ex-smokers (current non-smokers with >100 cigarettes in their lifetimes) or non-smokers (< 100 cigarettes in their lifetimes) [24].

*Exposure*

Due to the lack of ground-level air pollution monitoring from the start of the fire , mine fire-related $PM_{2.5}$ concentrations were retrospectively modelled by the Australian Commonwealth Scientific and Industrial Research Organisation (CSIRO) Oceans & Atmosphere using a chemical transport model that incorporated information on air monitoring, coal combustion and weather conditions [25]. The modelled 12-hourly concentrations were mapped to 12-hourly time-location diaries completed by participants as part of the Adult Survey, to estimate the level of exposure to mine-fire related $PM_{2.5}$ for each individual. Two exposure metrics were considered: mean and peak daily mine fire-related $PM_{2.5}$ exposure. Mean exposure was obtained by averaging the cumulative $PM_{2.5}$ exposure across all the locations that participants visited over the exposure period (9 February - 31 March 2014). Peak

exposure was obtained by assigning the highest 12-hourly $PM_{2.5}$ level from all locations, regardless of the time that the participant spent in that location.

*Outcomes*

Questionnaires

Self-reported respiratory health was assessed with standardised questionnaires, adapted from the European Community Respiratory Health Survey II (ECRHSII) [26] and administered by trained interviewers.

Respiratory function tests

Trained investigators conducted objective measurements of respiratory function. Standardisation was achieved through the use of Standard Operating Procedures (SOPs), validated equipment, temperature-controlled rooms and utilisation of the same staff and equipment at both clinic sites. Preparation instructions were provided to participants at the time of booking the clinic visit.

Respiratory function was measured using the EasyOne Pro™ LAB Respiratory Analysis System (nnd Medical Technologies AG, Zürich, Switzerland). Spirometry was performed according to American Thoracic Society/European Respiratory Society (ATS/ERS) guidelines [27]. Parameters measured included $FEV_1$, FVC and $FEV_1$/FVC. Z-scores for $FEV_1$, FVC and $FEV_1$/FVC were calculated using the Global Lung Initiative (GLI) 2012 spirometry reference equations [28]. This analysis focussed on spirometry measurements performed ten minutes after administration of 300 µg Salbutamol via pressurised Metered Dose Inhaler and spacer (pMDI)[29]

Carbon monoxide transfer factor ($T_Lco$) was measured using the single-breath testing technique according to ATS/ERS guidelines [30]. $T_Lco$ values were adjusted for haemoglobin (Hb) using standard equations from ATS/ERS guidelines [30]. Z-scores for $T_Lco$ were calculated using the GLI 2017 reference equations [31].

### *Definition of COPD and abnormal $T_Lco$*

COPD was defined by post bronchodilator $FEV_1$/FVC <5th percentile of predicted (lower limit of normal with $FEV_1$/FVC z-scores<-1.645) and abnormal $T_Lco$ was defined as haemoglobin adjusted $T_Lco$ <5th percentile of predicted ($T_Lco$ z-scores<-1.645). A similar definition for COPD was used in a recent population-based study on air pollution, lung function and COPD [32].

### *Statistical methods*

Descriptive statistics were used to summarise characteristics and clinical outcomes for the low, medium and high exposure groups in Morwell (tertiles of mean $PM_{2.5}$ exposure in Morwell) as well as the non-exposed Sale group. Crude statistical significance was assessed using Pearson chi-squared tests for categorical and t-tests for continuous measures. Multivariate linear regression models were used to analyse the association between mean and peak $PM_{2.5}$ exposure (controlling for key confounders) on continuous clinical outcomes. Logistic regression models were used for binary clinical outcomes. Oversampling of the asthmatic participants was corrected using the weighting method. Since smoking is the strongest risk factor for developing COPD, we tested whether the associations between $PM_{2.5}$ exposure and outcomes differed between smoking groups by including an

interaction between exposure and participants' smoking status (non-smoker, ex-smoker and current smoker) for all the outcome variables. Whether there was an overall interaction effect was tested using a Wald test of all interaction terms equal to zero. Sensitivity analyses were performed with unweighted and complete case models. Multiple imputation by chained equation (MICE) was used to deal with missing data. Statistical analyses were performed using Stata version 15 (Stata Corporation, College Station, Texas 2015).

*Ethical considerations*

The Monash University Human Research Ethics Committee (MUHREC) approved the Hazelwood Health Study: Cardiovascular and Respiratory Streams (approval number 1078). All participants provided written informed consent.

**Results**

As shown in Figure 1, 519 (39%) adults participated in the Respiratory Stream clinics; 346 from Morwell and 173 from Sale. Their characteristics were comparable across exposure groups (tertiles) and unexposed participants in terms of age, ethnicity, highest educational qualification, smoking status, years of smoking (for current and ex-smokers), occupational exposure and respiratory medication use in last 3 months, see Table 1. Distributions of exposure level for each group are provided in Supplementary Figure S1. However, there were higher proportions of male and obese participants and a lower proportion of employed participants in the higher exposure group compared with lower exposure group. Table 2 displays lung function and respiratory symptoms for participants in exposure groups. There were no substantial differences in post-bronchodilator spirometry or $T_{L}co$ between groups. However, respiratory symptoms in the last 12 months, including wheezing without an upper respiratory tract infection (URTI), chest tightness and chronic cough, were more prevalent in the higher exposure groups.

Adjusted associations between $PM_{2.5}$ exposure with respiratory symptoms and lung function are displayed in Table 3. A 10 µg/m$^3$ increase in mean $PM_{2.5}$ exposure was associated with a 30% (95%CI: 3% to 64%) increase in the odds of chest tightness in the last 12 months as well as a 24% (95%CI: 2% to 51%) increase in odds of chronic cough in the last 12 months. Figure 2 presents the results from logistic regression models for respiratory symptoms when interactions between exposure and smoking status were included. For ease of interpretation, we report the estimated OR associated with exposure (a 10 µg/m$^3$ increase in mean $PM_{2.5}$ or a 100 µg/m$^3$ increase in peak $PM_{2.5}$ exposure) in each smoking group separately. For chest tightness in the last 12 months, a dose-response relationship existed in non-smokers and current smokers, but not in ex-smokers. For chronic cough, there was strong evidence for a dose-response relationship in current smokers, but weaker evidence among non-smokers. There was also some evidence suggesting a dose response relationship among non-smokers for other respiratory symptoms including wheeze without URTI and chronic phlegm in the last 12 months (see Figure 2 and Table S1 in Supplementary Material).

Dose-response relationships between $PM_{2.5}$ exposure and post BD spirometry and $T_{L}co$ were not observed in the main model, however an interaction effect between exposure and smoking status was identified for spirometric COPD and post BD $FEV_1$/FVC z score but not $T_{L}co$ (see Figure 3 and Table S2 in the Supplementary Material). Among non-smokers, the interaction models estimated a 69% (95% CI: 11%, 158%) increase in the odds of spirometric COPD and a 0.13 (95% CI: 0.01, 0.26) reduction in post BD $FEV_1$/FVC z score per every 10 µg/m³ increase in mean exposure to $PM_{2.5}$, but no exposure effect was observed in smokers and ex-smokers (see Figure 3 and Table S2).

**Discussion**

This analysis found that COPD, as defined by a post bronchodilator $FEV_1/FVC$ ratio less than the lower limit of normal, was associated with both mean and peak $PM_{2.5}$ exposure from a coal mine fire in non-smoking individuals over three years after exposure. An association was also observed between mine-fire related $PM_{2.5}$ exposure and respiratory symptoms (chest tightness and chronic cough) consistent with airflow obstruction in all participants. Effect modification by smoking was apparent for the associations between respiratory symptoms and $PM_{2.5}$ exposure. This was clearest for chronic cough in the last 12 months, as current smokers had twice the odds of this symptom in association with increasing mean $PM_{2.5}$ exposure.

The exposure-response relationship between $PM_{2.5}$ and COPD in non-smokers in this study was consistent with findings from the European Studies of Cohorts for Air Pollution Effects (ESCAPE) which reported a positive association between proximity to traffic intensity and the incidence of GOLD-defined COPD in never smokers [33]. There are several possible explanations for this finding. Individuals more susceptible to the adverse effects of smoke may be less likely to continue cigarette smoking and may subsequently display a greater physiological response to smoke [34]. In addition, studies of lung cancer and cardiovascular disease provide evidence that relevant biological pathways for cardiovascular disease may be activated at low levels of $PM_{2.5}$ exposure and that further increasing exposure demonstrates a "saturation phenomenon". Diversely, a more linear relationship is found for lung cancer [34]. It is also possible that in smokers, ambient $PM_{2.5}$ exposure has an attenuated effect on spirometric COPD compared to non-smokers, although the biological pathways require further elucidation.

The lack of an effect on spirometry for smokers and ex-smokers exposed to $PM_{2.5}$ may equally relate to concurrent inhaled medication use, with these individuals potentially more likely to have used such medications compared to non-smokers. Lung function studies of asthmatics after a wildfire event have suggested a counter-intuitive preservation of lung function compared to non-asthmatics, with the possibility of a protective effect from the use of medications in this population [8]. Studies of COPD in non-smokers suggest a lower prevalence of respiratory symptoms, including chronic cough, reducing the likelihood of medication-seeking in this population [35].

From an epidemiological perspective, a quarter to a third of COPD occurs in non-smokers, with the proportion greater in low-middle compared to high income countries [35]. This may relate to an important role for indoor air pollution in the pathogenesis of COPD in non-smokers, with 45% of non-smokers reported to be exposed to biomass and 73% to coal smoke [35].

In general, there are limited data on the effects of particulate matter from coal mine fires on spirometry and our study provides a valuable addition to knowledge. Whilst comparison to other studies of coal mine fires is difficult for this reason, the similar emission profile between forest and coal mine fires allows comparison with the landscape fire literature [1]. Studies on the health impacts of wildfire smoke have found increasing evidence of a link with COPD using data on hospitalisations, physician visits and reliever medications [8]. However there were little data on effects on spirometry, with the exception of declines in peak expiratory flow among non-asthmatic children associated with wildfire exposure [8]. There was also little literature available on wild-fire smoke exposure lasting for up to six weeks, like the Hazelwood mine fire exposure. Wildfires often tend to be episodic and short in duration, and exposed populations from individual events are often small [8]. However, climate change may be contributing to increasing incidence of catastrophic wildfires globally, such as the

recent megafires in Australia which burned over a six month period and exposed more than 10 million people [36].

Declines in lung function (particularly $FEV_1$) after smoke exposure are commonly reported from studies of cross shift and cross season variations in wildfire fire fighters [37]. One study reported an association between levoglucosan (a woodsmoke marker) and decline in $FEV_1$ among wildland firefighters [38]. However, another study found no association between a decline in lung function and smoke components such as $PM_{3.5}$, acrolein, formaldehyde or carbon monoxide exposure [39]. The role of wildfire smoke causing sustained changes in lung function among adult fire fighters is unclear with contradictory results from existing studies [37] In some children with asthma, peak expiratory flow rates have been measured to assess response to wildfires [8]. One small study measured $FEV_1$ and peak flow prior to, and 1 month after, wildfires with high $PM_{2.5}$ levels and found stable lung function, but also elevated sputum eosinophils in two subjects tested during the fires, suggesting an acute inflammatory effect [40].

*Strengths & Limitations*

A major strength of our study was the measurement of lung function through spirometry, to enable an objective definition of COPD. To our knowledge, there were no comparable data available from other populations exposed to coal mine fires. Wildfire studies have largely focused on hospital and primary care presentations or medication dispensing [8]. This stream achieved the required sample size. The findings build upon previously reported respiratory health findings which have utilised administrative health datasets and self-reported symptoms.

Not relying on participants recalling their smoke exposure levels, an additional strength of our study was the estimation of individual exposures to $PM_{2.5}$ from a combination of detailed time-location diaries and spatially and temporally resolved modelled $PM_{2.5}$ concentrations using a chemical transport model. Whilst it is possible that some participants may have had difficulty recalling their precise locations and dates more than two years after the fire, considerable effort was made to manually review any detected inconsistencies in participant's time-location diaries.

However, a limitation of the study includes that it was essentially a cross-sectional analysis and it was not possible to imply causation between $PM_{2.5}$ exposure and COPD in non-smokers.

**Conclusions**

Our study found an association in non-smokers exposed to $PM_{2.5}$ from an open cut brown coal mine fire and increased risk of spirometry defined COPD. Conversely, we also found a pronounced association between respiratory symptoms and $PM_{2.5}$ exposure in smokers. These findings have important public health implications, as a better understanding of the exposure-response relationships across a range of $PM_{2.5}$ exposure levels and durations would help inform policy decisions including evidence-based exposure reduction strategies for communities. These may be particularly important in the event of future large-scale landscape fires similar to those recently in Australia and the United States. Our study also highlighted the importance of objective measurement of pulmonary impairment through spirometry, given that reliance on symptoms alone does not reliably identify disease, particularly in non-smokers. Long term followup of this exposed cohort may provide valuable insights into the incidence and prognosis of COPD in coal-mine fire $PM_{2.5}$ exposed non-smokers.

**Acknowledgments**

We gratefully acknowledge the contribution of all participating community members. We also thank Susan Denny, Kylie Sawyer, Shantelle Allgood and Kristina Thomas from the Monash University School of Rural Health who assisted with recruitment and data collection.

**Financial support**

The Hazelwood Health Study is funded by the Victorian Department of Health & Human Services. The paper presents the views of the authors and does not represent the views of the Department.

**References**

1.      Melody SM, Johnston FH. Coal mine fires and human health: What do we know? *Int J Coal Geol* 2015: 152: 1-14.

2.      Stracher G, Taylor T. Coal fires burning out of control around the world: Thermodynamic recipe for environmental catastrophe. *Int J Coal Geol* 2004: 59: 7-17.

3.      Song Z, Kuenzer C. Coal fires in China over the last decade: A comprehensive review. *Int J Coal Geol* 2014: 133: 72-99.

4.      Sinha PR. Mine fires in Indian coalfields. *Energy* 1986: 11(11): 1147-1154.

5.      Stracher G, Nolter M, Schroeder P, McCormack J, Blake D, Vice D. The great Centralia mine fire: a natural laboratory for the study of coalfires. *In:* FJ P, ed. Excursions in geology and history: field trips in the Middle Atlantic States: Geological Society of America field guide, 2006; pp. 33-45.

6.      Streufert S. Report on the Baseline Health Status for Centralia and Marion Heights, PA: Final Report. The Division, Pennsylvania State University, 1985.

7.      Streufert S, Baum A. Final Report on Health Status for Centralia and Marion Heights, Pa. Milton S. Hershey Medical Center, The Pennsylvania State University, 1987.

8.      Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical Review of Health Impacts of Wildfire Smoke Exposure. *Environmental Health Perspectives* 2016: 124(9): 1334-1343.

9.      Yao J, Eyamie J, Henderson SB. Evaluation of a spatially resolved forest fire smoke model for population-based epidemiologic exposure assessment. *J Expo Sci Environ Epidemiol* 2016: 26(3): 233-240.

10.     Henderson SB, Brauer M, Macnab YC, Kennedy SM. Three measures of forest fire smoke exposure and their associations with respiratory and cardiovascular health outcomes in a population-based cohort. *Environ Health Perspect* 2011: 119(9): 1266-1271.

11.     Reid CE, Jerrett M, Tager IB, Petersen ML, Mann JK, Balmes JR. Differential respiratory health effects from the 2008 northern California wildfires: A spatiotemporal approach. *Environ Res* 2016: 150: 227-235.

12.     Johnston FH, Purdie S, Jalaludin B, Martin KL, Henderson SB, Morgan GG. Air pollution events from forest fires and emergency department attendances in Sydney, Australia 1996–2007: a case-crossover analysis. *Environmental Health* 2014: 13(1): 105.

13.     Delfino RJ, Brummel S, Wu J, Stern H, Ostro B, Lipsett M, Winer A, Street DH, Zhang L, Tjoa T, Gillen DL. The relationship of respiratory and cardiovascular hospital admissions to the southern California wildfires of 2003. *Occup Environ Medi* 2009: 66(3): 189-197.

14.     Martin KL, Hanigan IC, Morgan GG, Henderson SB, Johnston FH. Air pollution from bushfires and their association with hospital admissions in Sydney, Newcastle and Wollongong, Australia 1994-2007. *Australian and New Zealand journal of public health* 2013: 37(3): 238-243.

15.     Elliott CT, Henderson SB, Wan V. Time series analysis of fine particulate matter and asthma reliever dispensations in populations affected by forest fires. *Environmental Health* 2013: 12(1): 11.

16.	Guo Y, Gao CX, Dennekamp M, Dimitriadis C, Straney L, Ikin J, Abramson MJ. The association of coal mine fire smoke with hospital emergency presentations and admissions: Time series analysis of Hazelwood Health Study. *Chemosphere* 2020: 253: 126667.

17.	Johnson AL, Dipnall JF, Dennekamp M, Williamson GJ, Gao CX, Carroll MTC, Dimitriadis C, Ikin JF, Johnston FH, McFarlane AC, Sim MR, Stub DA, Abramson MJ, Guo Y. Fine particulate matter exposure and medication dispensing during and after a coal mine fire: A time series analysis from the Hazelwood Health Study. *Environmental Pollution* 2019: 246: 1027-1035.

18.	Johnson AL, Gao CX, Dennekamp M, Williamson GJ, Carroll MTC, Dimitriadis C, Dipnall JF, Ikin JF, Johnston FH, McFarlane AC, Sim MR, Stub DA, Abramson MJ, Guo Y. Coal-mine fire-related fine particulate matter and medical-service utilization in Australia: a time-series analysis from the Hazelwood Health Study. *International Journal of Epidemiology* 2020: 49(1): 80-93.

19.	Johnson AL, Gao CX, Dennekamp M, Williamson GJ, Brown D, Carroll MTC, Ikin JF, Del Monaco A, Abramson MJ, Guo Y. Associations between Respiratory Health Outcomes and Coal Mine Fire $PM_{2.5}$ Smoke Exposure: A Cross-Sectional Study. *International Journal of Environmental Research and Public Health* 2019: 16(21): 4262.

20.	Ikin JF, Carroll M, Walker J, Borg B, Brown D, Cope M, Del Monaco A, Dennekamp M, Dimitriadis C, Gao CX, guo Y, Johnston FH, Liew D, Maybery D, Thompson BR, Sim MR, Abramson MJ. Cohort Profile: The Hazelwood Health Study Adult Cohort. *International Journal of Epidemiology* 2020: dyaa083.

21.	Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009: 42(2): 377-381.

22.	Cooper BG. An update on contraindications for lung function testing. *Thorax* 2011: 66(8): 714-723.

23.	WHO Consultation on Obesity. Obesity : preventing and managing the global epidemic : report of a WHO Consultation on Obesity, Geneva, 3-5 June 1997. World Health Organization, Geneva, 1998.

24.	World Health Organization. Guidelines for controlling and monitoring the tobacco epidemic. 1998.

25.	Luhar AK, Emmerson KM, Reisen F, Williamson GJ, Cope ME. Modelling smoke distribution in the vicinity of a large and prolonged fire from an open-cut coal mine. *Atmos Environ* 2020: 229: 117471.

26.	European Community Respiratory Health Survey Steering Committee. The European Community Respiratory Health Survey II. *Eur Respir J* 2002: 20(5): 1071-1079.

27.	Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CPM, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. Standardisation of spirometry. *European Respiratory Journal* 2005: 26(2): 319-338.

28.	Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, Enright PL, Hankinson JL, Ip MSM, Zheng J, Stocks J. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *European Respiratory Journal* 2012: 40(6): 1324.

29.	Borg BM, Reid DW, Walters EH, Johns DP. Bronchodilator reversibility testing: laboratory practices in Australia and New Zealand. *The Medical journal of Australia* 2004: 180(12): 610-613.

30.	Graham BL, Brusasco V, Burgos F, Cooper BG, Jensen R, Kendrick A, MacIntyre NR, Thompson BR, Wanger J. 2017 ERS/ATS standards for single-breath carbon monoxide uptake in the lung. *Eur Respir J* 2017: 49(1).

31.     Stanojevic S, Graham BL, Cooper BG, Thompson BR, Carter KW, Francis RW, Hall GL. Official ERS technical standards: Global Lung Function Initiative reference values for the carbon monoxide transfer factor for Caucasians. *Eur Respir J* 2017: 50(3).

32.     Doiron D, de Hoogh K, Probst-Hensch N, Fortier I, Cai Y, De Matteis S, Hansell AL. Air pollution, lung function and COPD: results from the population-based UK Biobank study. *Eur Respir J* 2019: 54(1).

33.     Schikowski T, Adam M, Marcon A, Cai Y, Vierkötter A, Carsin AE, Jacquemin B, Al Kanani Z, Beelen R, Birk M, Bridevaux P-O, Brunekeef B, Burney P, Cirach M, Cyrys J, de Hoogh K, de Marco R, de Nazelle A, Declercq C, Forsberg B, Hardy R, Heinrich J, Hoek G, Jarvis D, Keidel D, Kuh D, Kuhlbusch T, Migliore E, Mosler G, Nieuwenhuijsen MJ, Phuleria H, Rochat T, Schindler C, Villani S, Tsai M-Y, Zemp E, Hansell A, Kauffmann F, Sunyer J, Probst-Hensch N, Krämer U, Künzli N. Association of ambient air pollution with the prevalence and incidence of COPD. *European Respiratory Journal* 2014: 44(3): 614-626.

34.     Pope CA, Burnett RT, Turner MC, Cohen A, Krewski D, Jerrett M, Gapstur SM, Thun MJ. Lung cancer and cardiovascular disease mortality associated with ambient air pollution and cigarette smoke: shape of the exposure–response relationships. *Environmental Health Perspectives* 2011: 119(11): 1616-1621.

35.     Zeng G, Sun B, Zhong N. Non□smoking□related chronic obstructive pulmonary disease: A neglected entity? *Respirology* 2012: 17(6): 908-912.

36.     Walter, CM, Schneider□Futschik, EK, Knibbs, LD, Irving, LB. Health impacts of bushfire smoke exposure in Australia. *Respirology*. 2020; 25: 495– 501. https://doi.org/10.1111/resp.13798

37.     Groot E, Caturay A, Khan Y, Copes R. A systematic review of the health impacts of occupational exposure to wildland fires. *Int J Occup Med Environ Health* 2019: 32(2): 121-140.

38.     Gaughan DM, Piacitelli CA, Chen BT, Law BF, Virji MA, Edwards NT, Enright PL, Schwegler-Berry DE, Leonard SS, Wagner GR, Kobzik L, Kales SN, Hughes MD, Christiani DC, Siegel PD, Cox-Ganser JM, Hoover MD. Exposures and Cross-shift Lung Function Declines in Wildland Firefighters. *J Occup Environ Hyg* 2014: 11(9): 591-603.

39.     Slaughter J, Koenig J, Reinhardt T. Association between lung function and exposure to smoke among firefighters at prescribed burns. *J Occup Environ Hyg* 2004: 1(1): 45-49.

40.     Vora C, Renvall M, Chao P, Ferguson P, Ramsdell J. 2007 San Diego Wildfires and Asthmatics. *J Asthma* 2011: 48(1): 75-78.

Table 1. Participant characteristics by exposure groups (tertiles of mean $PM_{2.5}$ exposure in Morwell vs. Sale)

| Characteristics | Sale (no exposure) N=173 | Morwell low exposure N=109 | Morwell medium exposure N=113 | Morwell high exposure N=124 | p-value |
|---|---|---|---|---|---|
| **Age group, n (weighted %)** | | | | | |
| 18-44 years | 44 (22%) | 36 (26%) | 36 (30%) | 35 (25%) | 0.74 |
| 45-64 | 74 (42%) | 43 (44%) | 43 (37%) | 50 (37%) | |
| 65+ | 55 (36%) | 30 (29%) | 34 (34%) | 39 (38%) | |
| **Age, weighted mean (SD)** | 57.3 (20.0) | 54.7 (14.3) | 54.5 (15.3) | 56.7 (14.7) | 0.50 |
| **Male, n (weighted %)** | 62 (36%) | 43 (46%) | 46 (43%) | 62 (56%) | 0.023 |
| **Caucasian / White, n (weighted %)** | 171 (99%) | 108 (99%) | 112 (100%) | 123 (99%) | 0.92 |
| **Employment status, n (weighted %)** | | | | | |
| Employed | 89 (47%) | 44 (38%) | 46 (40%) | 50 (35%) | 0.021 |
| Retired | 57 (38%) | 32 (32%) | 33 (30%) | 37 (37%) | |
| Unable to work | 10 (6%) | 15 (16%) | 8 (5%) | 15 (12%) | |
| Other (unemployed, studying, home duties and other) | 17 (9%) | 18 (13%) | 26 (24%) | 22 (16%) | |
| **Highest educational qualification, n (weighted %)** | | | | | |
| Secondary up to year 10 | 34 (17%) | 28 (21%) | 29 (22%) | 28 (19%) | 0.79 |
| Secondary year 11-12 | 31 (19%) | 24 (23%) | 27 (24%) | 21 (17%) | |
| Certificate (trade/ apprenticeship/ technical) | 73 (42%) | 34 (37%) | 38 (37%) | 53 (48%) | |
| University or other Tertiary Institute degree | 34 (22%) | 22 (19%) | 16 (17%) | 21 (16%) | |
| **BMI, n (weighted %)** | | | | | |
| Underweight/Normal(BMI<25 kg/m$^2$) | 40 (24%) | 23 (20%) | 21 (18%) | 15 (11%) | 0.06 |
| Overweight (25≤BMI<30 kg/m$^2$) | 66 (38%) | 35 (33%) | 31 (29%) | 35 (30%) | |
| Obese (BMI≥30 kg/m$^2$) | 67 (38%) | 51 (47%) | 61 (53%) | 74 (59%) | |
| **Smoking status, n (weighted %)** | | | | | |
| Non-smoker | 82 (49%) | 58 (52%) | 60 (54%) | 49 (36%) | 0.10 |
| Ex-smoker | 66 (39%) | 35 (33%) | 34 (33%) | 51 (47%) | |
| Current smoker | 25 (12%) | 16 (15%) | 19 (13%) | 24 (17%) | |
| **Years of smoking among current and ex-smokers (N=270), weighted mean (SD)** | 23.6 (19.2) | 22.0 (11.0) | 20.5 (13.1) | 20.2 (12.3) | 0.54 |
| **Occupational exposure, n (weighted %)** | 64 (37%) | 43 (44%) | 44 (39%) | 54 (46%) | 0.47 |
| **Use any SAMA/SABA\* in the last 3 months, n (weighted %)** | 64 (31%) | 51 (33%) | 53 (36%) | 60 (36%) | 0.82 |
| **LAMA/LABA and/or ICS† use in the last 3 months, n (weighted %)** | 9 (4%) | 7 (3%) | 9 (6%) | 8 (6%) | 0.87 |

\* SAMA=short-acting muscarinic agonist, SABA=short-acting beta-agonist

† LAMA=long-acting muscarinic agonist, LABA=long-acting beta-agonist, ICS=inhaled corticosteroids

Table 2. Lung function and respiratory symptoms by exposure groups (tertiles of mean $PM_{2.5}$ exposure in Morwell vs. Sale)

| Outcome variables | Sale (no exposure) N=173 | Morwell low exposure N=109 | Morwell medium exposure N=113 | Morwell high exposure N=124 | p-value |
|---|---|---|---|---|---|
| | n (weighted%) | n (weighted%) | n (weighted%) | n (weighted%) | |
| Spirometric COPD | 24 (14%) | 9 (6%) | 14 (9%) | 12 (9%) | 0.27 |
| Abnormal $T_L$co | 18 (11%) | 9 (7%) | 15 (10%) | 14 (12%) | 0.50 |
| Spirometric COPD & abnormal $T_L$co | 7 (4%) | 2 (1%) | 6 (4%) | 6 (5%) | 0.27 |
| | Weighted mean (SD) | Weighted mean (SD) | Weighted mean (SD) | Weighted mean (SD) | |
| **Spirometry - post BD** | | | | | |
| $FEV_1$, Litres | 2.9 (1.1) | 3.0 (0.8) | 2.9 (0.8) | 2.9 (0.7) | 0.86 |
| $FEV_1$ z score | 0.0 (1.5) | 0.0 (1.0) | -0.3 (1.1) | -0.2 (1.0) | 0.22 |
| $FEV_1$ %Predicted | 99.3 (22.2) | 99.5 (13.9) | 95.9 (16.3) | 97.0 (14.9) | 0.29 |
| FVC, Litres | 3.7 (1.3) | 3.8 (1.1) | 3.7 (1.0) | 3.8 (0.8) | 0.94 |
| FVC z score | 0.2 (1.2) | 0.0 (0.9) | -0.1 (0.9) | -0.1 (0.8) | 0.10 |
| FVC %Predicted | 102.6 (18.4) | 100.7 (13.3) | 98.7 (13.5) | 99.2 (12.0) | 0.13 |
| $FEV_1$/FVC | 76.5 (12.7) | 78.7 (7.0) | 77.2 (9.2) | 77.0 (8.5) | 0.20 |
| $FEV_1$/FVC z score | -0.3 (1.4) | -0.1 (0.9) | -0.3 (1.0) | -0.3 (1.0) | 0.48 |
| **$T_L$co** | | | | | |
| $T_L$co z score | -0.2 (1.8) | -0.2 (1.0) | -0.2 (1.1) | -0.2 (1.2) | 0.99 |
| $T_L$co %Predicted | 98.6 (25.6) | 97.6 (15.0) | 97.8 (17.7) | 98.2 (19.8) | 0.98 |
| $T_L$co Hb corrected z score | -0.2 (1.8) | -0.2 (0.9) | -0.2 (1.1) | -0.2 (1.2) | 0.99 |
| $T_L$co Hb corrected %Predicted | 99.4 (25.5) | 98.2 (14.3) | 98.5 (17.6) | 99.0 (18.9) | 0.96 |
| **Respiratory symptoms in the last 12months** | | | | | |
| Wheeze | 66 (34%) | 53 (41%) | 52 (38%) | 71 (50%) | 0.08 |
| Wheeze and breathlessness | 46 (23%) | 43 (30%) | 36 (25%) | 46 (31%) | 0.47 |
| Wheeze without URTI | 49 (23%) | 38 (27%) | 36 (23%) | 61 (43%) | <0.001 |
| Chest tightness | 28 (15%) | 22 (15%) | 30 (19%) | 36 (27%) | 0.036 |
| Dyspnoea at rest | 26 (13%) | 20 (16%) | 22 (14%) | 28 (19%) | 0.60 |
| Dyspnoea after exercise | 76 (42%) | 65 (54%) | 49 (38%) | 68 (52%) | 0.049 |
| Woken with dyspnoea | 12 (6%) | 17 (12%) | 20 (14%) | 16 (10%) | 0.20 |
| Woken with cough | 56 (29%) | 52 (45%) | 42 (34%) | 54 (39%) | 0.11 |
| Chronic cough | 55 (29%) | 47 (39%) | 52 (45%) | 67 (48%) | 0.040 |
| Chronic phlegm | 25 (13%) | 14 (13%) | 21 (16%) | 28 (22%) | 0.22 |

Table 3. Adjusted associations (Odds Ratios and 95% Confidence Intervals) between $PM_{2.5}$ exposure, respiratory symptoms and lung function

| | Mean exposure model (10 μg/m³) | | Peak exposure model (100 μg/m³) | |
|---|---|---|---|---|
| | Adj OR (95% CI) | p-value | Adj OR (95% CI) | p-value |
| **Respiratory symptoms in the last 12 months**[*] | | | | |
| Wheeze | 1.13 (0.89,1.43) | 0.31 | 1.00 (0.89,1.13) | 0.99 |
| Wheeze and breathlessness | 1.00 (0.76,1.32) | 1.00 | 0.92 (0.80,1.06) | 0.26 |
| Wheeze without URTI | 1.19 (0.96,1.48) | 0.11 | 1.07 (0.96,1.20) | 0.22 |
| Chest tightness | 1.30 (1.03,1.64) | 0.026 | 1.08 (0.96,1.21) | 0.22 |
| Dyspnoea at rest | 1.02 (0.77,1.33) | 0.91 | 1.02 (0.87,1.19) | 0.84 |
| Dyspnoea after exercise | 1.08 (0.87,1.34) | 0.48 | 0.99 (0.89,1.11) | 0.88 |
| Woken with dyspnoea | 0.84 (0.63,1.12) | 0.23 | 0.92 (0.78,1.08) | 0.29 |
| Woken with cough | 0.97 (0.77,1.22) | 0.78 | 0.95 (0.84,1.07) | 0.39 |
| Chronic cough | 1.24 (1.02,1.51) | 0.035 | 1.06 (0.95,1.18) | 0.33 |
| Chronic phlegm | 1.23 (0.93,1.62) | 0.15 | 1.02 (0.88,1.17) | 0.83 |
| **Lung function**[†] | **Adj OR (95% CI)** | **p-value** | **Adj OR (95% CI)** | **p-value** |
| Spirometric COPD | 0.92 (0.67,1.26) | 0.60 | 1.02 (0.86,1.20) | 0.82 |
| Abnormal $T_Lco$ | 1.16 (0.83,1.61) | 0.39 | 1.05 (0.89,1.23) | 0.58 |
| | **Adj β (95% CI)** | **p-value** | **Adj β (95% CI)** | **p-value** |
| Post BD $FEV_1$ z score | 0.02 (-0.09,0.13) | 0.68 | 0.03 (-0.03,0.08) | 0.30 |
| Post BD $FEV_1$ %predicted | 0.04 (-1.71,1.78) | 0.97 | 0.28 (-0.56,1.11) | 0.51 |
| Post BD $FEV_1$/FVC z score | 0.04 (-0.06,0.13) | 0.44 | 0.00 (-0.05,0.05) | 0.99 |
| Corrected $T_Lco$ z score | -0.05 (-0.18,0.08) | 0.46 | 0.02 (-0.05,0.09) | 0.59 |

[*] Adjusted for smoking status, location of the participant (Morwell vs. Sale), BMI category, work place exposure, nasal allergies/hayfever, employed or not and having a certificate, university or other tertiary institute degrees.

[†] Adjusted for smoking status, location of the participant (Morwell vs. Sale), BMI category, employed or not and having a certificate, university or other tertiary institute degrees.

Figure 2. Forest plots of the estimated effects of PM$_{2.5}$ exposure on respiratory symptoms in the past 12 months for each smoking group.
*Note: The OR were estimated using multivariate logistic regression with an interaction between exposure and smoking status. All models were also adjusted for age, gender, location of the participant (Morwell vs. Sale), BMI category, work place exposure, nasal allergies/hayfever, employed or not and having a certificate university or other tertiary institute degrees. * indicates evidence of an overall interaction effect (p-value <0.1 from Wald test of all interaction terms equal to zero.*

Figure 3. Forest plots of the estimated effect of PM$_{2.5}$ exposure on lung function in each smoking group estimated using multivariate linear regression.
*Note: All models were adjusted for location of the participant (Morwell vs. Sale), BMI, work place exposure, employment and higher education.* indicates evidence of an overall interaction effect (p-value <0.1 from Wald test of all interaction terms equal to zero).*

**Adult Survey participants**
**n=4056**

Morwell = 3096      Sale = 960

**Participants eligible for Respiratory Stream**
**n=3846**

Morwell = 2948      Sale = 898

**Weighted random sample invited (40% with asthma)**
**n=1346**

Morwell = 842      Sale = 504

Target sample size was 339 Morwell and 170 Sale
Recruitment continued until target achieved.

Non participants: 398 declined, 4 deceased, 423 no response.

Respiratory Stream participants (first round)
n = 519

Morwell = 346      Sale = 173

|  | OR per 10 μg/m³ increase in mean PM₂.₅ | OR per 100 μg/m³ increase in peak PM₂.₅ |
|---|---|---|
| **Chest tightness** | 1.46 (1.07, 1.99) | 1.15 (0.96, 1.38) |
|  | 1.10 (0.75, 1.60) | 0.97 (0.80, 1.19) |
|  | 1.70 (1.05, 2.75) | 1.22 (0.98, 1.51) |
| **Chronic cough** | * 1.30 (0.97, 1.75) | * 1.04 (0.87, 1.24) |
|  | 1.10 (0.84, 1.45) | 1.00 (0.86, 1.17) |
|  | 2.18 (1.28, 3.71) | 1.32 (0.97, 1.79) |
| **Chronic phlegm** | 1.49 (0.99, 2.23) | 1.19 (0.94, 1.51) |
|  | 1.01 (0.68, 1.51) | 0.91 (0.73, 1.14) |
|  | 1.46 (0.78, 2.71) | 1.06 (0.84, 1.33) |
| **Dyspnoea after exercise** | 1.14 (0.85, 1.52) | 1.05 (0.89, 1.25) |
|  | 1.12 (0.86, 1.46) | 1.01 (0.87, 1.17) |
|  | 0.88 (0.54, 1.42) | 0.88 (0.70, 1.10) |
| **Dyspnoea at rest** | 1.20 (0.85, 1.68) | 1.09 (0.90, 1.33) |
|  | 0.93 (0.61, 1.43) | 1.06 (0.84, 1.34) |
|  | 0.97 (0.59, 1.60) | 0.91 (0.72, 1.14) |
| **Wheeze** | 1.26 (0.94, 1.69) | 1.08 (0.91, 1.28) |
|  | 1.07 (0.80, 1.44) | 0.94 (0.79, 1.11) |
|  | 1.08 (0.56, 2.10) | 1.09 (0.78, 1.51) |
| **Wheeze and breathlessness** | 1.25 (0.91, 1.72) | 1.06 (0.88, 1.27) |
|  | 0.85 (0.54, 1.32) | 0.85 (0.67, 1.09) |
|  | 1.03 (0.63, 1.69) | 0.91 (0.74, 1.13) |
| **Wheeze without URTI** | 1.36 (1.02, 1.81) | 1.17 (0.99, 1.38) |
|  | 1.05 (0.78, 1.42) | 0.97 (0.82, 1.14) |
|  | 1.36 (0.72, 2.57) | 1.25 (0.89, 1.75) |
| **Woken with cough** | * 0.87 (0.63, 1.21) | * 0.85 (0.71, 1.03) |
|  | 0.87 (0.62, 1.23) | 0.88 (0.73, 1.06) |
|  | 1.56 (0.97, 2.53) | 1.26 (1.01, 1.56) |
| **Woken with dyspnoea** | * 1.14 (0.79, 1.64) | * 1.09 (0.88, 1.35) |
|  | 0.46 (0.21, 1.00) | 0.57 (0.32, 1.01) |
|  | 0.96 (0.62, 1.49) | 1.05 (0.83, 1.35) |

Odds Ratio (95% Confidence Interval)

Smoking Status — Current smoker — Ex-smoker — Non-smoker