Life course air pollution exposure and cognitive decline: modelled historical air pollution data and the Lothian Birth Cohort 1936

RUNNING TITLE: Air pollution and cognitive decline

Tom C. **Russ**,[a-d] * Mark P. C. **Cherrie**,[e] Chris **Dibben**,[e,f] Sam **Tomlinson**,[g,h] Stefan **Reis**,[g,i]

Ulrike **Dragosits**,[g] Massimo **Vieno**,[g] Rachel **Beck**,[g] Ed **Carnell**,[g] Niamh K. **Shortt**,[k]

Graciela **Muniz-Terrera**,[a,c] Paul **Redmond**,[b] Adele M. **Taylor**,[b] Tom **Clemens**,[e]

Martie **van Tongeren**,[k] Raymond M **Agius**,[k] John M. **Starr**,[a,b †] Ian J. **Deary**,[b] Jamie R. **Pearce**[k]

† Professor Starr sadly died unexpectedly in December 2018 but made a great contribution to this project and would have fulfilled the ICJME criteria for authorship.

a. Alzheimer Scotland Dementia Research Centre, University of Edinburgh;

b. Lothian Birth Cohorts, Department of Psychology, University of Edinburgh

c. Edinburgh Dementia Prevention group, Centre for Clinical Brain Sciences, University of Edinburgh;

d. Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh;

e. School of GeoSciences, University of Edinburgh;

f. Scottish Centre for Administrative Data Research, University of Edinburgh;

g. UK Centre for Ecology & Hydrology (UKCEH), Bush Estate, Penicuik;

h. UK Centre for Ecology & Hydrology (UKCEH), Lancaster Environment Centre, Lancaster University

i. University of Exeter Medical School, Knowledge Spa, Truro

j. Centre for Occupational and Environmental Health, **School of Health Sciences, The University of** Manchester

k. Centre for Research on Environment, Society and Health, School of GeoSciences, University of Edinburgh

* Correspondence to: Dr Tom Russ, Alzheimer Scotland Dementia Research Centre, University of Edinburgh, 7 George Square, Edinburgh, EH8 9JZ, UK

Telephone: +44 (0)131 650 4340; Email: T.C.Russ@ed.ac.uk

Contributions: TCR drafted the article and all authors revised it for critical content.

Funding: This project was funded by the NERC-CSO-MRC call "Improving Health with Environmental Data" (NE/P010849/1).

Manuscript statistics: article body word count of 4103 plus an abstract of 246 words, one text box, five tables, three figures, two supplementary figures, and one supplementary table.

ABSTRACT

Background: Air pollution has been consistently linked with dementia and cognitive decline.

However, it is unclear whether risk is accumulated through long-term exposure or whether

there are sensitive/critical periods. A key barrier to clarifying this relationship is the dearth of

historical air pollution data. **Objective:** To demonstrate the feasibility of modelling historical

air pollution data and using them in epidemiological models. **Methods:** Using the EMEP4UK

atmospheric chemistry transport model, we modelled historical fine particulate matter ($PM_{2.5}$)

concentrations for the years 1935, 1950, 1970, 1980, and 1990 and combined these with

contemporary modelled data from 2001 to estimate life course exposure in 572 participants in

the Lothian Birth Cohort 1936 with lifetime residential history recorded. Linear regression

and latent growth models were constructed using cognitive ability (IQ) measured by the

Moray House Test at the ages of 11, 70, 76, and 79 years to explore the effects of historical air

pollution exposure. Covariates included sex, IQ at age 11 years, social class, and smoking.

Results: Higher air pollution modelled for 1935 (when participants would have been *in utero*)

was associated with worse change in IQ from age 11-70 years (β=-0.006, SE=0.002, P=0.03)

but not cognitive trajectories from age 70-79 years (P>0.05). There was no support for other

critical/sensitive periods of exposure or an accumulation of risk (all P>0.05). **Conclusions:**

The life course paradigm is essential in understanding cognitive decline and this is the first

study to examine life course air pollution exposure in relation to cognitive health.

KEYWORDS: Aging, dementia, Alzheimer disease, cognition, air pollution, epidemiologic

methods, atmosphere

INTRODUCTION

Dementia is a global public health crisis with almost 47 million people affected in 2015 and almost 10 million new cases every year, leading to a projected prevalence of over 130 million by 2050[1]. The brain changes which lead to many dementias – including the most common form, Alzheimer's dementia – begin in midlife and only manifest in later life[2]. Dementia prevention is now a worldwide priority and accepted risk factors include lower levels of educational attainment (in early life), cardiovascular disease risk factors (with hypertension and obesity particularly highlighted in mid-life), depression, hearing loss, and possession of the *APOE* ε4 allele[3]. A recent Lancet Commission report and other analyses have estimated that approximately a third of dementia risk can be explained by these common risk factors[4, 5]. With genetic factors (most prominently *APOE* ε4 carriage) explaining approximately another third[6], this leaves around a third of dementia risk unexplained. However, there is also evidence linking a number of environmental risk factors with dementia which might account for some of this unexplained risk[7]. The risk factor for which there is strongest evidence is air pollution[8, 9]. However, the field has been criticised since studies to date have lacked long-term (i.e. whole life) assessment of both exposure and outcome[10]. Thus, no light has yet been shed on the question of *when* in the life-course exposure to air pollution is most harmful to the brain. Recent papers describing "long-term exposure" to air pollution only estimated air pollution exposure at one time point[11, 12]. Indeed, answering this question from a life-course epidemiology perspective is hampered by both a dearth of available air pollution data from earlier than a few decades ago, when systematic long-term monitoring of atmospheric concentrations was implemented and limited information about the geographical location of study participants over their lives[13]. Therefore, we modelled air pollution data (fine particulate matter, with an aerodynamic diameter of 2.5μm or smaller; $PM_{2.5}$) for

multiple time periods and linked these with the Lothian Birth Cohort 1936 (LBC1936) — for

whom lifetime residential history is available — to investigate links between air pollution and

cognitive change over more than six decades.

MATERIALS & METHODS

Study participants

The LBC1936 is a well-established cohort study, originally comprising 1091 men and women

aged approximately 70 years at recruitment. Almost all sat the Moray House Test (MHT) of

general cognitive ability in the Scottish Mental Survey in 1947 when they were aged about 11

years[14]. Thus, general intelligence data are available for almost all participants at ages 11

years, and repeatedly from approximately 70 years onwards. In the present study, we used data

from waves 1, 3 and 4 when participants had mean ages of 69.5 (SD=0.8), 76.3 (0.7), and

79.3 (0.6) years respectively; the MHT was not administered in wave 2. We operationalised

cognitive function in the same way as previous studies, adjusting for age in days and

standardising to an IQ-type score with mean 100 and SD 15[15]. In line with previous

analyses, change in IQ score was computed as the standardised residual from a linear

regression model with age 11 IQ as the independent variable and age 70 IQ as the dependent

variable; this is superior to computing the arithmetic difference[16, 17].

In 2014, surviving LBC1936 participants were asked to complete a lifetime residential

questionnaire and 593 of 704 approached provided usable life grid data (full addresses) which

were geocoded to latitude and longitude[15]. Participants had a mean (SD) 11.3 (2.9)

separate locations throughout life, ranging from six to 27, with the years they lived there also

recorded. Each location was allocated to the closest time period for which air pollution data

were available: 1935 (location year 1942 or earlier); 1950 (1943-1959); 1970 (1960-1975);

1980 (1976-1985); 1990 (1986-1995); or 2001 (1995-2004); locations after 2004 were

excluded to avoid overlap with cognitive testing (wave 1 of the LBC1936 took place from

2004-2007[14]). Participants may have had more than one location allocated to each time

point — e.g., all locations between the years 1995 and 2004 would be allocated to the 2001

time point. Thus, participants had up to ten locations per time point (mean [SD] values

ranged from 1.11 [0.34] locations recorded in 2001 to 3.38 [1.33] locations in 1970). Twenty

one participants (3.5%) were missing location data for at least one time point, leaving 572 in

the final sample who had location (and therefore air pollution) data available for every time

point.

Other covariate data available in the LBC1936 and used in the models included sex, parental

occupational social class (using the Registrar General 1951 classification from I to V[18]), and

self-reported smoking status (current smoker at wave 1 or non-/ex-smoker).

Air pollution modelling

The EMEP4UK atmospheric chemistry transport model (rv4.3 for 1970-2010 and rv4.10 for

1935/50 [19]) was used to model historical ambient concentrations of fine particulate matter

($PM_{2.5}$) for the years 1935, 1950, 1970, 1980, and 1990 which were combined with

contemporary modelled data from 2000 onwards and residential histories to estimate life

course exposure. The EMEP4UK model setup, geographical coverage, and configuration used

here has been described previously[20, 21]. The model covers the European Union with a

horizontal resolution of 0.5° x 0.5° used to provide the boundary condition for a nested UK

domain (resolution of 0.055° x 0.055°). The modelled $PM_{2.5}$ and other key air pollutant

concentrations are routinely validated against observations across UK monitoring

networks[20-25] and have been used for the assessment of population exposure over longer

time scales in other studies for the period 1970 to 2010[26]. Emission data have been

identified as key sources of uncertainty in modelling historic air pollution. A detailed

assessment of sensitivity and uncertainty of the Atmospheric Chemistry Transport Models

(ACTM) applied in this study has been published elsewhere[27].

UK-specific gridded emissions of nitrogen oxides (NO_x), sulphur oxides (SO_x), ammonia

(NH_3), non-methane volatile organic compounds (NMVOCs), carbon monoxide (CO), and

coarse (PM_{10}) and fine ($PM_{2.5}$) particulate matter — all necessary for the atmospheric

composition calculations — were produced for the target years on a nominal 1km x 1km grid

covering the United Kingdom. Emissions data were internally re-projected and processed by

the EMEMP4UK model to provide output concentration data at the model grid resolution of

0.055° x 0.055° resolution (~5km x 6km) for the UK. The concentrations of $PM_{2.5}$ calculated

by the EMEP4UK model were used in conjunction with the residential history data (as

described above). The sources of primary emitted PM are varied but the main contributors are

essentially fuel combustion (from all sources) and the use of any mobile machinery, including

road traffic. This is in contrast to secondary produced PM — such as ammonium sulphate

which is formed by the interaction of ammonia gas and sulphur dioxide — which are strongly

linked to specific sectors, such as SO_x (energy) and NH_3 (agriculture). The PM components

included in the EMEP4UK model are; primary PM, secondary inorganic and organic aerosols,

sea salt, and mineral dust.[22] Although this work is focused on the UK the EMEP4UK

requires emission data for the whole of Europe to account for the transboundary

imports/export. EU data were kindly supplied at a 50km x 50km resolution[28, 29]. The

EMEP4UK model is driven by 3D hourly meteorology calculated by the weather and research

forecast model.[30] The meteorological year used for the 1935 and 1950 emission scenario

was the year 2014, for the 1970, 1980, 1990 emission scenario was the year 2012, and for the

2001 emission scenario the year was 2001.

For the years 1970, 1980 and 1990, emission data in the official UK inventory, the National

Atmospheric Emissions Inventory[31], were used to scale 2017 spatial distributions (1km x

1km resolution) of sectoral totals per pollutant, reported via the Selected Nomenclature for

sources of Air Pollution system (SNAP sectors). While the use of contemporary distributions

back to 1970 is imperfect, the majority of the time series had the best possible emissions

estimates per sector. Emissions for 1950 were estimated and distributed in the Long Term

Large Scale project[32], while the 1935 emissions were a scaled version of the 1950

distributions based upon activity data research, using the same spatial methods. Non-NH_3

activity data prior to 1970 are largely a reflection of the use of fossil fuels such as coal and of

oil-derived products such as diesel (DERV); coal usage in the UK had a double peak either

side of World War Two before a rapid decline in the 1960s. Agricultural activity data such as

animal numbers, principally associated with emissions of NH_3, were derived from the Vision of

Britain database.[33] For source strength emission factors (EFs), many were similar to those

used by the UK National Atmospheric Emission Inventory (NAEI) in 1970 while in terms of

the spatial distribution of pre-1970 data, the principal differences from the NAEI distributions

were: power stations relevant to the time period were mapped and the distribution of

industrial activity was tied to census employment data[32].

Raster files for each location year (1935, 1950, 1970, 1980, 1990, and 2001) were read into

the R statistical computing environment version 4.0.2 using the raster package[34]. The

latitude and longitude for each location were used to derive values from these raster files for

each participant at each time point. Since participants may have had multiple locations within

each time band, the unweighted mean of these multiple values per time point was calculated

and used in the analyses as the value for that participant at that time point. For example, the

mean of all the air pollution values corresponding to locations recorded between the years

1995 and 2004 would be allocated to the 2001 time point for an individual participant. For

the purposes of sensitivity analyses, we also computed the maximum value for each time point

per participant and the 90% percentile value.

Statistical modelling

Following the convention in previous LBC1936 analyses, we modelled change in IQ score

from age 11 to age 70 years separately from changes between the ages of 70, 76, and 79 years.

The former used a linear regression model of *in utero* air pollution exposure (i.e., using $PM_{2.5}$

data for 1935; this was the only measurement of air pollution which predated the MHT

administration at age 11 years) and residualised change in IQ score from age 11 to age 70

years in the R statistical computing environment version 4.0.2. We adjusted this model for

sex, parental occupation, and smoking status.

To estimate linear late life cognitive trajectories, we fitted latent growth models to IQ scores

from ages 70, 76, and 79 years to estimate the average population cognitive curves. These

longitudinal models permit estimation of the outcome's mean and individual trajectories while

permitting the inclusion of predictors (time-invariant or time-varying) to study their

association with curve parameters such as the intercept and slope parameters. Latent growth

models were estimated using maximum likelihood under a 'missing at random' missing data

assumption. All latent growth models were estimated using MPLUS[35].

We summarise the models fitted in **Figure 1**, in which observed data are represented within

rectangles, and latent variables such as the model's intercept and slope are represented within

circles. The one-way arrows indicate that the variable at the end of the arrow is explained in

the model by the variable at the beginning of the arrow. Often in such figures, two-way curved

arrows indicate covariances but as is commonly done – in order to simplify the figure – we

have omitted these arrows here as well as the arrows that indicate error terms.

Life course models

The two main models in life course epidemiology are critical/sensitive periods and

accumulation of risk[2]. To evaluate questions about any *critical/sensitive period(s)* of air

pollution exposure and maximise the use of pollution data collected over the life course, we

estimated late life trajectories of IQ scores at ages 70, 76 and 79 years, adjusting the intercept

and rate of change for air pollution measures collected at different ages in the life course, age

11 IQ scores, sex, parental social class, and smoking status. Specifically, we adjusted the

models separately for $PM_{2.5}$ in 1935, 1950, 1970, 1980, 1990, or 2001. **Figure 1a** depicts an

example of the critical/sensitive period model fitted here.

In order to evaluate an *accumulation of risk* model, we estimated a series of latent growth

models similar to the previous ones, adjusting the level and rate of change for variables

defined as the sum of air pollution to which the individual was exposed up to different stages

in life. That is, we added *in utero* air pollution measures (i.e., from 1935) to air pollution

measures collected in 1950 to derive an indicator of early life exposure; the sum of air

pollution measures from 1935 to 1970 covered early life to young adulthood; additionally

adding pollution from 1980 encompassed early life to mid-adulthood; the addition of air

pollution in 1990 covered early life to late adulthood; finally, adding air pollution from 2001

covered early life to later life. **Figure 1b** depicts an example of the accumulation period

model fitted here. **Text Box 1** summarises the models used in the present analyses.

RESULTS

A total of 572 LBC1936 participants were included in the present analyses. Their

characteristics are summarised in **Table 1**. Briefly, just under half were female, and had

completed more than compulsory education. Just over a quarter had parents from

occupational social classes I or II (i.e., less deprived), and about half were smokers at the time

of recruitment to the LBC1936. Comparing the 572 LBC1936 participants for whom we had

location (and therefore air pollution) data with the 519 participants excluded from these

analyses revealed no major differences.

Air pollution

Table 2 shows the average air pollution estimates for the LBC1936 participants and

Supplementary Figure 1 shows the distribution of air pollution exposure at each time period.

Supplementary Figure 2 shows participants' $PM_{2.5}$ exposure changes over time and

Supplementary Table 1 shows the correlations between individuals' $PM_{2.5}$ exposure ranking

at different time points. Rankings varied over time — likely due more to participants moving

than the relative ranking of areas changing — but were more closely correlated when closer in

time, suggesting it is feasible to explore critical/sensitive time periods. **Figure 2** shows the

modelled $PM_{2.5}$ values for Scotland in 1935; the urban centres are clearly visible.

Is in utero air pollution exposure associated with cognitive trajectories over the life course?

Results from this model are presented in **Table 3a**, where the potential impact of air pollution

measured in 1935 on the residualised change in IQ between the ages of 11 and 70 years,

controlling for sex, parental social, class, and smoking was explored. There was a small

association between higher levels of air pollution exposure in 1935 and *in utero* and a poorer

IQ trajectory in IQ from 11 to 70 years (β =-0.006 IQ point per 1 $\mu g/m^3$ increase in $PM_{2.5}$,

SE=0.002, P=0.03).

Is in utero exposure to air pollution associated with late life cognitive trajectories?

Figure 1a depicts the model that estimates linear changes in IQ scores between the ages of 70,

76, and 79 years. In this model, the intercept represents the average IQ score at age 70 for a

reference individual (a male whose father had a skilled job, who was exposed *in utero* to

average levels of pollution, and who had an average IQ score at age 11 years) and the slope,

the average rate of change of IQ scores from age 70 to 79 years. There was a small association

between air pollution values for 1935 and the intercept (IQ score at age 70 years), albeit only

of marginal statistical significance at conventional levels (P=0.06), but no association with

rate of change in IQ score from age 70 to 79 years (P=0.36; **Table 3b**).

Critical/sensitive period

For the sake of brevity, **Table 4** only contains estimates of the average value of IQ scores at

age 70, their rate of decline until age 79 (for a reference individual, as defined above) and

estimates of the association of air pollution exposure at each of the life course time points

(apart from 1935 which was reported above) with IQ level at age 70 and rate of change

(**Figure 1b**). No period of air pollution exposure had an effect on either the intercept or the

rate of change of the models which reached statistical significance at conventional levels (all

P>0.05).

Accumulation model

Summary results of the models are presented in **Table 5**, where estimates of average IQ level

at age 70, average IQ linear rate of change from that same age, and coefficients of the

association between air pollution exposure at different stages of life with these parameters are

presented. None of the risk periods had an effect on either the intercept or the rate of change

of the models which reached statistical significance at conventional levels (all P>0.05). Our

findings were robust to the sensitivity analyses varying the aggregation method used for

multiple air pollution values.

DISCUSSION

Our main finding is that it is feasible to model historical air pollutant concentration data and

incorporate them in epidemiological models to explore the influence of exposure to air

pollution across the life course. We found little evidence that exposure to air pollution at

different stages of the life course was associated with cognitive health and there was no

support for an accumulation of risk. There was some evidence of exposure to air pollution *in*

utero being associated with worse cognitive change between the ages of 11 and 70 years, but

the effect size was small (β=-0.006). In particular we would highlight that these results have

large degrees of uncertainties, considering the various methodologies used to produce the

different air pollution concentration estimates due the wide range of emissions estimates,

particularly for earlier estimates which have a lack of measured air quality data against which

to be evaluated.

Comparison with other literature

As noted in the introduction, there is a growing wealth of literature on the association

between air pollution and subsequent cognitive impairment and dementia, but the majority of

publications share the same shortcomings[10]: (1) an inability to explore *when* in the life

course exposure to air pollution has the most impact?; (2) which pollutant(s) or components

are most important?; and (3) since dementia describes a heterogeneous group of conditions,

which are most affected by exposure to air pollution?

Investigators from the Washington Heights–Inwood Community Aging Project (WHICAP)

and the Northern Manhattan Study (NOMAS) recently reported their findings of the impact

of "long-term" exposure to air pollution (nitrogen dioxide, PM_{10}, and $PM_{2.5}$), but these

participants (aged 65 years or older) were only recruited in the early 1990s and only their

residential address at the time of recruitment was used to estimate their exposure to air

pollution; air pollution values for the year before recruitment were used as the exposure[11,

12]. Our study was able to track migration and movement throughout the life course,

combined with modelled atmospheric concentration data covering most of the twentieth

century, to give a much better estimate of each person's exposure to air pollution at different

points in their lives.

Limitations and Strengths

Referring to the three criticisms of the air pollution literature described above,[10] the present

study could potentially shed some provisional light on the first (when in the life course is most

important), but not the second or third. A decision was taken early on to minimise the impact

of multiple testing by restricting the pilot modelling (of 1935 data) to a single pollutant; $PM_{2.5}$

was chosen since the majority of the literature linking air pollution and dementia focused on

that pollutant. Modelling other pollutants — such as smaller PM, NO_x etc. — is feasible and

we hope to do this in the future, now that the feasibility of this approach has been

demonstrated.

The collection of lifetime residential histories is rare and greatly augments the other data

available in the LBC1936. However, there are limitations to the approach taken

(retrospective collection of residential address history), including the fact that it is prone to

recall bias. Furthermore, only participants who were alive in 2014 were approached,

additionally introducing survivor bias. Finally, the accumulated $PM_{2.5}$ exposure was calculated

using an unweighted method – i.e. not taking into account the length of time an individual

lived at each address. Our main aim in the present analysis was to establish proof of concept

and hope that a more sophisticated weighted calculation – which could arguably be more

accurate – could be taken by future studies.

Almost all participants — until 2001 — were exposed to levels of $PM_{2.5}$ in excess of the World

Health Organization's guidelines of a maximum annual mean of $10\mu g/m^3$[36]. For

comparison, approximately half of UK Biobank participants were exposed to similarly

excessive values at baseline (mean [SD] 10.0 [1.1])[37], in line with most of the world[38].

There was a general reduction in air pollution over time — and marked step-changes between

some time points — but it is unclear how much of this is artefactual, relating to

methodological differences between the procedures used to generate these historical estimates.

There are inherent uncertainties at all stages of the emissions estimation process, even for the

present day, and many of these problems are magnified when trying to recreate an historical

context. The lack of measurement data to verify source strength, the lack of data regarding

the chemical composition of fuel and the behaviour of combustion technologies prior to

emissions mitigations are just some of the many issues that can influence the uncertainty.

From 1970 onwards, emissions uncertainties were estimated from the UK Inventory[39] while

for 1935 and 1950, uncertainty estimations are expert judgement.[40] Given the assumptions

that the fuel use data are lacking some detail (such as certain oil-based products, wood etc.),

combustion technology was a lot more polluting than in 1970 due to a lack of various

mitigating options such as scrubbers and particulate filters and that there are some missing

sources such as construction etc., it is very likely that the uncertainty range is asymmetric with

respect to the best estimate. To reflect this probable under estimation, the asymmetry was

estimated to be one order of magnitude centred on the mean, that is $\frac{1}{\sqrt{10}} < x < \sqrt{10}$. As an

example, the emissions of $PM_{2.5}$ in 1935 were 715kt (range 226-2261kt). Whilst the

uncertainties were not utilised within the EMEP4UK model, it is important to note these

qualitative estimates and the potential impacts on the final results. **Figure 3** shows the final

emissions estimates for all pollutants per year (2015 is displayed for context).

The focus of this paper is cognitive change rather than dementia. It is important to assess pre-

dementia cognitive change and its determinants in their own right, but dementia is inarguably

important. However, there were not sufficient LBC1936 participants who had developed

dementia to allow meaningful models to be constructed. A comprehensive programme of

dementia ascertainment in LBC1936 participants (who are currently in their mid-80s) is

currently underway and, once these data are available, similar models focused on dementia

will be possible.

Statistically, the life course model (change between ages 11 and 70 years) and late life models

(intercept and change from 70 to 79 years) are not comparable. Furthermore, a linear

assumption for the late life cognitive trajectories may be too strong. It may be necessary to

explore more complex models, such as quadratic trajectories, but the primary aim of this study

was proof of concept and so we have not taken that approach here. With additional time

points, we could have constructed a model that would permit estimation of piecewise

trajectories. This may become possible as further waves of data become available — wave 5 of

the LBC1936 was completed last year and wave 6 was due to begin in Spring 2020 but had to

be delayed because of the Covid-19 pandemic.

Future directions

This paper is the first step towards an understanding of the associations between air pollution

and cognitive decline and dementia from a life course epidemiology perspective. The modelled

historical air pollution data need to be refined and harmonised across different time points –

and these data used to provide a robust estimate of life course exposure – but we believe that

we have demonstrated the feasibility and value of this approach. However, these air pollution

data will be of little value without well-characterised cohort studies with full residential

histories for participants, such as are available for the LBC1936. All high quality longitudinal

cohort studies should explore whether it is possible to obtain such data through record linkage

or — as in the LBC1936 — self-report.

Conclusions

We have shown the feasibility of modelling historical air pollution data and incorporating

them in epidemiological models. This is the first step in a new area and we look forward to a

greater understanding of the life course effects of air pollution on the brain in coming years.

ACKNOWLEDGEMENTS

This study was funded by the "Improving Health with Environmental Data" call from the Natural Environment Research Council, the Chief Scientist Office, and the Medical Research Council (NE/P010849/1). It follows research undertaken under the Mobility, Mood, and Place research project (2013–2016), supported by Research Councils UK (EP/K037404/1) as part of the Cross-Council Lifelong Health and Wellbeing Program under Principal Investigator Catharine Ward Thompson and Co-Investigators Jamie Pearce and Niamh Shortt.

TCR and the late JMS are/were members of the Alzheimer Scotland Dementia Research Centre funded by Alzheimer Scotland. The Lothian Birth Cohort 1936 is funded by Age UK (Disconnected Mind grant). TCR, the late JMS, and IJD were members of the University of Edinburgh Centre for Cognitive Ageing & Cognitive Epidemiology (which recently closed), part of the cross council Lifelong Health and Wellbeing Initiative (MR/K026992/1). Funding from the Biotechnology and Biological Sciences Research Council and Medical Research Council is gratefully acknowledged for the latter.

Initial findings of this study were presented at the Alzheimer's Association International Conference in Chicago, 2018 (https://doi.org/10.1016/j.jalz.2018.06.2861).

CONFLICTS OF INTEREST

None

REFERENCES

[1]   Prince MJ, Wimo A, Guerchet M, Ali G-C, Wu Y-T, Prina AM (2015) *World Alzheimer Report 2015. The Global Impact of Dementia: an analysis of prevalence, incidence, cost and trends*, Alzheimer's Disease International, London.

[2]   Whalley LJ, Dick FD, McNeill G (2006) A life-course approach to the aetiology of late-onset dementias. *Lancet Neurology* **5**, 87-96.

[3]   Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, Ballard C, Banerjee S, Burns A, Cohen-Mansfield J, Cooper C, Fox N, Gitlin LN, Howard R, Kales HC, Larson EB, Ritchie K, Rockwood K, Sampson EL, Samus Q, Schneider LS, Selbaek G, Teri L, Mukadam N (2017) Dementia prevention, intervention, and care. *Lancet* **390**, 2673-2734.

[4]   Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C (2014) Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *Lancet Neurology* **13**, 788-794.

[5]   Tomata Y, Li X, Karlsson IK, Mosing M, Pedersen N, Hägg S (2020) Joint impact of common risk factors on incident dementia: A cohort study of the Swedish Twin Registry. *Journal of Internal Medicine*.

[6]   Van Cauwenberghe C, Van Broeckhoven C, Sleegers K (2016) The genetic landscape of Alzheimer disease: clinical implications and perspectives. *Genetics in Medicine* **18**, 421-430.

[7]   Killin LO, Starr JM, Shiue IJ, Russ TC (2016) Environmental risk factors for dementia: a systematic review. *BMC Geriatrics* **16**, 175.

[8]   Peters R, Ee N, Peters J, Booth A, Mudway I, Anstey KJ (2019) Air pollution and dementia: a systematic review. *Journal of Alzheimer's Disease*, 1-19.

[9]   Rizzuto D, Grande G, Ljungman P, Bellander T (2019) Long-term exposure to air pollution and the risk of dementia: the role of cardiovascular diseases. *Innovation in Aging* **3**, S119.

[10]   Russ TC, Reis S, van Tongeren M (2019) Air pollution and brain health: defining the research agenda. *Curr Opin Psychiatry* **32**, 97-104.

[11]   Kulick ER, Elkind MSV, Boehme AK, Joyce NR, Schupf N, Kaufman JD, Mayeux R, Manly JJ, Wellenius GA (2020) Long-term exposure to ambient air pollution, APOE-ε4 status, and cognitive decline in a cohort of older adults in northern Manhattan. *Environ Int* **136**, 105440.

[12]   Kulick ER, Wellenius GA, Boehme AK, Joyce NR, Schupf N, Kaufman JD, Mayeux R, Sacco RL, Manly JJ, Elkind MSV (2020) Long-term exposure to air pollution and trajectories of cognitive decline among older adults. *Neurology* **94**, e1782-e1792.

[13]   Pearce JR (2018) Complexity and Uncertainty in Geography of Health Research: Incorporating Life-Course Perspectives. *Annals of the American Association of Geographers* **108**, 1491-1498.

[14]   Taylor AM, Pattie A, Deary IJ (2018) Cohort Profile Update: The Lothian Birth Cohorts of 1921 and 1936. *Int J Epidemiol* **47**, 1042-1042r.

[15]   Cherrie MPC, Shortt NK, Mitchell RJ, Taylor AM, Redmond P, Thompson CW, Starr JM, Deary IJ, Pearce JR (2018) Green space and cognitive ageing: A retrospective life course analysis in the Lothian Birth Cohort 1936. *Soc Sci Med* **196**, 56-65.

[16]   Gow AJ, Whiteman MC, Pattie A, Whalley L, Starr J, Deary IJ (2005) Lifetime intellectual function and satisfaction with life in old age: longitudinal cohort study. *BMJ* **331**, 141-142.

[17]   Prochaska JJ, Velicer WF, Nigg CR, Prochaska JO (2008) Methods of quantifying change in multiple risk factor interventions. *Preventive Medicine* **46**, 260-265.

[18]   Office of Population Censuses and Surveys (1951) *Census of population 1951*, Her Majesty's Stationery Office, London.

[19]   Vieno M, Heal MR, Hallsworth S, Famulari D, Doherty RM, Dore AJ, Tang YS, Braban CF, Leaver D, Sutton MA, Reis S (2014) The role of long-range transport and domestic

emissions in determining atmospheric secondary inorganic particle concentrations across the UK. *Atmos. Chem. Phys.* **14**, 8435-8447.

[20]   Vieno M, Heal MR, Twigg MM, MacKenzie I, Braban CF, Lingard J, Ritchie S, Beck R, Móring A, Ots R (2016) The UK particulate matter air pollution episode of March–April 2014: more than Saharan dust. *Environmental Research Letters* **11**, 044004.

[21]   Vieno M, Heal M, Williams M, Carnell E, Nemitz E, Stedman J, Reis S (2016) The sensitivities of emissions reductions for the mitigation of UK PM2. 5. *Atmospheric Chemistry and Physics* **16**, 265-276.

[22]   Ots R, Vieno M, Allan JD, Reis S, Nemitz E, Young DE, Coe H, Di Marco C, Detournay A, Mackenzie IA (2016) Model simulations of cooking organic aerosol (COA) over the UK using estimates of emissions based on measurements at two sites in London. *Atmospheric Chemistry and Physics* **16**, 13773-13789.

[23]   Ots R, Heal MR, Young DE, Williams LR, Allan JD, Nemitz E, Di Marco C, Detournay A, Xu L, Ng NL (2018) Modelling carbonaceous aerosol from residential solid fuel burning with different assumptions for emissions. *Atmospheric Chemistry and Physics* **18**, 4497-4518.

[24]   Lin C, Heal MR, Vieno M, MacKenzie IA, Armstrong BG, Butland BK, Milojevic A, Chalabi Z, Atkinson RW, Stevenson DS (2017) Spatiotemporal evaluation of EMEP4UK-WRF v4. 3 atmospheric chemistry transport simulations of health-related metrics for NO2, O3, PM10, and PM2. 5 for 2001–2010. *Geoscientific Model Development* **10**, 1767-1787.

[25]   Twigg MM, Ilyinskaya E, Beccaceci S, Green DC, Jones MR, Langford B, Leeson SR, Lingard JJ, Pereira GM, Carter H (2016) Impacts of the 2014–2015 Holuhraun eruption on the UK atmosphere. *Atmospheric Chemistry and Physics* **16**, 11415-11431.

[26]   Carnell E, Vieno M, Vardoulakis S, Beck R, Heaviside C, Tomlinson S, Dragosits U, Heal MR, Reis S (2019) Modelling public health improvements as a result of air pollution control policies in the UK over four decades—1970 to 2010. *Environmental Research Letters* **14**, 074001.

[27]   Aleksankina K, Reis S, Vieno M, Heal MR (2019) Advanced methods for uncertainty assessment and global sensitivity analysis of an Eulerian atmospheric chemistry transport model. *Atmos. Chem. Phys.* **19**, 2881-2898.

[28]   Simpson D (2017) *Personal communication*.

[29]   Engardt M, Simpson D, Schwikowski M, Granat L (2017) Deposition of sulphur and nitrogen in Europe 1900–2050. Model calculations and comparison to historical observations. *Tellus B: Chemical and Physical Meteorology* **69**, 1328945.

[30]   Skamarock W, Klemp J, Dudhia J, Gill D, Liu Z, Berner J, Wang W, Powers J, Duda M, Barker D (2019) *A Description of the Advanced Research WRF Model. Version 4;(No. NCAR/TN-556+ STR)*, National Center for Atmospheric Research, Boulder, CO, USA.

[31]   NAEI, National Atmospheric Emissions Inventory: Data, https://naei.beis.gov.uk/data/,

[32]   Tipping E, Davies JAC, Henrys PA, Kirk GJD, Lilly A, Dragosits U, Carnell EJ, Dore AJ, Sutton MA, Tomlinson SJ (2017) Long-term increases in soil carbon due to ecosystem fertilization by atmospheric nitrogen deposition demonstrated by regional-scale modelling and observations. *Scientific Reports* **7**, 1890.

[33]   Vision of Britain Historic agricultural data, http://www.visionofbritain.org.uk,

[34]   Hijmans RJ, Etten Jv, raster: Geographic analysis and modeling with raster data. R package version 2.0-12., http://CRAN.R-project.org/package=raster,

[35]   Muthén LK, Muthén BO (1998-2017) *Mplus User's Guide*, Muthén & Muthén, Los Angeles, CA.

[36]   WHO (2006) *Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide. Global update 2005. Summary of risk assessment*, World Health Organization, Geneva.

[37]   UK Biobank, UK Biobank Data Showcase: Data-Field 24006 - Particulate matter air pollution (pm2.5), http://biobank.ctsu.ox.ac.uk/crystal/field.cgi?id=24006,

[38]   Shaddick G, Thomas ML, Mudu P, Ruggeri G, Gumy S (2020) Half the world's population are exposed to increasing air pollution. *NPJ Climate and Atmospheric Science* **3**, 23.

[39]   Richmond B, Misra A, Broomfield M, Brown P, Karagianni E, Murrells T, Pang Y, Passant N, Pearson B, Stewart R, Thistlethwaite G, Wakeling D, Walker C, Wiltshire J (2018) *UK Informative Inventory Report (1990 to 2017)*.

[40]   Smith SJ, van Aardenne J, Klimont Z, Andres RJ, Volke A, Delgado Arias S (2011) Anthropogenic sulfur dioxide emissions: 1850–2005. *Atmos. Chem. Phys.* **11**, 1101-1116.

Text Box. Summary of the models used in the present analyses

OUTCOME	EXPOSURE
Sensitive/critical period(s)	
Change in IQ from age 11 to age 70 years	In utero $PM_{2.5}$ exposure (1935)
Trajectories of IQ from age 70 to 79 years (intercept and rate of change)	*In utero* $PM_{2.5}$ exposure (1935) $PM_{2.5}$ exposure aged ~14 years (1950) $PM_{2.5}$ exposure aged ~34 years (1970) $PM_{2.5}$ exposure aged ~44 years (1980) $PM_{2.5}$ exposure aged ~54 years (1990) $PM_{2.5}$ exposure aged ~65 years (2001)
Accumulation of risk	
Trajectories of IQ from age 70 to 79 years (intercept and rate of change)	Early life (1935 + 1950) Early life to young adulthood (1935 + 1950 + 1970) Early life to mid-adulthood (1935 + 1950 + 1970 + 1980) Early life to late adulthood (1935 + 1950 + 1970 + 1980 + 1990) Early life to later life (1935 + 1950 + 1970 + 1980 + 1990 + 2001)

Table 1. Sample characteristics: life course air pollution exposure and cognitive decline in the

LBC1936

	Included[a]	Excluded[b]	P[c]	Total LBC1936 sample
N	572	519		1091
Age at SMS1947 (mean [SD] years)	10.92 (0.27)	10.96 (0.29)	0.027	10.94 (0.28)
Female (%)	46.9	53.0	0.0497	49.8
Age 11 IQ[d] (mean [SD])	101.6 (15.0)	98.2 (14.9)	<0.001	100.0 (15.0)
Parental occupational social class (% class I or II)	27.7	26.3	0.011	27.1
Current smoker at baseline (%)	49.3	42.2	0.022	45.9

[a] Participants were included if they had at least one location recorded for each time period.

[b] Excluded participants included 21 with missing location data for at least one time period, 111 who did not respond to the questionnaire requesting lifetime residential history, and 387 who were not approached, mainly because they had died or withdrawn from the study prior to the questionnaire being used in 2014.

[c] P-values from comparisons of included and excluded participants

[d] 31 participants were missing age 11 intelligence data

[e] Self-reported

LBC1936: Lothian Birth Cohort 1936 (N=1091);
SMS1947: Scottish Mental Survey 1947 (N=70,805, of which the LBC1936 is a subset)

Table 2. Annual average particulate matter ($PM_{2.5}$) values at different time points for all participants: life course air pollution exposure and cognitive decline in the LBC1936

Year	Mean (sd)	Range	N_{total}[a]	$>10\mu g/m^3$ [b]
1935	34.8 (16.0)	5.2-133.0	590	562 (95%)
1950	32.4 (12.8)	6.0-113.3	591	578 (98%)
1970	17.0 (1.5)	9.5-23.9	585	584 (100%)
1980	15.0 (1.5)	7.3-24.0	580	575 (99%)
1990	13.4 (1.2)	6.7-21.4	580	579 (100%)
2001	7.9 (0.6)	4.8-15.9	591	4 (0.7%)

[a] 593 participants provided lifetime residential histories; 572 had air pollution data from all time periods and were included in the present analyses
[b] The number (%) of participants whose $PM_{2.5}$ exposure exceeded the WHO guidelines of an annual mean of $\leq10\mu g/m^3$

Table 3. Results from (a) linear regression of residualised change in IQ from age 11 to age 70 years and (b) latent growth models fitted to IQ scores to estimate cognitive trajectories at ages 70, 76, and 79 years: life course air pollution exposure and cognitive decline in the LBC1936

(a) Change in IQ between ages 11 and 70 years		
	ß (SE)	P
In utero exposure to air pollution	-0.006 (0.002)	0.03

(b) IQ trajectories from age 70 to age 79 years					
Intercept (average IQ at 70 years)	97.74 (1.38)		Rate of change (in IQ from age 70 to 79 years)	-0.11 (0.31)	0.71
Random Intercept Variance	71.12 (5.61)		Random Slopes variance	2.36 (0.31)	<0.001
Intercept-slope correlation	-3.02 (0.06)	<0.001			
	ß (SE)	P		ß	P
In utero exposure to air pollution	0.05 (0.02)	0.06	*In utero* exposure to air pollution	-0.006 (0.006)	0.36

Model (a) is adjusted for sex, parental (father's) occupation, and smoking
Model (b) is adjusted for sex, age 11 IQ, parental (father's) occupation, and smoking
Coefficients (β) represent the change in IQ and rate of change per 1 $\mu g/m^3$ increase in $PM_{2.5}$

Table 4. Estimates of the association between air pollution exposures at different time points in the life course with mean IQ at age 70 and its rate of change from 70 to 79 years: life course air pollution exposure and cognitive decline in the LBC1936

Level and change in IQ between ages 70, 76, and 79 years						
	IQ	ß (SE)	P		ß (SE)	P
Age 70 IQ	102.14 (1.62)			Rate of change in IQ from age 70-79	-0.14 (0.33)	0.46
Pollution 1950		-0.027 (0.04)	0.52		-0.001 (0.006)	0.84
Age 70 IQ	105.14 (5.56)				0.21 (1.13)	0.85
Pollution 1970		-0.22 (0.04)	0.46		-0.03 (0.06)	0.65
Age 70 IQ	96.38 (4.94)				0.84 (1.51)	0.57
Pollution 1980		0.32 (0.32)	0.32		-0.07 (0.10)	0.45
Age 70 IQ	99.39 (7.34)				1.46 (1.51)	0.33
Pollution 1990		0.14 (0.54)	0.79		-0.13 (0.11)	0.24
Age 70 IQ	103.21 (8.84)				-0.91 (1.86)	0.62
Pollution 2001		-0.24 (1.10)	0.82		0.08 (0.23)	0.74

Models adjusted for sex, age 11 IQ, parental (father's) occupation, and smoking status
Coefficients (β) represent the change in IQ and rate of change per 1 $\mu g/m^3$ increase in $PM_{2.5}$

Table 5. Estimates of IQ intercept (at age 70 years) and rate of change from age 70 and of the association of cumulative air pollution exposure at various stages of life: life course air pollution exposure and cognitive decline in the LBC1936

Level and change in IQ between ages 70, 76, and 79 years						
	IQ	ß (SE)	P		ß (SE)	P
Age 70 IQ	100.49 (1.63)			Rate of change in IQ from age 70-79	-0.12 (0.34)	0.72
Early life (1935 + 1950)		0.01 (0.02)	0.54		-0.002 (0.003)	0.47
Age 70 IQ	100.42 (1.86)				-0.07 (0.39)	0.84
Early life to young adulthood (1935 + 1950 + 1970)		0.01 (0.02)	0.58		-0.002 (0.003)	0.46
Age 70 IQ	100.19 (2.08)				-0.02 (0.43)	0.96
Early life to mid-adulthood (1935 + 1950 + 1970 + 1980)		0.01 (0.02)	0.54		-0.003 (0.003)	0.42
Age 70 IQ	100.03 (2.27)				0.04 (0.47)	0.92
Early life to late adulthood (1935 + 1950 + 1970 + 1980 + 1990)		0.01 (0.02)	0.54		-0.003 (0.003)	0.38
Age 70 IQ	99.96 (2.39)				0.06 (0.49)	0.89
Early life to later life (1935 + 1950 + 1970 + 1980 + 1990 + 2001)		0.01 (0.02)	0.54		-0.003 (0.003)	0.38

Models adjusted for sex, age 11 IQ, parental (father's) occupation, and smoking status
Coefficients (β) represent the change in IQ and rate of change per 1 $\mu g/m^3$ increase in $PM_{2.5}$

FIGURE LEGENDS

Figure 1. Directed Acyclic Graphs representing the (a) critical/sensitive period and (b) accumulation models fitted to IQ scores: life course air pollution exposure and cognitive decline in the LBC1936

Figure 2. Modelled particulate matter ($PM_{2.5}$) values in 1935: life course air pollution exposure and cognitive decline in the LBC1936

Figure 3. Modelled emission totals (Gg) with uncertainty ranges for five air pollutants (CO, NH_3, NMVOCs, NO_x, and SO_x), plus $PM_{2.5}$, across five model years (2015 is included for context) for use in the EMEP4UK model: life course air pollution exposure and cognitive decline in the LBC1936

Figure 1. Figure representing the (a) critical/sensitive period and (b) accumulation models fitted to IQ scores: life course air pollution exposure and cognitive decline in the LBC1936

(a) Critical/sensitive period models   **(b) Accumulation models**

Figure 2. Modelled particulate matter (PM$_{2.5}$) values in 1935: life course air pollution exposure and cognitive decline in the LBC1936

PM$_{2.5}$ concentration in 1935 (µg/m^3)

25   50   75   100

The area displayed in the lower panel and enclosed in a box on the upper panel is the central belt of Scotland including Glasgow (left) and Edinburgh (right). Over half of the population of Scotland lives in this area.

Figure 3. Modelled emissions totals (Gg) with uncertainty ranges for five air pollutants (CO, NH$_3$, NMVOCs, NO$_x$, and SO$_x$), plus PM$_{2.5}$, across five model years (2015 is included for context) for use in the EMEP4UK model: life course air pollution exposure and cognitive decline in the LBC1936

Supplementary Figure 1a. Mean PM$_{2.5}$ exposure for each participant at each time point for which air pollution concentration data were modelled: life course air pollution exposure and cognitive decline in the LBC1936

Blue dotted line — mean PM$_{2.5}$ value; Solid red line — WHO guidelines (annual average $\leq$10µg/m^3)

Supplementary Figure 1b. Mean PM$_{2.5}$ exposure for each participant at each time point for which air pollution concentration data were modelled (all plotted on the same x and y scales): life course air pollution exposure and cognitive decline in the LBC1936

Blue dotted line — mean PM$_{2.5}$ value; Solid red line — WHO guidelines (annual average ≤10μg/m^3)

Supplementary Figure 2. Sankey diagram indicating the change in mean annual PM$_{2.5}$ exposure within individuals over time: life course air pollution exposure and cognitive decline in the LBC1936

Supplementary Table 1. Correlations between PM$_{2.5}$ exposure rankings at different time points: life course air pollution exposure and cognitive decline in the LBC1936

	1935	1950	1970	1980	1990	2001
1950	0.53	-				
1970	0.13	0.22	-			
1980	0.06	0.18	0.57	-		
1990	0.09	0.19	0.53	0.82	-	
2001	0.003	0.10	0.05	0.03	0.08	-