

1 **Minimum Dietary Diversity and Associated Factors among Lactating Mothers in Ataye**
2 **District, North Shoa Zone, Central Ethiopia: A Community-based Cross-sectional Study**

3 Lemma Getacher^{1*}, Gudina Egata², Tadesse Alemayehu², Agegnehu Bante³, and Abebaw
4 Molla⁴

5 ¹Department of Public Health, College of Health Sciences, Debre Berhan University, Debre
6 Berhan, Ethiopia

7 ²School of Public Health, College of Health and Medical Sciences, Haramaya University, Harar,
8 Ethiopia

9 ³Departments of Nursing, College of Medicine and Health Sciences, Arba Minch University,
10 Arba Minch, Ethiopia

11 ⁴Department of Nursing, College of Medicine and Health Sciences, Mizan Tepi University,
12 Mizan Tepi, Ethiopia

13 *Corresponding Author E-mail: lemmagetacher@gmail.com

14 **Institutional email:** lemmagetacher@dbu.edu.et

15 **Abstract**

16 **Background:** Low dietary diversity superimposed with poor quality monotonous diets is a
17 major problem that often results in undernutrition, mainly micronutrient deficiencies. However,
18 there is limited evidence on minimum dietary diversity and associated factors among lactating
19 mothers in resource-poor settings, including the study area. Therefore, the objective of the study
20 is to assess the prevalence of minimum dietary diversity and associated factors among lactating
21 mothers in Ataye District, Ethiopia.

22 **Methods:** A community-based cross-sectional study design was used among 652 lactating
23 mothers aged 15-49 years from January 25 to April 30, 2018. Dietary diversity was measured

24 by the minimum dietary diversity indicator for women (MDD-W) using the 24-hour dietary
25 recall method. Data were entered into EpiData version 4.2.0.0 and exported to the statistical
26 package for social science (SPSS) version 24 for analysis using the logistic regression model.

27 **Results:** The prevalence of minimum dietary diversity among lactating mothers was 48.8%,
28 (95% CI: (44.7%, 52.9%). Having formal education [(AOR=2.16, 95% CL: (1.14, 4.09)], a final
29 say on household purchases [(AOR=5.39, 95% CI: (2.34, 12.42)], home gardening practices [
30 (AOR=2.67, 95% CI: (1.49, 4.81)], a history of illness [(AOR=0.47, 95% CI: (0.26, 0.85), good
31 knowledge of nutrition [(AOR=5.11, 95% CI: (2.68, 9.78)], being from food-secure households
32 [(AOR=2.96, 95% CI: (1.45, 6.07)] and medium [(AOR=5.94, 95% CI: (2.82, 12.87)] and rich
33 wealth indices [(AOR= 3.55, 95% CI: (1.76, 7.13)] were significantly associated with minimum
34 dietary diversity.

35 **Conclusion:** The prevalence of minimum dietary diversity among lactating mothers was low in
36 the study area. It was significantly associated with mothers having a formal education, final say
37 on the household purchase, home garden, good knowledge of nutrition, history of illness, food-
38 secure households and belonging to medium and rich household wealth indices. Therefore,
39 efforts should be made to improve the mothers decision-making autonomy, nutrition knowledge,
40 household food security and wealth status.

41 **Keywords:** Ataye District, Minimum Dietary Diversity, Ethiopia, Lactating Mothers

42 **1. Background**

43 Appropriate and adequate maternal nutrition is a pivotal public health issue in resource-poor
44 countries throughout the globe, particularly in Africa, Southeast Asia, Latin America, and the
45 Caribbean. Women of reproductive age (WRA) are often nutritionally vulnerable, especially
46 lactating mothers [1]. Lactating mothers, particularly those who live in low-and middle-income

47 countries, are vulnerable to malnutrition due to increased physiological demand, lactogenesis
48 process, eating undiversified monotonous diet and increased nutrient needs during lactation.
49 Nutritional requirements during lactation are greater than during pregnancy, and they need at
50 least two extra meals per day. When these needs are not met, mothers may suffer from
51 malnutrition especially, from micronutrient deficiencies. To prevent and correct nutritional
52 vulnerability in women of reproductive age, especially during the most 1000 critical days, one
53 of the best nutritional interventions is dietary diversification. Dietary diversity (DD) refers to an
54 increase in the variety of foods across and within food groups capable of ensuring adequate
55 intake of essential nutrients that can promote good health, physical and mental development of
56 women. Moreover, it is a qualitative measure of food consumption that reflects household access
57 to a variety of foods and is also a proxy for the micronutrient adequacy diet of individuals [2-
58 6].

59 Micronutrient deficiency (aka hidden hunger) is one form of undernutrition that occurs due to
60 insufficient intake, poor knowledge about nutrition, inadequate dietary diversity or sufficient
61 intake combined with impaired absorption due to infection, disease or inflammation and
62 phytates (anti-nutrients) that can decrease the bioavailability of nutrients such as minerals [7].
63 The eleven micronutrients that are the most important nutrients during lactation are vitamin A,
64 vitamin C, thiamine, riboflavin, niacin, vitamin B6, folate, vitamin B12, calcium, iron, and zinc.
65 Undiversified diet during lactation leads to these important micronutrients being low in breast
66 milk and reflects the poor nutritional status and inadequate dietary intake of micronutrients [2,
67 8-10].

68 Low dietary diversity is one of the most important causes of both macronutrient and
69 micronutrient deficiencies [2, 13]. Studies conducted in some Asian countries showed that low
70 dietary diversity ranges from 55.2% to 87.8% [14, 15], whereas in African country studies, low

71 dietary diversity ranges from 42.1% to 88.7% [13, 16, 17]. In Ethiopia, the low dietary diversity
72 ranges from 56.4% to 99.9% [15, 18-20].

73 The major causes of inadequate dietary diversity stem from wider contextual factors such as
74 lack of education, poor socioeconomic status, cultural beliefs, poor agricultural practices,
75 poverty, disparities in households and food insecurity. Low dietary diversity has both short-term
76 consequences, such as mortality, morbidity, disability and long-term consequences, such as
77 stunting, impaired cognitive ability, poor economic productivity, poor reproductive
78 performance, increased metabolic and cardiovascular diseases, and intergenerational
79 consequences. Hence, the consequences of micronutrient malnutrition not affect only the health
80 and survival of women but also their children [7, 11].

81 The three main strategies that have been documented for the prevention and control of hidden
82 hunger are short-term supplementation, medium-term food fortification and a long-term focus
83 on dietary diversification [21]. According to different scientific pieces of evidence, there is a
84 growing worldwide consensus to increase the dietary diversity of women (consuming more food
85 groups), especially lactating mothers, which can help to improve their diet quality and infants.
86 Since no single food group can contain all nutrients required for the healthy functioning and
87 performance of the body, there is, therefore, the need for more food groups to be included in the
88 daily diet to meet their nutrient requirements [1, 3, 9]. The government of Ethiopia also has a
89 goal to reduce maternal undernutrition through dietary diversity intervention by launching a
90 revised national nutrition program (NNP II) among women of reproductive age, including
91 lactating mothers and their children in 2016 [22]. Therefore, assessing the prevalence of
92 minimum dietary diversity and associated factors among lactating mothers is vital to improving
93 the quality of diet taken by lactating mothers.

94 Despite having many studies done elsewhere in the globe on the prevalence of minimum dietary
95 diversity among pregnant women and household level, there is a paucity of data concerning the
96 nature and relative magnitude of minimum dietary diversity among lactating mothers in
97 Ethiopia, particularly there is a lack of study in central parts of the country including the study
98 area. In addition, most of the previous studies performed across the globe, including Ethiopia,
99 did not use the newly recommended indicator, which is called minimum dietary diversity-
100 women (MDD-W), except the South African study, to assess dietary diversity of mothers.
101 Furthermore, previous studies did not address the prevalence of minimum dietary diversity at
102 the community level using this indicator. Therefore, this study was aimed at assessing the
103 prevalence of minimum dietary diversity and associated factors among lactating mothers in
104 Ataye District, North Shoa Zone, Amhara Regional State, Central Ethiopia, including variables
105 that were not included in the previous studies.

106 **2. Methods**

107 **2.1. Study setting, period and design**

108 A community-based cross-sectional study was conducted from January 25 to April 30, 2018 in
109 Ataye District. Ataye is located 270 km away from Addis Ababa, the capital city of Ethiopia.
110 The District has 30 kebeles (lowest administrative unit in Ethiopia). The district has seven health
111 centers, 24 health posts, and one District hospital. The common agricultural practices in the
112 District are *teff*, wheat, sorghum, barley, maize, peas and beans, and some vegetables, such as
113 cabbage, potato, pumpkin, tomato, onion, carrots, and lettuce, and some fruits, such as orange,
114 papaya, banana, and mango, were among others. Lactating mothers aged 15-49 years and lived
115 for at least six months and above in the District during the study period were included in the
116 study, while mothers who were not able to respond to an interview and who had an unusual

117 dietary intake in the previous 24 hours, such as feasts and celebrations, were excluded from the
118 study.

119 **2.2. Sample Size Determination and Sampling Procedures**

120 The sample size was estimated using two approaches based on the objectives of the study. To
121 determine the prevalence of minimum dietary diversity, a single population proportions formula
122 was used with the following assumptions: the proportion of maternal dietary diversity among
123 lactating mothers was 43.6% [20], $Z_{\alpha/2}$ had a 95% confidence level of 1.96, a margin of error of
124 0.05, a non-response of 10%, and a design effect of 1.5. Accordingly, the calculated sample was
125 624. On the other hand, to assess the predictors of minimum dietary diversity using a double
126 proportion formula considering different predictors significantly associated with the outcome
127 variable, such as educational status, occupational status and meal frequency, were used.

128 During calculation, assumptions such as a two-sided confidence level of 95%, a margin of error
129 of 5% and power of 80%, a 1:1 proportion of exposed to unexposed ratio, a respective odds ratio
130 for each factor, a non-response of 10% and a design effect of 1.5 were considered using the Epi
131 Info 7 software program, and then the calculated sample size was 647. The latter sample size
132 was considered to increase the power of the study. However, the total number of mothers in the
133 selected kebeles was 652. Thus, due to the nature of the cluster sampling method, all mothers
134 were included in the selected cluster using a cluster sampling technique considering kebeles as
135 clusters.

136 This study was conducted in two urban kebeles and four kebeles of rural in the District. The
137 numbers of lactating mothers found in each kebele were taken from a family folder documented
138 by the Health Extension Workers (HEWs) with respect to their household. After that, all

139 lactating mothers in randomly selected clusters were included in the study through the house-
140 to-house visit.

141 **2.3. Data Collection Methods and Instruments**

142 A structured interviewer-administered questionnaire was first prepared in English and was
143 translated to Amharic (the language spoken in the study area) for data collection and then
144 translated back to English by an independent language expert to ensure its consistency.

145 Six female grade ten completed students who were fluent speakers in the local language were
146 recruited to participate in data collection. Two-degree health professionals were recruited for
147 the supervision of the data collection procedure. The data collectors underwent a community-
148 based face-to-face interview using a structured and pretested Amharic questionnaire. The
149 mothers were interviewed (on average 30 minutes) in a private setting inside their own house.

150 **2.4. Measurements**

151 In this study, the outcome variable (minimum dietary diversity) was measured by minimum
152 dietary diversity for a woman (MDD-W), which is a dichotomous indicator/tool that has been
153 developed by FAO. A total of 16 food groups were considered in this study (i.e. cereals, white
154 tubers and roots, vitamin A-rich vegetables and tubers, dark green leafy vegetables, other
155 vegetables, vitamin A-rich fruits, other fruits, organ meat, flesh meat, eggs, fish and seafood,
156 legumes, seeds and nuts, milk and milk products, oils and fats, sweets, spices, condiments, and
157 beverages). These food groups were further regrouped into ten food groups (i.e. all starchy
158 staples, pulses (beans, peas, and lentils), nuts and seeds, all dairy, flesh foods (including organ
159 meat), eggs, vitamin A-rich dark green leafy vegetables, other vitamin A-rich vegetables and
160 fruits, other vegetables and other fruits) during analysis [3, 4].

161 The Minimum Dietary Diversity Score (MDDS) was calculated for each lactating mother during
162 the previous 24 hours to classify mother’s dietary diversity as adequate (≥ 5 food groups) or not
163 adequate (< 5 food groups) from ten food groups. Then, the outcome variable was coded as
164 minimum dietary diversity score ≥ 5 food groups, as “1” and minimum dietary diversity score
165 < 5 food groups as “0” for logistic regression analysis.

166 Household food insecurity was measured with the Household Food Insecurity Access Scale
167 (HFIAS), a structured, standardized and validated tool that was developed mainly by FANTA,
168 to classify households as food secure or not [24, 25]. A scale is to be a valid tool in measuring
169 household food insecurity among both rural and urban areas of Ethiopia with Cronbach's alpha
170 values of 0.76 for round 1 and 0.73 for round 2 [26]. In this study, nine standard Household
171 Food Insecurity Access Scale (HFIAS) questions adapted from FANTA were computed. All
172 “Yes” responses were coded “1” and “No” responses were coded “0”, and the responses were
173 summed to obtain the household food insecurity status. The HFI status, which had a high internal
174 consistency (Cronbach’s alpha = 0.927), was further dichotomized as “food secure” and “food
175 insecure households which coded as “1” and as “0”, respectively, during analysis. Furthermore,
176 the household wealth index was analyzed using principal component analysis (PCA) by using
177 approximately 20 locally available household assets considering the assumptions.

178 The knowledge of mothers about nutrition was computed based on six questions that included
179 awareness of the mothers about nutrition and dietary diversity practice. The mothers who scored
180 above the mean cut-off point were considered to have good knowledge and coded as “1”,
181 whereas those who scored below this cut-off point were considered to have poor knowledge
182 and coded as “0”.

183 **2.5. Data Quality Control**

184 The adapted and further developed English language questionnaire was translated and
185 contextualized into the Amharic language by an Amharic language speaker who has attended a
186 Master of Arts in the Amharic language. It was translated back to the English language by a
187 person who attended Master of Arts in the English language, and comparison was made on the
188 consistency of the two versions. Data were collected using the translated and pretested Amharic
189 version structured questionnaire. The questionnaire was modified further after a pretest was
190 conducted. During data collection, the interviewers informed mothers about all the details of the
191 research. The mothers were encouraged to feel free and told that the confidentiality of their
192 responses was assured and that no information was shared with anybody without the
193 investigator. After this, women who were willing to participate and signed the informed consent
194 document were interviewed.

195 Two days of training (both theoretical and practical) was given for six female grade ten
196 completed students and the two supervisors. The training was focused on interview technique,
197 ethical issues, rights of the participants, reading through all the questions and understanding
198 them and ways of decreasing under/over-reporting and maintaining confidentiality.

199 Three weeks before the actual data collection, the questionnaire was pretested outside to the
200 selected kebele on 5% of the total sample size to ensure the validity of the tool. After the pretest
201 was performed, all the necessary adjustments were made. To minimize bias, interviews were
202 conducted in an area with adequate confidentiality, privacy and without the involvement of any
203 other person other than the respondent. On-site supervision was carried out during the whole
204 period of data collected daily by the supervisors and principal investigators. At the end of each
205 day, questionnaires were reviewed and cross-checked for completeness, accuracy, and

206 consistency by the supervisor and principal investigator, and in the end, corrective measures
207 were taken.

208 **2.6. Statistical Analysis**

209 All the interviewed questionnaires were checked visually by the principal investigator. Data
210 were coded, entered and cleaned using EpiData version 4.2.0.0 software. Double data entry was
211 performed by two data clerks to cross-check the data for completeness before analysis. The
212 entered data were exported and analyzed with Statistical Package for Social Science (SPSS)
213 version 24 software. Simple descriptive statistics, such as simple frequency distribution,
214 measures of central tendency, measures of variability and percentages, were performed to
215 describe the demographic, socioeconomic, and maternal-related characteristics of the
216 respondents. Then, the information was presented using tables and figures. The continuous
217 variables were tested for normal distribution using a histogram, Q-Q plot and some statistical
218 tests such as Kolmogorov-Smirnov and Shapiro-Wilk tests.

219 Bivariable analysis and crude odds ratio along with a 95% confidence interval (CI) were used
220 to see the association between each independent variable and the outcome variable by using
221 binary logistic regression. Independent variables with a p-value of ≤ 0.25 were included in the
222 multivariable analysis to control for confounding factors. Multicollinearity was checked to see
223 the linear correlation among the independent variables by using standard error (SE). Variables
224 with a standard error of ≥ 2 were dropped from the multivariable analysis. The fitness of the
225 model was tested by Hosmer-Lemeshow's goodness-of-fit test model coefficient with an enter
226 method, which was found to be insignificant with a large P-value ($P=0.374$) and the Omnibus
227 tests were significant ($P=0.0001$). The adjusted odds ratio along with 95% CI was estimated to
228 identify the factors associated with dietary diversity using multivariable logistic regression

229 analysis. All tests were two-sided, and the level of statistical significance was declared at a p-
230 value less than 0.05.

231 The ethical approval letter was obtained from Haramaya University, College of Health and
232 Medical Science Institutional Health Research Ethics Review Committee (IHRERC). The
233 approval letter was dated 15 January 2018 and numbered Ref C/AC/R/D/897/18. Before
234 informed consent was obtained, a clear description of the study title, purpose, procedure,
235 duration, possible risks and benefits of the study was explained for each study participant. Their
236 rights during the interview were guaranteed. Then written and signed informed consent was
237 obtained from each respondent before starting the interview. Questionnaire code numbers were
238 used to maintain the confidentiality of information gathered from each study participant
239 throughout the study.

240 **3. Results**

241 **3.1. Sociodemographic Characteristics of Study Participants**

242 Out of 652 planned lactating mothers, 631 of them aged 15-49 years participated in a response
243 rate of 96.8%. The mean age (\pm SD) of lactating mothers was 30.48 (\pm 6.36) years. From the
244 total of participating, respondents, 417 (66.1%) were rural by residence, 345 (54.7%) were in
245 the age group 15-30, 545 (86.4%) were Orthodox Christian by religion, 566 (89.7%) were
246 Amhara by ethnicity and 501 (79.1%) were married by their marital status (Table 1).

247 Table 1: Sociodemographic characteristics of lactating mothers in Ataye District, Central
248 Ethiopia, 2018 (n = 631)

Variables	Frequency	Percentage
-----------	-----------	------------

Residence		
Urban	214	33.9
Rural	417	66.1
Age		
15-30	345	54.7
31-40	237	37.6
>40	49	7.7
Religion		
Orthodox Christian	545	86.4
Muslim	47	7.4
Protestant	39	6.2
Ethnicity		
Amhara	566	89.7
Oromo	49	7.8
Tigray	11	1.7
Other*	5	0.8
Maternal education status		
No formal education	265	42.0
Read and write	136	21.5
Primary education	99	15.7
Secondary education	51	8.1
More than secondary education	80	12.7
Maternal occupation		

Housewife	393	62.3
Merchant	74	11.7
Daily laborer	54	8.6
Government employ	54	8.6
Self-employed	52	8.2
Others**	4	0.6
Marital status		
Single	39	6.2
Married	501	79.4
Divorced	54	8.6
Widowed	26	4.1
Separated	11	1.7
Sex of household head		
Male	513	81.3
Female	118	18.7
Maternal decision autonomy		
Involved	384	60.8
Not involved	250	39.6
Type of decision autonomy involved(n=384)		
Final say on household purchase	331	86.2
Final say on how to spend money	184	29.2
Other decision***	15	2.4
Income-generating activities		

Yes	384	60.8
No	247	39.2
Household wealth index		
Poor	267	42.4
Medium	170	26.9
Rich	194	30.7

249 *= Gurage, Argobba **=student, NGO employs ***=final say on total household activity

250 3.2. Maternal Health-care and Feeding Practice

251 More than half of lactating mothers 354 (56.1%) had one to three frequency of ANC visits in
 252 their previous/last pregnancy. Among those who had ANC visits, 297 (47%) delivered in the
 253 health center and 119 (18.9%) delivered in a hospital (Table 2).

254 Table 2: Maternal healthcare and feeding practices of lactating mothers in Ataye district, Central
 255 Ethiopia, 2018 (n = 631)

Variables	Minimum Dietary Diversity		Frequency (%)
	≥5 FGs (%)	<5 FGs (%)	
Number of ANC visit			
No visit	32 (5.1)	87 (13.8)	119 (18.9)
1-3	174 (27.6)	180 (28.5)	354 (56.1)
≥4	102 (16.2)	56 (8.8)	158 (25.0)
Place of delivery			
Home	78 (12.4)	137 (21.7)	215 (34.1)
Health center	144 (22.8)	153 (24.2)	297 (47.0)

Hospital	86 (13.6)	33 (5.3)	119 (18.9)
Postnatal follow-up			
Yes	189 (57.3)	141 (42.7)	330 (52.3)
No	119 (39.5)	182 (60.5)	301 (47.7)
Source of food			
Own production	146 (23.1)	219 (34.7)	365 (57.8)
Purchasing	157 (24.9)	99 (15.7)	256 (40.6)
Food aid/relief	5 (0.8)	3 (0.5)	8 (1.3)
Other sources*	1 (0.15)	1 (0.15)	2 (0.3)
Home gardening practice			
Yes	179 (28.4)	81 (12.8)	260 (41.2)
No	129 (20.4)	242 (38.4)	371 (58.8)
Meal frequency			
Two meals only or below	21 (3.3)	32 (5.1)	53 (8.4)
Three meals	197 (31.2)	237 (37.6)	434 (68.8)
Three meals above	90 (62.5)	54 (37.5)	144 (22.8)
Illness in the last 2 weeks			
Yes	74 (11.7)	169 (26.8)	243 (38.5)
No	234 (37.1)	154 (24.4)	388 (61.5)

256 * = no formal source like helping from others/relatives, FGs = food groups

257 3.3. Minimum Dietary Diversity Score of lactating mothers

258 The prevalence of the minimum dietary diversity score (≥ 5 food groups) among lactating
 259 mothers was 48.8 (95% CI: (44.7%, 52.9%)) during the preceding 24 hours of the survey. The

260 majority of the study participants had consumed all starchy staple foods (99.4%) and pulses
261 (beans and peas) (79.1%). The least consumed food group was other fruits (8.7%) (**Figure 1**).

262 **3.4. Knowledge about nutrition**

263 Regarding knowledge about nutrition, of the total participants in this study, 54.8% (95% CI:
264 50.9, 58.5) had good knowledge of nutrition.

265 **3.5. Household food security status**

266 The household food security status of the study participants was 75.9%, 95% CI: (72.4, 79.3)
267 food secure, while 24.1%, 95% CI: (20.7, 27.6) were food insecure households.

268 **3.6. Factors Associated with Minimum Dietary Diversity among Lactating Mothers**


269 In the bivariable logistic regression analysis model, educational status of lactating mothers,
270 mothers who had the final say on a household purchase, mothers who had income-generating
271 activities, increased number of ANC visits, place of delivery, postnatal follow-up, the main
272 source of food, home gardening practice, history of illness, daily eating pattern, knowledge of
273 nutrition, household food insecurity, and household wealth indices were significantly associated
274 with minimum dietary diversity of mothers and became candidates for further multivariable
275 analysis.

276 In the multivariable logistic regression analysis model, those variables with a P-value of ≤ 0.25
277 in the bivariable analysis were entered into a multivariable analysis using the enter method to
278 identify the independent predictors of the minimum dietary diversity score. After adjusting for
279 all possible confounders, the formal educational status of mothers, maternal decision-making
280 autonomy related to a final say on a household purchase, home gardening practice, good
281 knowledge of nutrition, illness in the last two weeks, food-secure households and medium and

282 rich household wealth indices remained significant and independent predictors of the minimum
283 dietary diversity score of mothers.

284 Lactating mothers who had formal education were more than two times [(AOR=2.16, 95% CL:
285 (1.14, 4.09)] more likely to have minimum dietary diversity scores than those who had no formal
286 education. Regarding maternal decision-making autonomy, mothers who had the final say on
287 household purchases were more than five times [(AOR=5.39, 95% CI: (2.34, 12.42)] more likely
288 to achieve minimum dietary diversity than their counterparts. The odds of minimum dietary
289 diversity scores were nearly three times higher among mothers who had home gardening
290 practice [(AOR=2.67, 95% CI: (1.49, 4.81)] than their counterparts.

291



292 Figure 1: The ten grouped food groups consumed by lactating mothers in Ataye District, Central
293 Ethiopia, 2018 (n=631)

294 Mothers who had a history of illness in the last two weeks were 53% [(AOR=0.47, 95% CI:
295 (0.26, 0.85)] less likely to reach a minimum dietary diversity score than their counterparts. The
296 odds of minimum dietary diversity score were more than five times [(AOR=5.11, 95% CI: (2.68,
297 9.78)] more common among mothers who had good knowledge of nutrition than those who had
298 poor knowledge of nutrition. Food-secure households had nearly three times the minimum
299 dietary diversity scores than food-insecure households [(AOR=2.96, 95% CI: (1.45, 6.07)].

300 Mothers from households who had medium and rich wealth indices were nearly six times and
 301 more than three times more likely to have minimum dietary diversity experience than poor
 302 households [(AOR=5.94, 95% CI: (2.82, 12.87)] and [(AOR= 3.55, 95% CI: (1.76, 7.13)],
 303 respectively (Table 3).

304 Table 3: Bivariable and multivariable logistic regression¹ analysis of factors associated with
 305 MDD among lactating mothers in Ataye district, Central Ethiopia, 2018 (n=631).

Variables	Minimum DD		COR (95% CI)	AOR (95% CI)
	≥5 FGs	<5 FGs		
	N ₀ (%)	N ₀ (%)		
Maternal education				
Formal education	182(66.2)	93(33.8)	3.57(2.56,4.97) **	2.16(1.14,4.09) *
No formal education	126(35.4)	230(64.6)	1.00	1.00
Final say on household purchase (n=384)				
Yes	211(63.7)	120(36.3)	3.72(2.01,6.91) **	5.39(2.34,12.42) **
No	17(32.1)	36(67.9)	1.00	1.00
Meal frequency				
Two meals only or below	21(39.8)	32(60.4)	1.00	1.00
Three meals	197(45.4)	237(54.6)	1.27(0.7,.2.27)	0.59(0.19,1.85)
Three meals above	90(62.5)	54(37.5)	2.54(1.33,4.84) *	0.68(0.20,2.31)
Income-generating activities				
Yes	205(53.4)	179(46.6)	1.60(1.16,2.21)	0.56(0.26,1.22)
No	103(41.7)	144(58.3)	1.00	1.00

Place of delivery				
Health institutions	230(55.2)	187(44.8)	2.15(1.53, 3.01) **	0.53(0.21,1.35)
Home	78(36.4)	136(63.6)	1.00	1.00
Postnatal follow-up				
Yes	189(57.3)	141(42.7)	2.05(1.49,2.82) **	1.56(0.81,2.99)
No	119(39.5)	182(60.5)	1.00	1.00
Number of ANC visits				
≥4	102(64.6)	56(35.4)	4.95(2.94,8.33) **	3.32(0.85, 12.87)
1-3	174(49.2)	180(50.8)	2.63(1.67,4.15) **	2.43(0.74,7.93)
No visit	32(26.9)	87(73.1)	1.00	1.00
Home gardening practice				
Yes	179(68.2)	81(31.2)	4.15(2.95,5.82) **	2.67(1.49,4.81) **
No	129(34.8)	242(65.2)	1.00	1.00
Source of food				
Another source	162(60.9)	104(39.1)	1.00	1.00
Own source	146(40.0)	219(60.0)	2.34(1.69,3.23) **	1.29(0.59,2.81)
History of illness				
Yes	74(30.5)	169(69.5)	0.28(0.21,0.41)	0.47(0.26,0.85) **
No	234(60.5)	154(39.5)	1.00	1.00
Knowledge about nutrition				
Good	234(67.6)	112(32.4)	5.96(4.21,8.43) **	5.11(2.68,9.78) **
Poor	74(26.0)	211(74.0)	1.00	1.00
Household food insecurity				

Food secure	269(56.2)	210(43.8)	3.71(2.47,5.57) **	2.96(1.45,6.07) **
Food insecure	39(25.7)	113(74.3)	1.00	1.00
Household wealth index				
Rich	117(60.3)	77(39.7)	3.62(2.45,5.34) **	3.55(1.76,7.13) **
Medium	112(65.9)	58(34.1)	4.59(3.04,6.94) **	5.94(2.82,12.87) **
Poor	79(29.6)	188(70.4)	1.00	1.00

306 * = P-value < 0.05, ** = P-value < 0.001, CI = Confidence Interval, COR = Crude Odds Ratio,

307 AOR = Adjusted odds ratio, FGs = food groups, ¹Hosmer and Lemeshow test (P-value = 0.374)

308 **4. Discussion**

309 In this study, approximately 48.8% of lactating mothers achieved a minimum dietary diversity
 310 score, and they were more likely to have higher (more adequate) micronutrient intakes than
 311 51.2% of lactating mothers who did not. This result indicated that the prevalence of minimum
 312 dietary diversity in the study area is low. The 2016 FAO guideline recommended that all of the
 313 mothers should achieve the minimum dietary diversity score (≥ 5 food groups) to have adequate
 314 micronutrient adequacy (3). The present study also examined the demographic and
 315 socioeconomic characteristics, household-related characteristics, maternal healthcare and
 316 feeding practice-related characteristics associated with minimum dietary diversity among
 317 lactating mothers in Ataye District. Accordingly, attending formal education, having decision-
 318 making autonomy on household purchases, having home gardening practices, having good
 319 knowledge of nutrition, having an illness in the last two weeks, being from food secure
 320 households and being in medium and rich household wealth indices were independent predictors
 321 of minimum dietary diversity.

322 The prevalence of minimum dietary diversity is relatively higher than the study findings carried
323 out elsewhere in the world such as Bangladesh (12.8%) and Vietnam (22%) [15], South Africa
324 (25%) [16], Ghana (43.1%) [13], Aksum Town (43.6%) [20] and Jimma (32.8%) [18]. In
325 contrast, this prevalence is lower than that in the study performed in Uganda (57.9%) [17]. This
326 inconsistency could be due to the difference in the sample size because some of the studies were
327 large surveys, they had a relatively large sample size (study done in Bangladesh and Vietnam)
328 and some of the studies also had a lower sample size than the present study (study done in South
329 Africa and Aksum). The study period and seasonal variation (since this study was done during
330 the postharvest season) may also be another explanation for the observed discrepancy in which
331 it can result in food security status change, while most of the previous studies had a high
332 prevalence of food insecurity compared to the present study, which is nearly one-fourth.
333 Methodological (measurement tool) differences may be another possible explanation because
334 the current study uses the newly recommended indicator of ten food groups rather than the nine
335 food groups' indicators. The feeding culture, educational status and socioeconomic status of the
336 study participants were also other explanations for this observed variation.

337 The educational status of mothers was associated with a minimum dietary diversity score of
338 lactating mothers. Lactating mothers who had formal education were more than two times highly
339 dietary diversified mothers than those who had no formal education. This finding is similar to
340 that of prior studies done in Ghana [13] and three countries (Bangladesh, Vietnam, and Ethiopia)
341 [15]. This could be related to the fact that educated mothers may obtain sufficient information
342 about varieties of food during their health service utilization and understanding the importance
343 of increasing dietary diversity and meal frequency during lactation than non-educated mothers
344 in case they would have the probability of taking a variety of food groups.

345 Minimum dietary diversity was associated with maternal decision-making autonomy practice.
346 Mothers who had the final say on household purchase were more than five times more likely to
347 achieve minimum DDS compared to those who had no final say on household purchases. This
348 figure is in line with the finding of Ghana, women who had a final say were almost twice as
349 likely to achieve a higher DD score [13]. This consistency may be because as the mother's
350 involvement is increased in the family on a nutrition-related decision, they have the probability
351 of purchasing and deciding on an important variety of foods for the households, their child and
352 themselves, as a result, they may have increased diversify diets.

353 In the present study, the home gardening practice of lactating mothers was found to be an
354 independent predictor of the minimum DDS of mothers. Lactating mothers who practice home
355 gardening showed nearly three-fold higher likelihood of consuming highly diversified diets than
356 their counterparts. This is comparable with the study conducted in Aksum town, Ethiopia [20],
357 which demonstrated that home gardening was strongly associated with diet diversity. This might
358 be because mothers from households with gardens would grow vegetables and fruits and become
359 beneficiary from gardens to diversify their daily food intake; therefore, they may obtain
360 additional diet options that enhance the diversity of the household's food consumption of
361 different food groups in their diets.

362 Good maternal knowledge of nutrition predicted the minimum dietary diversity score of
363 lactating mothers in the current study area. Mothers who scored above the mean cut-off point
364 upon responding to nutrition and food diversity knowledge questions were more than five times
365 more likely to achieve a minimum dietary diversity score compared to their counterparts. This
366 result is comparable with the study done in three countries (Bangladesh, Vietnam, and Ethiopia)
367 and Uganda [5, 17]. This may be because mothers who had increased knowledge about nutrition

368 may know the importance of varieties of foods and have a greater likelihood of taking these
369 varieties of food groups. It could also be related to mothers getting enough nutrition and dietary
370 diversity-related information and education when they have good access to health services
371 during their antenatal and postnatal follow up. This helps them to increase their knowledge about
372 varieties of food groups, and as a result, they may have adequate intake and consumption.

373 Maternal history of illness was associated with minimum DDS of mothers. Mothers who had a
374 history of illness were 53% less likely to have the probabilities of achieving minimum DD scores
375 when compared to their counterparts. A possible explanation could be that illness may decrease
376 appetite and intake of the diet which in turn affects the dietary intake and dietary diversity of
377 mothers. In addition, most of the respondents (nearly two-thirds) in the present study were rural
378 residents and they may not receive enough health service in the nearby health facility
379 immediately as they worsened, leading to a decrease in the minimum DDS mothers.

380 Another important predictor of the minimum dietary diversity score of mothers was household
381 food insecurity. Lactating mothers from food-secure households had nearly three-fold higher
382 probabilities of achieving a high level of MDDS than food-insecure households. This is
383 supported by previous studies in three countries Bangladesh, Vietnam, and Ethiopia [15]. This
384 could be explained in such a way that food security promotes the intake of adequate quantity
385 and quality of diet that in turn contributes to having a better minimum dietary diversity score
386 and nutritional status.

387 The odds of MDD score was more common among mothers found in medium and rich wealth
388 indices. Mothers belonging to medium and rich socioeconomic status households were nearly
389 six times and four times more likely to have minimum dietary diversity scores compared to poor
390 socioeconomic status households respectively. This finding is consistent with previous results

391 in three countries (Bangladesh, Vietnam, and Ethiopia) [15], South Africa [16] and Ghana [13].
392 The possible explanation behind this may be that as the socioeconomic status of the family
393 increases, they would have stable food security status in case their probabilities of eating
394 diversified food groups would also increase.

395 In summary, this finding will also support one of the strategic objectives of the National
396 Nutrition Program II (NNP-II) in Ethiopia, which was launched to improve the nutritional status
397 of women of reproductive age, including adolescent girls, pregnant women, and lactating
398 mothers, through appropriate dietary diversity to alleviate the current burden of micronutrient
399 deficiencies [22].

400 Regarding the strength and limitations of the study, the major strength of the current study is
401 that using the newly recommended 10-point food group indicator, MDD-W, instead of the
402 previous 9-point food group using a community-based cross-sectional study design was the
403 limitation of other studies. However, recall bias was one of the limitations of the study since
404 some of the questions we asked about the event that occurred 24 hours and 4 weeks back. This
405 was minimized by probing the respondents about the event. Seasonal variation was also another
406 limitation of the study. This study was also affected by reporting and social desirability bias.
407 Therefore, this study could not reject the possibility of reporting bias from lactating mothers
408 who did not eat. To minimize these biases lactating mothers were asked in a private setting.

409 **5. Conclusion**

410 Nearly half of the lactating mothers achieved the minimum dietary diversity score in the current
411 study. Regarding the predictors of minimum dietary diversity score, attending formal education,
412 having decision autonomy on the household purchase, having home gardening practice, having

413 good knowledge of nutrition, having illness in the last two weeks, being food secure household
414 and being in medium and rich household wealth indices were the significant and independent
415 predictors of minimum dietary diversity. Therefore, the government of Ethiopia, particularly the
416 Ataye District Health Office, should be prioritizing, planning, designing and initiating dietary
417 diversity intervention programs aimed at improving maternal nutrition through appropriate
418 food-based approaches. Strengthen nutrition education programs on proper maternal MDD
419 practices and adequate dietary intake during lactation to improve the nutrition outcomes of
420 lactating mothers. Cooperatively work with health extension workers to increase the awareness
421 of lactating mothers on how to improve the dietary diversity practice of mothers. Create strong
422 multisectoral collaboration targeted at improving the mother's educational status, decision-
423 making autonomy (women empowerment) and family socioeconomic status. District Health
424 Extension Workers (HEWs) should also facilitate and promote nutrition education about dietary
425 diversification for lactating mothers and the community at large. Educate mothers about visiting
426 health service institutions when they become sick and about increasing their intake of varieties
427 of food. Work in cooperation with developmental agents to increase the minimum dietary
428 diversity score of lactating mothers in the District. Local NGOs should actively work in
429 collaborating with the woreda health office to increase the awareness and knowledge of lactating
430 mothers about food diversity and nutrition. Collaboratively work with another stakeholder to
431 increase the stability of the HFS and socioeconomic status of the mother's family and the whole
432 community to prevent inadequate DD practice of the mother's, thereby preventing poor
433 nutritional status. Finally, the scientific community should study a different study design, such
434 as a prospective cohort study design, that is recommended to address seasonal variability during
435 the preharvest and postharvest seasons. Since dietary diversity is multifactorial and

436 multifaceted, further study is needed to identify other independent predictors for lactating
437 mothers' dietary diversity.

438 **Abbreviations**

439 DDS: Dietary Diversity Score

440 EDHS: Ethiopian Demographic and Health Survey

441 FAO: Food and Agricultural Organization of the United Nation

442 FANTA: Food and Nutrition Technical Assistant

443 HFIAS: Household Food Insecurity Access Scale

444 MDDS: Minimum Dietary Diversity Score

445 MDD-W: Minimum Dietary Diversity for Women

446 WHO: World Health Organization

447 **Data Availability**

448 The data used to support the findings of this study are available from the corresponding author
449 upon reasonable request.

450 **Competing interests**

451 Authors are declared that they have no competing interests.

452 **Disclosure**

453 This manuscript was prepared from the thesis work of Mr. Lemma Getacher. The cost of
454 conducting this research was covered by him.

455 **Author's contribution**

456 LG: - participated in the design of the study, performed the data collection, performed the
457 statistical analysis and served as the lead author of the manuscript. GE: - participated in the
458 design of the study, revised subsequent drafts of the paper and contributed to the finalization of

459 the manuscript. TA, AB, and AM: - participated in the design of the study, statistical analysis
460 and finalization of the manuscript. All authors read and approved the final manuscript.

461 **Acknowledgments**

462 We would like to express our deepest gratitude to Haramaya University, College of Health and
463 Medical Science for ethical clearance. Our great thanks also deserve to our study participants,
464 data collectors, supervisors and language translators for their invaluable contribution to this
465 study.

466 **References**

467 [1] USAID (United States Agency for Development). Maternal dietary diversity and the
468 implications for children's diets in the context of food security. USAID 'S Infant and Young
469 Child Nutrition Project. 2012. Available at: [http://iycn.wpengine.netdna-](http://iycn.wpengine.netdna-cdn.com/files/IYCN_Brief_Maternal_Dietary_Diversity0112.pdf)
470 [cdn.com/files/IYCN_Brief_Maternal_Dietary_Diversity0112.pdf](http://iycn.wpengine.netdna-cdn.com/files/IYCN_Brief_Maternal_Dietary_Diversity0112.pdf) Accessed on 20,
471 December 2017.

472 [2] Arimond, M., Wiesmann, D., Becquey, E., Alicia, C., Melissa, D., Megan, D. et al. Dietary
473 diversity as a measure of the micronutrient adequacy of women's diets in resource-poor
474 areas: Summary of results from five sites. Washington, DC: FANTA-2 Bridge, FHI360:14-
475 15. 2011. Available at:
476 [https://www.fantaproject.org/sites/default/files/resources/WDDP_Summary_Report_Jul20](https://www.fantaproject.org/sites/default/files/resources/WDDP_Summary_Report_Jul2011.pdf)
477 [11.pdf](https://www.fantaproject.org/sites/default/files/resources/WDDP_Summary_Report_Jul2011.pdf) Accessed on November 21, 2017.

478 [3] FAO(Food and Agriculture Organization). Guidelines for measuring household and
479 individual dietary diversity. FAO. Rome. 2011. Available at:

480 http://agrobiodiversityplatform.org/files/2011/05/guidelines_MeasuringHousehold.pdf

481 Accessed on December 21, 2018.

482 [4] FAO (Food and Agriculture Organization). Minimum Dietary Diversity for Women A Guide
483 to Measurement. Rome. FAO and FANTA. 2016. Available at:
484 <https://index.nutrition.tufts.edu/printpdf/160> Accessed on 21, December 2017.

485 [5] FAO (Food and Agriculture Organization) and FANTA (Food and Nutrition Technical
486 Assistant). Introducing the Minimum Dietary Diversity–Women (MDD-W) Global Dietary
487 Diversity Indicator for Women. FAO. Washington, DC. 2014. Available at:
488 [https://www.fantaproject.org/sites/default/files/resources/Introduce-MDD-W-indicator-](https://www.fantaproject.org/sites/default/files/resources/Introduce-MDD-W-indicator-brief-Sep2014_0.pdf)
489 [brief-Sep2014_0.pdf](https://www.fantaproject.org/sites/default/files/resources/Introduce-MDD-W-indicator-brief-Sep2014_0.pdf) Accessed on 21, December 2017.

490 [6] Hailelassie, K., Afework, M. and Meron, G. Feeding practices, nutritional status and
491 associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray,
492 Ethiopia. *Nutrition Journal*. 2013.12(28).

493 [7] Bailey, R. L., Keith, B. P., West, J. B. and Robert, E. B. The Epidemiology of Global
494 Micronutrient Deficiencies. *Ann Nutr Metab*. 2015. 66(2): 22–33.

495 [8] Allemand, P. Reaching Consensus on a Global Dietary Diversity Indicator for Women,
496 Washington, DC: 2014.1-6. Available at:
497 [http://www.fao.org/fileadmin/templates/nutrition_assessment/Dietary_Diversity/AG2NUT](http://www.fao.org/fileadmin/templates/nutrition_assessment/Dietary_Diversity/AG2NUT_JULY30_ballard.pdf)
498 [_JULY30_ballard.pdf](http://www.fao.org/fileadmin/templates/nutrition_assessment/Dietary_Diversity/AG2NUT_JULY30_ballard.pdf) Accessed on November 21, 2017.

499 [9] Martin-Prévela, Y., Allemanda, P., Wiesmannb, D., Arimond, M., Ballardd, T., Megan, D.
500 et al. Moving forward on choosing a standard operational indicator of women’s dietary
501 diversity. *FAO*. 2015. Available at: <https://cgspace.cgiar.org/bitstream/handle>

502 [/10568/72450/FAO_2015_MDD-W.PDF?sequence=1&isAllowed=y](#). Accessed on 21,
503 December 2017.

504 [10] WHO (World Health Organization). Nutrition of women in the preconception period, during
505 pregnancy and the breastfeeding period. Geneva, WHO. Geneva. 2012. Available at:
506 https://apps.who.int/iris/bitstream/handle/10665/78900/A65_12en.pdf?sequence=1&isAllowed=y
507 Accessed on 20, December 2017.

508 [11] IFPRI (International Food Policy Research Insititute). Global Nutrition Report 2015.
509 Actions and accountability to advance nutrition & sustainable development. Washington,
510 DC. 2015. Available at: <http://www.ifpri.org/publication/global-nutrition-report-2015>
511 Accessed on 21, December 2017.

512 [12] CSA(Central Statistical Agency). Ethiopia Demographic and Health Survey 2016. CSA
513 and ICF International. 2016. Addis Ababa, Ethiopia, and Rockville, Maryland, USA.
514 Available at: <https://dhsprogram.com/pubs/pdf/FR328/FR328.pdf> Accessed on 21,
515 December 2017

516 [13] Amugsi, D. A., Lartey, A., Kimani, E., and Mberu, B. U. Women's participation in
517 household decision-making and higher dietary diversity: findings from nationally
518 representative data from Ghana. *J Health Popul Nutr*. 2016. 35(1): 16.

519 [14] Henjum, S., Torheim, L. E., Thorne-Lyman, A. L., Chandyo, R., Fawzi, W. W.,
520 Shrestha, P. S. et al. Low dietary diversity and micronutrient adequacy among lactating
521 women in a peri-urban area of Nepal. *Public Health Nutr*. 2015.18(17): 3201-10.

- 522 [15] Nguyen, P. H., Avula, R., Ruel, M. T., Saha, K. K., Ali, D., Tran, L. M. et al. Maternal
523 and child dietary diversity are associated with child dietary diversity in Bangladesh,
524 Vietnam, and Ethiopia. *J Nutr.* 2013.143(7): 1176-83.
- 525 [16] Chakona, G. and Shackleton, C. Minimum Dietary Diversity Scores for Women Indicate
526 Micronutrient Adequacy and Food Insecurity Status in South African Towns. *Nutrients.*
527 2017.9(8).
- 528 [17] Ray, O., Wamani, H., and Bukenya, J. Factors associated with dietary diversity among
529 women of reproductive age (15-49 years) in agago district. *Nurs Times.* 2014.1(2).
- 530 [18] Alemayehu, M., Alemayehu, A. and Abebe, G. Factors Associated with Malnutrition
531 among Lactating Women in Subsistence Farming Households from Dedo and Seqa-
532 Chekorsa Districts, Jimma Zone. *Developing Country Studies*, 2015. 5(21): 11-12.
- 533 [19] Sitotaw, I. K., Hailesslasie, K., and Adama, Y. Comparison of nutritional status and
534 associated factors of lactating women between lowland and highland communities of
535 District Raya, Alamata, Southern Tigray, Ethiopia. *BMC Nutrition.* 2017. 3(61).
- 536 [20] Weldehaweria, N. B., Misgina, K. H., Weldu, M. G., Gebregiorgis, Y. S., Gebrezgi, B.
537 H., Zewdie, S. W. et al. Dietary diversity and related factors among lactating women visiting
538 public health facilities in Aksum town, Tigray, Northern Ethiopia. *BMC Nutrition*, 2016.
539 2(1): 10-12.
- 540 [21] Nair, M. K., Augustine, L. F. and Konapur, A. Food-Based interventions to Modify Diet
541 Quality and Diversity to Address Multiple Micronutrient Deficiency. *Environmental Health,*
542 *a section of the Journal of Frontiers in Public Health*, 2016. 3(277): 10-13.

- 543 [22] NNP II (National Nutrition Programme II). The government of the Federal Democratic
544 Republic of Ethiopia National Nutrition Programme (NNP) 2016-2020. NNP Implementing
545 Sectors' Declaration. 2016. Available at: [https://unicefethiopia.org/2016/12/20/new-](https://unicefethiopia.org/2016/12/20/new-national-nutrition-programme-ii-envisions-an-ethiopia-free-of-malnutrition/)
546 [national-nutrition-programme-ii-envisions-an-ethiopia-free-of-malnutrition/](https://unicefethiopia.org/2016/12/20/new-national-nutrition-programme-ii-envisions-an-ethiopia-free-of-malnutrition/) Accessed on
547 21, December 2017.
- 548 [23] CSA(Central Statistical Agency). 2008. Population and housing census atlas of Ethiopia:
549 Central Statistical Agency. Available at: [http://www.csa.gov.et/census-report/complete-](http://www.csa.gov.et/census-report/complete-report/census-2007)
550 [report/census-2007](http://www.csa.gov.et/census-report/complete-report/census-2007) Accessed on 21, December 2017
- 551 [24] Coates, J., Swindale, A. and Blinisky, P. Household Food Insecurity Access Scale
552 (HFIAS) for Measurement of Food Access: Indicator Guide VERSION 3 Washington, D.C.:
553 Food and Nutrition Technical Assistance Project, Academy for Educational Development.
554 2007. http://www.fao.org/fileadmin/user_upload/eufao-fsi4dm/doc-training/hfias.pdf
555 Accessed on 15, November 2017.
- 556 [25] Deitchle, M., Terri, B., Swindale, A. and Jennifer, C. Validation of a Measure of
557 Household Hunger for Cross-Cultural Use. Washington, DC: Food and Nutrition Technical
558 Assistance II Project (FANTA-2), AED. 2010. Available at
559 [https://www.fantaproject.org/sites/default/files/resources/HHS_Validation_Report_May20](https://www.fantaproject.org/sites/default/files/resources/HHS_Validation_Report_May2010_0.pdf)
560 [10_0.pdf](https://www.fantaproject.org/sites/default/files/resources/HHS_Validation_Report_May2010_0.pdf) accessed on November 30, 2017.
- 561 [26] Hagos, S., Lunde, T., Damen, H., Tasew, W. and Lindtjorn, B. Is the adapted Household
562 Food Insecurity Access Scale (HFIAS) developed internationally to measure food insecurity
563 valid in urban and rural households of Ethiopia?, *BMC Nutrition*, 2015. 1(2).

564 [27] Ongosi, A. N., Gericke, G., Mbutia , E. and Oelofse, A. Food variety, dietary diversity,
565 and perceived hunger among lactating women (0-6 months postpartum) in a low socio-
566 economic area in Nairobi, Kenya. *African Journal of Food, Agriculture, Nutrition and*
567 *Development*, 2014.14(2).