

Clinical characteristics of 4490 COVID-19 patients in Africa: A meta-analysis

Testimony Jesupamilerin Olumade^{1,2,4} and Leonard Ighodalo Uzairue^{3,4*}

1-Department of Biological Sciences, Redeemer's University, Ede, Osun State.

2 -African Centre of Excellence for Genomics of Infectious Diseases, Redeemer's University, Ede, Osun state.

3-Department of Microbiology, College of Biosciences, Federal University of Agriculture, Abeokuta, Ogun-State Nigeria.

4-Research Hub Nigeria, Kuwba, Abuja, FCT, Nigeria

* - Corresponding author

Email address: uzairue.leonard@gmail.com or uzairue.leonard@researchhub.ng

Full postal address: Department of Microbiology, College of Biosciences, Federal University of Agriculture, Abeokuta, Ogun-State Nigeria.

Running Head: Clinical Characteristics of COVID-19 Patients in Africa.

Word Count: 1764 words

ABSTRACT

Introduction: The novel coronavirus disease-2019 (COVID-19) pandemic that started in December 2019 has affected over 39 million people and killed over 1.1 million people. While more studies are published to help us understand the virus, there is a dearth of studies on the clinical characteristics and associated outcomes of the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) on the African continent.

Methods: We evaluated evidence from previous studies in Africa available in six databases between January 1 and October 6, 2020. Meta-analysis was then performed using Open-Meta Analyst software.

Results: A total of seven studies including 4490 COVID-19 patients were included. The result of the meta-analysis showed 68.8% of infected patients were male. Common symptoms presented (with their incidences) were fever (42.8%), cough (33.3%), headache (11.3%), breathing problems (16.8%). Other minor occurring symptoms included diarrhea (7.5%), and rhinorrhea (9.4%). Fatality rate was 5.6%.

Conclusion: This study presents the first description and analysis of the clinical characteristics of COVID-19 patients in Africa. The most common symptoms are fever, cough and breathing problems.

INTRODUCTION

A disease that was first reported in early December 2019¹ has led to a pandemic in 2020 that has resulted in about 39 million cases and over 1.1 million deaths as of 18th October, 2020². Nearly every continent of the earth, except Antarctica, has felt the impact of the novel Coronavirus Disease 2019 (COVID-19) caused by the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)^{3,4}. It was predicted that Africa will be the worst hit with the pandemic, having no less than 223 million cases and more than 150,000 fatalities^{3,5}. However, eight months into the pandemic, this is not the case. The WHO African Region, comprising only 47 member states on the continent (without Djibouti, Egypt, Libya, Morocco, Somalia, Sudan and Tunisia), has reported only 3.2% of all COVID-19 cases (1,259,192 out of 39,596,858) and 2.6% of deaths (28,313 out of 1,107,374) globally as of October 18th, 2020^{2,6}.

Nevertheless, to improve our understanding and management of this novel disease, it remains pertinent to define and analyze the clinical characteristics and outcomes of the disease in patients⁷. Several studies have described the clinical and epidemiological characteristics, risk factors, case management, and associated outcomes of different patient cohorts with COVID-19 globally^{1,7-15}. Clinical features, diagnosis and treatment of COVID-19 have also been reviewed elsewhere¹⁶. However, there is limited information on the clinical characteristics of COVID-19 in patients in Africa. In this article, we describe a meta-analysis of the clinical characteristics of COVID-19 patients in Africa.

METHODOLOGY

Search databases and search strategy

PubMed, Google scholar, Scopus, The Cochrane Library, EMBASE, and Africa journal online (AJOL) were electronically searched to collect clinical studies on the clinical characteristics of COVID-19 from January 1, 2020 to October 6, 2020. We also performed a manual search of the reference lists of included studies to avoid omitting any eligible study. Only studies written in English language and published online were included. The following terms were used in search alone OR in combination: “Coronavirus”

OR“2019-nCoV”OR“COVID-19”OR“SARS-CoV-2” AND “Africa” OR "Comoros" OR "Djibouti" OR "Madagascar" OR "Malawi" OR "Seychelles" OR "Cameroon" OR "Central African Republic" OR "Chad" OR "Congo" OR "Equatorial Guinea" OR "Atlantic Islands" OR “Gabon” OR "Morocco" OR "South Sudan" OR "Sudan" OR "Botswana" OR "Lesotho" OR "Swaziland" OR "Benin" OR "Burkina Faso" OR "Cape Verde" OR "Ghana" OR "Guinea" OR "Guinea-Bissau" OR "Mauritania" OR "Niger" OR "Senegal" OR "Sierra Leone" OR "Togo" OR “Burundi” OR “Eritrea” OR “Ethiopia” OR “Kenya” OR “Mozambique” OR “Rwanda” OR “Somalia” OR “Tanzania” OR “Uganda” OR “Zambia” OR “Zimbabwe” OR “Angola” OR “Algeria” OR “Egypt” OR “Tunisia” OR “Namibia” OR “South Africa” OR “Gambia” OR “Liberia” OR “Mali” OR “Nigeria” AND English[lang].
Figure 1 shows the flow chart of the literature screening process.

Inclusion and exclusion criteria

The inclusion criteria were as follows: (1) cohort studies, case-control studies, and case series studies; (2) the study population included individuals diagnosed with COVID-19; (3) the primary outcomes were: clinical symptoms, signs, demographics and fatality rate, etc. The exclusion criteria were as follows: (1) overlapping or duplicate studies; (2) The epidemiological analysis with only secondary outcomes such as fatality rate, without the primary outcomes; (3) had no clinical indicators or lacking necessary data.

Data extraction and quality assessment

Two reviewers, using the inclusion and exclusion criteria, independently selected the literature, and extracted data to an Excel database and any disagreement was resolved by consensus. Data extraction included the first author's surname and the date of publication of the article, study region/country, study design, sample size, age and outcome measurement data such as clinical symptoms. The included studies of this meta-analysis were observational case series studies, so the British National Institute for Clinical Excellence (BNICE) was used to evaluate the study quality by the independent reviewers. The evaluation included 8 items and the total score was 8. Studies with a score greater than 4 were seen as high-quality.

Statistical analyses

All the meta-analyses were performed by using Open-Meta Analyst¹⁷ Single-arm meta-analyses were carried out. The heterogeneity was quantified using the I^2 statistic. The random model was utilized for statistical heterogeneity between the results of each study.




Figure 1: Flow chart of the literature screening process

RESULTS

Literature retrieval and articles characteristics

A total of 1065 Records were identified during literature retrieval from databases. A total of 7 studies involving 4490 COVID-19 patients were included in this meta-analysis^{8,9,15,18-21} (Figure 1 and Table 1). All studies included in this meta-analysis were conducted in African countries. The characteristics of the studies included in this meta-analysis were published between June 1, 2020 to October 2, 2020. The quality score of the included studies ranged from 5 to 7 on a maximum scale of quality score of 8 using the NICE criteria²² (Table 1).

Gender distribution

The random effect model was used in the meta-analysis. Gender distribution showed that the proportion of male was 68.8% (95% CI, 64.6%-72.9%) with significant heterogeneity $I^2 = 65.16\%$, $p = 0.014$ (Figure 2).

Clinical symptoms

The incidence of commonly encountered symptoms were as follows: Fever (42.8%, 95% CL 30.7%-54.9%), Cough (33.3%, 95% CL, 24.0%-42.5%), Headache (11.3%, 95%CL, 2.9%-25.4%), Breathing problem (16.8%, 95% CL, 4.5%-29.1%) Diarrhea (7.5%, 95%CL, 2.8%-12.2%) and Rhinorrhea (9.4%, 95% CL, 8.5%-10.4%) as show in Table 2.

Fatality rate

The fatality rate in the 4490 patients included in the meta-analysis was 5.6% (95% CL, 2.7%-8.6%) with significant heterogeneity ($I^2=85.9\%$, $p= <0.001$) (Figure 3).

Table 1: Basic characteristics and quality score of included studies

S/N	Studies	Countries	Number of patients	Quality Score according to NICE Criteria	Date Published
1	Cekhlabi <i>et al.</i>	Morocco	15	6	June 1, 2020
2	Abimbola-Bowale <i>et al.</i>	Nigeria	32	5	May 6, 2020
3	Abdoul Aziz-Diouf <i>et al.</i>	Senegal	9	5	

4	Nachegea <i>et al.</i>	Congo	766	6	October 2, 2020
5	Kirenga <i>et al.</i>	Uganda	56	7	August 24, 2020
6	Elimian <i>et al.</i>	Nigeria	3467	6	August 26, 2020
7	Otuonye <i>et al</i> (Pre-Print)	Nigeria	154	5	September 24, 2020
Total			4490		




Figure 2: Proportion of males in COVID-19 patients

Table 2: Meta-analysis of different clinical symptoms in COVID-19 patients

Symptoms	No. of studies	No. of Patients	Heterogeneity		Model	Meta-analysis	
			P	I ²		R (95% CL)	P
Fever	6	1108	<0.001	88.25%	Random	0.428(0.307, 0.549)	<0.001
Cough	6	1003	<0.001	79.99%	Random	0.332(0.240, 0.425)	<0.001
Headache	3	31	<0.001	85.99%	Random	0.113(-0.029, 0.254)	<0.001
Rhinorrhea	5	354	0.430	0.0%	Random	0.094(0.085, 0.104)	0.430
Diarrhea	6	163	0.003	72.15%	Random	0.075(0.028, 0.104)	0.003

						0.122)	
Breathing Problem	4	427	<0.001	95.83%	Random	0.168(0.045, 0.291)	<0.001




Figure 3: Proportion of death rate in COVID-19 patients examined

DISCUSSION

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-COV-2) is a positive-stranded single-stranded RNA virus within the β -coronavirus cluster¹¹. Humans have experienced two prior outbreaks of coronavirus infections - Severe Acute Respiratory Syndrome (SARS) in 2002, and Middle East Respiratory Syndrome (MERS) in 2012, all within the past 2 decades²³. Since the outbreak of COVID-19 in china, and the first infection record on African soil, identifying the clinical characteristics and its dynamics in COVID-19 patients has been the key effort of most clinical studies. In this study, we evaluated evidence from previous studies in Africa and conducted this meta-analysis to systematically review the clinical characteristics of COVID-19 patients on the continent between June 1 and October 2, 2020. Our analysis included 4490 COVID-19 patients from seven studies conducted in five countries (Morocco, Senegal, Uganda, Congo and Nigeria). To the best of our knowledge,

this is the first description and analysis of the clinical characteristics of COVID-19 patients in Africa.

From our analysis, males were the most affected subjects. This finding agrees with the finding of several studies conducted on other continents such as China^{1,10,11,13,14}, Kuwait⁷, and the United States¹². Although, during the SARS-CoV pandemic in 2002, there was a female dominance²⁴, and in a study of 869 patients confirmed with COVID-19 in Wuhan, China, there were less men (43.4%)¹³. The most common symptoms of patients with COVID-19 encountered were fever, cough, breathing problem and headache, while diarrhea and rhinorrhea were least encountered. Fever as a clinical symptom of SARS-CoV-2 infection has been found to be associated with other previous coronaviruses infections as documented by SARS-CoV, MERS-CoV, and SARS-CoV-2^{24,25}. Our findings of fever, cough and breathing problems as clinical symptoms of COVID-19 were in tandem with the findings of other studies, with up to 99% of patients experiencing fever²⁴. However, the proportion of patients that developed fever in our analysis was lower than the reported from meta-analysis done in China and other countries where over 80% of the patients examined had fever^{1,10,12,26}. Our study also reported low proportion in cough and breathing problems as compared with studies done outside Africa^{11,12}. The case fatality rate from the analyzed data for COVID-19 patients in Africa was lower than reported in other studies,^{10,12,14} but was higher than reports reviewed by Zhu *et al.*²⁴ and Rodriguez-Morales *et al.*²⁷

This study is not without its strengths and weaknesses. The strengths of this study include its large sample size and the high-quality score of studies included in the analysis. Some limitations were also conducting our meta-analysis. First, most studies included studies were retrospective and conducted at single centers. This may have introduced admission bias and selection bias, so we cannot rule out the influence of other confounding factors. The sample sizes for two studies^{9,18} were relatively small, so the test efficiency may be insufficient. Also, data collected in most of the included studies did not include laboratory findings, hence, it was difficult to analyze the clinicopathological characteristics of the disease in patients across Africa. This meta-analysis showed a significant heterogeneity between the studies, due to too many outcomes, and there was no subgroup analysis, which could affect the accuracy of the results presented.

CONCLUSION

Currently, documented evidence shows that the most commonly encountered symptoms of COVID-19 patients in Africa were fever, cough, breathing problem and headache. The case-

fatality was low, contrary to predictions and modeling results by experts. More high-quality prospective studies are required to verify the above conclusions and to gain more insights into the clinical characteristics of COVID-19 patients in Africa. However, in the absence of any vaccine, relevant measures such as those that address infectious disease outbreak response²⁸ should be put in place to slow down the spread of the virus.

Consent for Publishing

All authors gave their consent for publishing

Availability of Supporting Data

Besides those presented in the study, there are no other supporting data for this manuscript.

Competing Interests

All authors declare no competing interests.

Funding

None

Authors' contributions

TJO and LIU were both involved in the conceptualization, design, analysis, and writing of the manuscript. Both authors approved the final draft of the manuscript.

REFERENCES

1. Guan W et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382(18):1708-1720. doi:10.1056/NEJMoa2002032
2. World Health Organisation (WHO). *WHO Coronavirus Disease (COVID-19) Dashboard.*; 2020. <https://covid19.who.int/>
3. Cabore JW et al. The potential effects of widespread community transmission of SARS-CoV-2 infection in the World Health Organization African Region: a predictive model. *BMJ Glob Heal.* 2020;5(5):e002647. doi:10.1136/bmjgh-2020-002647
4. Rahman S, Bahar T. COVID-19: The New Threat. *Int J Infect.* 2020;7(1). doi:10.5812/iji.102184
5. El-Sadr WM, Justman J. Africa in the Path of Covid-19. *N Engl J Med.* 2020;383(3):e11. doi:10.1056/NEJMp2008193
6. Amimo F, Lambert B, Magit A. What does the COVID-19 pandemic mean for HIV, tuberculosis, and malaria control? *Trop Med Health.* 2020;48(1):32. doi:10.1186/s41182-020-00219-6
7. Almazeedi S, Al-Youha S, Jamal MH, Al-Haddad M, Al-Muhaini A, Al-Ghimlas F, Al-Sabah S. Characteristics, risk factors and outcomes among the first consecutive 1096 patients diagnosed with COVID-19 in Kuwait. *EClinicalMedicine.* 2020;24:100448. doi:10.1016/j.eclinm.2020.100448
8. Bowale A et al. Clinical presentation, case management and outcomes for the first 32

- COVID-19 patients in Nigeria. *Pan Afr Med J.* 2020;35(Supp 2):1-4. doi:10.11604/pamj.supp.2020.35.2.23262
9. Diouf AA et al. Clinical characteristics and outcomes of COVID-19 infection in nine pregnant women: A report from a Sub-Saharan African Country, Senegal. *Pan Afr Med J.* 2020;35(2):1-3. doi:10.11604/pamj.2020.35.2.23736
 10. Zhu J, Ji P, Pang J, Zhong Z, Li H, He C, Zhang J, Zhao C. Clinical characteristics of 3,062 COVID-19 patients: a meta-analysis. *J Med Virol.* Published online 2020:0-1. doi:10.1002/jmv.25884
 11. Zhu J, Zhong Z, Ji P, Li H, Li B, Pang J, Zhang J, Zhao C. Clinicopathological characteristics of 8697 patients with COVID-19 in China: A meta-analysis. *Fam Med Community Heal.* 2020;8(2):1-11. doi:10.1136/fmch-2020-000406
 12. Suleyman G et al. Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. *JAMA Netw open.* 2020;3(6):e2012270. doi:10.1001/jamanetworkopen.2020.12270
 13. Zhang J et al. The Clinical Characteristics and Prognosis Factors of Mild-Moderate Patients With COVID-19 in a Mobile Cabin Hospital: A Retrospective, Single-Center Study. *Front Public Heal.* 2020;8(June):1-11. doi:10.3389/fpubh.2020.00264
 14. Chen N et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020;395(10223):507-513. doi:10.1016/S0140-6736(20)30211-7
 15. Otuonye N et al. Clinical and Demographic Characteristics of COVID-19 Patients in Lagos, Nigeria: A Descriptive Study. *MedRxiv.* doi:<https://doi.org/10.1101/2020.09.15.20195412>
 16. Hassan SA, Sheikh FN, Jamal S, Ezeh JK, Akhtar A. Coronavirus (COVID-19): A Review of Clinical Features, Diagnosis, and Treatment. *Cureus.* 2020;12(3). doi:10.7759/cureus.7355
 17. Wallace BC, Dahabreh IJ, Trikalinos TA, Lau J, Trow P, Schmid CH. Closing the gap between methodologists and end-users: R as a computational back-end. *J Stat Softw.* 2012;49(5):1-15.
 18. Chekhlabi N, Kettani C El, Haoudar A, Bahlaoui A, Mahi M, Ettair S, Dini N. The epidemiological and clinical profile of covid-19 in children: Moroccan experience of the cheikh khalifa university center. *Pan Afr Med J.* 2020;35(2):1-5. doi:10.11604/pamj.2020.35.2.23571
 19. Nachega JB et al. Clinical Characteristics and Outcomes of Patients Hospitalized for COVID-19 in Africa: Early Insights from the Democratic Republic of the Congo. *Am J Trop Med Hyg.* 2020;(October). doi:10.4269/ajtmh.20-1240
 20. Kirenga B et al. Characteristics and outcomes of admitted patients infected with SARS-CoV-2 in Uganda. *BMJ Open Respir Res.* 2020;7(1). doi:10.1136/bmjresp-2020-000646
 21. Elimian KO et al. Descriptive epidemiology of coronavirus disease 2019 in Nigeria , 27 February -. 2020;148(e208).
 22. NICE. Appendix 4 Quality of case series form. Accessed October 12, 2020.

<https://www.nice.org.uk/guidance/cg3/documents/appendix-4-quality-of-caseseries-form2>

23. Hui DSC, Zumla A. Severe Acute Respiratory Syndrome: Historical, Epidemiologic, and Clinical Features. *Infect Dis Clin N Am.* 2020;33:869-889.
24. Zhu Z, Lian X, Su X, Wu W, Marraro GA, Zeng Y. From SARS and MERS to COVID-19: a brief summary and comparison of severe acute respiratory infections caused by three highly pathogenic human coronaviruses. *Respir Res.* 2020;21(1):224. doi:10.1186/s12931-020-01479-w
25. Hui DS, Chan PK. Clinical features, pathogenesis and immunobiology of severe acute respiratory syndrome. *Curr Opin Pulm Med.* 2008;14(3):241-247. doi:10.1097/MCP.0b013e3282fb81b2
26. Sun P, Qie S, Liu Z, Ren J, Li K, Xi J. Clinical characteristics of hospitalized patients with SARS-CoV-2 infection: A single arm meta-analysis. *J Med Virol.* 2020;92(6):612-617. doi:10.1002/jmv.25735
27. Rodriguez-Morales AJ et al. Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel Med Infect Dis.* 2020;34:101623. doi:10.1016/j.tmaid.2020.101623
28. Olumade TJ, Adesanya OA, Fred-Akintunwa IJ, Babalola DO, Oguzie JU, Ogunsanya OA, George UE, Akin-Ajani OD, Osasona DG. Infectious disease outbreak preparedness and response in Nigeria: history, limitations and recommendations for global health policy and practice. *AIMS Public Heal.* 2020;7(4):736-757. doi:10.3934/publichealth.2020057