Engineered phage endolysin eliminates *Gardnerella* in bacterial vaginosis without

damaging the healthy vaginal microbiome

Christine LANDLINGER, Dr. [a], Lenka TISAKOVA, Dr. [a], Vera OBERBAUER, Ms. [a], Timo

SCHWEBS, Mr. [a], Agnieszka LATKA, Dr. [b], Leen VAN SIMAEY, Ms. [b], Mario

VANEECHOUTTE, Prof. [b], Alexander GUSCHIN, Dr. [c], Sonja SWIDSINSKI, Dr. [d],

Alexander SWIDSINSKI, Dr. [e,f], Lorenzo CORSINI, Dr. [a]

[a] PhagoMed Biopharma GmbH, Vienna Biocenter, Austria

[b] Laboratory Bacteriology Research, Department of Diagnostic Sciences, Faculty of Medicine

& Health Sciences, Ghent University, Flanders, Belgium

[c] Moscow Scientific and Practical Center of Dermatovenerology and Cosmetology.Moscow,

Russia

[d] MDI Limbach Berlin GmbH, Berlin, Germany

[e] Medizinische Klinik, Charité CCM, Humboldt Universität, Berlin, Germany

[f] Institute of Molecular Medicine, Sechenov First Moscow State Medical University, Moscow,

Russia

Correspondence to:

Lorenzo Corsini

PhagoMed Biopharma GmbH

Leberstraße 20, 1110 Wien, Austria

Tel: +43 6765797380

e-mail: Lorenzo.corsini@phagomed.com

Disclosure

Conflicts of interests have been declared by the following authors: CL, LT, VO, are employees of PhagoMed Biopharma GmbH at the time of the study and LC holds shares in PhagoMed Biopharma GmbH and is the inventor on a patent application related to the work described here. The research of AS, AL, LVS and MV has been supported financially by PhagoMed Biopharma GmbH.

Funding

This study was supported by the Austrian Research Promotion Agency (FFG).

Ethics approval

The study was approved by the Charité ethical committee EA1 /088110.

Word count of the abstract

343 words

Word count of the main text

5434 words

Keywords: *Gardnerella*, prophage, endolysin, bacterial vaginosis, polymicrobial biofilm, genus-specificity, antimicrobial resistance, alternative to antibiotic treatment

Background - Bacterial vaginosis is characterized by an imbalance of the vaginal microbiome in which the normally predominant lactobacilli are replaced by other bacterial species. Initiated by strains of the bacterium *Gardnerella,* a characteristic biofilm forms on the vaginal epithelium, explaining the typical presence of clue cells. This biofilm contributes to the resilience of the bacteria to antibiotic treatment, which may explain the frequent recurrence of BV.

Objective - In this study, we investigate whether a therapy based on bacteriophage endolysins which specifically lyse *Gardnerella*, in particular the drug candidate PM-477, might be a promising alternative to broad-spectrum antibiotics and antiseptics.

Study design - To identify *Gardnerella* specific endolysins, we searched for endolysin-encoding sequences in regions of *Gardnerella* genomes that are of prophage origin. This search identified fourteen homologous genes predicted to encode 1,4-beta-N-acetylmuramidase-type endolysins. When expressed in *Escherichia coli* and purified, the recombinant proteins demonstrated strong bactericidal activity against four different *Gardnerella* species. By shuffling the N-terminal catalytic domains and C-terminal cell wall-binding domains between the homologues, we produced 81 chimeric endolysins. These endolysins were tested for their activity and specificity *in vitro* and *ex-vivo* on vaginal samples from fifteen BV positive patients. Fluorescence in situ hybridization was used for visualization.

Results - Several engineered endolysins were 10-fold more active than the most active wild-type enzymes. When tested against a panel of 20 *Gardnerella* strains, the most active endolysin, called PM-477, showed minimum inhibitory concentrations of 0.13–8 µg/ml. PM-477 had no effect on *Lactobacillus* strains or other species of vaginal bacteria. Furthermore, the efficacy of PM-477 was tested on vaginal samples from fifteen patients with either first time or recurring bacterial vaginosis. In fourteen cases, PM-477 killed the *Gardnerella*

bacteria and physically dissolved the biofilms without affecting the remaining vaginal

microbiome.

Conclusion - The high selectivity and effectiveness in eliminating *Gardnerella*, both in

cultures of isolated strains as well as in clinically derived samples of natural polymicrobial

biofilms, makes PM-477 a promising drug candidate and an alternative to antibiotics for the

treatment of bacterial vaginosis, especially in patients with frequent recurrence.

Background

Bacterial vaginosis (BV) is a very common disorder in women of reproductive age, with a prevalence estimated at 10–30 % worldwide [1,2]. It is caused by an imbalance in the normal, healthy microbiome of the vagina that results mostly in discharge, odor and irritation. BV is associated with an increased risk of preterm delivery and low birthweight [3,4], infertility and early spontaneous abortion [5–7], and it is also a high risk factor for contracting sexually transmitted diseases, including HIV [8,9]. Besides these potentially severe physiological consequences, frequently recurring and strongly symptomatic BV can have a huge negative impact on some women's quality of life and psychological wellbeing.

The healthy vaginal microbiome is characterized by a low diversity and uniform colonization by a few species of *Lactobacillus* [10]. In BV, the imbalance in the vaginal microbiome is represented predominantly by the loss of beneficial lactobacilli and overgrowth of bacteria such as *Gardnerella*, *Prevotella*, *Atopobium*, and *Sneathia* [11] and *Lactobacillus iners* [12]. The current model of the etiology of BV focuses on the importance of *Gardnerella* species. It postulates that virulent strains of *Gardnerella* form an adherent biofilm on the vaginal epithelium in which other species can proliferate, resulting in a polymicrobial biofilm [13–17]. This adherent biofilm is only weakly affected by the innate immune response [18], and the interplay with the host immune system is only partially understood [19]. BV development may be triggered by sexual transmission of a mix of bacteria including *Gardnerella*, which can adhere to epithelial cells in the presence of lactobacilli. It has been suggested that, unlike strictly anaerobic bacteria associated with BV, *Gardnerella*, a facultative anaerobe, can tolerate the high redox potential created by the *Lactobacillus*-dominated healthy vaginal microbiome [15,20,21]. The metabolism of *Gardnerella spp.*, in turn, results in a local increase in pH and decrease of the redox potential, favoring growth of iron-dependent anaerobes [10] and suppression of lactobacilli [22]. Recently, the species previously known as *Gardnerella vaginalis* was shown to comprise at least 13 different species of which four were named [23].

Clinical studies indicated that various *Gardnerella* species may contribute differently to the

pathogenesis of BV. A higher abundance of *G. vaginalis* and *G. swidsinskii* was found to be

related with vaginal symptoms of abnormal odor and discharge and the relative abundances of

G. vaginalis, *G. swidsinskii*, and *G. piotii* but not *G. leopoldii* were strongly associated with

BV microbiome [24,25].

The therapies used currently to treat BV are broad-spectrum antibiotics, mainly metronidazole

(MDZ), tinidazole (TDZ), or clindamycin (CLI), antiseptics (such as Octenisept®, based on

the detergent-like octenidin), probiotics (preparations of lactobacilli), and prebiotics (lactate

gels). Although all of these treatments may have some beneficial effects, none has proven to

have satisfactory efficacy in preventing recurrence. Antibiotics are the most effective in

quickly reducing the symptoms, but they are associated with a recurrence rate of up to 60 %

within six months of treatment [26]. One reason may be the presence of an antibiotic-tolerant

biofilm in BV that shelters a reservoir of persister cells [26,27]. Antibiotic treatment not only

often fails to eradicate the biofilm, but can also lead to further dysbiosis of the vaginal

microbiome and can promote candidiasis [28]. Moreover, CLI can trigger pseudomebrannous

colitis, cause antibiotics associated diarrhea and other gastro-intestinal side effects [29,30]. Also,

Gardnerella strains frequently develop resistance to MDZ and TDZ [31]. Antiseptic treatments

for BV have been less well studied than antibiotics but they are similar with regard to their

effectiveness and are also associated with high recurrence rates and deterioration of the

healthy vaginal microbiome [16,32]. Studies of probiotics, and prebiotics are generally

inconclusive: they appear not to result in lasting benefits [26]. Vaginal microbiome

transplantation after a course of antibiotic treatment had a promising effect in one study of

five patients, but the approach is very resource-intense and hardly scalable [33]. In conclusion,

novel treatments that offer an alternative to antibiotics or that can be used in combination with

antibiotics, are urgently needed.

Bacteriophage-encoded peptidoglycan hydrolases – also called endolysins or 'enzybiotics' –

are a promising alternative to antibiotics [34,35]. Produced towards the end of the lytic cycle in

phage-infected bacteria, these enzymes cleave peptidoglycan in the bacterial cell wall, thus

lysing the cells and releasing the progeny phages. They have been shown to be particularly

effective against Gram-positive bacteria, which lack the Gram-negative outer membrane that

might limit accessibility to the cell wall peptidoglycan. Endolysins have several advantages

over antibiotics; not least, their very narrow host spectrum, which is usually limited to a single

genus or even a single species [36], and their low propensity to generate resistance in their hosts

[37]. Bacteriophages that invade Gram-positive bacteria encode a variety of highly diverse

endolysins. In general, they have a modular structure consisting of one or more enzymatically

active domain (EAD)s connected by a flexible interdomain linker to at least one cell wall-

binding domain (CBD), typically located at the C-terminus of the protein. Both domains can

contribute to the specificity for a given genus or species of bacteria [38].

Although no bacteriophages that infect *Gardnerella* species have yet been isolated, prophages

encoding for endolysins are present in the *Gardnerella* genomes. Phage endolysins are a

potentially novel and promising approach to treat BV because *Gardnerella*, now generally

acknowledged as the central agent in the etiology of BV, has a thin cell wall. To develop

endolysins specific for *Gardnerella*, we used an *in silico* approach to identify the genes

encoding endolysins. We cloned and sequenced 14 such genes, demonstrated the lytic activity

and specificity of purified recombinant proteins for *Gardnerella* spp. and engineered an

enzyme with enhanced activity by domain shuffling. Moreover, we demonstrate the efficacy

and specificity of this engineered endolysin on biofilms of *Gardnerella* in samples from

patients with BV.

Material and Methods

Bacterial strains and culture conditions

Atopobium vaginae, *Gardnerella* spp., *Lactobacillus iners*, *Mobiluncus spp*, *Streptococcus*

agalactiae, *Prevotella bivia*, and and were obtained from the Laboratory of Bacteriology,

University of Ghent, Belgium. Other *Lactobacillus* spp. were obtained from the German

Collection of Microorganisms (DSMZ). *Gardnerella* strains were either grown on chocolate

(Choc) agar plates (Becton Dickinson) under anaerobic conditions in an anaerobic chamber

equipped with anaerobic atmosphere generation bags (Sigma Aldrich) for 24–48 h.

Lactobacillus spp. were cultured on Schaedler plates supplemented with vitamin K1 and 5 %

sheep blood (Becton Dickinson), anaerobically at 37 °C for 24–48 h. *A. vaginae, M. curtisii,*

M. mulieris, S. agalactiae, and P. bivia were either grown on Columbia blood agar plates,

supplemented with 5 % sheep blood (Becton Dickinson) or Schaedler agar plates

supplemented with vitamin K1 and 5 % sheep blood, anaerobically at 37 °C for 48 h.

Escherichia coli BL21(DE3) cells (New England Biolabs) were used for protein expression

and were grown on Luria Bertani (LB) agar plates (Becton Dickinson) or in LB (Luria/Miller)

broth (Carl Roth) supplemented with appropriate selection antibiotic, aerobically at 37 °C.

Liquid cultures of all species were done in New York City (NYC) broth III (10 mM HEPES

(Sigma Aldrich), 15 g/L Proteose Peptone (Sigma Aldrich), 3.8 g/L yeast extract (Thermo

Fisher Scientific), 86 mM sodium chloride (Carl Roth), 28 mM α-D-glucose (Sigma

Aldrich)), supplemented with 10 % horse serum (HS) (Thermo Fisher Scientific).

Identification of* Gardnerella *genes encoding endolysin within prophage regions

Prophage regions in *Gardnerella* genomic sequences were identified by using PHASTER

(https://phaster.ca/) to scan translated nucleotide sequences of *Gardnerella* genome entries in

the NCBI database (https://blast.ncbi.nlm.nih.gov). Protein domain searches for endolysins

were performed using INTERPRO (https://www.ebi.ac.uk/interpro/) to identify predicted

enzymatically active domains in the corresponding amino acid sequence for each identified

phage endolysin.

Gene cloning and overexpression of phage endolysins

Briefly, DNA of genes encoding wild-type or genetically engineered endolysins fused to a

His$_6$-tag was synthesized, cloned into the expression vector pET29b(+) and expressed in *E.*

coli BL21(DE3). Protein expression was induced by TB (Terrific Broth Medium; 24 g/L

Bacto Yeast Extract (Becton Dickinson), 12 g/L Bacto Tryptone (Becton Dickinson), 0.4 %

glycerol (87%; Applichem), 0.02 mM potassium phosphate monobasic (Sigma Aldrich), 0.07

mM di-potassium hydrogen phosphate (Merck) and 1.5 % α-lactose monohydrate (Carl

Roth)), by incubating at 25 °C for 24 h, with shaking at 250 rpm/min. The overexpressed

proteins were purified by affinity chromatography on a nickel–nitrilotriacetic acid (Ni–NTA)

affinity matrix (HISTrap column) eluted with 50 mM MES (Carl Roth) pH 7, 150 mM NaCl

(Carl Roth), 250 mM imidazole (Carl Roth) and followed by size exclusion chromatography.

Fractions were pooled and dialyzed against MES buffer (50 mM MES pH 5.5, 100 mM NaCl,

8 mM MgSO$_4$ (Sigma Aldrich)). Protein concentration was determined at OD 260/280 nm or

by using the Pierce^{TM} BCA (bicinchoninic acid) protein assay kit (Thermo Fisher Scientific).

Culture-based assessment of bactericidal activity

Bacterial suspensions (OD$_{600}$ of 0.1, corresponding to approximately 10^7–10^8 CFU/ml) were

prepared by scraping the cells from confluently grown agar plates and diluting them into

NYCIII+ 10 % HS, pH 5.5. Reactions were performed in triplicate by mixing 10 µl of

endolysin (200 µg/ml) with 90 µl bacterial suspension in the wells of a 96-well plate. Ten µl

MES buffer without the endolysin was used as a control. The 96-well reaction plate was

incubated anaerobically at 37 °C for 5 h. Tenfold dilution series (10^{-1} to 10^{-6}) of the cell

reaction mixtures were prepared in NYCB+ 10 % HS and 2 µl of each dilution were spotted

onto Choc agar plates. After anaerobic incubation at 37 °C for 48 h, colonies were counted,

CFU/ml calculated, and the log$_{10}$ reduction compared to MES buffer treated control was

determined.

MIC assessment

The minimum inhibitory concentration (MIC), which is a standard measure of the activity of

antimicrobials, was determined according to the Clinical and Laboratory Standards Institute

protocol (2018) *Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria* [39].

Bacterial suspensions of 10^5 to 10^6 CFU/ml in NYCIII+ 10 % HS were treated with a 1:2

dilution series of either PM-477 (starting concentration 64 µg/ml) or the antibiotics

metronidazole (Gatt-Koller), tinidazole (Sigma Aldrich), or clindamycin (Clindamycin

hydrochloride, Sigma Aldrich), starting with a concentration of 128 µg/ml and twofold

dilution down to 0.0625 µg/ml. Controls for growth in the absence of antimicrobials were also

included. OD_{620} was recorded by microplate reader (Tecan) after incubation at 37 °C for 48 h

or, for some fast-growing strains, after incubation for 24 h, and compared to t=0 values.

Phase contrast microscopy

Dense bacterial suspensions were prepared in NYCIII+ 10 % HS either as *Gardnerella*

monocultures or as a co-cultures of *Gardnerella* and *Lactobacillus* strains at final $OD_{600}= 4$

for each strain. Bacterial mixtures were centrifuged at 3,800 x g for 7 min at room

temperature in a benchtop centrifuge (Heraeus Instruments, Biofuge Pico). Supernatants were

removed and cell pellets were resuspended in 50 µl of PM-477 (460 µg/ml) or in MES buffer

alone as a negative control and incubated for 5 h at 37 □C under anaerobic conditions.

Subsequently, 5 µl of each bacterial suspension were placed on a glass slide, covered with a

glass coverslip and observed by phase contrast microscopy (Olympus BX41 microscope,

magnification 1000x).

For the comparison of bacteriolytic and bactericidal effects, bacterial suspensions of

Gardnerella swidsinskii and *Gardnerella vaginalis* were prepared in NYCIII+ 10 % HS at a

final $OD_{600}= 1$. Bacterial suspensions of 50 µl were centrifuged at 3,800 x g for 7 min at room

temperature in a benchtop centrifuge (Heraeus Instruments, Biofuge Pico). Supernatants were

removed and pellets were resuspended in 50 µl of MES buffer containing PM-477 at 400

µg/ml, 200 µg/ml, 100 µg/ml, 50 µg/ml or 25 µg/ml, or in MES buffer alone as a negative

control, and incubated for 5 h at 37 □C in anaerobic conditions. Subsequently, 5 µl of each

bacterial suspension were observed by phase contrast microscopy as above. To determine

viability, 10 µl of the reaction mixture was serially diluted and spotted onto Choc agar plates.

After incubation at 37 °C for 48 h, the colonies were counted and CFU/ml was calculated.

Lytic effects of PM-477 and Octenisept® on* ex vivo *BV patient samples as detected by

fluorescence in situ hybridization (FISH) microscopy

Vaginal samples were collected from randomly selected 24–49 year-old Caucasian women

who were previously diagnosed with BV according to the Amsel criteria [40]. In addition, the

presence of clue cells (epithelial cells covered with cohesive *Gardnerella* bacteria) was

determined by FISH microscopy [16]. Fresh vaginal smears were collected with a swab and put

in ESwab™ 493C02 pre-filled vials (COPAN Diagnostics). The vials were vortexed gently,

resulting in a suspension that contained epithelial cells and vaginal bacteria. Four consecutive

vaginal smears from each patient were pooled. The effects of 0.2, 2, 20 and 200 µg/ml of PM-

477 after 2, 6, and 24 h, respectively, were tested either at room temperature or 36 °C. Since

the most pronounced lytic effect on *Gardnerella* was observed after 24 h exposure to 200

µg/ml PM-477 at 36 °C (**Supplementary Table 1**), this condition was used for further

studies. Subsequently, vaginal swabs from 15 BV patients were collected, aliquoted, and

mixed 1:1 (v/v) with 400 µg/ml PM-477 (final concentration of 200 µg/ml) or with MES

buffer as a negative control, or mixed 1:20 (v/v) with the disinfectant Octenisept® (Schülke)

and incubated for 24 h at 36 °C. Untreated samples were also included in the analyses as

baseline controls. The samples were fixed with Carnoy solution (alcohol/chloroform/acetic

acid 6/3/1 by volume) [41]. Fields of 10 mm x 10 mm were marked on SuperFrost slides

(Langenbrinck, Emmendingen, Germany) with a PAP pen (Kisker-Biotech, Steinfurt,

Germany). Five µl aliquots of vortexed, fixed vaginal smear sample were dropped onto the

marked field. The slides were dried for 60 min at 50 °C before FISH analysis.

For multicolor FISH analyses, we used the Gard662 *Gardnerella*-specific DNA hybridization probe [42] and the universal bacterial probe Eub338 [43]. Oligonucleotide probes were synthesized with a fluorescent dye (Cy3 or Cy5). The hybridization was performed at 50 °C as previously described [41]. The probe Ato291 was used for *Atopobium*, Liner23-2 for *L. iners*, and Lcrisp16-1 for *L. crispatus* [42]. DAPI was used to visualize the vaginal epithelial cells (microphotographs not shown). Change**s** in density and distribution of microbial species were monitored by using a Nikon Eclipse 80i fluorescence microscope, a Nikon SHG1 camera and accompanying software (Nikon, Tokyo, Japan). To avoid dispersion related biases (the thickness of smears on the glass slide and the microbial distribution is uneven), bacteria were enumerated for representative areas containing 10 epithelial cells as follows. For bacteria in low densities (e.g. lactobacilli), larger areas, including at least 10 microscopic fields were evaluated and mean cell numbers were expressed in relation to 10 epithelial cells. In areas with high local bacterial concentrations (e.g. in patches of biofilms), where cells are adjacent and in part overlapping, each 10 µm x 10 µm area covered with bacteria was counted as 500 bacterial cells. The numbers were rounded to full 100s.

Statistical analysis

Where appropriate, data were log-normalized prior to applying statistical tests (e.g. for CFU/ml values, and as indicated in the figure legends). When only two groups were compared, the unpaired two-tailed Student's *t*-test was used as indicated in the respective figure legends. Multiple groups were compared by two-tailed one-way ANOVA tests. The software used for statistical analyses was GraphPad Prism8. Differences between groups were considered statistically significant when $P \leq 0.05$.

Results

Gardnerella genomes contain prophage genes that encode active endolysins

Although no bacteriophages of *Gardnerella* species are known, multiple *Gardnerella*

genomes that have been sequenced contain DNA regions that are predicted to be of prophage

origin [44,45]. To identify endolysins that might target specifically *Gardnerella*, we used Basic

Local Alignment Search Tool (BLAST) to search for sequences with similarities to known

endolysin genes in the prophage regions of published *Gardnerella* genomes. A total of 14

genes were identified that encoded 1,4-beta-N-acetylmuramidases, with sequence similarity

between 87 to 98 %, when compared to each other. Twelve genes encode a 306 residues

protein whereas two encode similar proteins that lack 55 residues at the C-terminus

(**Supplementary Table 2**). The sequences all encode domain structures common to all known

endolysins: an N-terminal EAD of the glycoside hydrolase family 25 (GH25) and a CBD,

comprising two CW_7 motifs, homologous to the CBD of Cpl-7 lysozyme, encoded by the

Streptococcus pneumoniae bacteriophage Cp-7. The sequences of these putative endolysins

are 90–94% identical to some proteins encoded by the genomes of *Atopobium* and

Bifidobacteriaceae but < 70 % identical to other endolysins, which may already indicate a

high specificity for the genus *Gardnerella* (data not shown).

We expressed the 14 predicted endolysin genes (EL1–EL14) as recombinant proteins in *E.*

coli, and ten (EL 1–7 and EL 10–12) yielded sufficient protein to perform further

experiments. Their lytic activities were tested on representatives of the four *Gardnerella*

species: *G. vaginalis* (ATCC 14018^T), *G. leopoldii* (UGent 09.48), *G. piotii* (UGent 18.01^T),

and *G. swidsinskii* (GS10234). Bacterial suspensions of 10^7-10^8 CFU/ml were incubated with

each of the 10 endolysins (at 20 µg/ml) or with buffer as a negative control. After a 5 h

incubation, we plated the suspensions and quantified the surviving bacteria, measured as

colony-forming units (CFU)/ml (**Figure 1**). Most of the endolysins reduced the CFU/ml of all

four *Gardnerella* species by multiple $\log_{10}$ units. Only EL5 and EL6 (one of the two truncated

proteins mentioned above) were completely inactive in this assay. EL3 and EL10 were the

most active against all four *Gardnerella* species tested, *G. swidsinskii* GS10234 being the

most susceptible with a > 5 $\log_{10}$-fold reduction in CFU/ml (**Figure 1**).

Engineering of a* Gardnerella *endolysin with enhanced bactericidal activity

To increase the lytic activity of the endolysins identified above, we used domain shuffling [46,47]

to recombine systematically ten of the EADs encoded by the newly identified endolysin genes

EL 1–7 and EL 10–12 (annotated as H1–7 and H10–12) with nine CBDs encoded by EL1–5,

EL7 and EL10–12 (annotated as B1–5, B7, and B10–12; since EL6 was inactive, we did not

use its truncated CBD). All of the 81 resulting chimeric proteins were expressed in *E. coli* and

purified by means of their N-terminal His tags. We tested these chimeric endolysins for their

bactericidal activity on the abovementioned *Gardnerella* strains and compared their activity to

that of the 10 wild-type endolysins (**Figure 2**). Overall, *G. swidsinskii* GS10234 was most

sensitive to most endolysins – wild-type and chimeric. Two of the chimeric endolysins,

H2B10 and H2B11, were more active than any wild-type enzyme against all four *Gardnerella*

strains, whereas others were more active than any wild-type endolysin against some strains

(e.g. H2B12, H5B3, H5B4, H7B3 and H7B11) (**Figure 2**). Overall, chimeric endolysins

combining domains H2, H7 or H10 with domains B10, B11 or B12 showed highest

bactericidal activity. The combination of H2 and B10 was most effective against any of the

four strains and reduced their viability by an average of 4.3 $\log_{10}$ CFU/ml. We designated this

construct PM-477 and chose it for further development as a candidate for the treatment of BV.

***Specificity of PM-477 for the genus* Gardnerella**

We characterized the activity and specificity of PM-477 against panels of 13 *Gardnerella*

strains of the four species described above, 12 strains of vaginal lactobacilli (including *L.*

crispatus, *L. jensenii* and *L. gasseri,* selected as representative of the healthy vaginal

microbiome) and nine other species typically found in BV, by using the quantitative assay for

bacterial viability described above. The engineered endolysin caused a 4–5 $\log_{10}$-fold loss of

viability in three *Gardnerella* strains (Gs_GS10234, Gs_GS9838-1[T], Gp_UGent 18.01[T]), a 2–

3 $\log_{10}$-fold loss of viability in five strains (Gv_ATCC 14018[T], Gv_BV50.1, Gl_UGent 09.48,

Gp_P80275, Gs_BV7.1), and a 1–2 $\log_{10}$-fold loss of viability in a further five strains

(Gv_UGent 09.07, Gv_UGent 09.01, Gl_BV13.2, Gl_BV86.5, Gp_UGent 21.28) (**Figure**

3A). The same treatment of lactobacilli, by contrast, had no statistically significant effect on

the viability of any of the 12 strains tested (**Figure 3B**). It also had no effect on strains of

Atopobium vaginae, *Mobiluncus mulieris*, *Prevotella bivia* and *Streptococcus agalactiae*, and

had only a minor effect one of three strains tested of *Mobiluncus curtisii* (**Figure 3C**). (In the

cases of *Atopobium vaginae* and *Prevotella bivia* the cells either grew poorly or died rapidly

upon transfer from agar plates to suspension, even without treatment.) Thus, the engineered

endolysin PM-477 is highly selective for *Gardnerella*: it kills strains of each of the four main

species [23], without harming beneficial lactobacilli or other species typical of the vaginal

microbiome.

The genus-specific bacteriolytic effect of PM-477 was confirmed by microscopy in mixed

cultures of *Gardnerella* and lactobacilli. These two genera can be distinguished by their

distinct morphologies: *Gardnerella* cells are small coccoidal rods, whereas lactobacilli form

mostly long and thick rods. PM-477 (at 460 µg/ml for 5 h) lysed *G. vaginalis* and *G.*

swidsinskii cells in mono-culture and selectively lysed them in mixed cultures of *G. vaginalis*

and *L. crispatus* and of *G. swidsinskii* and *L. gasseri*, respectively, while the lactobacilli in

these mixed cultures were unaffected (**Figure 4A**).

We compared the bactericidal and bacteriolytic effects of PM-477 on *G. vaginalis* and *G.*

swidsinskii by treating suspensions of the bacteria with 25–400 µg/ml PM-477 for 5 h and

then counting the cell numbers by microscopy and determining their viability by quantitative

plating (**Figure 4B–E**). The highest concentration of PM-477 (400 µg/ml) decreased the

number of cell-like structures by only 80–90 % (**Figure 4B, C**), whereas the same treatment

resulted in a ≥ 5.6 $\log_{10}$ reduction in viable cells. This indicates that PM-477 may kill before

fully dissolving bacterial cells, resulting in visible but dead cell-like structures seen under the microscope (**Figure 4D, E**).

Efficacy of PM 477 compared to standard antimicrobials

To compare the efficacy of PM-477 in suspension to the standard antimicrobials used to treat BV, we determined the minimum inhibitory concentration (MIC)s of PM-477, CLI, MDZ and TDZ for 20 strains of *Gardnerella* by following the Clinical and Laboratory Standards Institute protocol [39]. All the *Gardnerella* species tested were highly susceptible to PM-477 with a MIC_{90} value (the MIC value which inhibits 90 % of strains) of 8 µg/ml (**Table 1**). The *G. swidsinskii* strains had MICs in the range of 0.25–1 µg/ml, whereas the *G. vaginalis* strains had MICs of 0.13–8 µg/ml, and *G. leopoldii and G. piotii* strains had MICs of 1–8 µg/ml. Among this small number of strains tested for each species, there was no evidence that the four *Gardnerella* species differed with regard to their susceptibility to PM-477. Whereas 12 of the 20 *Gardnerella* strains were resistant to MDZ and 15 were resistant TDZ, CLI was effective against all strains at MICs ≤ 1 µg/ml. All tested strains of three *Lactobacillus* species (*L. crispatus*, *L. gasseri*, and *L. jensenii*) were resistant to the highest concentrations of PM-477, MDZ, and TDZ (MICs > 128 µg/ml), whereas only one strain was resistant to CLI.

Efficacy of PM-477 against biofilms of* Gardnerella *in human BV samples

To analyze the efficacy of PM-477 in a physiological environment that resembles closely the *in vivo* situation, we treated vaginal swabs from 15 BV patients, and analyzed them by fluorescence in-situ hybridization (FISH).

Areas of strong yellow fluorescence in the untreated (baseline) and buffer control-treated samples indicated the presence of *Gardnerella*-dominated biofilm on exfoliated vaginal epithelial cells (visible due to their pale autofluorescence and clearly corresponding to DAPI counterstain), called clue cells. At baseline and after buffer treatment, the yellow fluorescence

of the *Gardnerella* probe outshines the red fluorescence of the probe for all bacteria,

indicating that the biofilm is composed in large part of *Gardnerella* cells. **Figure 5A** shows a

total resolution of the dense yellow staining on the epithelial cells upon treatment with PM-

477. Therefore, PM 477 proved strong bacteriolytic activity also in this *ex-vivo* setting. The

specificity of PM-477 was demonstrated by the remaining red signal from the universal

bacterial probe, indicating mostly non-*Gardnerella* cells. Only very few individual

Gardnerella cells seemed to remain associated with or in between the epithelial cells.

However, the comparison of bacteriolytic and bactericidal effect in **Figure 4B-E** would

indicate that these might be dead cell-like structures with sufficient DNA associated to them

to hybridize the FISH probes. In contrast to PM-477, Octenisept killed and lysed both the

Gardnerella and other bacterial cells non-specifically, as indicated by the visible reduction of

both types of fluorescence.

The effect of treatment was also quantified by counting the number of bacterial cells per 10

epithelial cells in representative fields of vision (**Figure 5B**). When compared with the

baseline samples, no significant changes were observed upon incubation with the buffer

control, indicating that the vaginal microbiome was generally stable over the time-course of

the experiment. PM-477 significantly reduced *Gardnerella* cells in the patient samples

(median of 300 cells/10 epithelial cells) when compared to baseline (median of 1200 cells/10

epithelial cells, *p*=0.02) and when compared to the buffer control-treated samples (median of

900 cells/ 10 epithelial cells, *p*=0.007): 3 of the 15 samples showed a >500-fold reduction in

Gardnerella load, 4 showed a >100-fold reduction, 7 a >10-fold reduction and 11 a >2-fold

reduction (data for individual patient samples are presented in **Supplementary Table 3**).

The effects of PM-477 on *Gardnerella* were similar to those of Octenisept[®]**.** Unlike PM-477,

however, the lytic activity of the disinfectant was not specific for *Gardnerella* but affected all

bacteria (**Figure 5A, Supplementary Table 3**).

Two other potential BV associated bacteria, *A. vaginae* and *L. iners*, as well as *L. crispatus*,

were visualized by using selective FISH probes and their cell numbers per 10 epithelial cells

were counted (**Supplementary Table 3**). *A. vaginae*, *L. crispatus,* and *L. iners* were detected

in 10, 3 and 8 out of the 15 samples, respectively. Treatment with PM-477 reduced the

median cell counts of *L. iners* when compared to treatment with the buffer control in all

samples whereas the numbers of *L. crispatus* and *A. vaginae* bacteria were unaffected.

Octenisept® strongly reduced the cell counts of all three species.

In summary, these experiments provide a proof of the efficacy and selectivity of PM-477 for

the genus *Gardnerella in vitro* and also in the native BV biofilm.

Discussion

This is the first study to show that prophage-derived endolysins active on *Gardnerella* could

be used as an innovative treatment for bacterial vaginosis. Although no Gardnerella phages

were ever isolated in a lab, we were able to identify putative endolysins *in silico*. We

expressed them as recombinant proteins, improved their potency by systematic domain

shuffling, demonstrated their efficacy and selectivity *in vitro*, and, ultimately, on the native

polymicrobial biofilms in vaginal swabs from BV patients.

The currently available therapy of bacterial vaginosis is unsatisfactory. Antibiotic-centered

'standard-of-care' treatment for BV is associated with a recurrence rate of ~60 % within 6

months, which may be explained by low efficacy of antibiotics on bacteria growing as

biofilms, but also by selective destruction of beneficial lactobacilli, which are thereby

prevented from re-establishing a protective environment [26]. Bacteriophage-derived endolysins

have been described as a potential alternative to antibiotics; they are both highly active on

mucosal biofilms and selective for individual genera or even species of bacteria [35]. Neither

endolysins nor whole phages specific to *Gardnerella* have been isolated previously, although

the existence of *Gardnerella*-specific phages has been postulated on the basis of genomic

analyses [44]. We identified *Gardnerella*-specific endolysin genes in the prophage-like genome

sequences of strains of *Gardnerella*. A similar approach was reported earlier, in which the

phage lysin PlyCD was identified by searching prophage-like sequences in the genomes of

Clostridium difficile strains [48]. The fact that it is possible to predict and generate endolysins

from prophage encoded sequences in the genomes of unrelated species (*C. difficile* and

Gardnerella) may indicate a wide application of this approach, which is easily adaptable to

other pathogenic bacteria.

We could show that the wild-type endolysins we identified and, even more so, our engineered

constructs, are highly bactericidal and selective for the genus *Gardnerella*.

Our engineered endolysin PM-477 is superior to the current standard of care antibiotics in that

it is active in the low µg/ml range and effective against all 20 of the *Gardnerella* strains we

tested, covering all four named species. In total, 60 % of the 20 strains were resistant to MDZ

and 75 % to TZD, reinforcing the notion that PM-477 may be more effective against BV than

currently used standard antibiotics.

The high resistance rate for MDZ, which was also reported elsewhere [31], does not support the

finding that MDZ treatment temporarily relieves the symptoms in a majority of cases [26].

However, long-term cure is mostly not achieved with MDZ indicating that their effectiveness

may be reduced due to the high resistance rate of *Gardnerella* strains-[49].

In our study, CLI was also effective against all the strains of different *Gardnerella* genotypes

we tested. Curiously, however, CLI has not shown superior efficacy when compared to MDZ

in clinical trials [26], which may indicate that BV biofilms are more tolerant to CLI than they

are of MDZ, thus compensating the higher rate of resistance against MDZ.

In conclusion, PM-477 is superior to all tested antibiotics because, unlike nitroimidazoles, it is

active on all *Gardnerella* strains and, unlike CLI, it does not kill lactobacilli. Antibiotics may

have the advantage over PM-477 of being active against a broader spectrum of pathogens.

But, unlike *Gardnerella*, these pathogens may not be pivotal in vaginal biofilm formation as

is the case in BV. Also, antibiotics typically have low activity on bacteria in biofilms [27]

whereas our *ex-vivo* data indicate that PM-477 works well on *Gardnerella* in vaginal biofilms.

Progress has been made in developing animal models for BV. For example, it was possible to

model some of the clinical characteristics of BV, such as epithelial exfoliation and clue cells

upon infection of mice with *Gardnerella* [50]. However, it was not possible so far to model the

increase in pH and the diverse microbiome composition of human BV in animals. Therefore,

the next best and, arguably, more biologically relevant model to test the efficacy of PM-477 is

to use *ex-vivo* vaginal samples from BV patients, which contain clue cells, *i.e.* exfoliated

epithelial cells covered with a polymicrobial biofilm. The microbial populations of all the *ex*

vivo samples that we collected from 15 individual BV patients were clearly dominated by

Gardnerella and a substantial part of the *Gardnerella* cells adhered to the epithelial cells.

Treatment with PM-477 fully resolved the dense staining due to *Gardnerella* cells on the

epithelial cells while not affecting non-*Gardnerella* cells. Thus, we conclude that PM-477

quantitatively removes *Gardnerella* cells inside the polymicrobial biofilm on vaginal

epithelial cells *ex vivo*. Furthermore, it is effective in the environment of the vaginal fluid

despite the putative presence of proteases [51] and other substances that might reduce its

activity.

PM-477 shows promising results *in vitro* and *ex vivo* but we do not know yet whether

selectively removing *Gardnerella* will be sufficient to resolve the clinical symptoms of BV,

which are caused by a polymicrobial biofilm, and to allow lactobacilli to re-colonize the

vagina. The increasing evidence of the central role that *Gardnerella* plays in the etiology of

BV [15] and the observation that BV seems not to be possible without *Gardnerella* [22], indicates

that this might well be the case. The biomass of the polymicrobial biofilm consists largely of

Gardnerella cells [52,53], so selective removal of *Gardnerella* should decimate the biofilm.

Moreover, *Gardnerella* mediates adhesion to the epithelial cells, whereas other bacteria,

especially in smaller numbers, may not be able to initiate biofilm formation on these cells [54].

Other pathogens, especially *Prevotella bivia*, synergize with *Gardnerella*, so when the latter is

removed the viability of the other pathogens is likely reduced as well [55,56]. Removal of

Gardnerella (and the concomitant reduction in the viability of *P. bivia*) would reduce the

levels of sialidase produced by *Gardnerella* and *P. bivia* and allow the mucus layer to reform,

in turn reducing the adherence of the remaining BV pathogens and depriving them of carbon

sources [57]. Finally, the active suppression of lactobacilli by *Gardnerella* [22] would be released,

allowing recolonization by benign bacteria. This hypothesis has been investigated with *in*

vitro biofilm models [17,20], and PM-477 as a selective antimicrobial agent may be a very useful

tool to further investigate the role of *Gardnerella* in *ex vivo* as well as in *in vivo* studies.

In summary, we report here that an engineered, phage-derived endolysin has the potential to

completely disrupt the *Gardnerella*-dominated biofilm in vaginal swabs from patients with

BV. PM-477 is, thus a promising candidate for an effective treatment of recurrent BV, which

remains, despite decades of research, a major public health concern.

Acknowledgements

We thank Grégory Resch and Life Science Editors for scientific editing assistance during the

preparation of the manuscript.

References

1.   Bautista CT, Wurapa E, Sateren WB, Morris S, Hollingsworth B, Sanchez JL. Bacterial

vaginosis: A synthesis of the literature on etiology, prevalence, risk factors, and

relationship with chlamydia and gonorrhea infections. *Mil Med Res*. 2016;3(1):4.

doi:10.1186/s40779-016-0074-5

2.   Kenyon C, Colebunders R, Crucitti T. The global epidemiology of bacterial vaginosis:

a systematic review. *Am J Obstet Gynecol*. 2013;209(6):505-523.

doi:10.1016/j.ajog.2013.05.006

3. Tabatabaei N, Eren A, Barreiro L, et al. Vaginal microbiome in early pregnancy and

subsequent risk of spontaneous preterm birth: a case-control study. *BJOG An Int J*

Obstet Gynaecol. 2019;126(3):349-358. doi:10.1111/1471-0528.15299

4. Leitich H, Kiss H. Asymptomatic bacterial vaginosis and intermediate flora as risk

factors for adverse pregnancy outcome. *Best Pract Res Clin Obstet Gynaecol*.

2007;21(3):375-390. doi:10.1016/j.bpobgyn.2006.12.005

5. Haahr T, Zacho J, Bräuner M, Shathmigha K, Skov Jensen J, Humaidan P.

Reproductive outcome of patients undergoing in vitro fertilisation treatment and

diagnosed with bacterial vaginosis or abnormal vaginal microbiota: a systematic

PRISMA review and meta-analysis. *BJOG An Int J Obstet Gynaecol*. 2019;126(2):200-

207. doi:10.1111/1471-0528.15178

6. Mania-Pramanik J, Kerkar SC, Salvi VS. Bacterial vaginosis: A cause of infertility? *Int*

J STD AIDS. 2009;20(11):778-781. doi:10.1258/ijsa.2009.009193

7. van Oostrum N, De Sutter P, Meys J, Verstraelen H. Risks associated with bacterial

vaginosis in infertility patients: a systematic review and meta-analysis. *Hum Reprod*.

2013;28(7):1809-1815. doi:10.1093/humrep/det096

8. Unemo M, Bradshaw CS, Hocking JS, et al. Sexually transmitted infections: challenges

ahead. *Lancet Infect Dis*. 2017;17(8):e235-e279. doi:10.1016/S1473-3099(17)30310-9

9. Hoang T, Toler E, DeLong K, et al. The cervicovaginal mucus barrier to HIV-1 is

diminished in bacterial vaginosis. Silvestri G, ed. *PLOS Pathog*. 2020;16(1):e1008236.

doi:10.1371/journal.ppat.1008236

10. Vaneechoutte M. The human vaginal microbial community. *Res Microbiol*.

2017;168(9-10):811-825. doi:10.1016/j.resmic.2017.08.001

11.   Ceccarani C, Foschi C, Parolin C, et al. Diversity of vaginal microbiome and

532   metabolome during genital infections. *Sci Rep*. 2019;9(1):1-12. doi:10.1038/s41598-

533   019-50410-x

534   12.   Petrova MI, Reid G, Vaneechoutte M, Lebeer S. Lactobacillus iners: Friend or Foe?

535   *Trends Microbiol*. 2017;25(3):182-191. doi:10.1016/j.tim.2016.11.007

536   13.   Swidsinski A, Mendling W, Loening-Baucke V, et al. Adherent biofilms in bacterial

537   vaginosis. *Obstet Gynecol*. 2005;106(5):1013-1023.

538   doi:10.1097/01.AOG.0000183594.45524.d2

539   14.   Machado A, Cerca N. Influence of Biofilm Formation by *Gardnerella vaginalis* and

540   Other Anaerobes on Bacterial Vaginosis. *J Infect Dis*. 2015;212(12):1856-1861.

541   doi:10.1093/infdis/jiv338

542   15.   Muzny CA, Taylor CM, Swords WE, et al. An Updated Conceptual Model on the

543   Pathogenesis of Bacterial Vaginosis. - PubMed - NCBI. *J Infect Dis*.

544   2019;220(9):1399-1405. doi:10.1093/infdis/jiz342

545   16.   Swidsinski A, Loening-Baucke V, Mendling W, et al. Infection through structured

546   polymicrobial Gardnerella biofilms (StPM-GB). *Histol Histopathol*. 2014;29(5):567-

547   587. doi:10.14670/HH-29.10.567

548   17.   Castro J, Alves P, Sousa C, et al. Using an in-vitro biofilm model to assess the

549   virulence potential of bacterial vaginosis or non-bacterial vaginosis Gardnerella

550   vaginalis isolates. *Sci Rep*. 2015;5:11640. doi:10.1038/srep11640

551   18.   Castro J, Jefferson KK, Cerca N, Biomédicas E, Biomédicas B, Salazar A. Innate

552   immune components affect growth and virulence traits of bacterial-vaginosis-

553   associated and non-bacterial-vaginosis-associated Gardnerella vaginalis strains

554   similarly. *Pathog Dis*. 2019;76(9):89. doi:10.1093/femspd/fty089

19.   Murphy K, Mitchell CM. The Interplay of Host Immunity, Environment and the Risk

of Bacterial Vaginosis and Associated Reproductive Health Outcomes. *J Infect Dis*.

2016;214(Suppl 1):S29-S35. doi:10.1093/infdis/jiw140

20.   Castro J, Machado D, Cerca N. Unveiling the role of Gardnerella vaginalis in

polymicrobial Bacterial Vaginosis bio fi lms□: the impact of other vaginal pathogens

living as neighbors. *ISME J*. 2019:1306-1317. doi:10.1038/s41396-018-0337-0

21.   Machado A, Jefferson KK a., Cerca N. Interactions between Lactobacillus crispatus

and bacterial vaginosis (BV)-associated bacterial species in initial attachment and

biofilm formation. *Int J Mol Sci*. 2013;14(6):12004-12012.

doi:10.3390/ijms140612004

22.   Schwebke JR, Muzny CA, Josey WE. Role of Gardnerella vaginalis in the pathogenesis

of bacterial vaginosis: A conceptual model. *J Infect Dis*. 2014;210(3):338-343.

doi:10.1093/infdis/jiu089

23.   Vaneechoutte M, Guschin A, Van Simaey L, Gansemans Y, Van Nieuwerburgh F,

Cools P. Emended description of Gardnerella vaginalis and description of Gardnerella

leopoldii sp. nov., Gardnerella piotii sp. nov. and Gardnerella swidsinskii sp. nov., with

delineation of 13 genomic species within the genus Gardnerella. *Int J Syst Evol*

Microbiol. 2019;69(3):679-687. doi:10.1099/ijsem.0.003200

24.   Hill JE, Albert AYK. Resolution and cooccurrence patterns of gardnerella leopoldii, G.

Swidsinskii, G. Piotii, and G. Vaginalis within the vaginal microbiome. *Infect Immun*.

2019;87(12). doi:10.1128/IAI.00532-19

25.   Shipitsyna E, Krysanova A, Khayrullina G, et al. Quantitation of all Four Gardnerella

vaginalis Clades Detects Abnormal Vaginal Microbiota Characteristic of Bacterial

Vaginosis More Accurately than Putative G. vaginalis Sialidase A Gene Count. *Mol*

Diagnosis Ther. 2019;23(1):139-147. doi:10.1007/s40291-019-00382-5

26.   Bradshaw CS, Sobel JD. Current Treatment of Bacterial Vaginosis-Limitations and

Need for Innovation. *J Infect Dis ®*. 2016;214(S1):14-20. doi:10.1093/infdis/jiw159

27.   Swidsinski A, Mendling W, Loening-Baucke V, et al. An adherent Gardnerella

vaginalis biofilm persists on the vaginal epithelium after standard therapy with oral

metronidazole. *Am J Obstet Gynecol*. 2008;198(1):97.e1-6.

doi:10.1016/j.ajog.2007.06.039

28.   Sobel JD. Vulvovaginal candidosis. *Lancet*. 2007;369(9577):1961-1971.

doi:10.1016/S0140-6736(07)60917-9

29.   Buffie CG, Jarchum I, Equinda M, et al. Profound alterations of intestinal microbiota

following a single dose of clindamycin results in sustained susceptibility to Clostridium

difficile-induced colitis. *Infect Immun*. 2012;80(1):62-73. doi:10.1128/IAI.05496-11

30.   Vardakas KZ, Trigkidis KK, Boukouvala E, Falagas ME. Clostridium difficile

infection following systemic antibiotic administration in randomised controlled trials: a

systematic review and meta-analysis. *Int J Antimicrob Agents*. 2016;48(1):1-10.

doi:10.1016/j.ijantimicag.2016.03.008

31.   Petrina MAB, Cosentino LA, Rabe LK, Hillier SL. Susceptibility of bacterial vaginosis

(BV)-associated bacteria to secnidazole compared to metronidazole, tinidazole and

clindamycin HHS Public Access. *Anaerobe*. 2017;47:115-119.

doi:10.1016/j.anaerobe.2017.05.005

32.   Swidsinski A, Loening-Baucke V, Swidsinski S, Verstraelen H. Polymicrobial

Gardnerella biofilm resists repeated intravaginal antiseptic treatment in a subset of

women with bacterial vaginosis: a preliminary report. *Arch Gynecol Obstet*.

2015;291(3):605-609. doi:10.1007/s00404-014-3484-1

33.   Lev-Sagie A, Goldman-Wohl D, Cohen Y, et al. Vaginal microbiome transplantation in

604   women with intractable bacterial vaginosis. *Nat Med*. 2019;25(10):1500-1504.

605   doi:10.1038/s41591-019-0600-6

606   34.   Schmelcher M, Donovan DM, Loessner MJ. Bacteriophage endolysins as novel

607   antimicrobials. *Future Microbiol*. 2012;7(10):1147-1171. doi:10.2217/fmb.12.97

608   35.   Fischetti VA. Development of phage lysins as novel therapeutics: A historical

609   perspective. *Viruses*. 2018;10(6):310. doi:10.3390/v10060310

610   36.   Fischetti VA. Bacteriophage endolysins: a novel anti-infective to control Gram-positive

611   pathogens. *Int J Med Microbiol*. 2010;300(6):357-362.

612   doi:10.1016/j.ijmm.2010.04.002

613   37.   Schuch R, Lee HM, Schneider BC, et al. Combination therapy with lysin CF-301 and

614   antibiotic is superior to antibiotic alone for treating methicillin-resistant staphylococcus

615   aureus-induced murine bacteremia. *J Infect Dis*. 2014;209(9):1469-1478.

616   doi:10.1093/infdis/jit637

617   38.   Oliveira H, Melo LDR, Santos SB, et al. Molecular Aspects and Comparative

618   Genomics of Bacteriophage Endolysins. *J Virol*. 2013;87(8):4558-4570.

619   doi:10.1128/jvi.03277-12

620   39.   Carpenter DE, Karen Anderson Diane Citron CM, JoAnn Dzink-Fox BL, et al.

621   *Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria.*; 2018.

622   www.clsi.org. Accessed June 9, 2020.

623   40.   Amsel R, Totten PA, Spiegel CA, Chen KCS, Eschenbach D, Holmes KK. Nonspecific

624   vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med*.

625   1983;74(1):14-22. doi:10.1016/0002-9343(83)91112-9

41.   Swidsinski A, Loening-Baucke V. Evaluation of Polymicrobial Involvement Using

Fluorescence In Situ Hybridization (FISH) in Clinical Practice. 2017:531-543.

doi:10.1007/978-3-662-52959-1_52

42. Swidsinski A, Guschin A, Tang Q, et al. Vulvovaginal candidiasis: histologic lesions

are primarily polymicrobial and invasive and do not contain biofilms. *Am J Obstet*

Gynecol. 2019;220(1):91.e1-91.e8. doi:10.1016/j.ajog.2018.10.023

43. Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. Combination

of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed

microbial populations. *Appl Environ Microbiol*. 1990;56(6).

44. Malki K, Shapiro JW, Price TK, et al. Genomes of Gardnerella Strains Reveal an

Abundance of Prophages within the Bladder Microbiome. Kuo C-H, ed. *PLoS One*.

2016;11(11):e0166757. doi:10.1371/journal.pone.0166757

45. Bohr LL, Mortimer TD, Pepperell CS. Lateral Gene Transfer Shapes Diversity of

Gardnerella spp. *Front Cell Infect Microbiol*. 2020;10. doi:10.3389/fcimb.2020.00293

46. Schmelcher M, Tchang VS, Loessner MJ. Domain shuffling and module engineering of

Listeria phage endolysins for enhanced lytic activity and binding affinity. *Microb*

Biotechnol. 2011;4(5):651-662. doi:10.1111/j.1751-7915.2011.00263.x

47. Nelson DC, Schmelcher M, Rodriguez-Rubio L, et al. Endolysins as Antimicrobials.

In: *Advances in Virus Research*. Vol 83. Academic Press Inc.; 2012:299-365.

doi:10.1016/B978-0-12-394438-2.00007-4

48. Wang Q, Euler CW, Delaune A, Fischetti VA. Using a novel lysin to help control

Clostridium difficile infections. *Antimicrob Agents Chemother*. 2015;59(12):7447-

7457. doi:10.1128/AAC.01357-15

49. Schuyler JA, Mordechai E, Adelson ME, Sobel JD, Gygax SE, Hilbert DW.

Identification of intrinsically metronidazole-resistant clades of Gardnerella vaginalis.

Diagn Microbiol Infect Dis. 2016;84(1):1-3. doi:10.1016/j.diagmicrobio.2015.10.006

50.   Morrill S, Gilbert NM, Lewis AL. Gardnerella vaginalis as a Cause of Bacterial

Vaginosis: Appraisal of the Evidence From in vivo Models. *Front Cell Infect*

Microbiol. 2020;10:168. doi:10.3389/fcimb.2020.00168

51.   Muytjens CMJ, Vasiliou SK, Oikonomopoulou K, Prassas I, Diamandis EP. Putative

functions of tissue kallikrein-related peptidases in vaginal fluid. *Nat Rev Urol*.

2016;13(10):596-607. doi:10.1038/nrurol.2016.161

52.   Swidsinski A, Mendling W, Loening-Baucke V, et al. Adherent Biofilms in Bacterial

Vaginosis. *Obs Gynecol*. 2005;106(5):1013-1022.

53.   Swidsinski A, Doerffel Y, Loening-Baucke V, et al. Gardnerella biofilm involves

females and males and is transmitted sexually. *Gynecol Obstet Invest*. 2010;70(4):256-

263. doi:10.1159/000314015

54.   Alves P, Castro J, Sousa C, Cereija TB, Cerca N. Gardnerella vaginalis Outcompetes

29 other bacterial species isolated from patients with bacterial vaginosis, using in an in

vitro biofilm formation model. *J Infect Dis*. 2014;210(4):593-596.

doi:10.1093/infdis/jiu131

55.   Pybus V, Onderdonk AB. Evidence for a commensal, symbiotic relationship between

Gardnerella vaginalis and Prevotella bivia involving ammonia: Potential significance

for bacterial vaginosis. *J Infect Dis*. 1997;175(2):406-413.

doi:10.1093/infdis/175.2.406

56.   Chen KCS, Forsyth PS, Buchanan TM, Holmes KK. Amine content of vaginal fluid

from untreated and treated patients with nonspecific vaginitis. *J Clin Invest*.

1979;63(5):828-835. doi:10.1172/JCI109382

57.   Briselden AM, Moncla BJ, Stevens CE, Hillier SL. Sialidases (neuraminidases) in

675  bacterial vaginosis and bacterial vaginosis-associated microflora. *J Clin Microbiol*.

676  1992;30(3).

677 **Table 1:** Minimum inhibitory concentration (MIC) of PM-477 and the antibiotics CLI, MDZ,

678 and TDZ for various *Gardnerella* spp. and *Lactobacillus* spp. (R) Resistance, defined as ≥ 32

Gardnerella spp.			MIC of Antimicrobials [µg/ml]			
Name	Strain	Clade	PM-477	CLI	MDZ	TDZ
G. vaginalis (Gv9)	ATCC 14018T	IB	1	0.25	8	128 (R)
G. vaginalis (Gv1)	UGent 09.07	IA	2	0.25	>128 (R)	>128 (R)
G. vaginalis (Gv5)	UGent 09.01	IB	0.25	0.13	8	4
G. vaginalis (Gv8)	UGent 25.49	IB	0.13	<0.06	8	4
G. vaginalis (BV50)	SJHBruges isolate		8	0.25	32	64 (R)
G. vaginalis (BV111)	SJHBruges isolate		2	0.125	8	4
G. vaginalis (FB049)	SJHBruges isolate		4	0.5	16	8
G. vaginalis (FB061)	SJHBruges isolate		8	0.25	8	4
G. leopoldii (Gl11)	UGent 09.48	II	4	0.5	128 (R)	128 (R)
G. leopoldii (BV13)	SJHBruges isolate		4	0.5	>128 (R)	>128 (R)
G. leopoldii (BV86)	SJHBruges isolate		1	0.25	>128 (R)	>128 (R)
G. piotii (Gp17)	UGent 18.01T	III	8	0.5	16	64 (R)
G. piotii (Gp22)	UGent 21.28	III	2	0.25	64 (R)	>128 (R)
G. piotii (P80275)	UGent isolate		1	0.5	16	32
G. piotii (FB041)	UGent isolate		4	1	32	64 (R)
G. piotii (VMF1800 SVT21)	SJHBruges isolate		8	1	32	64 (R)
G. swidsinskii (Gs23)	GS 10234	IV	0.5	0.25	64 (R)	>128 (R)
G. swidsinskii (Gs24)	GS 9838-1T	IV	0.25	<0.06	>128 (R)	>128 (R)
G. swidsinskii (BV7)	SJHBruges isolate		1	0.5	>128 (R)	128 (R)
G. swidsinskii (BV112)	SJHBruges isolate		1	0.06	64 (R)	64 (R)
L. crispatus	DSM 20584		>128	4	>128	>128
L. gasseri	DSM 20077		>128	64	>128	>128
L. jensenii	PB2003-073-T2-3		>128	2	>128	>128

679 µg/ml for MDZ and TDZ, and ≥ 8 µg/ml for CLI, according to international standards.

680 (R) Resistance, defined as ≥ 32 µg/ml for MDZ and TDZ and ≥ 8 µg/ml for CLI according to

681 international standards.

Figure Legends:

Figure 1: Wild-type endolysins are highly active against four different *Gardnerella* species. Suspensions of *G. vaginalis* (ATCC 14018T), *G. leopoldii* (UGent 09.48), *G. piotii* (UGent 18.01T), and *G. swidsinskii* (GS10234) were treated with recombinant endolysins prepared by expression of the naturally occurring genes (EL1–EL7 and EL10–EL12) (see Methods) and the $\log_{10}$ reduction in viable CFU/ml was determined. The $\log_{10}$ reduction was calculated by comparing the CFU/ml after endolysin treatment with that of the buffer control (w/o). LOD indicates the limit of detection.

Figure 2: Domain swapping increases the lytic activity of engineered endolysins. Eighty-one constructs comprising all possible combinations of the EAD (H) and CBD (B) domains of 10 wild-type endolysins were tested on four different *Gardnerella* species as in Figure 1. The values represent the $\log_{10}$ of the ratio of viable cells (CFU/ml; mean of triplicate assays) in endolysin-treated suspensions of *Gardnerella* compared to buffer control-treated suspensions of the species, as indicated. Blue shades indicate higher bactericidal effect (high reduction in viable CFU/ml) compared to the wild-type endolysin, red shades indicate lower bactericidal effect, and white coloring indicates equal reduction compared to wild-type endolysin (boxed in black). The endolysin candidate with the highest average reduction over four *Gardnerella* species is boxed in yellow.

Figure 3: PM-477 is highly specific for *Gardnerella* and spares lactobacilli and other BV associated microbiome components. (a) *Gardnerella* strains, (b) *Lactobacillus spp.*, (c) and *Mobiluncus curtisii*, *Mobiluncus mulieris*, *Streptococcus agalactiae*, *Atopobium vaginae* and *Prevotella bivia* were tested for their susceptibility to PM-477 (20 µg/ml for 5h; treated, black squares) and compared to buffer control-treated (untreated, white circles) bacteria suspensions. For statistical purposes CFU/ml values from the untreated and the treated groups

were log-normalized prior to applying the unpaired two-tailed multiple Student's *t*-test. *P*-

values ≤ 0.05 were considered statistically significant; ** *P*≤0.01, *** *P*≤0.001, ****

P≤0.0001.

Figure 4: PM-477 selectively kills *Gardnerella* in a co-culture with lactobacilli. **A**, Phase

contrast microscopy of dense cultures of *G. vaginalis* (ATCC14018T) (first row) and *G.*

swidsinskii (GS10234) (second row), or co-cultures of *G. vaginalis* (ATCC14018T) and *L.*

crispatus (LABCRI FB020-08c) (third row), as well as *G. swidsinskii* (GS10234) and *L.*

gasseri (LAB GAS LMG 9203T) (fourth row) treated with PM-477 (460 µg/ml for 5h) or

with buffer (control). **B** and **C**, Cell counts of *Gardnerella* cells determined by phase contrast

microscopy after treatment with 25–400 µg/ml PM-477 for 5h or with buffer (Ctr). **D** and **E**,

Viable cells remaining ($\log_{10}$ reduction CFU/ml compared to the buffer-treated control) in

suspensions of *Gardnerella* treated with 25–400 µg/ml PM-477 for 5h.

Figure 5: PM-477 dissolves the *Gardnerella*-dominated biofilm on exfoliated vaginal

epithelial cells from BV patients. **A**, Representative examples of multi-color FISH with Gard

Cy3 probe (*Gardnerella,* yellow fluorescence) and universal Eub 338 Cy5 probe (all bacteria,

red fluorescence) in 400x magnification (where not stated otherwise) for patients 11, 12 and

15. *Baseline (prior to treatment): Gardnerella* bacteria are cohesive and adherent to epithelial

cells forming dense biofilm patches on the surface of clue cells which is apparent due to

background fluorescence. The attachment of *Gardnerella* biofilm to epithelial cells can be

seen especially well in DAPI stains and on zoomed original microphotographs (DAPI

microphotographs are added to supplementary material together with all other original

microphotographs of Figure 5) *Control*: *Gardnerella* dominated biofilms in buffer-treated

controls after 24 h incubation are still dense in patient 11 and 12 and start to partially

disintegrate in patient 15. *PM-477*: After incubation with PM-477, the *Gardnerella* biofilm is

completely disintegrated in all patients. The disappearance of intense yellow fluorescent

Gardnerella bacteria unmasks the red fluorescence of still intact bacterial mass. Octenisept:

similar to PM-477, Octenisept massively reduces *Gardnerella* (yellow) but additionally nearly

completely eliminates any other non-*Gardnerella* bacteria (red). **B**, *Gardnerella* cell density

in samples of patients 1-15 expressed as the number of yellow bacteria counted in areas

containing 10 epithelial cells (see methods for a detailed description of the counting method).,

p-values were calculated with ANOVA in GraphPad Prism.

Effect of natural ELs on *Gardnerella spp.*

	B1				B2				B3				B4				B5				B7				B10				B11				B12			
	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$	$Gv\_ATCC\ 14018^T$	$Gl\_UGent\ 09.48$	$Gp\_UGent\ 18.01^T$	$Gs\_GS10234$
H1	-1.9	-1.3	-0.9	-3.5	-0.4	-0.3	-0.2	-1.4	-0.7	-0.7	-0.3	-1.9	-0.8	-0.7	-0.7	-2.3	-1.8	-0.8	-0.8	-3.3	-1.6	-1.2	-0.9	-3.3	-2.5	-2.3	-1.5	-4.9	-1.5	-1.6	-0.8	-3.8	-0.7	-0.6	-0.4	-2.3
H2	-1.6	-1.2	-1.0	-3.5	-1.2	-1.1	-2.5	-5.4	-2.9	-2.1	-1.6	-5.4	-2.3	-1.8	-1.6	-4.5	-1.1	-1.5	-2.7	-4.7	-1.0	-1.3	-2.6	-6.7	-3.6	-3.3	-3.8	-6.7	-3.3	-3.0	-3.7	-6.7	-2.8	-3.3	-3.6	-6.7
H3	-1.1	-1.3	-2.9	-4.3	-1.6	-1.4	-3.1	-4.1	-3.0	-2.0	-1.6	-5.4	-1.9	-1.4	-1.2	-4.0	-1.4	-1.2	-0.6	-3.5	-0.9	-1.1	-0.3	-2.8	-1.5	-2.2	-0.6	-3.8	-1.6	-2.0	-0.8	-4.2	-1.3	-1.9	-0.5	-3.7
H4	0.1	0.7	0.4	0.0	0.1	0.5	0.4	0.0	0.0	0.4	0.2	-0.3	-0.4	-0.5	-0.4	-2.0	0.1	0.3	0.5	0.1	0.1	0.1	-0.4	-1.3	-3.0	-2.9	-1.9	-4.8	-2.9	-3.3	-1.8	-4.8	-0.3	-1.1	-0.3	-3.9
H5	-1.3	-1.5	-1.1	-3.8	-1.2	-1.3	-0.9	-3.3	-2.8	-3.0	-1.8	-4.8	-1.8	-3.0	-2.2	-4.5	0.2	0.3	-0.1	-0.1	-1.6	-2.2	-1.2	-4.2	-2.5	-3.6	-1.1	-4.7	-2.4	-1.6	-1.6	-4.4	-2.3	-1.6	-1.6	-4.3
H6	-0.3	-0.1	-0.4	-1.8	-0.8	-0.4	-0.7	-2.4	-0.3	-0.5	-0.3	-1.6	-0.4	-0.7	-0.3	-1.6	-0.4	-0.5	-0.4	-1.7	-0.4	-0.8	-0.6	-2.0	-0.5	-0.7	-0.6	-1.8	-1.0	-1.5	-0.7	-2.7	-0.4	-0.6	-0.3	-1.3
H7	-0.9	-1.4	-0.9	-3.1	-2.0	-2.2	-1.3	-4.2	-3.2	-3.3	-2.0	-4.8	-2.5	-3.1	-1.7	-4.0	-1.6	-2.1	-1.2	-3.2	-1.6	-2.1	-1.2	-3.2	-3.4	-3.0	-1.8	-5.1	-3.5	-3.3	-1.9	-3.7	-2.6	-3.3	-1.7	-4.1
H10	-1.7	-1.1	-1.4	-3.2	-1.6	-1.2	-1.2	-3.9	-2.7	-1.9	-2.1	-4.9	-2.3	-1.5	-1.7	-4.2	-1.2	-0.8	-1.2	-3.5	-1.4	-0.8	-1.2	-3.5	-3.5	-1.6	-2.4	-5.5	-3.2	-1.7	-1.8	-4.6	-3.6	-2.8	-1.6	-5.0
H11	-0.8	-1.3	-0.8	-2.3	-0.2	-0.4	-0.4	-0.3	-2.5	-2.2	-1.7	-4.2	-1.7	-1.7	-1.1	-2.8	-1.5	-1.7	-0.9	-3.2	-0.1	-0.9	-0.7	-1.9	-2.8	-2.2	-1.4	-4.1	0.1	-1.6	-1.0	-3.5	-3.0	-2.3	-1.3	-4.1
H12	0.0	-0.8	-0.2	-1.5	-0.2	-0.9	-0.3	-2.1	-0.2	-1.2	-0.6	-2.5	0.0	-0.7	-0.3	-2.3	-0.2	-1.1	-0.4	-2.1	-0.8	-0.6	-0.4	-2.6	-1.3	-0.3	-0.4	-2.9	-2.4	-1.0	-1.0	-3.9	-0.2	-1.0	-0.3	-3.9

Effect of PM-477 on *Gardnerella spp.*

Effect of PM-477 on *Lactobacillus spp.*

Effect of PM-477 on other vaginal bacteria

Bacteriolytic effect of PM-477 on *G.vaginalis*

Bacteriolytic effect of PM-477 on _G.swidsinskii_

Bactericidal effect of PM-477 on *G.vaginalis*

Bactericidal effect of PM-477 on *G.swidsinskii*

a

	Baseline	Control	PM-477	Octenisept
Patient 11				
Patient 12			x 1000	
Patient 15				

Gardnerella