Simulation and prediction of spread of COVID-19 in The Republic of Serbia by SEIRDS model of disease transmission

*Slavoljub STANOJEVIC[a] , Mirza PONJAVIC[b] , Slobodan STANOJEVIC[c] , Aleksandar STEVANOVIC[d], Sonja RADOJICIC[e]

[a]Directorate of National Reference Laboratories, Batajnicki drum 10, 11080 Zemun, Serbia

[b]International Burch University, Francuske revolucije bb, Ilidza 71210, Sarajevo, Bosnia and Herzegovina, e-mail: mirza.ponjavic@gis.ba

[c]Veterinary Scientific Institute of Serbia, Janisa Janulisa 14, 11107 Belgrade, Serbia, e-mail: slobodan.vet@gmail.com

[d]University of Pittsburgh, Department of Civil and Environmental Engineering, 3700 O'Hara Street, Pittsburgh, PA 15261, USA, e-mail: stevanovic@pitt.edu

[e]Belgrade University, Faculty of Veterinary Medicine – Department of Infectious Diseases and Epidemiology, Bulevar Oslobodenja 18, 11000 Belgrade, Serbia, e-mail: sonjar@vet.bg.ac.rs

*Corresponding author: Slavoljub Stanojevic, e mail: slavoljub.stanojevic@minpolj.gov.rs

Abstract

As a response to the pandemic caused by SARS-Cov-2 virus, on 15 March, 2020, the Republic of Serbia introduced comprehensive anti-epidemic measures to curb COVID-19. After a slowdown in the epidemic, on 6 May, 2020, the regulatory authorities decided to relax the implemented measures. However, the epidemiological situation soon worsened again. As of 7 February, 2021, a total of 406,352 cases of SARSCov-2 infection have been reported in Serbia, 4,112 deaths caused by COVID-19. In order to better understand the epidemic dynamics and predict possible outcomes, we have developed an adaptive mathematical model SEAIHRDS (S-susceptible, E-exposed, A-asymptomatic, I-infected, H-hospitalized, R-recovered, D-dead due to COVID-19 infection, S-susceptible). The model can be used to simulate various scenarios of the implemented intervention measures and calculate possible epidemic outcomes, including the necessary hospital capacities. Considering promising results regarding the

development of a vaccine against COVID-19, the model is extended to simulate vaccination among

different population strata. The findings from various simulation scenarios have shown that,

with implementation of strict measures of contact reduction, it is possible to control COVID-19 and

reduce number of deaths. The findings also show that limiting effective contacts within the most

susceptible population strata merits a special attention. However, the findings also show that the

disease has a potential to remain in the population for a long time, likely with a seasonal pattern. If

a vaccine, with efficacy equal or higher than 65%, becomes available it could help to significantly slow

down or completely stop circulation of the virus in human population.

The effects of vaccination depend primarily on: 1. Efficacy of available vaccine(s), 2. Prioritization

of the population categories for vaccination, and 3. Overall vaccination coverage of the population,

assuming that the vaccine(s) develop solid immunity in vaccinated individuals. With expected basic

reproduction number of R_o=2.46 and vaccine efficacy of 68%, an 87% coverage would be sufficient to

stop the virus circulation.

Keywords: COVID-19, SEAIHRDS mathematical model, prediction, vaccination

1. Introduction

On 11 March, 2020, the World Health Organization characterised the disease caused by the

novel SARS-Cov-2 virus as a pandemic [1]. The Initial epidemic outbreak in China spread outside the

Wuhan area, and subsequently on a global scale. On 6 March, 2020, the first case of the novel

coronavirus infection was reported in the Republic of Serbia. Taking into consideration the escalation of

the disease and limited effects of the initially implemented measures, the state of emergency was

declared throughout the country on 15 March, 2020. Comprehensive anti-epidemic measures (e.g.

lockdown of entire country) were introduced in the entire country [2].

Due to the absence of specific pharmaceutical intervention, Serbia, like other countries,

implemented an anti-epidemic strategy based on physical distancing, school and university closures,

reduced number of workers present in the workplaces, closure of places of worship for public religious

services, reduced working hours of cafés and restaurants, avoiding mass gatherings, events, sports

games, tracing and identification of infected people's contacts, etc. After a slowdown in the epidemic, as

shown in the relevant officially published data, the regulatory authorities decided to relax the introduced

measures on 6 May, 2020. However, the epidemiological situation soon worsened again, resulting in the

reinstatement of some measures, as well as the introduction of new measures [2]. Although the return of

extensive measures has yielded favourable results, the further development of the epidemic is not clear.

For these reasons, mathematical modelling has a crucial role in understanding the epidemic and

predicting possible outcomes. Modelling is a particularly useful tool for devising strategies for combating

the epidemic, capacity planning, and selection of efficient measures, especially in the absence of specific

pharmaceutical treatments [3, 4, 5]. Mathematical modelling based on differential equations dates back to

the first half of 20^{th} century. In 1927, Kermack and McKendrick developed the basic model of disease

transmission consisting of three compartments: susceptible (S), infected (I) and removed (R). The model

was based on a connected system of nonlinear differential equations as a special case of the general

epidemiological model [6, 7]. Subsequent models, became more complex and adapted to the needs of

research [5].

Since the outbreak of COVID-19, many published papers have dealt with the implementation of

mathematical modelling and prediction of possible outcomes of COVID-19 epidemics. Most of these

research efforts have been based on the implementation of the SIR (susceptible-infected-removed)

model. Many of the other models provide a clear picture of dynamics of COVID-19 spreading, including

the overloading of the relevant health systems. For example, Ferguson et al., developed one of the first

models for COVID19 simulation, which was, among other things, used to plan the health care resources

[8]. Wu et al., developed the SEIR model to examine the dynamics of SARS-Cov-2 transmission from

person to person. This model was also used to calculate the basic reproduction number R_o, which we use

in this paper as one of the key parameters [9]. The classical SIR model assumes that there is

homogenous mixing of infected and susceptible populations and that the total population is constant and

does not change over time. Moreover, according to the classical SIR model, there is a monotonous

decline in susceptible population towards zero [10]. However, such assumptions are not objective in the

case of COVID-19 spreading and they are the basic problem in the modelling of this pandemic. In reality,

the human population fluctuates constantly [11]. In order to account for such fluctuation, and better

understand the COVID-19 epidemic in the Republic of Serbia, we have employed mathematical modelling

of the epidemic using the available data on the characteristics of the disease, such as incubation period,

latent period, recovery period, severity of clinical signs, and mortality rate caused by COVID-19.

Unlike the classic SIR model, the SEAIHRDS (S-susceptible, E-exposed, A-asymptomatic, I-

infected, H-hospitalized, R-recovered, D-dead due to COVID-19 infection, S-susceptible) epidemic model,

developed for this research, simulates the spreading of COVID-19 in an open population. Taking into

account that the population is constantly changing and that various measures are applied for different

strata or subgroups of the population (such as preschool children, children attending primary school, high

school students, students, employees, the unemployed and retirees), as well as changes in the intensity

of applied measures, we have proposed the use of a model that takes these circumstances into account.

Based on input disease parameters taken from scientific literature and specific data related to Serbia, this

model simulates daily disease occurrence, including the number of hospitalized patients and cases which

require intensive care. The model also predicts the expected number of deaths, as well as hospital

capacities necessary to accommodate the patients. It provides a possibility to simulate different scenarios

of disease control and intervention measures. Considering the expectations of successful development of

the vaccine against COVID-19 in the near future, we added an option to model vaccination of different

strata of the population as a set of disease control strategies.

2. Methodology

This section presents the research methodology and the proposed model, which were used to

predict the further dynamics of the epidemic in Serbia. We also presented the data that were used to

model the epidemic, a simulated strategy to combat COVID-19, and a sensitivity analysis.

2.1. SEAIHRDS mathematical model

Classical SEIRDS model divides the population into compartments, i.e. groups, and follows the

disease dynamics at all times. The population is divided into the following compartments: the portion of

the population susceptible to the infection is denoted by S, those latently infected with SARSCov-2

(exposed to) are denoted by E, the infected individuals who are able to spread the disease are denoted

by I, the ones recovered from the infection are denoted by R, and those who died due to disease with D.

Assuming that individuals mix homogenously, the force of infection λ (the rate at which susceptible

persons are infected) is related to *per capita* contact rate β. Also, the risk of infection is closely related to

the number of infectious individuals in the population I_t. It depends on the number of infectious individuals

(I_t) and how frequently they make contacts with other persons. In a situation of homogenous mixing

among the population, the force of infection λ can be express as follows:

$\lambda(t) = \beta I(t)$ (1)

The change of rates in every compartment per unit time in SEIRDS model is presented in the following

series of differential equations:

$S_{t+1} = bN_t - \beta S_t I_t - mS_t + \omega R_t$ (2)

$E_{t+1} = \beta S_t I_t - fE - mE_t$ (3)

$I_{t+1} = fE_t - (r+m)I_t - \delta I_t$ (4)

$R_{t+1} = rI_t - (m+\omega)R_t$ (5)

$D_{t+1} = \delta I_t$ (6)

where f is rate of onset of infectiousness expressed as the reciprocal of the latent infection period, r is

the rate at which infectious individuals are recovered, δ is the rate at which infectious individuals die from

COVID-19 infection and ω is rate of waning of immunity. The total population at any particular interval of

time t is:

$N_t = S_t + E + I_t + R_t + D_t + bN_{t-1} - mS_{t-1}$ (7)

where parameters b and m are *per capita* daily birth rate and death rate unrelated to COVID-19.

However, considering that implemented anti-epidemic measures against COVID-19 do not have

an identical impact on the population's age subgroups and that COVID-19 pathogenesis varies in different

age subgroups, we propose the use of multi-compartment version of standard SEIRDS model. The

model, named SEAIHRDS, monitors the dynamics of following compartments: susceptible individuals (*S*),

latently infected with SARSCov-2 (*E*) (exposed to/presymptomatic), asymptomatic infectious individuals

(*A),* infectious individuals with symptoms/clinically ill (*I*), hospitalized patients (*H*), recovered individuals

(*R*), and those who died due to disease (*D).* In this model the susceptible population was further stratified

within the compartment S according to age and occupations. Grouping into various strata was done

according to the real age structure of the Republic of Serbia population as follows: pre-school children

(S_{ps}), elementary school children (S_{es}), high school children (S_{hs}), college students (S_{cs}), unemployed

population (S_{ua}), employed population (S_{ea}), and elderly/retired (S_r) (Table 1). To simulate the epidemic

progression through different population strata-subgroups, we used appropriate, stratum-specific, model

parameters and factors of effective contact reduction (anti-epidemic intervention measures - ρ), which

were adapted to the relevant population groups: lockdown of the entire country, closures of pre-school,

school and college sessions closures, reduced number of workers allowed in the workplaces, work from

home, restrictions of mobility of elderly people, etc. During the simulation, we monitored the effects of

various levels of contact reduction, ranging from 20% to 75%, taking into account the realistic possibilities

of maintaining a minimum work process, functioning of the society and feasibility of such measures.

Given that intervention measures, applied in response to the emergence of COVID-19, are not

the same for all population strata, homogeneous mixing can be expected only within same population

stratum. The rate of effective contacts β, after the application of intervention measures, is no longer

identical at the level of all strata of the population. Effective contacts are limited by the intensity and types

of measures applied and are identical only when it comes to individuals within the same population strata.

Furthermore, persons in different population strata become infected at different rates depending on how

frequently they interact with other persons in their own subgroup and other subgroups. If we assume that

force of infection differs between different strata of population, the equation for the force of infection would

be as follows: $\lambda_i(t) = \sum_{j=1}^{n} \beta_{ij} I_j(t)$ \hfill (8)

where $\lambda_i(t)$ is force of infection in the i^{th} population strata, β_{ij} is the rate at which susceptible persons in the

i^{th} population strata and infectious persons in j^{th} population strata come into effective contact per unit of

time, *and $I_j(t)$* is the number of infectious persons in j^{th} population strata. Also, in this model the number of

recovered and dead is conditioned with different ages and genders.

Now our model will be expressed as follows:

$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) - \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) + \omega R_i(t) - m S_i(t)$ \hfill (9)

$\frac{dE_i(t)}{dt} = \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) + \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) - (m + f) E_i(t)$ \hfill (10)

$\frac{dA_i(t)}{dt} = f\alpha E_i(t) - (\gamma + m) A_i(t)$ \hfill (11)

$\frac{dI_i(t)}{dt} = f(1 - \alpha) E_i(t) - \sum_{j=1}^{n} \sigma_{ij} I_j(t) - (r + m) I_i(t)$ \hfill (12)

$\frac{dH_i(t)}{dt} = \sum_{j=1}^{n} \sigma_{ij} I_j(t) - \sum_{j=1}^{n} \delta_{ij} H_i(t) - (\varepsilon + m) H_i(t)$ (13)

$\frac{dR_i(t)}{dt} = \sum_{j=1}^{n} r_{ij} I_j(t) + \sum_{j=1}^{n} \gamma_{ij} A_j(t) + \varepsilon H_i(t) - (m + \omega) R_i(t)$ (14)

$\frac{dD_i(t)}{dt} = \sum_{j=1}^{n} \delta_{ij} I_i(t)$ (15)

where α is the proportion of asymptomatic cases, η accounts for the relative infectiousness of

asymptomatic carriers (in comparison to symptomatic carriers), r is the rate at which infectious individuals

whit symptoms are recovered, γ is the rate at which asymptomatic infectious individuals are recovered, σ

is rate at which infectious individuals are hospitalized, ε is the rate at which hospitalized patients are

recovered, δ is the rate at which infected individuals die from COVID-19 infection and ω is rate of waning

of immunity (Supplementary material).

2.2. Determining the herd immunity threshold and control of COVID-19 by vaccination policy

Considering the undergoing worldwide efforts to develop a vaccine against COVID-19 and

promising results, we extended the model to simulate and analysed the effects of a hypothetical

vaccination on the epidemic dynamics, and to estimate the extent of coverage of vaccination which can

interrupt the chain of infection. The control of COVID-19 by vaccination means targeting the entire

susceptible population with mass vaccination until critical herd immunity achieved. In such situation there

is a "race" between the exponential growth of epidemic and mass vaccination. The level of herd immunity

threshold (HIT) can be calculated by the following formula: $HIT = 1 - 1/R_o = (R_o - 1)/R_o$ (16)

and the critical vaccination coverage required to achieve herd immunity can be obtained by multiplying

herd immunity threshold with reciprocal value of vaccine efficacy, v_e:

$V_c = \frac{1}{v_e}(1 - \frac{1}{R_o})$ (17)

Most people infected with SARS-CoV-2 develop an immune response followed by the

development of specific antibodies between 10 and 21 days after getting infected [12]. Specific IgM and

IgG antibodies against SARS-CoV-2 develop 6 to 15 days after the onset of the disease [13-17].

According to some studies, the presence of antibodies has been confirmed in less than 40% of the

patients within 1 week after the onset of the disease, whereas percentage reaches 100% of subjects 15

days after the onset of disease [18]. Although duration of the immune response against CVOVID 19 is still

unknown, comparing with other coronaviruses, where immunity wane within 12 to 52 weeks after the first

symptoms appear [19, 20], while in the case of SARS-CoV-1 infection the presence of IgG antibodies was

confirmed in 90% and 50% of infected patients, respectively, over two and three years, respectively [29],

we assumed that durable immunity against COVID-19 is possible [20, 21]. Immunity to HCoVOC43 and

HCoV-HKU1 appears to wane appreciably within 1 year [21, 22], whereas SARS-CoV-1 infection can

induce longer-lasting immunity [23]. S. F. Lumley et. in a study conducted on 1,246 persons recovered

from COVID-19 found no symptomatic re-infections over 6 months [24].

Based on these findings, and fact that SARS-Cov-2 virus is also beta coronavirus, we assumed

that in the event of the development of a successful vaccine, immunity against the SARS-Cov-2 virus

could last for a year, as well as after recovery after a natural infection. In a study conducted to determine

the dynamics of SARS-Cov-2 transmission in the post pandemic period, Kissler et al. applied a similar

approach in defining the possible length of the immunity period [20].

For the purpose of modelling a control strategy based on vaccination, additional compartment to

the model was added denoted with V(t), in which there are vaccinated persons who have successfully

developed protective immunity after vaccination. The vaccination parameter, u, is the daily rate of

vaccination of susceptible population and it represents the proportion of susceptible population

immunized per unite time. The critical daily rate of vaccination, u_o is $u_c = (b+\omega)(R_o -1)$, required to

interrupt the infection [5]. The basic reproduction number under the vaccination is $R_{op} = (1-p)R_o$. The

proportion of effectively protected persons, p, is conditioned by parameter the vaccine efficacy, v_e. This

parameter represents a proportion of person who successfully developed protective immunity after

vaccination, whereas total number of actively protected individuals in time t is V(t) = number of vaccinated

x v_e [5]. In this compartment the daily rate of waning of immunity at which immunity of vaccinated

population fades out is ω, and it is reciprocal to the period of lasting of immunity. Vaccinated persons,

after losing their immunity, become sensitive again and removed to the compartment S. The change of

rate in this compartment per unit time is as follows:

$\frac{dV_i(t)}{dt} = v_e S_i(t) - mV_i - \omega V_i.$ (18)

The compartment S(t) is slightly modified as follows:

$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) + \omega R\_i(t) - mS_i(t) - v_e S_i(t) + \omega V_i.$ (19)

The other of compartments of SERIDS model remain unchanged.

2.3.   Model parametrisation

In proposed model, β is *per capita* effective contact rate at which specific persons come into

effective contact per unit of time. An effective contact is defined as a contact sufficient to cause disease

transmission [10, 24, 26]. We calculated the parameter β by using the formula:

$\beta = \mathrm{R_o}/\mathrm{NT_R}$ (20)

where $\mathrm{R_o}$ is a basic reproduction number of the disease, i.e. the average number of newly infected people

with COVID-19 (secondary infection cases), infected by one infectious individual in a totally susceptible

population, N is total population, and $\mathrm{T_R}$ is the average duration of infectious period [10, 25, 26]. The R

values of 2.46 and 3.1 are adopted from the relevant literature. The $\mathrm{R_o}$ values were based on the data

obtained during the initial phase of the epidemic in Italy [27]. Since the implemented measures and

disease transmission were simulated through various population strata, we corrected the β parameter

with a relevant, stratum-specific contact reduction factor ρ_i. In this way, we obtained the *per capita* contact

rate specific for each separate stratum based on following formula

$\beta_i = \beta(1-\rho_i)$.   (21)

The values of the ρ_i factor in different population strata ranged from 0.25 to 0.75 (effective contact

reduction ranged from 25% to 75%).

Parameters such as daily birth and death rates were calculated based on the data published by

the Office of Statistics of the Republic of Serbia, and data published by the World Bank regarding the life

expectancy in the Republic of Serbia [28, 29]. The latter study reported that the life expectancy in Serbia

was 79.06 years in 2017 [29]. By using this figure, we expressed the daily death rate as a value reciprocal

to life expectancy m = 0.000036. We calculated the daily *per capita* birth rate of b= *0.000025* based on

the figure of 9.2 births in the Republic of Serbia per 1000 people in a year. These estimates were needed

to realistically simulate fluctuations of the total population. To simplify the calculations, we assumed that

the general morality rate m is applicable for all population strata.

The infectivity rate, i.e. the rate of transfer from compartment E to I, was derived as a value

reciprocal to the COVID-19's average latent period. The data on the average duration of latent infection

(f^{-1}) and the average period during which an infected person is shading the SARS-Cov-2 virus (T_R) were

adopted from the relevant literature as f^{-1} = 3.5 days [30, 31] and T_R = 9.3 days, respectively [30]. Also

the data on the percentage of hospitalized patients and those whose therapy requires intensive care,

used for prediction of required hospital capacities, as shown in Table 1, were taken from literature [30, 8].

Parameters such as δ and r are related to the infectious fatality rate (IFR) for COVID-19 and

average times taken from onset of symptoms to death (T_D) or recovery (T_R). These parameters were

calculated using the following formulas:

$\delta = IFR_{stratum}/T_D$ and   (22)

$r = (1\text{-}IFR_{stratum})/T_R$ (23)

The IFRs, shown in Table 2, were taken from literature and compared with local IFR value which

was calculated based on officially registered deaths published by the health system of the Republic of

Serbia [2]. The Calculation of local IFR is presented in section 2.4. Population data, (e.g. total population,

age structure, and stratification) are presented in Tables 1 and 2. A summary of all model parameters is

given in Table 3.

2.4. Setting disease control scenarios

Five different scenarios were developed for simulating the COVID-19 epidemic control based on

non-pharmaceutical interventions. SC1 implies a base-case scenario where the epidemic spreads in

susceptible population without any anti-epidemic measures being implemented. In the other scenarios,

the extent of contacts was reduced, for each population stratum separately, according to objective

possibilities and measures which were implemented during the actual epidemic in the Republic of Serbia.

Scenarios are presented in Table 4.

The timing of the simulation of anti-epidemic measures, i.e. the reduction of individual contacts

corresponds to the actual date when the implementation of measures in the real epidemic in Serbia

began (March 15, 2020). Considering that it is not realistic to expect the desired level of reduction of

physical contacts to be achieved in one day, in each scenario, a period of gradual introduction of

measures was simulated (lag period of 7 days). Since each individual scenario was simulated at R_0 of

2.46 and 3.10, different contact reduction rates were applied accordingly.

Additional four scenarios of control of COVID-19 were simulated based on vaccination policy. We

assumed that vaccine efficacy was 50%, 68%, 80%, and 85%. The initial conditions assumed that all

other anti-epidemic measures are excluded from the model and replaced with mass vaccination.

Indicators of epidemic dynamics were monitored, such as: CI, hospitalized patients, patient in intensive

care units and deaths.

2.5.   Model sensitivity analysis and calibration

Considering the world experience with detection of COVID-19 cases, as well as the unreliable

data on COVID-19 infections which are currently available worldwide, model calibration is very

challenging, and can result in obtaining inaccurate values for the parameters [40]. This is especially due

to the facts that a significant percentage of the infected individuals do not exhibit any symptoms. The

other issue is small percentage of tested population [40].

As part of the national seroepidemiological study, 1,006 subjects were tested in Serbia from May

11[th] to June 25[th], 2020, for the purpose of estimating the extent of COVID-19 infection. According to the

published data, seroconversion was confirmed in 6,4% of the subjects. On the other hand, a total of

13,372 cases of the infection were reported, which means that those who were infected constitute around

0.19% of the overall population. However, it is our opinion that the data on reported deaths caused by

COVID-19 infection is more reliable for use in model calibration, e.g. infection fatality rate. Alex et al.

reached a similar conclusion when simulating COVID-19 by using the SEIRD model with heterogeneous

diffusion [40]. When we compared the data recorded during the beginning of the epidemic in Serbia with

the results obtained during the simulation, such as the initial doubling time, the two data series matched

well. However, later, the obtained results did not match well the officially registered data on the number of

infected, especially after the beginning of the implementation of measures in Serbia. We attribute these

differences to the methodology by which official authorities register cases of infection, and collect the

data.

The parameter that determines the number of deaths is the IFR. It is the number of persons who

die of the COVID-19 among all infected individuals regardless of whether the infected show symptoms of

the disease or not. As with many diseases, IFR is not always equivalent to the number of reported deaths

caused by COVID-19. This is because a significant number of deaths, although caused by COVID-19, will

not be recognized as deaths caused by COVID-19 [41]. Also, there are many asymptomatic cases of

infection which are never detected [42, 43, 44, 45].

However, according to new findings, the overall estimate of the proportion of people who become

infected with SARS-CoV-2 and remain asymptomatic throughout infection was 20% (95% confidence

interval) with a prediction interval of 3%–67% in 79 studies that addressed this question [46, 47]. Michael

A. Johansson et al. reported that 30% of individuals with infection never develop symptoms and are 75%

as infectious as those who do develop symptoms, and concluded that persons with infection who never

develop symptoms may account for approximately 24% of all transmission [39].

Due the fact that there is a lag in time between when people are infected and when they die,

patients who die on any given day were infected much earlier, and thus the denominator of the mortality

rate should be the total number of patients infected at the same time as those who died [41]. David et al.

estimated mortality rate by dividing of deaths on a given date by the total number of persons confirmed as

COVID 19 cases 14 days before [29, 41]. It is based on the assumption that maximum incubation period

is14 days [34]. However, if we take into account that the number of registered cases of COVID-19

infection is usually significantly lower than the actual number, assuming that the data on deaths are

accurate, the real IFR value is significantly lower than the calculated value [40]. If we apply this to the

situation in Serbia, the daily value of IFR on July 10[th], 2020, when the largest number of deaths was

registered in one day, was 9.33%, considering that 18 deaths were registered on July 10[th] and 14 days

earlier 193 confirmed cases of COVID-19 infection. The raw values of IFR for the period between March

6[th] and August 10[th] were as follows: median of IFR = 2.11%, and average value of IFR = 7.15% bounded

in interval 4.17%-10.13%. When we compared these values with those published by the WHO, CDC and

other authors [43, 44, 48] we concluded that they differ significantly. Considering these findings, the IFR

values adopted in the model (for various population groups and genders) were primarily taken from the

literature, with a remark that the selection of IFR values was based on preliminary comparison of the

overall Serbian IFR with similar IFRs found in the literature, taking into account the registered deaths and

most probable number of infected individuals [33]. To make this possible, the first step was to correct the

local raw IFR value mentioned above. Based on real data we first calculated the population at risk of

dying from COVID-19 infection for each individual day since the outbreak, ending on August 10[th], 2020.

The number at risk on a given day should correspond to the number of deaths from COVID-19 infection,

considering the lag period from infection to death. For this calculation, we used the data on the number of

deaths D_t in Serbia registered on daily bases [2]. We hypothesized that the distribution of time periods t_n

from the moment of COVID19 infection to death follows the lognormal distribution defined by the

parameters $m = 26.8$ and $\sigma = 12.4$ days [48, 49].

Based on the formula: $I_{r(t)} = \sum_{j=1}^{n} I_{(t-n)} * m_{(t-n)}$ by reverse, we calculated the population at risk of

dying from COVID-19 infection for each individual day, where m_t is the probability that the time between

infection and death is t days and follows the lognormal distribution ($m = 26.8$, $\sigma = 12.4$) [48, 49]. After that,

the daily IFR values were calculated according to the formula $IFR_{(t)} = D_t / I_{r(t)}$ [50]. Based on IFR values

calculated in this way, we made descriptive statistics and obtained the mean value of IFR = 0.70%,

bounded in the interval 0.46-0.94% and a median of 0.19%. It is important to note here that this value

corresponds with the COVID-19 IFR values found in Eastern European countries and Spain [50, 33].

Taking these findings into account, we decided to take the IFR values specific to certain population strata

recorded in Spain as the most appropriate for our case [33]. The adopted IFR values are listed in Table 3.

In this section, we used sensitivity analysis to estimate the amount of change on outcomes when

varying the input values used in the model. Sensitivity analysis is one of the methods most frequently

used for the evaluation of disease spread models [51, 52]. A sensitivity analysis is carried out to

characterise the impact of uncertainty of input parameters on model outputs. Sensitivity analysis consists

of assessing the impact that changes in input parameters have on model outcomes. We evaluate two

aspects of the model: the global behavior of the model when perturbing a group of key parameters

together, and the impact of changes when perturbing the key parameters used in the model individually.

The model sensitivity analysis was conducted by changing the most sensitive model parameters:

$R_0, \beta, f\text{-}1, r, b, m, \gamma, \varepsilon, \eta, \omega, \bar{\tau}_d$. The values of these parameters were increased by 5%, 10% and 25%

relative to the base scenario and changes in output indicators were observed.

Finaly, validation of the model was performed by comparing the historical data of the real

epidemic in Serbia with the data obtained with the SEAIHRDS model. For validation purposes, the current

epidemic of COVID-19 was simulated, along with actual anti-epidemic measures. Taking into account the

risks described above related to the accuracy of historical data from the COVID-19 epidemic in Serbia

and significantly higher confidence in the accuracy of data related to the number of deaths from COVIDA-

19 compared to data on the number of infected, for the purpose of model validation, only data on the

cumulative number of deaths were used. The reason for this assumption is that there is still uncertainty

about the proportion of the infected population that is not reported due to a mild form of the disease or

the patients are asymptomatic.

In statistical analysis the coefficient of determination, R^2, was used to check goodness of fit of

SEAIHRD model with COVID-19 data recorded during the real epidemic. The regression coefficient

compares predicted values (y) against actual data (x). To address model uncertainties, bias, mean

absolute error (MAE), mean square error (MSE), the root mean square error (RMSE), maximum deviation

(MaxDev) and normalized root mean square error (NRMSE) were also estimated (*Supplementary*

material).

3. Results

3.1. Predicting the number of sick, hospitalized patients and deaths caused by COVID-19 in the

absence of any intervention measures

After the simulation, the model predicted that with $R_o = 2.46$, and without the implementation of

any anti-epidemic measures, the initial doubling time of the infection could be five days. The epidemic

wave could peak 219 days after the outbreak, and it could yield 99,819 infected individuals in a day.

Afterwards, the infection rate could decline for 215 days, eventually reaching the daily incidence of 492

newly infected, after which the next epidemic wave could start. The second wave could peak 706 days

after the onset of epidemic and yielding 25,232 infected individuals in a day. The third epidemic wave

could peak 429 days later, with 13,709 infected individuals in a day. The true cumulative incidence in the

first year of the epidemic could be 6,229,144 infected people with SARS-Cov-2 virus, while the apparent

cumulative incidence could be 4,360,401 infected. A total of 20,894 patients could die due to COVID-19

consequences.

With R_o = 3.1, the following results were obtained: the initial doubling time of the infection was five

days, true cumulative incidence 7,133,221, apparent cumulative incidence 4,993,254, and the total

deaths of 23,951. Fig. 1, panels a) and b) show daily variations in the number of susceptible, latently

infected, infected and recovered patients, at basic reproduction numbers of R_o=2.46 and R_o=3,

respectively. Panels c) and d) of the same figure show daily fluctuations in susceptible, recovered and net

reproduction rates R_n for R_o=2.46 and R_o=3, respectively. Panels e) and f) of Fig. 1 show daily variations

in R_n, true and apparent disease incidences at basic reproduction numbers of R_o=2.46 and R_o=3. The

shaded area corresponds to the period when the daily number of new COVID-19 infected individuals

increasing, and therefore all of the following hold: R_n>1, proportion susceptible >1/R_o and the proportion

of population that is recovered (immune) is below the herd immunity threshold. Fig. 2 panels a) and b)

show a prediction of necessary hospital capacities. Panels c) and d) of Fig. 2 show predicted numbers of

sick and dead due to COVID-19 at R_o=2.46 and R_o=3 and age structure of hospitalized patients and

deaths.

3.2.   Predicting the number of sick, hospitalized patients and deaths caused by COVID-19 in the

conditions of application of restrictive anti-epidemic measures

When the spread of COVID-19 epidemic through totally susceptible population in the Republic of

Serbia is simulated, under an assumption of only incidental movement among the population, basic

reproduction number of R_o = 2.46, and with lock-down of entire country, a significant slowdown of the

epidemic was observed. Initial infection doubling time was 6 days. The peak of the epidemic wave could

occur 702 days after the epidemic onset, when there could be 2,848 infected in one day. In the first year

of the epidemic 308,581 people could be infected and 1,031 people could die.

When the basic reproductive number was increased to R_0 = 3.1, and for certain segments of the

population the contact reduction factor ρ_i increased compared to the values from the scenario with R_0 =

2.46 (reduction of contacts in public places and contacts of persons over 65 years by 55% and 65%,

respectively), the results changed significantly. The model predicted that with R_0 = 3.10, the initial

infection doubling time could be 5 days, the peak of epidemic wave could occur after 246 days and it yield

8,110 infected people in one day. In the first year of the epidemic 2,219,251 people could be infected and

7,194 could die.

Table 5 provides the overview of epidemic indicators obtained from the simulations of all five

scenarios with R_o = 2.46 and R_o = 3.1. Figures 3. and 4 provide a comparative overview of results of all

five simulated scenarios. Panels a) and b) of Fig. 3 show the values of cumulative incidences on a daily

basis for R_o=2.46 and R_o=3.1, respectively. Panels b) and c) of Fig. 3 show the expected total number of

hospitalized patients and patients in intensive care units (ICU) for R_o=2.46 and R_o=3.1, respectively. Fig.

4 provides overview of required hospital capacities e.g. hospital bed occupancy and the occupancy of

beds in ICU on daily bases for R_o=2.46 and R_o=3.1, respectively. The results show that after applying

various measures to slow down the circulation of SARS-Cov-2, the number of newly infected people,

hospitalized patients and the occupancy of hospital capacities are the lowest in the situations where

rigorous anti-epidemic measures are applied to all population strata (Scenario 2 in Table 5 and in Fig. 3

and Fig. 4). Openings of pre-school and elementary school's facilities leads to a visible jump in the

number of infected and hospitalized in all strata. This finding clearly shows that children, although least

susceptible to developing more severe clinical pictures, are important when transmitting SARS-Cov-2

(Scenario 3 in Table 5 and Fig. 3). Opening of the high schools and colleges also leads to a visible

increase in the number of newly infected and hospitalized patients, including an increase in the number of

deaths (Scenario 5 in Table 5 and Fig. 3). Without the application of any intervention measures, the

greatest burden on the health system could be expected 228 days from the beginning of the epidemic at

R_o=2.46 or 168 days at R_o=3.1.

Depending on the R_o value used in simulation, it could be necessary to provide 42,351 (61,739)

hospital beds and an additional 20,173 (28,740) in intensive care units. In the case of Scenario 2, there is

a significant slowdown in the epidemic. According to the predictions obtained by the simulation of

Scenario 2, after 713 (259) days at the moment of maximum occupancy of hospital capacities, it might be

necessary to provide 1,387 (3,940) beds in COVID-19 hospitals and 784 (2,162) beds in intensive care

units (Fig. 4).

3.3. COVID-19 simulation and disease control by implementing a hypothetical vaccine

Assuming that the disease is spreading at the basic reproduction number of R_o=2.46, the herd immunity threshold (when the disease can be expected to slow down and the chain of infection is expected to be broken) would be 59.35%, while at R_o=3.1, it would be 67.74%. Depending on the efficacy of the potential vaccine, the required vaccination coverage should be 87% (v_e =68%), 74.19% (v_e =80%), 69.82% (v_e=85%), 65.94% (v_e=90%) and 95.41% (v_e=71%), 84.68% (v_e=80%), 79.70% (v_e=85%), 75.27% (v_e=90%) for R_o=2.46 and R_o=3.1, respectively. The different ways of including vaccination in the SEAIHRD model are detailed in the supplementary material. Fig. 5 shows different scenarios of COVID-19 control strategies based solely on vaccination.

3.4. Results of the model sensitivity analysis

The results of the sensitivity analysis are presented in tables 6 and 7. The tables show increased values of input parameters and the percentage of the parameter increase relative to the basic scenario, as well as the values of output results obtained after the simulation of the changed scenario.

3.5. Results of the model validation

The model was validated on the most recent historical data (from January 1, 2021 to February 1, 2021). As shown in Table 8. all measures of the prediction quality of deceased due to COVID-19 are low. The average difference between the actual number of people who died of COVID-19 and predicted one is only 2.05% and the maximum deviation between the predicted and actual number will not exceed 4.82% with a probability of 95%.

Based on Fig. 6 and Table 9, we can conclude that the SEAIHRD model fits historical data quite well. Pearson's r and coefficient of determination (R^2) have shown strong the linear relationship between real deceased and the number of deaths predicted by SEAIHRD model.

More information on validation results are detailed in the supplementary material.

4. Conclusions and discussion

For the needs of this research, we augmented the classic deterministic model by adding the compartments of vaccinated, asymptomatic, hospitalized and latently infected subjects. By adding birth and death rates, we enabled daily fluctuations of the overall simulated population, which brought us closer to the real conditions in which the disease is transmitted. When we assumed that the recovered lose immunity over time, we obtained dynamic oscillations of epidemic waves through susceptible population.

The input values for the parameters used to simulate the COVID-19 were obtained either from literature review or were calculated on the basis of data have taken from the literature and other official sources. Whereas some of these inputs are well documented; other input values are either not so well documented or based on potential subjective opinions (i.e., expert opinion, historical data from epidemic etc.). In any case, well documented or not input values have potential to impact results and, therefore, should be carefully evaluated. The results of conducted sensitivity analysis show that the sources of uncertainty are different for each output considered and it is necessary to consider multiple output variables for a proper assessment of the model. The most influential parameter is r than R_0 and f^{c-1} .

The model was tested on Serbian COVID-19 statistic data and obtained validation results allow us to conclude that the proposed model has good prediction ability and performance. However, although we obtained satisfactory results during the validation of the model, worth noticing also that some of the model parameters were estimated based on available data that might be less precise due to the difficulty of being measured. That could be the reason why the values of some parameters e.g. recovery rate, contact rate, daily infection rate, that are estimated in hospitals may differ from those acquired by this study. For that reasons SEAIHRD model can be used for the long-term rough predictions of the epidemic. Obtained long-term predictions reflect the general dynamic of the outbreak and are especially useful for the healthcare system workers and governmental officials.

When we simulated different disease control scenarios of the COVID-19 epidemic based on non-pharmaceutical intervention measures, scenario number 2 proved to be the most effective approach to the disease control, because it implemented the most comprehensive anti-epidemic measures (entire

country lock down). However, the basic problem of this approach is the feasibility and practicability to

maintain the measures in the long term.

The model predicted that students, children and younger school-age generations have an

important role in transmitting COVID-19, especially if they come into contact with a more vulnerable

population. The model showed that, in the case of returning school children of all ages to schools, an

increase of 10.48% in the estimated deaths and 12.16% of the number of infected is possible, when

compared to the conditions before opening of the schools (Scenarios 2 and 4). However, most dead and

seriously diseased people are found in the older population. This is particularly important when planning

intervention measures, especially when deciding on which restrictions to be lifted and how (opening

schools, students' return to faculties etc.). The model predicted that COVID-19 has a potential to spread

rapidly and linger in population. Due to a large number of the infected persons and duration of the

disease, there are significant needs for hospital capacities, especially in the conditions when the disease

is suppressed by implementing partly relaxed anti-epidemic measures, or in the case of absence of any

measures. According to the prediction, without the application of any intervention measures, at the

moment of the greatest load, depending of actual R_o, the health system should provide 42,351 (R_o=2.46)

hospital beds for the care of the patients and an additional 20,733 in intensive care units. On the other

hand, in the case of the application of the strictest anti-epidemic measures, the needs decrease to only

1,387 beds in COVID-19 hospitals and additional 784 beds in intensive care units. In the case of

continued implementation of current measures, which are significantly less intense than the measures

applied at the beginning of the epidemic, it is necessary to provide 3,537 beds in COVID-19 hospitals and

1,945 beds in intensive care units in the entire country. The model also shows that the needs for hospital

capacities decline with the ending of the first epidemic wave, since daily incidence decreases and during

the second and third waves it never reaches the initial peaks, but these needs still remain substantial. For

example, in the case of Scenario 1, at the top of the second epidemic wave, it is necessary to provide

11,845 beds in COVID-19 hospitals and 6,378 in intensive care units, which makes 27.96% and 30,76%

of the required capacities of the first wave of the epidemic.

Based on the cyclical patterns of the epidemic waves and duration of simulated epidemics, the

model predicted that the disease has a potential to linger in population and that it will most probably have

a seasonal pattern. Therefore, potential vaccines can have an enormous potential and significance for

COVID-19 control. Depending on the efficacy of future vaccines, the disease can be stopped and curbed

almost solely by implementing the measure of vaccination. However, the necessary conditions for these

predictions and expectations are the efficacy of potential vaccines and the ability of a health systems to

implement vaccination to a satisfactory extent and rapidly, especially with regards to the most sensitive

categories of population. Depending on the R_o, a vaccine that would have an efficacy $\geq$ 68-71% could

stop the pandemic and break the chain of infection. However, even vaccines with lower efficacy could be

useful as they would significantly reduce the number of cases and deaths, especially if used in

combination with the other disease control measures. The effects of vaccination depend primarily on: 1.

Efficacy of available vaccine(s), 2. Prioritization of the population categories for vaccination, and 3.

Overall vaccination coverage of the population, assuming that the vaccine(s) develop solid immunity in

vaccinated individuals. With expected basic reproduction number of Ro=2.46 and vaccine efficacy of

68%, an 87%- coverage would be sufficient to stop the virus circulation. The required minimal vaccination

coverage should be 87% (v_e =68%), 74.19% (v_e =80%), 69.82% (v_e=85%) and 95.41% (v_e=71%), 84.68%

(v_e=80%), 79.70% (v_e=85%) for R_o=2.46 and R_o=3.1, respectively. The minimum daily vaccination rate

should be 0.47% for vaccines with an efficiency of 85%, and 0.59% for vaccines with an efficiency of

68%.

Based on the obtained results, we can conclude that at this point, without the application of

specific pharmaceutical products, COVID-19 suppression is highly dependent on the basic reproduction

number (R_o), and that more intensive contacts and relaxed measures can result in a dramatic spread of

the virus. The choice of intervention measures depends on the feasibility of their implementation and their

efficacy in different social contexts. COVID-19 will likely have to be suppressed in this way for a certain

period of time. This will most probably last until sufficient quantities of a reliable and effective vaccine are

available, and thereafter until optimal vaccination coverage is achieved.

Funding: This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

References

[1] World Health Organization (WHO), Coronavirus disease (COVID-19) outbreak situation. 2020.

https://www.who.int/emergencies/diseases/ novel-coronavirus-2019, [accessed 18 August 2020].

[2] https://covid19.rs, 2020, [accessed 18 August 2020].

[3] Wangping J, Ke H, Yang S, Wenzhe C, Shengshu W, Shanshan Y, Miao L. Extended SIR prediction

of the epidemics trend of COVID-19 in Italy and compared with Hunan, China. Frontiers in Medicine 2020;

7:169.

[4] Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, Davies N. Early dynamics of

transmission and control of COVID-19: a mathematical modelling study. The Lancet infectious diseases

2020. Doi: 10.1016/S1473-3099(20) 30144-4.

[5] Matt J. Keeling, Pejman Rohani. Modelling Infectious Diseases in Humans and Animals, Princeton

University Press, 3 Market Place, Woodstock, Oxfordshire OX20 1SY, p. 8-10.

[6] Kermack, W. and A. McKendrick (1927). A contribution to the mathematical theory of epidemics.

Proceedings of the Royal Society. A 115, p. 700–721.

[7] Barnes B. and G. R. Fulford. Mathematical modelling with case studies Using Maple™ and MATLAB,

CRC Press Taylor & Francis Group 6000 Broken Sound Parkway NW, Suite 300, Boca Raton, FL 33487-

2742, p. 100-102.

[8] Imperial College COVID-19 Response Team. Report 9: Impact of non-pharmaceutical interventions

(NPIs) to reduce COVID-19 mortality and healthcare demand; 2020.

https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-

COVID19-NPI-modelling-16-03-2020.pdf.

[9] Wu Z, Googan JMM. Characteristics of and important lessons from the coronavirus disease 2019

(COVID-19) outbreak in China: Summary of a Report of 72□314 Cases from the Chinese Center for

Disease Control and Prevention. JAMA. 2020;323(13):1239-1242. Doi:10.1001/jama.2020.2648.

[10] Emilia Vynnycky, Richard G. White with an introduction by Paul E.M. Fine. An Introduction to

Infectious Disease Modelling. Oxford University Press Inc., New York © Oxford University Press; 2010.

[12] European Centre for Disease Prevention and Control. Immune responses and immunity to SARS-

CoV-2; 2020. https://www.ecdc.europa.eu/en/covid-19/latest-evidence/immune-responses; [accessed 18

August 2020].

[13] Woelfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Mueller MA, et al. Clinical presentation

and virological assessment of hospitalized cases of coronavirus disease 2019 in a travel-associated

transmission cluster. medRxiv. 2020; https://doi.org/10.1101/2020.03.05.20030502.

[14] Liu W, Liu L, Kou G, Zheng Y, Ding Y, Ni W, et al. Evaluation of Nucleocapsid and Spike Protein-

based ELISAs for detecting antibodies against SARS-CoV-2. medRxiv. 2020.

https://doi.org/10.1101/2020.03.16.20035014.

[15] Long Q-x, Deng H-j, Chen J, Hu J, Liu B-z, Liao P, et al. Antibody responses to SARS-CoV-2 in

COVID-19 patients: the perspective application of serological tests in clinical practice. medRxiv.

https://doi.org/10.1101/2020.03.18.20038018.

[16] Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-CoV-2 in patients

of novel coronavirus disease 2019. medRxiv. 2020; https://doi.org/10.1101/2020.03.02.20030189.

[17] Okba NMA, Muller MA, Li W, Wang C, Geurtsvan Kessel CH, Corman VM, et al. SARS-CoV-2

specific antibody responses in COVID-19 patients. medRxiv. 2020;

https://doi.org/10.1101/2020.03.18.20038059.

[18] Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-CoV-2 in patients

of novel coronavirus disease 2019. medRxiv. 2020; https://doi.org/10.1101/2020.03.02.20030189.

[19] Kellam P, Barclay W. The dynamics of humoral immune responses following SARS-CoV-2 infection

and the potential for reinfection. The Journal of general virology. 2020 May 20.

[20] Kissler et al. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period,

Science 10.1126/science.abb5793 (2020).

[21] K. A. Callow et al. The time course of the immune response to experimental coronavirus infection of

man. Epidemiol. Infect. 105, 435–446 (1990)

[22] J. M. Dan et al. Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection

Science 371, eabf4063 (2021). DOI: 10.1126/science.abf4063.

[23] K.-H. Chan et al. Cross-reactive antibodies in convalescent SARS patients' sera against the

emergingnovel human coronavirus EMC (2012) byboth immunofluorescent and neutralizingantibody tests

J. Infect. 67, 130–140 (2013).

[24] Sheila F Lumley et al. Antibodies to SARS-CoV-2 are associated with protection against reinfection.

medRxiv preprint doi: https://doi.org/10.1101/2020.11.18.20234369.

[25] Gerardo Chowell, James M. Hyman, Lu´ıs M. A. Bettencourt, Carlos Castillo-Chavez. Mathematical

and Statistical Estimation Approaches in Epidemiology, ISBN 978-90-481-2312-4, Doi 10.1007/978-90-

481-2313-1, Springer Dordrecht Heidelberg London New York; 2009.

[26] Michael Thrusfield. Veterinary epidemiology. Third edition. Blackwell Science Ltd, a Blackwell

Publishing company; 2007.

[27] Marco D'Arienzo, Angela Coniglio. Assessment of the SARS-CoV-2 basic reproduction number, R0,

based on the early phase of COVID-19 outbreak in Italy. Biosafety and Health 2(2020) 57-59.

[28] Demografski pregled 2017. Broj: 68/2018, https://www.minrzs.gov.rs/sites/default/files/2019-

01/demografski_pregled_68.pdf; [accessed 15 July 2020].

[29] The World Bank. World Developments Indicators;

https://www.google.com/publicdata/explore?ds=d5bncppjof8f9_&met_y=sp_dyn_le00_in&idim=country:S

RB:HRV:BIH&hl=en&dl=en#!ctype=l&strail=false&bcs=d&nselm=h&met_y=sp_dyn_le00_in&scale_y=lin&

ind_y=false&rdim=world&idim=country:SRB&ifdim=world&hl=en_US&dl=en&ind=false; [accessed 25

August 2020].

[30] Byrne AW et al. Inferred duration of infectious period of SARS-CoV-2: rapid scoping review and

analysis of available evidence for asymptomatic and symptomatic COVID-19 cases. BMJ Open

2020;10:e039856. doi:10.1136/ bmjopen-2020-039856].

[31] Shujuan Ma et al., Epidemiological parameters of coronavirus disease 2019: a pooled analysis of

publicly reported individual data of 1155 cases from seven countries. medRxiv 2020;preprint doi:

https://doi.org/10.1101/2020.03.21.20040329].

[32] Robert Verity et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis.

Lancet Infect Dis 2020; 20: 669–77, https://doi.org/10.1016/ S1473-3099(20)30243-7

[33] Roberto Pastor-Barriuso et al. Infection fatality risk for SARS-CoV-2: a nationwide

seroepidemiological study in the noninstitutionalized population of Spain. medRxiv. 2020; preprint doi:

https://doi.org/10.1101/2020.08.06.20169722.

[34] Marco D'Arienzo, Angela Coniglio. Assessment of the SARS-CoV-2 basic reproduction number, R0,

based on the early phase of COVID-19 outbreak in Italy. Biosafety and Health 2(2020) 57-59.

[35] McAloon C. et al. Incubation period of COVID-19: a rapid systematic review and meta-analysis of

observational research. BMJ Open 2020;10:e039652. doi:10.1136/ bmjopen-2020-039652.

[36] Pierre-Yves Boëlle at al. Trajectories of Hospitalization in COVID-19 Patients: An Observational

Study in France. J. Clin. Med. 2020, 9, 3148; doi:10.3390/jcm9103148.

[37] Natalie M. Linton at al. Incubation Period and Other EpidemiologicalCharacteristics of 2019 Novel

Coronavirus Infections with Right Truncation: A Statistical Analysis of Publicly Available Case Data. J.

Clin. Med. 2020, 9(2), 538; https://doi.org/10.3390/jcm9020538.

[38] Ma S, Zhang J, Zeng M, et al. Epidemiological parameters of coronavirus disease 2019: a pooled

analysis of publicly reported individual data of 1155 cases from seven countries. medRxiv 2020.

[39] Michael A. Johansson et al. SARS-CoV-2 Transmission From People Without COVID-19 Symptoms.

JAMA Network Open. 2021;4(1):e2035057. doi:10.1001/jamanetworkopen.2020.35057.

[40] Alex Viguerie, Guillermo Lorenzo, Ferdinando Auricchio, Davide Baroli, Thomas J.R. Hughes, Alessia

Patton, Alessandro Reali, Thomas E. Yankeelov, Alessandro Veneziani. Simulating the spread of COVID-

19 via a spatially-resolved susceptible–exposed–infected–recovered–deceased (SEIRD) model with

heterogeneous diffusion. Applied Mathematics Letters 111 (2021) 106617.

[41] David Baud, Xiaolong Qi, Karin Nielsen-Saines, Didier Musso, Léo Pomar, Guillaume Favre. Real

estimates of mortality following COVID-19 infection. The Lancet. 2020 Jul; 20(7): 773.

Doi: 10.1016/S1473-3099(20)30195-X.

[42] Charles F. Manski, Francesca Molinari. Estimating the COVID-19 infection rate: Anatomy of an

inference problem. Journal of Econometrics. https://doi.org/10.1016/j.jeconom.2020.04.041.

[43] WHO. Q&A: Influenza and COVID-19 - similarities and differences.

https://www.who.int/westernpacific/news/q-a-detail/q-a-similarities-and-differences-covid-19-and-

influenza. [accessed 18 August 2020].

[44] CDC. https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html; [accessed 18 August

2020].

[45] BMJ 2020; 368. Covid-19: identifying and isolating asymptomatic people helped eliminate virus in

Italian village. doi: https://doi.org/10.1136/bmj.m1165.

[46] Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, Ipekci AM, et al. (2020)

Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A

living systematic review and metaanalysis. PLoS Med 17(9): e1003346. https://doi.

org/10.1371/journal.pmed.1003346.

[47] Eric A Meyerowitz et al. Towards an accurate and systematic characterisation of persistently

asymptomatic infection with SARS-CoV-2. Lancet Infect Dis 2020. https://doi.org/10.1016/ S1473-

3099(20)30837-9.

[48] CEBM- The Centre for Evidence-Based Medicine develops, promotes and disseminates better

evidence for healthcare. Estimating the infection fatality ratio in England. https://www.cebm.net/covid-

19/estimating-the-infection-fatality-ratio-in-england/; [accessed 9 October 2020].

[49] Wood, S. N. Did COVID-19 infections decline before UK lockdown? ArXiv:2005.02090 [q-Bio, Stat].

http://arxiv.org/abs/2005.02090.

[50] Selene Ghisolfi, Ingvild Almås, Justin Sandefur, Tillmann von Carnap, Jesse Heitner, Tessa Bold.

Predicted COVID-19 Fatality Rates Based on Age, Sex, Comorbidities, and Health System Capacity.

2020Jun.https://www.cgdev.org/sites/default/files/predicted-covid-19-fatality-rates-based-age-sex-

comorbidities-and-health-system-capacity.pdf.

[51] B. Martınez-Lo´pez et al. A novel spatial and stochastic model to evaluate the within and between

farm transmission of classical swine fever virus: II Validation of the model. / Veterinary Microbiology 155

(2012) 21–32 doi:10.1016/j.vetmic.2011.08.008.

[52] Karsten et al. Monte Carlo simulation of classical swine feverepidemics and control II. Validation of

the model. Veterinary Microbiology 108 (2005) 199–205. doi:10.1016/j.vetmic.2005.04.008.

Tables

Table 1. Structure of different population strata in the Republic of Serbia [23]

Stratum	Population	Percentage of total population
Younger than 7 years	356,377	5.10%
Elementary school	550,527	7.88%
Secondary school	249,455	3.57%
Students	241,698	3.46%
Employed	2,197,065	31.46%
Pensioners	1,715,152	24.56%
Others	1,672,330	23.95%

689 Table 2. Age structure of the population of the Republic of Serbia and expected percentage of

690 hospitalized patients, patients in intensive care, and death rate caused by COVID-19.

Population age groups	*Population	Percentage of total population	**Expected % of hospitalized patients (σ)	***Expected % of patients whose treatment requires intensive care	****Infection fatality rate IFR (male/female)
0-9	458,199	6.56%	0.00%	5.00%	0.04%;0.01%
10-19	445,481	6.38%	0.04%	5.00%	0.00%;0.02%
20-29	1,028,226	14.73%	1.04%	5.00%	0.00%;0.01%
30-39	951,615	13.63%	3.43%	5.00%	0.00%;0.05%
40-49	968,854	13.88%	4.25%	6.30%	0.08%;0.04%
50-59	963,229	13.79%	8.16%	12.20%	0.33%;0.20%
60-69	815,244	11.68%	11.80%	27.40%	1.62%;0.62%
70-79	696,045	9.97%	16.60%	43.20%	6.11%;2.68%
80-	655,711	9.39%	18.40%	70.90%	16.40%;6.49%

691 *[28], **[32], ***[8], ***[33]

693　Table 3. SEAIHRD model parameters

Input parameters	Mark	Value	Source
Population	N_{t0}	6,982,604	[28]
Initial number of cases	I_{t0}	1	Assumed
Initially immune	I_{mm0}	0	Assumed
Basic reproduction number	R_0	2.46 (3.1)	[34]
Effective contact rate	C_e	0.38	Estimated
Per capita contact rate	β	0.0000000378	Estimated
Daily infection rate (transfer E→I)	f	0.294118	Estimated
Recovery rate of symptomatic cases	r	0.107527	Estimated
Daily rate of waning of immunity	ω	0.002739726	Estimated
Per capita birth rate	b	0.000025205	[28]
Per capita death rate unrelated to COVID-19	m	0.000036006	Estimated
Life expectancy in years	L	76.09	[29]
Duration of latent infection in days	f^{-1}	3.5	[30]
Duration of infectious period in days (clinical cases)	T_R	9.3	[30]
Duration of immunity in days	I_{mm}	365.00	Assumed
Incubation period in days	Inc	5.8	[35]
Time period (day)	dt	1.00	-
Average treatment duration in hospital	*days*	15,9	[36]
Average time spent in intensive care	*days*	27	[36]
Recovery rate of hospitalized cases	ε	0.062893	Estimated
Average times taken from onset of symptoms to death	*days*	17	[37]
Infectious period for asymptomatic cases	*days*	7.25	[30, 38]
Recovery rate of asymptomatic cases	γ	0.137931	Estimated
Expected percentage of asymptomatic cases	-	30%	[39]
Infectiousness of asymptomatic cases in relation to symptomatic cases	η	75%	[39]

694   Table 4. Description of different simulated non-pharmaceutical intervention scenarios

Mark	Scenario	Scenario description
SC1	Base-case scenario	The population relies on development of herd immunity. No anti-epidemic measures are implemented.
SC2	Lock down of the entire country	Pre-schools, schools, and colleges are fully closed – reduction in contacts at educational institutions by 75%; reduction in contacts in workplaces by 50%; reduction in contacts of the elderly (older than 65) by 50% at $R_0 = 2.46$ or by 65% at $R_0 = 3.10$; physical distancing of the unemployed population and in public places – reduction in contacts by 45% at $R_0 = 2.46$) or by 55% at $R_0 = 3.10$.
SC3	Partial lockdown of the country - I	Elementary and pre-school educational institutions are open. High schools and colleges are closed. Reduction in contacts by 75% at colleges and high-schools; reduction in contacts in workplaces by 50%; reduction in contacts of the elderly (older than 65) by 60% at $R_0 = 2.46$ or by 65% at $R_0 = 3.10$; social distancing of the unemployed population and in public places – reduction in contacts by 40% at $R_0 = 2.46$ or by 55% at $R_0 = 3.10$.
SC4	Partial lockdown of the country - II	Colleges are closed – reduction in contacts by 75%; reduction in contacts in the workplace by 50%; reduction in contacts of the elderly (older than 65) by 60% at $R_0 = 2.46$ or by 65% at $R_0 = 3.10$; social distancing of the unemployed population and in public places – reduction in contacts by 40% at $R_0 = 2.46$ or by 55% at $R_0 = 3.10$.
SC5	Partial lockdown of the country - III	Reduction in contacts in the workplace by 50%; reduction in contacts of the elderly (older than 65) by 60% at $R_0 = 2.46$ or by 65% at $R_0 = 3.10$; social distancing of the unemployed population and in public places – reduction in contacts by 40% at $R_0 = 2.46$ or by 55% at $R_0 = 3.10$.

697 Table 5. Results of different simulated scenarios (Ro = 2.46 and Ro = 3.1). The data refers to the period of 365 days from epidemic onset.

Scenario mark	SC1		SC2		SC3		SC4		SC5	
Basic reproductive number	$R_o=2.64$	$R_o=3.1$	$R_o=2.64$	$R_o=3.1$	$R_o=2.64$	$R_o=3.1$	$R_o=2.64$	$R_o=3.1$	$R_o=2.64$	$R_o=3.1$
Cumulative incidence (CI)	6,229,144	7,133,221	308,581	2,219,251	1,286,227	2,419,079	1,375,416	2,489,197	1,408,262	2,514,936
Apparent CI	4,360,401	4,993,254	216,007	1,553,476	900,359	1,693,355	962,791	1,742,438	985,783	1,760,456
Overall hospitalized	278,781	320,567	13,970	98,337	56,631	105,930	60,319	108,554	61,685	109,538
Overall in intensive care	85,633	98,333	4,260	29,753	17,085	32,058	18,202	32,855	18,612	33,147
Overall deaths	20,894	23,951	1,031	7,194	4,113	7,754	4,383	7,948	4,483	8,018

699 Table 6. Results of the model sensitivity analysis of individual parameters used in the model

Model parameter mark	Change relative to the basic scenario	CI	Deaths (D_{th})	Change in CI relative to the basic scenario	Change in D_{th} relative to the basic scenario
R_0	5%	6,398,486	21,470	2.72%	2.75%
R_0	10%	6,553,978	21,998	5.21%	5.28%
R_0	25%	6,982,604	23,782	12.10%	13.82%
f^{-1}	5%	6,224,056	19,885	0.08%	4.83%
f^{-1}	10%	6,219,502	18,968	0.15%	9.22%
f^{-1}	25%	6,208,481	16,666	0.33%	20.24%
r	5%	6,068,299	20,351	2.58%	2.60%
r	10%	5,908,223	19,811	5.15%	5.18%
r	25%	5,425,721	18,182	12.90%	12.98%

702 Table 7. Results of the model sensitivity analysis of group of parameters using a perturbation up to 25%

Model parameter mark	Change relative to the basic scenario	CI	Deaths (D_{th})	Change in CI relative to the basic scenario	Change in D_{th} relative to the basic scenario
R_0, f^{-1}, r	5%	6,242,433	19,948	-0.21%	4.53%
R_0, f^{-1}, r	10%	6,257,685	19,093	-0.46%	8.62%
R_0, f^{-1}, r	25%	6,321,339	16,984	-1.48%	18.71%
$R_0, \beta, f^{-1}, r, b, m, \gamma, \varepsilon, \eta, \omega, \bar{\tau}_d$	25%	6,147,725	13,203	1.31%	36.81%

704 Table 8. Measures of the prediction quality

	MAE	%Error	MSE	RMSE	Normalized MAE	Normalized MSE	Max Deviation
Deceased	73	2.04%	7,212.20	84.92	2.05%	0.06%	4.82%

706 Table 9. Regression statistics

Multiple R	R Square	Adjusted R Square	Standard Error	Observations
0.981678289	0.963692263	0.963124954	86.24143528	66

721 **Figures**

a) Distribution of the total number of the susceptible, latently infected, asymptomatic infectious individuals, infected and recovered on a daily basis since the epidemic onset ($R_o=2.46$)

b) Distribution of the total number of the susceptible, latently infected, asymptomatic infectious individuals, infected and recovered on a daily basis since the epidemic onset ($R_o=3.1$)

c) Daily fluctuations of the susceptible, recovered and net disease transmission rates ($R_o=2.46$)

d) Daily fluctuations of the susceptible, recovered and net disease transmission rates ($R_o=3.1$)

e) Daily fluctuations of apparent incidence, true incidence and net disease transmission rates ($R_o=2.46$)

f) Daily fluctuations of apparent incidence, true incidence and net disease transmission rates ($R_o=2.46$)

722 Fig.1 Model prediction of latently infected, asymptomatic infectious individuals, infected, recovered and
723 daily fluctuations of R_n.

a) Necessary hospital capacities (R_o=2.46)

b) Necessary hospital capacities (R_o=3.1)

c) Age structure of hospitalized patients (R_o=2.46)

d) Age structure of hospitalized patients (R_o=3.1)

e) Expected age structure of the dead (R_o=2.46)

f) Expected age structure of the dead (Ro=3.1)

725   Fig.2 Model prediction of required hospital capacities under the assumption of different intervention

726   measures.

a) Comparative overview of cumulative incidences (CI). Results obtained from simulated scenarios 1-5 (R_o=2.46)

b) Comparative overview of cumulative incidences (CI). Results obtained from simulated scenarios 1-5 (R_o=3.1)

c) Comparative overview of hospitalized patients on a daily basis. Results obtained from simulated scenarios 1-5 (R_o=2.46)

d) Comparative overview of hospitalized patients on a daily basis. Results obtained from simulated scenarios 1-5 (R_o=3.1)

e) Comparative overview of expected number of patients in intensive care on a daily basis. Results obtained from simulated scenarios 1-5 with R_o=2.46

f) Comparative overview of expected number of patients in intensive care on a daily basis. Results obtained from simulated scenarios 1-5 with R_o=3.1

729   Fig.3 Model prediction of expected number of hospitalized patient and patient in intensive care.

a) Comparative overview of hospital bed occupancy (R_o=2.46)　　b Comparative overview of hospital bed occupancy (R_o=3.1)

c) Comparative overview of ICU bed occupancy (Ro=2.46)　　d) Comparative overview of ICU bed occupancy (Ro=3.1)

731　Fig. 4 Model prediction of required hospital capacities needed to treat patients with COVID-19.

a) Comparative overview of daily incidences (R_o=2.46).

b) Comparative overview of daily incidences (R_o=3.1).

c) Comparative overview of hospitalized patients (R_o=2.46).

d) c) Comparative overview of hospitalized patients (R_o=3.1).

e) Comparative overview of deaths (Ro=2.46)

f) Comparative overview of deaths (Ro=3.1)

734   *NV-without vaccination and any non-pharmaceutical intervention*
735
736   Fig. 5 Results of simulated COVID-19 control based solely on vaccination, scenarios 6-9 (R_o=2.46 and
737   R_o=3.1).

739   Fig. 6. The observed number of deceased individuals (blue), number of deaths modeled with SEAIHRD

740   model (yellow) and predicted number of deaths (green).

741   *Figures 1 to 6 should be printed in color.*

$$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) - \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) + \omega R_i(t) - mS_i(t)$$

$$\frac{dE_i(t)}{dt} = \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) + \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) - (m + f)E_i(t)$$

$$\frac{dA_i(t)}{dt} = f\alpha E_i(t) - (\gamma + m)A_i(t)$$

$$\frac{dI_i(t)}{dt} = f(1 - \alpha)E_i(t) - \sum_{j=1}^{n} \sigma_{ij} I_j(t) - (r + m)I_i(t)$$

$$\frac{dH_i(t)}{dt} = \sum_{j=1}^{n} \sigma_{ij} I_j(t) - \sum_{j=1}^{n} \delta_{ij} H_i(t) - (\varepsilon + m)H_i(t)$$

$$\frac{dR_i(t)}{dt} = \sum_{j=1}^{n} r_{ij} I_j(t) + \sum_{j=1}^{n} \gamma_{ij} A_j(t) + \varepsilon H_i(t) - (m + \omega)R_i(t)$$

$$\frac{dD_i(t)}{dt} = \sum_{j=1}^{n} \delta_{ij} I_i(t)$$

Fig. 1. SEAIHRDS model with demography

$$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) - \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) + \omega R_i(t) - mS_i(t)$$

$$\frac{dE_i(t)}{dt} = \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) + \sum_{j=1}^{n} \beta_{ij} \eta A_j(t) S_i(t) - (m + f)E_i(t)$$

$$\frac{dA_i(t)}{dt} = f\alpha E_i(t) - (\gamma + m)A_i(t)$$

$$\frac{dI_i(t)}{dt} = f(1 - \alpha)E_i(t) - \sum_{j=1}^{n} \sigma_{ij} I_j(t) - (r + m)I_i(t)$$

$$\frac{dH_i(t)}{dt} = \sum_{j=1}^{n} \sigma_{ij} I_j(t) - \sum_{j=1}^{n} \delta_{ij} H_i(t) - (\varepsilon + m)H_i(t)$$

$$\frac{dR_i(t)}{dt} = \sum_{j=1}^{n} r_{ij} I_j(t) + \sum_{j=1}^{n} \gamma_{ij} A_j(t) + \varepsilon H_i(t) - (m + \omega)R_i(t)$$

$$\frac{dD_i(t)}{dt} = \sum_{j=1}^{n} \delta_{ij} I_i(t)$$

$$\frac{dV_i(t)}{dt} = v_e \psi S_i(t) - mV_i - \omega V_i.$$

Fig. 2. SEAIHRDVS model with demography and vaccination

a) Prediction of the total number of the susceptible, latently infected, asymptomatic infectious individuals, infected and recovered on a daily basis since the epidemic onset. No intervention measures applied (R_o=2.46)

b) Prediction of the total number of the susceptible, latently infected, asymptomatic infectious individuals, infected and recovered on a daily basis since the epidemic onset. No intervention measures applied (R_o=3.1)

c) The results obtained during the simulation of the application of various contact reduction measures (scenarios 1-5, R_o=2.46)

d) The results obtained during the simulation of the application of various contact reduction measures (scenarios 1-5, R_o=3.1)

e) Results of simulated COVID19 control based solely on vaccination, scenarios 6-9 (R_o=2.46).

f) Results of simulated COVID19 control based solely on vaccination, scenarios 6-9 (R_o=3.1).