Simulation and prediction of further spread of COVID-19 in The Republic of Serbia by SEIRDS model of disease transmission

*Slavoljub STANOJEVIC[a] , Mirza PONJAVIC[b] , Slobodan STANOJEVIC[c] , Aleksandar STEVANOVIC[d], Sonja RADOJICIC[e]

[a]Directorate of National Reference Laboratories, Batajnicki drum 10, 11080 Zemun, Serbia

[b]International Burch University, Francuske revolucije bb, Ilidza 71210, Sarajevo, Bosnia and Herzegovina, e-mail: mirza.ponjavic@gis.ba

[c]Veterinary Scientific Institute of Serbia, Janisa Janulisa 14, 11107 Belgrade, Serbia, e-mail: slobodan.vet@gmail.com

[d]University of Pittsburgh, Department of Civil and Environmental Engineering, 3700 O'Hara Street, Pittsburgh, PA 15261, USA, e-mail: stevanovic@pitt.edu

[e]Belgrade University, Faculty of Veterinary Medicine – Department of Infectious Diseases and Epidemiology, Bulevar Oslobodenja 18, 11000 Belgrade, Serbia, e-mail: sonjar@vet.bg.ac.rs

*Corresponding author: Slavoljub Stanojevic, e mail: slavoljub.stanojevic@minpolj.gov.rs

Abstract

As a response to the pandemic caused by SARSCov-2 virus, on 15 March, 2020, the Republic of Serbia introduced comprehensive anti-epidemic measures to curb COVID 19. After a slowdown in the epidemic, on 6 May, 2020, the regulatory authorities decided to relax the implemented measures. However, the epidemiological situation soon worsened again. As of 15 October, 2020, a total of 35,454 cases of SARSCov-2 infection have been reported in Serbia, including 770 deaths caused by COVID19. In order to better understand the epidemic dynamics and predict possible outcomes, we have developed a mathematical model SEIRDS (S-susceptible, E-exposed, I-infected, R-recovered, D-dead due to COVID19 infection, S-susceptible). When developing the model, we took into account the differences between different population strata, which can impact the disease dynamics and outcome. The model can be used to simulate various scenarios of the implemented intervention measures and calculate possible epidemic

outcomes, including the necessary hospital capacities. Considering promising results regarding the

development of a vaccine against COVID19, the model is enabled to simulate vaccination among different

population strata. The findings from various simulation scenarios have shown that, with implementation

of strict measures of contact reduction, it is possible to control COVID19 and reduce number of deaths.

The findings also show that limiting effective contacts within the most susceptible population strata merits

a special attention. However, the findings also show that the disease has a potential to remain in the

population for a long time, likely with a seasonal pattern. If a vaccine, with efficacy equal or higher than

65%, becomes available it could help to significantly slow down or completely stop circulation of the virus in

human population. The effects of vaccination depend primarily on: 1. Efficacy of available vaccine(s), 2.

Prioritization of the population categories for vaccination, and 3. Overall vaccination coverage of the

population, assuming that the vaccine(s) develop solid immunity in vaccinated individuals. With expected

basic reproduction number of R_o=2.46 and vaccine efficacy of 68%, an 87%–coverage would be sufficient

to stop the virus circulation.

Keywords: COVID19, SEIRDS mathematical model differential equation, prediction, vaccination

44   **1.   Introduction**

On 11 March, 2020, the World Health Organization characterised the disease caused by the novel SARS

Cov-2 virus as a pandemic [1]. Initial epidemic outbreak in China spread outside the Wuhan area, and

subsequently on a global scale. On 6 March, 2020, the first case of the novel corona virus infection was

reported in the Republic of Serbia. Taking into consideration the escalation of the disease and limited effects

of the initially implemented measures, the state of emergency was declared throughout the country on 15

March, 2020. Comprehensive anti-epidemic measures (e.g. lockdown of entire country) were introduced in

the entire country [2]. Due to the absence of specific pharmaceutical intervention, Serbia, like other

countries, implemented an anti-epidemic strategy based on social distancing, school and university

closures, reduced number of workers present in the workplaces, closure of places of worship for public

religious services, reduced working hours of cafés and restaurants, avoiding mass gatherings, events,

sports games, tracing and identification of infected people's contacts, etc. After a slowdown in the epidemic,

as shown in the relevant officially published data, the regulatory authorities decided to relax the introduced

measures on 6 May, 2020. However, the epidemiological situation soon worsened again, resulting in the

reinstatement of some measures, as well as the introduction of new measures [2].   Although the return of

extensive measures has yielded favourable results, the further development of the epidemic is not clear.

For these reasons, mathematical modelling has a crucial role in understanding the epidemic and

predicting possible outcomes. Modelling is a particularly useful tool for devising strategies for combating

the epidemic, capacity planning, and selection of efficient measures, especially in the absence of specific

pharmaceutical treatments [3, 4, 5]. Mathematical modelling based on differential equations dates back to

the first half of 20th century. In 1927, Kermack and McKendrick developed the basic model of disease

transmission consisting of three compartments: susceptible (S), infected (I) and removed (R). The model

was based on a connected system of nonlinear differential equations as a special case of the general

epidemiological model [6, 7]. Subsequent models, became more complex and adapted to the needs of

research [5].

Since the outbreak of COVID19, many published papers have dealt with the implementation of

mathematical modelling and prediction of possible outcomes of COVID19 epidemics. Most of these

research efforts have been based on the implementation of the SIR (susceptible-infected-removed) model.

Many of the other models provide a clear picture of dynamics of COVID19 spreading, including the

overloading of the relevant health systems. For example, Ferguson et al., developed one of the first models

for COVID19 simulation, which was, among other things, used to plan the health care resources [8]. Wu et

al., developed the SEIR model to examine the dynamics of SARS-Cov-2 transmission from person to

person. This model was also used to calculate the basic reproduction number R_o, which we use in this

paper as one of the key parameters [9]. The classical SIR model assumes that there is homogenous mixing

of infected and susceptible populations and that the total population is constant and does not change over

time. Moreover, according to the classical SIR model, there is a monotonous decline in susceptible

population towards zero [10]. However, such assumptions are not objective in the case of COVID-19

spreading and they are the basic problem in the modelling of this pandemic. In reality, human population

fluctuates constantly [11]. In order to account for such fluctuation, and better understand the COVID 19

epidemic in the Republic of Serbia, we have employed mathematical modelling of the epidemic using the

available data on the characteristics of the disease, such as incubation period, latent period, recovery

period, severity of clinical signs, and mortality rate caused by COVID19. Unlike the classic SIR model,

SEIRDS (S-susceptible, E-exposed, I-infected, R-recovered, D-dead due to COVID19 infection, S-

susceptible) epidemic model, developed for this research, simulates the spreading of COVID 19 in an open

population. Taking into account that the population is constantly changing and that various measures are

applied for different strata or subgroups of the population (such as preschool children, children attending

primary school, high school students, students, employees, the unemployed and retirees), as well as

changes in the intensity of applied measures, we have proposed the use of a model that takes these

circumstances into account. Based on input disease parameters taken from scientific literature and specific

data related to Serbia, this model simulates daily disease occurrence, including the number of hospitalised

patients and cases which require intensive care. The model also predicts the expected number of deaths,

as well as hospital capacities necessary to accommodate the patients. It provides a possibility to simulate

different scenarios of disease control and intervention measures. Considering the expectations of

successful development of the vaccine against COVID 19 in the near future, we added an option to model

vaccination of different strata of the population as a set of disease control strategies.

2. Methodology

This section presents the research methodology and the proposed model, which was used to predict the

further dynamics of the epidemic in Serbia. We also presented the data that were used to model the

epidemic, a simulated strategy to combat COVID 19, and a sensitivity analysis.

2.1. SEIRDS mathematical model

Classical SEIRDS model divides the population into compartments, i.e. groups, and follows the disease

dynamics at all times. The population is divided into the following compartments: the portion of the

population susceptible to the infection is denoted by S, those latently infected with SARSCov-2 (exposed

to) are denoted by E, the infected individuals who are able to spread the disease are denoted by I, the ones

recovered from the infection are denoted by R, and those who died due to disease with D. Assuming that

individuals mix homogenously, the force of infection λ (the rate at which susceptible persons are infected)

is related to *per capita* contact rate β. Also, the risk of infection is closely related to the number of infectious

individuals in the population I_t. It depends on the number of infectious individuals (I_t) and how frequently

they make contacts with other persons. In a situation of homogenous mixing among the population, the

force of infection λ can be express ae follows:

$\lambda(t) = \beta I(t)$ (1)

The change of rates in every compartment per unit time in SEIRDS model is presented in the following

series of differential equations:

$S_{t+1} = bN_t - \beta S_t I_t - mS_t + \omega R_t$ (2)

$E_{t+1} = \beta S_t I_t - fE - mE_t$ (3)

$I_{t+1} = fE_t - (r+m)I_t$ (4)

$R_{t+1} = rI_t - (m+\omega)R_t$ (5)

$D_{t+1} = \delta I_t$ (6)

where f is rate of onset of infections expressed as the reciprocal of the latent infection period, r is the rate

at which infectious individuals are recovered, δ is the rate at which infectious individuals die from COVID

19 infection and ω is rate of waning of immunity. The total population at any particular interval of time t is:

$N_t = S_t + E_t + I_t + R_t + D_t + bN_{t-1} - mS_{t-1}$ (7)

where parameters b and m are *per capita* daily birth and death rates unrelated to COVID19.

However, considering that implemented anti-epidemic measures against COVID 19 do not have an identical

impact on population's age subgroups and that COVID19 pathogenesis varies in different age subgroups,

we propose the use of multi-compartment version of standard SEIRDS model. In this model susceptible

population was further stratified within the compartment S according to age and occupations. Grouping into

various strata was done according to the real age structure of the Republic of Serbia population as follows:

pre-school children (S_{ps}), elementary school children (S_{es}), high school children (S_{hs}), college students (S_{cs}),

unemployed population (S_{ua}), employed population (S_{ea}), and elderly/retired (S_r) (Table 1). To simulate the

epidemic progression through different population strata-subgroups, we used appropriate, stratum-specific,

model parameters and factors of effective contact reduction (anti-epidemic intervention measures - ρ),

which were adapted to the relevant population groups: lockdown of the entire country, closures of pre-

school, school and college sessions closures, reduced number of workers allowed in the workplaces, work

from home, restrictions of mobility of elderly people, etc. During the simulation, we monitored the effects of

various levels of contact reduction, ranging from 25% to 75%, taking into account the realistic possibilities

of maintaining a minimum work process, functioning of the society and feasibility of such measures.

Given that intervention measures, applied in response to the emergence of COVID19, are not the

same for all population strata, homogeneous mixing can be expected only within same population stratum.

The rate of effective contacts β, after the application of intervention measures, is no longer identical at the

level of all strata of the population. Effective contacts are limited by the intensity and types of measures

applied and are identical only when it comes to individuals within the same population strata. Furthermore,

persons in different population strata become infected at different rates depending on how frequently they

interact with other persons in their own subgroup and other subgroups. If we assume that force of infection

differs between different strata of population, the equation for the force of infection would be as follows:

$\lambda_i(t) = \sum_{j=1}^{n} \beta_{ij} I_j(t)$ (8)

where $\lambda_i(t)$ is force of infection in the i^{th} population strata, β_{ij} is the rate at which susceptible persons in the

i^{th} population strata and infectious persons in j^{th} population strata come into effective contact per unite of

time, *and* $I_i(t)$ is the number of infectious persons in j^{th} population strata. Also, in this model the number of

recovered and dead is conditioned with different ages and genders.

Now our model will be expressed as follows:

$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) - mS_i(t)$ (9)

$\frac{dE_i(t)}{dt} = \sum_{j=1}^{n} \beta_{ij} I_j(t) S_i(t) - (m + f) E_i(t)$ (10)

$\frac{dI_i(t)}{dt} = f E_i(t) - (r + m) I_i(t) - \sum_{j=1}^{n} \delta_{ij} I_j(t)$ (11)

$\frac{dR_i(t)}{dt} = \sum_{j=1}^{n} r_{ij} I_j(t) - (m + \omega) R_i(t)$ (12)

$\frac{dD_i(t)}{dt} = \sum_{j=1}^{n} \delta_{ij} I_i(t)$ (13)

In this model susceptible, exposed, infectious, recovered, deaths and total population are:

$S(t)=S_{ps}(t)+S_{es}(t)+S_{hs}(t)+S_{cs}(t)+S_{ua}(t)+S_{ea}(t)+S_p(t)$ (14)

$E(t)=E_{ps}(t)+E_{es}(t)+E_{hs}(t)+E_{cs}(t)+E_{ua}(t)+E_{ea}(t)+E_p(t)$ (15)

$I(t)=I_{ps}(t)+I_{es}(t)+I_{hs}(t)+I_{cs}(t)+I_{ua}(t)+I_{ea}(t)+I_p(t)$ (16)

$R(t)=R_{ps}(t)+R_{es}(t)+R_{hs}(t)+R_{cs}(t)+R_{ua}(t)+R_{ea}(t)+R_p(t)$ (17)

$D(t)=D_{ps}(t)+D_{es}(t)+D_{hs}(t)+D_{cs}(t)+D_{ua}(t)+D_{ea}(t)+D_p(t)$ (18)

N(t)=S(t)+E(t)+I(t)+R(t)-D(t)   (19)

2.2.   Determining the herd immunity threshold and control of CIVID 19 by vaccination policy

Considering the undergoing worldwide efforts to develop a vaccine against COVID 19 and promising

results, we extended the model to simulate and analysed the effects of a hypothetical vaccination on the

epidemic dynamics, and to estimate the extent of coverage of vaccination which can interrupt the chain of

infection. The control of COVID 19 by vaccination means targeting the entire susceptible population with

mass vaccination until critical herd immunity achieved. In such situation there is a "race" between the

exponential growth of epidemic and mass vaccination. The level of herd immunity threshold (HIT) can be

calculated by the following formula:

HIT = 1− 1/R_o = (R_o−1)/R_o   (20)

and the critical vaccination coverage required to achieved herd immunity can be obtained by multiplying

herd immunity threshold with reciprocal value of vaccine efficacy, v_e:

$V_c = \frac{1}{v_e}(1 - \frac{1}{R_o})$ (21)

Most people infected with SARS-CoV-2 develop an immune response followed by the development

of specific antibodies between 10 and 21 days after getting infected [12]. Specific IgM and IgG antibodies

against SARS-CoV-2 develop 6 to 15 days after the onset of the disease [13-17]. According to some

studies, the presence of antibodies has been confirmed in less than 40% of the patients within 1 week after

the onset of the disease, whereas percentage reaches 100% of subjects 15 days after the onset of disease

[18]. Although duration of the immune response against CVOVID 19 is still unknown, comparing with other

coronaviruses, immunity wane within 12 to 52 weeks after the first symptoms appear [19], while in the case

of SARS-CoV-1 infection, the presence of IgG antibodies was confirmed in 90% and 50% of infected

patients, respectively, over two and three years, respectively [29]. Based on these findings, we assumed

that in the event of the development of a successful vaccine, immunity against the SARRSCov-2 virus could

last for a year, as well as after recovery after a natural infection.

For the purpose of modelling, additional compartment to the model was added denoted with V(t),

in which there are vaccinated persons who have successfully developed protective immunity after

vaccination. The vaccination parameter, *u*, is the daily rate of vaccination of susceptible population and it

represents the proportion of susceptible population immunized per unite time. The critical daily rate of

vaccination, u_c, is $\psi_c = (b+\omega)(R_o -1)$, required to interrupt the infection [5]. The basic reproduction number

under the vaccination is $R_{op} = (1-p)R_o$. The proportion of effectively protected persons, p, is conditioned by

parameters the vaccine efficacy, v_e. This parameter represents a proportion of person who successfully

developed protective immunity after vaccination, whereas total number of actively protected individuals in

time t is V(t) = number of vaccinated x v_e [5]. In this compartment the daily rate of waning of immunity at

which immunity of vaccinated population fades out is ω, and it is reciprocal to the period of lasting of

immunity. Vaccinated persons, after losing their immunity, become sensitive again and removed to the

compartment S. The change of rate in this compartment per unit time is as follow:

$\frac{dV_i(t)}{dt} = v_e\psi S_i(t) - mV_i - \omega V_i.$ (22)

The compartment S(t) is slightly modified as follow:

$\frac{dS_i(t)}{dt} = bN(t) - \sum_{j=1}^{n} \beta_{ij}I_j(t) S_i(t) - mS_i(t) - v_e\psi S_i(t).$ (23)

The other of compartments of SERIDS model remain unchanged.

2.3.   **Model parametrisation**

In proposed model, β is *per capita* effective contact rate at which specific persons come into

effective contact per unit of time. An effective contact is defined as a contact sufficient to cause disease

transmission [**10, 20, 21**]. We calculated the parameter β by using the formula:

β = R_o/NT_R (24)

where R_o is a basic reproduction number of the disease, i.e. the average number of newly infected people

with COVID 19 (secondary infection cases), infected by one infectious individual in a totally susceptible

population, N is total population, and T_R is the average duration of infectious period [10, 20, 21]. The R

values of 2.46 and 3.1 are adopted from the relevant literature. The R_o values were based on the data

obtained during the initial phase of the epidemic in Italy [22]. Since the implemented measures and disease

transmission were simulated through various population strata, we corrected the β parameter with a

relevant, stratum-specific contact reduction factor ρ_i. In this way, we obtained the *per capita* contact rate

specific for each separate stratum based on following formula. $\beta_i = \beta(1-\rho_i)$. The values of the ρ_i factor in

different population strata ranged from 0.25 to 0.75 (effective contact reduction ranged from 25% to 75%).

Parameters such as daily birth and death rates were calculated based on the data published by the

Office of Statistics of the Republic of Serbia, and data published by the World Bank regarding the life

expectancy in the Republic of Serbia [23, 24]. The latter study reported that the life expectancy in Serbia

was 79.06 years in 2017 [23]. By using this figure, we expressed the daily death rate as a value reciprocal

to life expectancy m = 0.000036. We calculated the daily *per capita* birth rate of *b= 0.000025* based on the

figure of 9.2 births in the Republic of Serbia per 1000 people in a year. These estimates were needed to

realistically simulate fluctuations of the total population. To simplify the calculations, we assumed that the

general morality rate m is applicable for all population strata.

The infectivity rate, i.e. the rate of transfer from compartment E to I, was derived as a value

reciprocal to the COVID 19's average latent period. The data on the average duration of latent infection (f^{-1}

) and the average period during which an infected person is shading the SARSCov-2 virus (T_R) were

adopted from the relevant literature as f^{-1} = 4.6 days and T_R = 6.5 days, respectively [8]. The same study

is used as a source of the data: on the percentage of hospitalised patients and those whose therapy requires

intensive care, used for prediction of required hospital capacities, as shown in Table 1 [8].

Parameters such as δ and r are related to the infectious fatality rate (IFR) for COVID 19 and

average times taken from infection to death (T_D) or recovery (T_R). These parameters were calculated using

the following formulas:

$\delta = IFR_{stratum}/T_D$ and   (25)

$r = (1-IFR_{stratum})/T_R$ (26)

The IFRs, shown in Table 3, were taken from literature and compared with local IFR value which was

calculated based on officially registered deaths published by the health system of the Republic of Serbia

[2]. The Calculation of local IFR is presented in section 2.4. Population data, (e.g. total population, age

structure, and stratification) are presented in Tables 1 and 2. A summary of all model parameters is given

in Table 3.

251   Table 1. Structure of different population strata in the Republic of Serbia [23]

Stratum	Population	Percentage of total population
Younger than 7 years	356,377	5.10%
Elementary school	550,527	7.88%
Secondary school	249,455	3.57%
Students	241,698	3.46%
Employed	2,197,065	31.46%
Pensioners	1,715,152	24.56%
Others	1,672,330	23.95%

254   Table 2. Age structure of the population of the Republic of Serbia and expected percentage of hospitalised

255   patients, patients in intensive care, and death rate caused by COVID19.

Population age groups	*Population	Percentage of total population	**Excepted % of hospitalised patients	**Expected % of patients whose treatment requires intensive care	***Infection fatality rate IFR (male/female)
0-9	458,199	6.56%	0.10%	5.00%	0.04%;0.01%
10-19	445,481	6.38%	0.30%	5.00%	0.00%;0.02%
20-29	1,028,226	14.73%	1.20%	5.00%	0.00%;0.01%
30-39	951,615	13.63%	3.20%	5.00%	0.00%;0.05%
40-49	968,854	13.88%	4.90%	6.30%	0.08%;0.04%
50-59	963,229	13.79%	10.20%	12.20%	0.33%;0.20%
60-69	815,244	11.68%	16.60%	27.40%	1.62%;0.62%
70-79	696,045	9.97%	24.30%	43.20%	6.11%;2.68%
80-	655,711	9.39%	27.30%	70.90%	16.40%;6.49%

256   *[23], **[8], ***[25]

259 Table 3. SEIRD model parameters

Input parameters	Mark	Value	Source
Population	N_{t0}	6,982,604	[23]
Initial number of cases	I_{t0}	1	
Initially immune	I_{mm0}	0	
Basic reproduction number	R_0	2.46 (3.1)	[22]
Effective contact rate	C_e	0.38	
Per capita contact rate	β	0.000000054	
Daily infection rate (transfer E→I)	f	0.217391304	
Recovery rate	r	0.153846154	
Daily rate of waning of immunity	ω	0.002739726	
Per capita birth rate	b	0.000025205	[23]
Per capita death rate unrelated to COVID19	m	0.000036006	
Waning immunity rate	v	0.002777778	
Life expectancy in years	L	76.09	[24]
Duration of latent infection in days	f^{-1}	4.60	[8]
Duration of infectious period in days (clinical cases)	T_R	6.50	[8]
Duration of immunity in days	I_{mm}	365.00	
Incubation period in days	Inc	5.10	[8]
Time period (day)	dt	1.00	
Average treatment duration in hospital	*days*	10,4	[8]
Average time spent in intensive care	*days*	16	[8]

2.4.   Setting disease control scenarios

Five different scenarios were developed for simulating the COVID19 epidemic control based on non-

pharmaceutical interventions. SC1 implies a base-case scenario where the epidemic spreads in susceptible

population without any anti-epidemic measures being implemented. In the other scenarios, the extent of

contacts was reduced, for each population stratum separately, according to objective possibilities and

measures which were implemented during the actual epidemic in the Republic of Serbia. Scenarios are

presented in table 4.

Table 4. Description of different simulated non-pharmaceutical intervention scenarios

Mark	Scenario	Scenario description
SC1	Base-case scenario	The population relies on development of herd immunity. No anti-epidemic measures are implemented.
SC2	Lock down of the entire country	Pre-schools, schools, and colleges are fully closed – reduction in contacts at educational institutions by 75%; reduction in contacts in workplaces by 50%; reduction in contacts of the elderly (older than 65) by 25%; social distancing of the unemployed population and in public places – reduction in contacts by 25%.
SC3	Partial lockdown of the country - I	Elementary and pre-school educational institutions are open. High schools and colleges are closed. Reduction in contacts by 75% at colleges and high-schools; reduction in contacts in workplaces by 50%; reduction in contacts of the elderly (older than 65) by 25%; social distancing of the unemployed population and in public places – reduction in contacts by 25%.
SC4	Partial lockdown of the country - II	Colleges are closed – reduction in contacts by 75%; reduction in contacts in the workplace by 50%; reduction in contacts of the elderly (older than 65) by 25%; social distancing of the unemployed population and in public places – reduction in contacts by 25%.
SC5	Partial lockdown of the country - III	Reduction in contacts in the workplace by 50%; reduction in contacts of the elderly (older than 65) by 25%; social distancing of the unemployed population and in public places – reduction in contacts by 25 %

Additional four scenarios of control of COVID 19 were simulated based on vaccination policy. We assumed

that vaccine efficacy was 50%, 68%, 80%, and 85%. The initial conditions assumed that all other anti-

epidemic measures are excluded from the model and replaced with mass vaccination. Indicators of

epidemic dynamics were monitored, such as: CI, hospitalized patients, patient in intensive care units and

deaths.

2.5.   Model sensitivity analysis and calibration

Considering the world experience with detection of COVID19 cases, as well as the unreliable data on

COVID19 infections which are currently available worldwide, model calibration is very challenging, and can

result in obtaining inaccurate values for the parameters [26]. This is especially due to the facts that a

significant percentage of the infected individuals do not exhibit any symptoms. The other issue is small

percentage of tested population [26].

As part of the national seroepidemiological study, 1,006 subjects were tested in Serbia from May

11[th] to June 25[th], 2020, for the purpose of estimating the extent of COVID-19 infection. According to the

published data, seroconversion was confirmed in 6,4% of the subjects. On the other hand, a total of 13,372

cases of the infection were reported, which means that those who were infected constitute around 0.19%

of the overall population. However, it is our opinion that the data on reported deaths caused by COVID19

infection is more reliable for use in model calibration, e.g. infection fatality rate. Alex et al. reached a similar

conclusion when simulating COVID19 by using the SEIRD model with heterogeneous diffusion [26]. When

we compared the data recorded during the beginning of the epidemic in Serbia with the results obtained

during the simulation, such as the initial doubling time, the two data series matched well. However, later,

the obtained results did not match well the officially registered data on the number of infected, especially

after the beginning of the implementation of measures in Serbia. We attribute these differences to the

methodology by which official authorities register cases of infection, and collect the data. However, on

October 15[th], 2020 official authorities have published new epidemiological estimates of COVID19

population infection, which is 20% of the total population, which approximately coincides with the results

obtained from our simulation, (e.g. 1,396,521 infected cases obtained from simulation compared to

1,120,460 cases estimated by the Serbian official authorities).

The parameter that determines the number of deaths is the IFR. It is the number of persons who

die of the COVID 19 among all infected individuals regardless of whether the infected show symptoms of

the disease or not. As with many diseases, IFR is not always equivalent to the number of reported deaths

caused by COVID 19. This is because a significant number of deaths, although caused by COVID 19, will

not be recognized as deaths caused by COVID 19 [27]. Also, there are many asymptomatic cases of

infection which are never detected [28, 29, 30].

Due the fact that there is a lag in time between when people are infected and when they die,

patients who die on any given day were infected much earlier, and thus the denominator of the mortality

rate should be the total number of patients infected at the same time as those who died [27]. David et al.

estimated mortality rate by dividing of deaths on a given date by the total number of persons confirmed as

COVID 19 cases 14 days before [27]. It is based on the assumption that maximum incubation period is14

days [34]. However, if we take into account that the number of registered cases of COVID 19 infection is

usually significantly lower than the actual number, assuming that the data on deaths are accurate, the real

IFR value is significantly lower than the calculated value [26]. If we apply this to the situation in Serbia, the

daily value of IFR on July 10th, 2020, when the largest number of deaths was registered in one day, was

9.33%, considering that 18 deaths were registered on July 10th and 14 days earlier 193 confirmed cases of

COVID 19 infection. The raw values of IFR for the period between March 6th and August 10th were as

follows: median of IFR = 2.11%, and average value of IFR = 7.15% bounded in interval 4.17%-10.13%.

When we compared these values with those published by the WHO, CDC and other authors [29, 30, 31]

we concluded that they differ significantly. Considering these findings, the IFR values adopted in the model

(for various population groups and genders) were primarily taken from the literature, with a remark that the

selection of IFR values was based on preliminary comparison of the overall Serbian IFR with similar IFRs

found in the literature, taking into account the registered deaths and most probable number of infected

individuals [8]. To make this possible, the first step was to correct the local raw IFR value mentioned above.

Based on real data we first calculated the population at risk of dying from COVID 19 infection for each

individual day since the outbreak, ending on August 10th, 2020. The number at risk on a given day should

correspond to the number of deaths from COVID 19 infection, considering the lag period from infection to

death. For this calculation, we used the data on the number of deaths D_t in Serbia registered on daily bases

[2]. We hypothesized that the distribution of time periods t_n from the moment of COVID 19 infection to death

follows the lognormal distribution defined by the parameters m = 26.8 and σ = 12.4 days [32]. Taking into

account the published data on the expected percentages of asymptomatic cases in the population, we

assumed that the registered number of cases of infection is 20% of the total number of actually infected

[29, 33, 34]. Based on the formula: $I_{r(t)} = \sum_{j=1}^{n} I_{(t-n)} * m_{(t-n)}$ by reverse, we calculated the population at risk

of dying from COVID 19 infection for each individual day, where m_t is the probability that the time between

infection and death is t days and follows the lognormal distribution ($m = 26.8$, $\sigma = 12.4$) [35]. After that, the

daily IFR values were calculated according to the formula $IFR_{(t)} = D_t / I_{r(t)}$ [35]. Based on IFR values calculated

in this way, we made descriptive statistics and obtained the mean value of IFR = 0.70%, bounded in the

interval 0.46-0.94% and a median of 0.19%. It is important to note here that this value corresponds with the

COVID 19 IFR values found in Eastern European countries and Spain [35, 36]. Taking these findings into

account, we decided to take the IFR values specific to certain population strata recorded in Spain as the

most appropriate for our case [36]. The adopted IFR values are listed in Table 3.

The model sensitivity analysis was conducted by changing the most sensitive model parameters:

β, f, r. The values of these parameters were increased by 5%, 10% and 25% relative to the base scenario

and changes in output indicators were observed.

3. Results

3.1.   Predicting the number of sick, hospitalized patients and deaths caused by COVID 19 in

the absence of any intervention measures

After the simulation, the model demonstrated that with $R_o = 2.46$, and without the implementation

of any anti-epidemic measures, the initial doubling time of the infection was five days. The epidemic wave

would peak 158 days after the outbreak, and it would yield 157,424 infected individuals in a day. Afterwards,

the infection rate would decline for 180 days, eventually reaching the daily incidence of 225 newly infected,

after which the next epidemic wave would start. The second wave would peak 569 days after the onset of

epidemic and yielding 36,922 infected individuals in a day. The third epidemic wave would peak 914 days

later, with 18,791 infected individuals in a day. The true cumulative incidence in the first year of the epidemic

would be 6,633,834 infected people with SARSCov-2 virus, while the apparent cumulative incidence would

be 1,326,767 infected. A total of 5,525 patients would die due to COVID19 consequences. With $R_o = 3.1$,

the following results were obtained: the initial doubling time of the infection was five days, true cumulative

incidence 7,035,433, apparent cumulative incidence 1,407,089, and the total deaths of 5,868. Fig. 1, panels

a) and b) show daily variations in the number of susceptible, latently infected, infected and recovered

patients, at basic reproduction numbers of R_o=2.46 and R_o=3, respectively. Panels c) and d) of the same

figure show daily fluctuations in susceptible, recovered and net reproduction rates R_n for R_o=2.46 and R_o=3,

respectively. Panels e) and f) of Fig. 1 show daily variations in R_n, true and apparent disease incidences at

basic reproduction numbers of R_o=2.46 and R_o=3. The shaded area corresponds to the period when the

daily number of new COVID19 infected individuals increasing, and therefore all of the following hold: R_n>1,

proportion susceptible >1/R_o and the proportion of population that is recovered (immune) is below the herd

immunity threshold. Fig. 2 panels a) and b) show a prediction of necessary hospital capacities. Panels c)

and d) of Fig. 2 show predicted numbers of sick and dead due to COVID19 at R_o=2.46 and R_o=3 and age

structure of hospitalized patients and deaths.

a) Distribution of the total number of the susceptible, latently infected, infected and recovered on a daily basis since the epidemic onset (R_o=2.46)

b) Distribution of the total number of the susceptible, latently infected, infected and recovered on a daily basis since the epidemic onset (R_o=3.1)

c) Daily fluctuations of the susceptible, recovered and net disease transmission rates (R_o=2.46)

d) Daily fluctuations of the susceptible, recovered and net disease transmission rates (R_o=3.1)

e) Daily fluctuations of apparent incidence, true incidence and net disease transmission rates (R_o=2.46)

f) Daily fluctuations of apparent incidence, true incidence and net disease transmission rates (R_o=2.46)

380   Fig.1 Model prediction of latently infected, diseased, recovered and daily fluctuations of R_n.

a) Necessary hospital capacities (R_o=2.46)

b) Necessary hospital capacities (R_o=3.1)

c) Age structure of hospitalised patients (R_o=2.46)

d) Age structure of hospitalised patients (R_o=3.1)

e) Expected age structure of the dead (R_o=2.46)

f) Expected age structure of the dead (Ro=3.1)

Fig.2 Model prediction of required hospital capacities under the assumption of different intervention

measures.

3.2. Predicting the number of sick, hospitalized patients and deaths caused by COVID 19 in

the conditions of application of restrictive anti-epidemic measures

When the spread of COVID19 epidemic through totally susceptible population in the Republic of Serbia is

simulated, under an assumption of only incidental movement among the population, basic reproduction

number of R_o = 2.46, and with lock-down of entire country, a significant slowdown of the epidemic was

observed. Initial infection doubling time was 11 days. The peak of the epidemic wave was reached 269

days after the epidemic onset and it was 25,754 infected in one day. In the first year of the epidemic

3,283,691 people would get infected and 3,391 people would die. When the basic reproduction number

was increased to R_o=3.1, while keeping the same conditions of locking down the entire country, results

changed significantly. The initial infection doubling time was 9 days, the epidemic wave peak was reached

after 164 days and it yield 65,327 infected people in one day. In the first year of the epidemic 5,021,561

people would get infected and 4,923 would die. Table 6 provides the overview of epidemic indicators

obtained from the simulations of all five scenarios with R_o = 2.46 and R_o = 3.1. Figures 3. and 4 provide a

comparative overview of results of all five simulated scenarios. Panels a) and b) of Fig. 3 show the values

of cumulative incidences on a daily basis for R_o=2.46 and R_o=3.1, respectively. Panels b) and c) of Fig. 3

show the expected total number of hospitalised patients and patients in intensive care units (ICU) for

R_o=2.46 and R_o=3.1, respectively. Fig. 4 provides overview of required hospital capacities e.g. hospital bed

occupancy and the occupancy of beds in ICU on daily bases for R_o=2.46 and R_o=3.1, respectively. The

results show that after applying various measures to slow down the circulation of SARSCov-2, the number

of newly infected people, hospitalized patients and the occupancy of hospital capacities are the lowest in

the situations where rigorous anti-epidemic measures are applied to all population strata (Scenario 2 in

Table 6 and in Fig. 3 and Fig. 4). Openings of pre-school and elementary school's facilities leads to a visible

jump in the number of infected and hospitalized in all strata. This finding clearly shows that children,

although least susceptible to developing more severe clinical pictures, are important when transmitting

SARSCov-2 (Scenario 3 in Table 6 and Fig. 3). Opening of the high schools and colleges also leads to a

visible increase in the number of newly infected and hospitalized patients, including an increase in the

number of deaths (Scenario 5 in Table 6 and Fig. 3). Without the application of any intervention measures,

the greatest burden on the health system could be expected 196 days from the beginning of the epidemic

at R_o=2.46 or 144 days at R_o=3.1. Depending on the R_o value used in simulation, it would be necessary to

provide 12,740 (19,380) hospital beds and an additional 8,486 (12,355) in intensive care units. In the case

of Scenario 2, there is a significant slowdown in the epidemic, so that after 621 (238) days at the moment

the highest occupancy of hospital facilities would be to provide 461 (4,181) beds in COVID hospitals and

358 (3,120) beds in intensive care units (Fig. 4).

Table 6. Results of different simulated scenarios (R_o = 2.46 and R_o = 3.1). The data refers to the period of 365 days from epidemic onset.

Scenario mark	SC1		SC2		SC3		SC4		SC5	
Basic reproductive number	R_o=2.64	R_o=3.1	R_o=2.64	R_o=3.1	R_o=2.64	R_o=3.1	R_o=2.64	R_o=3.1	R_o=2.64	R_o=3.1
Cumulative incidence (CI)	6,633,834	7,035,443	3,283,691	5,021,561	4,969,942	5,990,015	5,171,901	6,087,288	5,408,422	6,257,345
Apparent CI	1,326,767	1,407,089	656,738	1,004,312	993,988	1,198,003	1,034,380	1,217,458	1,081,684	1,251,469
Overall hospitalized	115,612	122,782	67,142	98,973	88,529	106,442	90,361	106,945	92,727	108,242
Overall in intensive care	37,212	39,512	22,353	32,690	29,288	34,975	29,865	35,121	30,545	35,452
Overall deaths	5,525	5,868	3,391	4,923	4,424	5,260	4,510	5,281	4,609	5,328

a) Comparative overview of cumulative incidences (CI). Results obtained from simulated scenarios 1-5 (R_o=2.46)

b) Comparative overview of cumulative incidences (CI). Results obtained from simulated scenarios 1-5 (R_o=3.1)

c) Comparative overview of hospitalised patients on a daily basis. Results obtained from simulated scenarios 1-5 (R_o=2.46)

d) Comparative overview of hospitalised patients on a daily basis. Results obtained from simulated scenarios 1-5 (R_o=3.1)

e) Comparative overview of expected number of patients in intensive care on a daily basis. Results obtained from simulated scenarios 1-5 with R_o=2.46

f) Comparative overview of expected number of patients in intensive care on a daily basis. Results obtained from simulated scenarios 1-5 with R_o=3.1

421   Fig.3 Model prediction of expected number of hospitalised patient and patient in intensive care.

a) Comparative overview of hospital bed occupancy (R_o=2.46)

b Comparative overview of hospital bed occupancy (R_o=3.1)

c) Comparative overview of ICU bed occupancy (Ro=2.46)

d) Comparative overview of ICU bed occupancy (Ro=3.1)

Fig. 4 Model prediction of required hospital capacities needed to treat patients with COVID 19.

3.3.   COVID19 simulation and disease control by implementing a hypothetical vaccine

Assuming that the disease is spreading at the basic reproduction number of R_o=2.46, the herd

immunity threshold (when the disease can be expected to slow down and the chain of infection is expected

to be broken) would be 59.35%, while at R_o=3.1, it would be 67.74%. Depending on the efficacy of the

potential vaccine, the required vaccination coverage should be 87% (v_e =68%), 74.19% (v_e =80%), 69.82%

(v_e=85%) and 95.41% (v_e=71%), 84.68% (v_e=80%), 79.70% (v_e=85%) for R_o=2.46 and R_o=3.1, respectively.

Fig. 5 shows different scenarios of COVID19 control strategies based solely on vaccination.

a) Comparative overview of daily incidences (R_o=2.46).

b) Comparative overview of daily incidences (R_o=3.1).

c) Comparative overview of hospitalised patients (R_o=2.46).

d) c) Comparative overview of hospitalised patients (R_o=3.1).

e) Comparative overview of deaths (Ro=2.46)

f) Comparative overview of deaths (Ro=3.1)

NV-without vaccination and any non-pharmaceutical intervention
Fig. 5 Results of simulated COVID19 control based solely on vaccination, scenarios 6-9 (R_o=2.46 and
R_o=3.1).

3.4. Results of the model sensitivity analysis

The results of the sensitivity analysis are presented in table 9. The table shows increased values of input

parameters and the percentage of the parameter increase relative to the basic scenario, as well as the

values of output results obtained after the simulation of the changed scenario.

Table 9. Results of model sensitivity analysis.

Model parameter mark	Change relative to the basic scenario	CI	Deaths (D_{th})	Change in CI relative to the basic scenario	Change in D_{th} relative to the basic scenario
β	5%	6,733,914	5,610	1.51%	1.54%
β	10%	6,819,454	5,683	2.80%	2.86%
β	25%	7,022,923	5,857	5.87%	6.01%
f^{-1}	5%	6,642,443	5,532	0.13%	0.13%
f^{-1}	10%	6,651,063	5,539	0.26%	0.25%
f^{-1}	25%	6,676,747	5,560	0.65%	0.63%
r	5%	6,508,468	5,420	-1.89%	-1.90%
r	10%	6,376,497	5,309	-3.88%	-3.90%
r	25%	5,938,474	4,943	-10.48%	-10.54%

4. Conclusions and discussion

For the needs of this research, we augmented the classic deterministic model by adding the compartments

of vaccinated and latently infected subjects. By adding birth and death rates, we enabled daily fluctuations

of the overall simulated population, which brought us closer to the real conditions in which the disease is

transmitted. When we assumed that the recovered lose immunity over time, we obtained dynamic

oscillations of epidemic waves through susceptible population. When we simulated different disease control

scenarios of COVID19 epidemic based on non-pharmaceutical intervention measures, scenario number 2

proved to be the most effective approach to the disease control, because it implemented the most

comprehensive anti-epidemic measures (entire country lock down). However, the basic problem of this

approach is feasibility and possibility to maintain the measures in the long term.

The model demonstrated that students, children and younger school-age generations have an

important role in transmitting COVID19, especially if they come into contact with a more vulnerable

population. The model showed that, in the case of returning school children of all ages to schools, an

increase of 32.98% of the estimated deaths and 57.50% of the number of infected is possible, when

compared to the conditions before opening of the schools (Scenarios 2 and 4). However, most dead and

seriously diseased people are found in the older population. This is particularly important when planning

intervention measures, especially when deciding on which restrictions to be lifted and how (opening

schools, students' return to faculties etc.). The model demonstrated that COVID19 has a potential to spread

rapidly and linger in population. Due to a large number of the infected and duration of the disease, there

are significant needs for hospital capacities, especially in the conditions when the disease is suppressed

by implementing partly relaxed anti-epidemic measures, or in the case of absence of any measures. Without

the application of any intervention measures, at the moment of the greatest load, depending of actual R_o,

the health system would have to provide 12,740 (R_o=2.46) hospital beds for the care of the patients and an

additional 8,486 in intensive care units. On the other hand, in the case of the application of the strictest anti-

epidemic measures, the needs decrease to only 461 beds in COVID hospitals and additional 358 beds in

intensive care units. In the case of continued implementation of current measures, which are significantly

less intense than the measures applied at the beginning of the epidemic (Scenario 4), it is necessary to

provide 3,508 beds in COVID hospitals and 2,617 beds in intensive care units in the entire country. The

model also shows that the needs for hospital capacities decline with the ending of the first epidemic wave,

since daily incidence decreases and during the second and third waves it never reaches the initial peaks,

but these needs still remain substantial. For example, in the case of Scenario 1, at the top of the second

epidemic wave, it is necessary to provide 3,643 beds in COVID hospitals and 2,575 in intensive care units,

which makes 30% and 25% of the required capacities of the first wave of the epidemic.

Based on the cyclical patterns of the epidemic waves and duration of simulated epidemics, the

model demonstrated that the disease has a potential to linger in population and that it will most probably

have a seasonal pattern. Therefore, potential vaccines can have an enormous potential and significance

for COVID19 control. Depending on the efficacy of future vaccines, the disease can be stopped and curbed

almost solely by implementing the measure of vaccination. However, the necessary conditions for these

predictions and expectations are the efficacy of potential vaccines and the ability of a health systems to

implement vaccination to a satisfactory extent and rapidly, especially with regards to the most sensitive

categories of population. Depending on the R_o, a vaccine that would have an efficacy ≥ 68-71% could stop

the pandemic and break the chain of infection. However, even vaccines with lower efficacy could be useful

as they would significantly reduce the number of cases and deaths, especially if used in combination with

the other disease control measures. The required minimal vaccination coverage should be 87% (v_e =68%),

74.19% (v_e =80%), 69.82% (v_e=85%) and 95.41% (v_e=71%), 84.68% (v_e=80%), 79.70% (v_e=85%) for

R_o=2.46 and R_o=3.1, respectively. The minimum daily vaccination rate should be 0.47% for vaccines with

an efficiency of 85%, and 0.59% for vaccines with an efficiency of 68%.

Based on the obtained results, we can conclude that at this point, without the application of specific

pharmaceutical products, COVID19 suppression is highly dependent on the basic reproduction number

(R_o), and that more intensive contacts and relaxed measures can result in a dramatic spread of the virus.

The choice of intervention measures depends on the feasibility of their implementation and their efficacy in

different social contexts. COVID19 will likely have to be suppressed in this way for a certain period of time.

This will most probably last until sufficient quantities of a reliable and effective vaccine are available, and

thereafter until optimal vaccination coverage is achieved.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial,

or not-for-profit sectors.

References

[1] World Health Organization (WHO), Coronavirus disease (COVID-19) outbreak situation. 2020.

https://www.who.int/emergencies/diseases/ novel-coronavirus-2019, [accessed 18 August 2020].

[2] https://covid19.rs, 2020, [accessed 18 August 2020].

[3] Wangping J, Ke H, Yang S, Wenzhe C, Shengshu W, Shanshan Y, Miao L. Extended SIR prediction of

the epidemics trend of COVID-19 in Italy and compared with Hunan, China. Frontiers in Medicine 2020;

7:169.

[4] Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, Davies N. Early dynamics of

transmission and control of COVID-19: a mathematical modelling study. The Lancet infectious diseases

2020. Doi: 10.1016/S1473-3099(20) 30144-4.

[5] Matt J. Keeling, Pejman Rohani. Modelling Infectious Diseases in Humans and Animals, Princeton

University Press, 3 Market Place, Woodstock, Oxfordshire OX20 1SY, p. 8-10.

[6] Kermack, W. and A. McKendrick (1927). A contribution to the mathematical theory of epidemics.

Proceedings of the Royal Society. A 115, p. 700–721.

[7] Barnes B. and G. R. Fulford. Mathematical modelling with case studies Using Maple™ and MATLAB,

CRC Press Taylor & Francis Group 6000 Broken Sound Parkway NW, Suite 300, Boca Raton, FL 33487-

2742, p. 100-102.

[8] Imperial College COVID-19 Response Team. Report 9: Impact of non-pharmaceutical interventions

(NPIs) to reduce COVID-19 mortality and healthcare demand; 2020.

https://www.imperial.ac.uk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-

COVID19-NPI-modelling-16-03-2020.pdf.

[9] Wu Z, Googan JMM. Characteristics of and important lessons from the coronavirus disease 2019

(COVID-19) outbreak in China: Summary of a Report of 72 314 Cases from the Chinese Center for Disease

Control and Prevention. JAMA. 2020;323(13):1239-1242. Doi:10.1001/jama.2020.2648.

[10] Emilia Vynnycky, Richard G. White with an introduction by Paul E.M. Fine. An Introduction to Infectious

Disease Modelling. Oxford University Press Inc., New York © Oxford University Press; 2010.

[12] European Centre for Disease Prevention and Control. Immune responses and immunity to SARS-CoV-

2; 2020.   https://www.ecdc.europa.eu/en/covid-19/latest-evidence/immune-responses; [accessed 18

August 2020].

[13] Woelfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Mueller MA, et al. Clinical presentation

and virological assessment of hospitalized cases of coronavirus disease 2019 in a travel-associated

transmission cluster. medRxiv. 2020; https://doi.org/10.1101/2020.03.05.20030502.

[14] Liu W, Liu L, Kou G, Zheng Y, Ding Y, Ni W, et al. Evaluation of Nucleocapsid and Spike Protein-based

ELISAs for detecting antibodies against SARS-CoV-2. medRxiv. 2020.

https://doi.org/10.1101/2020.03.16.20035014.

[15] Long Q-x, Deng H-j, Chen J, Hu J, Liu B-z, Liao P, et al. Antibody responses to SARS-CoV-2 in COVID-

19 patients: the perspective application of serological tests in clinical practice. medRxiv.

https://doi.org/10.1101/2020.03.18.20038018.

[16] Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-CoV-2 in patients

of novel coronavirus disease 2019. medRxiv. 2020; https://doi.org/10.1101/2020.03.02.20030189.

[17] Okba NMA, Muller MA, Li W, Wang C, Geurtsvan Kessel CH, Corman VM, et al. SARS-CoV-2 specific

antibody responses in COVID-19 patients. medRxiv. 2020; https://doi.org/10.1101/2020.03.18.20038059.

[18] Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-CoV-2 in patients

of novel coronavirus disease 2019. medRxiv. 2020; https://doi.org/10.1101/2020.03.02.20030189.

[19] Kellam P, Barclay W. The dynamics of humoral immune responses following SARS-CoV-2 infection

and the potential for reinfection. The Journal of general virology. 2020 May 20;

[20] Gerardo Chowell, James M. Hyman, Lu´ıs M. A. Bettencourt, Carlos Castillo-Chavez. Mathematical

and Statistical Estimation Approaches in Epidemiology, ISBN 978-90-481-2312-4, Doi 10.1007/978-90-

481-2313-1, Springer Dordrecht Heidelberg London New York; 2009.

[21] Michael Thrusfield. Veterinary epidemiology. Third edition. Blackwell Science Ltd, a Blackwell

Publishing company; 2007.

[22] Marco D'Arienzo, Angela Coniglio. Assessment of the SARS-CoV-2 basic reproduction number, R0,

based on the early phase of COVID-19 outbreak in Italy. Biosafety and Health 2(2020) 57-59.

[23] The World Bank. World Developments Indicators;

https://www.google.com/publicdata/explore?ds=d5bncppjof8f9_&met_y=sp_dyn_le00_in&idim=country:S

RB:HRV:BIH&hl=en&dl=en#!ctype=l&strail=false&bcs=d&nselm=h&met_y=sp_dyn_le00_in&scale_y=lin&

ind_y=false&rdim=world&idim=country:SRB&ifdim=world&hl=en_US&dl=en&ind=false; [accessed 25

August 2020].

[24] Demografski pregled 2017. Broj: 68/2018, https://www.minrzs.gov.rs/sites/default/files/2019-

01/demografski_pregled_68.pdf; [accessed 15 July 2020].

[25] Roberto Pastor-Barriuso et al. Infection fatality risk for SARS-CoV-2: a nationwide seroepidemiological

study in the noninstitutionalized population of Spain. medRxiv. 2020; preprint doi:

https://doi.org/10.1101/2020.08.06.20169722.

[26] Alex Viguerie, Guillermo Lorenzo, Ferdinando Auricchio, Davide Baroli, Thomas J.R. Hughes, Alessia

Patton, Alessandro Reali, Thomas E. Yankeelov, Alessandro Veneziani. Simulating the spread of COVID-

19 via a spatially-resolved susceptible–exposed–infected–recovered–deceased (SEIRD) model with

heterogeneous diffusion. Applied Mathematics Letters 111 (2021) 106617.

[27] David Baud, Xiaolong Qi, Karin Nielsen-Saines, Didier Musso, Léo Pomar, Guillaume Favre. Real

estimates of mortality following COVID-19 infection. The Lancet. 2020 Jul; 20(7): 773. Doi: 10.1016/S1473-

3099(20)30195-X.

[28] Charles F. Manski, Francesca Molinari. Estimating the COVID-19 infection rate: Anatomy of an

inference problem. Journal of Econometrics. https://doi.org/10.1016/j.jeconom.2020.04.041.

[29]   WHO.   Q&A: Influenza   and   COVID-19   -   similarities   and   differences.

https://www.who.int/westernpacific/news/q-a-detail/q-a-similarities-and-differences-covid-19-and-

influenza. [accessed 18 August 2020].

[30] CDC. https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html; [accessed 18 August

2020].

[31] CEBM- The Centre for Evidence-Based Medicine develops, promotes and disseminates better

evidence for healthcare. Estimating the infection fatality ratio in England. https://www.cebm.net/covid-

19/estimating-the-infection-fatality-ratio-in-england/; [accessed 9 October 2020].

[32] Wood, S. N. Did COVID-19 infections decline before UK lockdown? ArXiv:2005.02090 [q-Bio, Stat].

http://arxiv.org/abs/2005.02090.

[33] CEBM- The Centre for Evidence-Based Medicine develops, promotes and disseminates better

evidence for healthcare. COVID-19: What proportion are asymptomatic? https://www.cebm.net/covid-

19/covid-19-what-proportion-are-asymptomatic/; [accessed 18 August 2020].

[34] BMJ 2020; 368 doi: https://doi.org/10.1136/bmj.m1165. https://www.bmj.com/content/368/bmj.m1165

[35] Selene Ghisolfi, Ingvild Almås, Justin Sandefur, Tillmann von Carnap, Jesse Heitner, Tessa Bold.

Predicted COVID-19 Fatality Rates Based on Age, Sex, Comorbidities, and Health System Capacity. 2020

Jun.https://www.cgdev.org/sites/default/files/predicted-covid-19-fatality-rates-based-age-sex-

comorbidities-and-health-system-capacity.pdf.

[36] Roberto Pastor-Barriuso et al. Infection fatality risk for SARS-CoV-2: a nationwide seroepidemiological

study   in   the   noninstitutionalized   population   of   Spain.   medRxiv.   2020;   preprint   doi:

https://doi.org/10.1101/2020.08.06.20169722.