Title: Rate of Antibiotic Use and Associated Risk Factors in COVID-19 Hospitalized Patients

Running title: COVID-19 antibiotic risk factors

Authors: Alysa J. Martin, PharmD[1]; Stephanie Shulder, PharmD[1]; David Dobrzynski, MD[2];
Katelyn Quartuccio, PharmD[3]; Kelly E. Pillinger, PharmD[1]

[1] University of Rochester Medical Center Strong Memorial Hospital, Department of Pharmacy, Rochester, NY, USA

[2] University of Rochester Medical Center, Division of Infectious Diseases, Rochester, NY, USA

[3] University of Rochester Medical Center Highland Hospital, Department of Pharmacy, Rochester, NY, USA

Acknowledgements:

The authors thank Raquel Roberts, PharmD for her assistance with data collection. There was

no funding for the study.

Conflict of Interest Statement: The authors have no personal conflicts of interests to disclose.

The authors would like to disclose that K. Quartuccio's spouse consults for AstraZeneca.

Key words: COVID-19, antimicrobial stewardship, antibiotics, community-acquired pneumonia,

bacterial infection

Abstract

Background- Literature suggests that antibiotic prescribing in COVID-19 patients is high, despite

low rates of confirmed bacterial infection. There are little data on what drives prescribing

habits.

Objective- This study sought to determine antibiotic prescribing rates and risk factors for

antibiotic prescribing in hospitalized patients. It was the first study to assess risk factors for

receiving more than one course of antibiotics.

Methods- This was a retrospective, multi-center, observational study. Patients admitted from

March 1, 2020 to May 31, 2020 and treated for PCR-confirmed COVID-19 were included. The

primary endpoint was the rate of antibiotic use during hospitalization. Secondary endpoints

included risk factors associated with antibiotic use, risk factors associated with receiving more

than one antibiotic course, and rate of microbiologically confirmed infections.

Results- A total of 208 encounters (198 patients) were included in the final analysis. Eighty-

three percent of patients received at least one course of antibiotics, despite low rates of

microbiologically confirmed infection (12%). Almost one-third of patients (30%) received more

than one course of antibiotics. Risk factors identified for both antibiotic prescribing and

receiving more than one course of antibiotics were more serious illness, increased hospital

length of stay, intensive care unit admission, mechanical ventilation, and acute respiratory

distress syndrome.

Conclusion and relevance – There were high rates of antibiotic prescribing with low rates of

bacterial co-infection. Many patients received more than one course of antibiotics during

hospitalization. This study highlights the need for increased antibiotic stewardship practices in

COVID-19 patients.

Introduction

The novel coronavirus disease identified in 2019 (COVID-19) is caused by a betacoronavirus known as SARS-CoV-2. Despite the fact that COVID-19 is a viral infection, clinical manifestations may present similarly to a bacterial pneumonia.[1] Patients often present with respiratory symptoms including fever, cough, dyspnea, and bilateral changes on chest imaging.[1] It can be difficult to determine if patients have a bacterial respiratory coinfection along with COVID-19 infection. This may lead to over-prescribing of antibiotics in patients with COVID-19.

Despite bacterial coinfection being reported in 2-8% of patients, rates of antibiotic prescribing have ranged from 57-95% in hospitalized patients.[1-7] Additionally, there are reports of increased consumption of antibiotics during the COVID-19 pandemic.[6-9] Over-prescribing of antibiotics may increase the risk of adverse side effects, nosocomial infection, and antibiotic resistance. Several organizational guidelines recommend starting empiric antibiotic therapy if concern for bacterial pneumonia or sepsis, but to re-evaluate daily and de-escalate or discontinue if no evidence of bacterial infection.[10-14] Furthermore, they emphasize the importance of antimicrobial stewardship programs which may help to avoid starting antibiotics reflexively or continuing unnecessary courses. The World Health Organization (WHO) has released guidance which discourages the use of antibiotic therapy for patients with COVID-19 infections unless signs and symptoms of a bacterial infection exist.[11]

Unfortunately, there are limited data to date describing the need or role for antibiotics in COVID-19 or outcomes for patients who receive them. Additionally, little is known regarding

risk factors associated with antibiotic use. There have only been two previous studies which

evaluated prescribing trends and risk factors of antibiotic prescribing. Both studies focused on

prescribing rates, coinfections, and risk factors associated with antibiotic prescribing within the

first 48 hours of admission.[4, 5]

The purpose of this study was to evaluate the rate of antibiotic use in COVID-19 patients

and risk factors associated with antibiotic use throughout hospitalization. A better

understanding of prescribing practices can be used to evaluate opportunities for antibiotic

stewardship for future patients with COVID-19.

Methods

Study design

This was a multi-center, retrospective, observational study of antibiotic prescribing

trends in patients with confirmed COVID-19 at three hospitals within the University of

Rochester Medical Center.

Patients 18 years and older admitted from March 1, 2020 to May 31, 2020 and treated

for PCR-confirmed COVID-19 were eligible for inclusion. Patients did not meet inclusion criteria

if they were only treated in the emergency department without subsequent admission. Patients

could be included more than once if they had multiple admissions within the study period.

Patients were excluded if they were asymptomatic. Data were collected until June 30, 2020.

This study was reviewed and deemed exempt by the Institutional Review Board.

Data Collection

Patients were identified from the electronic health record based on a reported positive

SARS-CoV-2 PCR test. Baseline data including age, gender, race, weight, height, comorbidities,

date of first positive SARS-CoV-2 PCR, and admission and discharge dates were collected. Labs

on admission (or at time of positive test if nosocomial transmission), microbiologic data, clinical

symptoms, imaging, antibiotic data, COVID-19 investigational or off-label treatment, death and

re-admission within 30 days were also collected.

Outcomes

The primary endpoint of the study was the rate of antibiotic use in patients admitted for

COVID-19. Secondary endpoints included risk factors associated with antibiotic use, risk factors

associated with receiving more than one antibiotic course, and rate of microbiologically-

confirmed infections. Microbiologically-confirmed infection was defined as a positive result

from positive urinary antigens or respiratory and blood cultures not suggestive of contaminant

or colonization.

Statistical Analysis

Data were analyzed using R statistical software version 4.0.2 (Boston, MA) with a p-

value < 0.05 considered statistically significant. Continuous variables were described using a

median with interquartile range (IQR). Categorical variables were analyzed using Fisher's exact

testing. Two-sample Wilcoxon test was used for continuous variables. To evaluate risk factors

associated with antibiotic use, a univariate analysis was conducted and variables with a p-value

≤ 0.05 were included in a multivariate logistic regression.

Results:

A total of 228 encounters met inclusion criteria, with twenty excluded due to asymptomatic disease, leaving 208 encounters included in the final analysis (n=198 patients). The median age was 69 years (IQR 60-80) and 50% (105/208) were male. The most common comorbidities were hypertension (143/208, 69%), obesity (81/208, 39%), and diabetes (81/208, 39%). Of the 208 encounters, 83% (172/208) of patients received at least one course of antibiotics (Table 1). Antibiotic prescribing did not differ based on gender, age, race, or comorbidities. Even though a small subset of patients (n=11), all immunocompromised patients received antibiotics

While rates of antibiotic prescribing overall were high, we saw a temporal decrease in antibiotic prescribing. When consolidating encounters by month of admission, rates of antibiotic prescribing were highest in March, with 88% (42/49) of encounters receiving antibiotics, compared to 82% (127/155) in April and 75% (3/4) in May.

Rates of antibiotic prescribing were significantly higher in patients admitted to the intensive care unit (ICU), requiring O2 support (including ventimask, BiPAP and mechanical ventilation), and with acute respiratory distress syndrome (ARDS) ($p < 0.01$). Median length of stay was 6.5 days longer in those receiving antibiotics ($p < 0.01$). Patients who received investigational or off-label therapy for COVID-19 were also more often prescribed antibiotics ($p = 0.03$). Patients who died within 28 days were more often prescribed antibiotics (92%, 47/51, $p = 0.02$). Procalcitonin (PCT) at baseline was higher in those who were prescribed antibiotics ($p < 0.01$). In a subgroup analysis, PCT > 0.5 was associated with a higher rate of antibiotic

prescribing [94% (44/47) prescribed antibiotics vs. 6% (3/47) did not receive antibiotics, OR

3.76, CI 1.09-20.1, p = 0.03]. Of patients with a PCT > 0.5, 43/47 had respiratory or blood

cultures obtained. Microbiologically confirmed bacterial infection was present in 12% of

patients with PCT >0.5 (5/43).

Respiratory cultures were obtained in 32% (67/208) of patients. In total, 19% (13/67)

were positive (Table 2). *Staphylococcus aureus* was identified in 5 of 13 positive respiratory

culture results (3% of total patients). Blood cultures were obtained in the majority of patients

(149/208, 72%), but few were positive (12/149, 7%). All influenza PCR and *Legionella* urinary

antigen (UAg) tests were negative (n= 88 and 93 respectively). Two *Streptococcus pneumoniae*

UAg tests were positive (2.4%, 2/84). In addition, 75 patients (36%) were screened for MRSA

nares colonization. Overall, there were 24 microbiologically-confirmed infections and all were

treated with antibiotics [12% (24/208)]. In a subgroup analysis, patients with microbiologically-

confirmed infection were more likely to receive additional courses of antibiotics (20/24, 83% p

< 0.01), be admitted to the ICU (23/24, 96%, p < 0.01), and have a longer length of stay (median

29.5 days, p < 0.01).

The most frequently prescribed initial antibiotics were ceftriaxone in 113 (66%) patients

and azithromycin in 99 (57%) patients for community acquired pneumonia (CAP), followed by

an anti-pseudomonal beta lactam (23%) and vancomycin (23%) for hospital acquired

pneumonia (HAP) or ventilator associated (VAP) (Figures 1 and 2). The median time from SARS-

CoV-2 collection to antibiotic initiation was one day (IQR 1-4.5 days). Median duration of

therapy for the initial course of antibiotics was five days (IQR 2-6 days). In 41 (24%) patients,

the initial regimen was changed; 17 patients were escalated to broader therapy (42%) and 15

patients were de-escalated to narrower therapy (37%). Nine (22%) had another regimen change

such as intravenous to oral conversion or transition to another agent with similar spectrum of

activity and not based on positive culture result.

Fifty-two patients (30%) received a second course of antibiotics and 19 patients (11%)

received more than 2 antibiotic courses during hospitalization. The most frequently prescribed

antibiotics included an anti-pseudomonal beta lactam (58%) and vancomycin (58%) for

HAP/VAP (Figures 1 and 2). The median duration of second course antibiotics was 6 days (IQR 3-

8.25). Duration of admission, ICU admission, ICU length of stay (LOS), mechanical ventilation,

microbiologically-confirmed infection, and duration of initial course of antibiotics were all risk

factors for receiving more than one course of antibiotics (Table 3). Of patients with

microbiologically-confirmed infection who received additional antibiotic courses, 14 (70%)

received treatment for HAP/VAP, 4 (20%) received treatment for bloodstream infection and 2

(10%) received treatment for other infections unrelated to cultures.

Discussion

In this multi-center study of patients hospitalized with COVID-19, we found an overall

high rate of antibiotic prescribing, with 83% of patients receiving antibiotics. Of these patients,

30% received at least one additional course of antibiotics within the same hospitalization,

increasing the overall antibiotic exposure. Despite the high rate of antibiotic prescribing, there

were low rates of bacterial coinfection, with only 12% of patients having microbiologically-

confirmed infection for initial antibiotic use. These findings are similar to the results of previous

studies, suggesting the need for strategies to help clinicians judiciously prescribe antibiotics in

patients with COVID-19.[1-5, 15] Previous studies conducted in the United States have shown that

concomitant community acquired infections in COVID-19 patients are low based on

microbiological data within the first 48 hours of admission.[4, 5] In our study, we found only

slightly higher rates of microbiologically-confirmed infections than in other US studies, despite

the fact that we reported all microbiologic data throughout hospitalization. The higher rate may

be explained by bacterial infections acquired during hospitalization. Patients with

microbiologically-confirmed infections were more likely to receive more than one course of

antibiotics, be admitted to the ICU, and have a longer length of stay. Additional courses of

antibiotics were often prescribed for empiric or targeted treatment of nosocomial infection.

Early in the pandemic, little was known regarding COVID-19 presentation, disease

progression and rates of bacterial co-infection. Thus, guideline recommendations were evolving

throughout the study period. Initial recommendations often encouraged antibiotic use. Within

the University of Rochester Medical Center, our practices and institutional guidelines

recommended initiating empiric antibiotics in the majority of COVID-19 patients. As information

became more available, both national and institutional guidance changed, recommending

judicious use of antibiotics. Subsequently, our antibiotic prescribing decreased. This was

reflected in the temporal decline in antibiotic prescribing we observed from March to May.

There were several indicators that serious illness increased the risk of antibiotic

prescribing. Initial antibiotic prescribing was more likely in patients with elevated PCT, longer

length of stay, ICU admission, longer ICU length of stay, and administration of off-label or

investigational therapy for COVID-19. Additionally, all patients with ARDS or requiring more

invasive O2 support received at least 1 course of antibiotics. These findings are consistent with

previous studies which found patients were more likely to receive antibiotics if they had severe

disease upon presentation.[4, 5] To our knowledge, this is the first study to report rates of

additional antibiotic prescribing beyond initial empiric therapy. Patients with COVID-19 often

have a prolonged hospitalization due to the underlying disease process. Our median LOS was

significantly longer in patients who received at least one antibiotic course (12 days vs 5.5 days,

p < 0.001), and prolonged hospitalization may also increase the potential for additional

antibiotic exposure. Thus, it is important to understand antibiotic prescribing trends and

associated risk factors beyond the first course. Our study found that nearly one-third of all

patients received two or more courses of antibiotics. Variables associated with receiving two or

more antibiotic courses included: longer length of stay, ICU admission, longer ICU length of

stay, and mechanical ventilation. This is likely explained as these are all risk factors for hospital-

acquired illness and antibiotics were commonly prescribed for nosocomial infection (e.g.

HAP/VAP).

Procalcitonin has been suggested as a potentially useful biomarker to differentiate

bacterial and viral infections and assist with clinical decision-making. However, the ATS/IDSA

CAP guidelines strongly advise initiating antibiotics based off clinical judgement rather than PCT

alone.[13] In alignment with these guidelines, our institution does not recommend routinely using

PCT to guide antibiotic initiation for CAP or sepsis. In COVID-19 patients specifically, there are

conflicting data regarding baseline PCT in COVID-19 patients.[4, 16] One two-center study found

that patients with PCT available prior to antibiotic initiation were less likely to be prescribed

antibiotics.[17] At our sites, the patients prescribed antibiotics had significantly higher baseline

PCT, however the median was only 0.28. In a subgroup analysis focusing on patients who had a

PCT of > 0.5, 94% of patients (41/44) received antibiotics. Rates of microbiologically confirmed

infection were similar between patients with a PCT > 0.5 and the entire cohort (12%). It is

unclear at this time the utility of using PCT to determine if there is bacterial co-infection with

COVID-19, and in our study it did not appear to be predictive of confirmed infection.

Previous studies have shown high rates of broad-spectrum antibiotics for initial empiric

therapy in COVID-19 patients.[1, 3-6, 8, 15] However, in our patient population, most patients were

treated initially for CAP. The most common agents included ceftriaxone plus azithromycin,

which is consistent with the 2019 ATS/IDSA recommendations for CAP.[13] It is worth noting that

our institution did not routinely recommend use of azithromycin off-label for anti-inflammatory

effects in COVID-19. During the initial antibiotic course, 37% of patients were de-escalated to

narrower therapy. Not surprisingly, patients who received a second course of antibiotics were

more likely to receive broad-spectrum antimicrobials; over half of patients who received more

than one course of antibiotics were treated for HAP/VAP and prescribed an anti-pseudomonal

beta lactam. Respiratory cultures were obtained in only 30.2% of patients, with a large number

of patients receiving empiric treatment for CAP, HAP or VAP. In part, the low number of

respiratory cultures obtained may be due to the symptomatology of disease, with the majority

of patients experiencing dry, non-productive cough.[1] Additionally, limited PPE early in the

pandemic created additional challenges in obtaining respiratory cultures due to concerns of

transmission and aerosolization. Also, the 2019 ATS/IDSA CAP guidelines do not recommend

routine sputum gram stain and culture in patients with non-severe CAP or those without multi-

drug resistant organism risk factors due to overall poor yield and detection of organisms.[13]

However, respiratory cultures are recommended in patients being treated empirically for MDR

organisms, and in those being treated for HAP/VAP.[13, 14] Respiratory cultures were not obtained

in all patients who met these criteria. Obtaining respiratory cultures and other microbiologic

data, when possible, on patients treated with broad spectrum antibiotics can potentially help

with de-escalation of therapy.

One limitation of this study is its retrospective design. Laboratory and microbiologic

data collected were not consistent amongst all patients. We found duration of admission and

ICU length of stay to be associated with higher antibiotic use, however it is difficult to say that

these are true risk factors for antibiotic prescribing. Instead, this difference was likely due to

antibiotic prescribing being higher in patients with more serious illness, and perhaps longer

admission leading to increased risk of nosocomial infections and subsequent antibiotic courses.

Additionally, we did not perform a time adjusted analysis. Another limitation is that the vast

majority of patients within our study received antibiotics which could have affected our ability

to detect differences between the two groups. Early in the pandemic, most COVID-19 patients

within our institution received antibiotics. While we saw a temporal decrease over time, it is

difficult to assess risk factors for antibiotic use in our patients treated for COVID-19 early in the

study when antibiotic use was routine.

While there were limitations of this study, there were also several strengths. This study

was multi-site, including three hospitals within the University of Rochester Medical Center: two

community hospitals and a tertiary academic medical center. Therefore, the results of this

study may be generalizable to similar sites. Finally, as previously mentioned, this is the first

study to assess overall antibiotic use throughout hospitalization, and determine risk factors for

additional antibiotic courses. Many patients with COVID-19 have prolonged hospitalization,

increasing the risk for antibiotic exposure. Our study examined antibiotic use that is more

representative of hospitalized COVID-19 patients. Based on the results of this study, our

institution plans to examine how other factors may increase or decrease the risk of antibiotic

prescribing, as well as trend our use of antibiotics throughout the pandemic.

Conclusions

Our study found high rates of antibiotic prescribing, despite low rates of respiratory

cultures and confirmed microbiologic infections. Nearly one-third of patients were treated with

more than one antibiotic course during hospitalization. These data highlight a role for

antimicrobial stewardship during the COVID-19 pandemic. Further studies are needed to

examine the impact of antimicrobial stewardship initiatives in patients with COVID-19 and

assess if antibiotic prescribing practices have changed with increasing information related to

bacterial co-infection that wasn't available at the beginning of the pandemic.

References:

1. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet (London, England) 2020;10229:1054-62.

2. Crotty MP, Dominguez EA, Akins R, et al. Investigation of subsequent and co-infections associated with SARS-CoV-2 (COVID-19) in hospitalized patients. medRxiv 2020.

3. Rawson TM, Moore LSP, Zhu N, et al. Bacterial and fungal co-infection in individuals with coronavirus: A rapid review to support COVID-19 antimicrobial prescribing. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America 2020.

4. Vaughn VM, Gandhi T, Petty LA, et al. Empiric Antibacterial Therapy and Community-onset Bacterial Co-infection in Patients Hospitalized with COVID-19: A Multi-Hospital Cohort Study. 2020.

5. Wei W, Ortwine JK, Mang NS, Joseph C, Hall BC, Prokesch BC. Limited Role for Antibiotics in COVID-19: Scarce Evidence of Bacterial Coinfection. Available at SSRN 3622388 2020.

6. Beović B, Doušak M, Ferreira-Coimbra J, et al. Antibiotic use in patients with COVID-19: a 'snapshot' Infectious Diseases International Research Initiative (ID-IRI) survey. Journal of Antimicrobial Chemotherapy 2020.

7. Buehrle DJ, Decker BK, Wagener MM, et al. Antibiotic consumption and stewardship at a hospital outside of an early Coronavirus disease 2019 epicentre. 2020;AAC.01011-20.

8. Abelenda-Alonso G, Padullés A, Rombauts A, et al. Antibiotic prescription during the COVID-19 pandemic: A biphasic pattern. Infection Control & Hospital Epidemiology 2020;1-2.

9. Nestler M, Godbout E, Lee K, et al. Impact of COVID-19 on Pneumonia-Focused Antibiotic Use at an Academic Medical Center. Infection control and hospital epidemiology 2020;1-9.

10. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. "Government Document"

11. Clinical management of COVID-19 Interim Guidance - May 2020, Available from https://www.who.int/publications-detail/clinical-management-of-covid-19. 2020.

12. Alhazzani W, Møller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). Intensive care medicine 2020;5:854-87.

13. Metlay JP, Waterer GW, Long AC, et al. Diagnosis and Treatment of Adults with Community-acquired Pneumonia. An Official Clinical Practice Guideline of the American Thoracic Society and Infectious Diseases Society of America. 2019;7:e45-e67.

14. Kalil AC, Metersky ML, Klompas M, et al. Management of Adults With Hospital-acquired and Ventilator-associated Pneumonia: 2016 Clinical Practice Guidelines by the Infectious Diseases Society of America and the American Thoracic Society. Clinical Infectious Diseases 2016;5:e61-e111.

15. Lansbury L, Lim B, Baskaran V, Lim WS. Co-infections in people with COVID-19: a systematic review and meta-analysis. The Journal of infection 2020;2:266-75.

16. Bass J, Singh A, Williams R, Lindner DH. Procalcitonin and COVID-19: A Reliable Clinical Tool. 2020.

17. Pulia MS, Wolf I, Schwei RJ, et al. Antibiotic Prescribing Patterns for COVD-19 in Two Emergency Departments with Rapid Procalcitonin. 2020;1-7.

Tables

Table 1. Baseline Characteristics of COVID-19 patients and Risk factors for Antibiotic Prescribing				
Baseline Characteristic	**Total**	**Antibiotics n=172**	**No Antibiotics n=36**	**P-value**
Male gender	105 (50.5)	84 (48.8)	21 (55.6)	0.43
Age, years (median, IQR)	69.0 (60-80)	70.0 (61-81)	65.5 (47-77)	0.12
Race				0.56
Caucasian	116 (55.7)	98 (56.9)	18 (47.2)	
Black/African American	69 (33.2)	53 (30.8)	16 (44.4)	
Asian	2 (1.0)	2 (1.2)	0 (0)	
Other	6 (2.9)	6 (3.5)	0 (0)	
Unknown	15 (7.2)	13 (7.6)	2 (5.6)	
Comorbidities				
Hypertension	143 (68.8)	116 (67.4)	27 (72.2)	0.43
Diabetes	81 (38.9)	70 (23.2)	11 (27.8)	0.35
Obesity	80 (38.5)	66 (38.4)	14 (38.9)	>0.99
Cardiac Disease	56 (26.9)	50 (29.1)	6 (16.7)	0.15
Chronic Lung Disease	43 (20.7)	37 (21.5)	6 (16.7)	0.65
CKD	45 (21.7)	40 (23.2)	5 (13.9)	0.20
None	21 (10.1)	17 (9.9)	4 (11.1)	0.77
Immunocompromised[a]	11 (5.3)	11 (6.4)	0 (0)	0.22
Labs (median, IQR)				
AST, U/L	43 (31-65)	46 (33.25-65)	35 (26.75-49)	0.02
ALT, U/L	30 (20-46)	29.5 (20-45)	32.5 (20-49.5)	0.83
Scr, mg/dL	1.130 (0.84-1.64)	1.14 (0.84-1.725)	1.06 (0.81-1.3)	0.14
WBC, /mm3	7.30 (5.50-10.50)	7.4 (5.5-10.7)	6.8 (4.8-9.2)	0.11
ALC, K/uL	1.0 (0.7-1.3)	1.0 (0.7-1.3)	1.1 (0.5-1.4)	0.10
CRP, mg/L	93.0 (39.5-134.5)	93 (39.5-134.5)	73 (37.5-118.5)	0.37
LDH, U/L	344 (264.5-467)	348 (285-474)	294 (239-387)	0.03
Ferritin, ng/mL	627 (304-1086)	627 (309-1133.5)	589 (235-980)	0.45
D-dimer, ug/mL	1.26 (0.63-2.08)	1.37 (0.71-2.1)	0.86 (0.6-1.6)	0.24
PCT, ng/mL	0.215 (0.120-0.620)	0.28 (0.13-0.63)	0.13 (0.09-0.20)	<0.01
Symptoms Prior to Admission				
Fever	120 (57.7)	101 (58.7)	19 (52.8)	0.59
Respiratory Symptoms	180 (86.5)	152 (88.4)	28 (75)	0.10

Gastrointestinal symptoms	76 (36.5)	60 (34.9)	16 (41.7)	0.34
Anosmia/Hyperosmia/Hypogeusia	11 (5.3)	10 (5.8)	1 (2.8)	0.69
Symptom duration prior to admission, days (median, IQR)	6.0 (3-9)	6.0 (3-9)	7.5 (4.75-9)	0.16
Location prior to admit				0.08
Home	140 (67.3)	110 (63.9)	30 (80.5)	
SNF	60 (28.8)	54 (31.4)	6 (16.7)	
Transfer from OSH	8 (3.8)	8 (4.7)	0 (0)	
Duration of admission, days (median, IQR)	11 (5-22)	12 (5.75-25)	5.5 (4-11.3)	<0.01
ICU admission	104	102 (58.9)	2 (5.6)	<0.01
ICU LOS, days (median, IQR)	10 (3-18)	10 (3.25-18)	6.5 (6.25-6.75)	0.59
O2 support				<0.01
Room Air	40 (19.2)	24 (13.9)	16 (44.4)	
Standard Nasal Cannula	67 (32.2)	49 (28.5)	18 (50.0)	
HFNC/Venti Mask	19 (9.1)	17 (9.9)	2 (5.6)	
BiPAP	5 (2.4)	5 (2.9)	0 (0)	
Non-rebreather	7 (3.4)	7 (4.1)	0 (0)	
Mechanical ventilation	70 (33.7)	70 (40.7)	0 (0)	
ARDS	78 (37.5)	78 (45.3)	0 (0)	<0.01
Received investigational or off-label treatment for COVID-19[b]	37 (17.8)	35 (20.2)	2 (5.6)	0.03
28-day mortality	51 (24.5)	48 (27.9)	3 (8.3)	0.02
30-day readmission	16 (7.7)	12 (6.9)	4 (11.1)	0.75
More than 1 admission for COVID-19	12 (5.8)	8 (4.7)	4 (11.1)	0.23
Time from discharge to COVID-19 readmission, days (median, IQR)	7 (4.5-12.25)	5 (2.75-11.5)	10 (7.75-12.5)	0.22

()= Percentages unless otherwise specified
a = immunocompromised patients were defined as those with previous solid organ transplant, active cancer receiving chemotherapy, patients on biologic therapy, and uncontrolled HIV
b = Investigational or off-label therapy included remdesivir, remdesivir + baricitinib, hydroxychloroquine, tocilizumab or convalescent plasma
IQR = Interquartile Range
CKD = chronic kidney disease
AST = Aspartate aminotransferase
ALT = Alanine aminotransferase
SCr= Serum Creatinine
WBC = White blood cell count
ALC = Absolute lymphocyte count
CRP = C-reactive protein
LDH= Lactate dehydrogenase
PCT = procalcitonin
SNF = Skilled nursing facility
HFNC = high flow nasal cannula
Respiratory symptoms were defined as hypoxia, SOB, cough
GI symptoms were defined as nausea, vomiting, diarrhea

Table 2: Microbiologic Data			
Micro Data	Antibiotics n=172	No antibiotics n=36	P-value
Microbiologically-confirmed infection	24 (14.0)	0 (0)	0.02
Respiratory cultures obtained	66 (38.3)	1 (2.7)	<0.01
No growth	53 [80.3]	1 [100]	>0.99
Positive	13 [19.7]	0 [0]	>0.99
Blood cx obtained	135 (78.5)	14 (38.9)	<0.01
No growth	123 [91.1]	14 [100]	0.61
Positive	12 [8.9]	0 [0]	0.61
Influenza PCR obtained	75 (43.6)	13 (36.1)	0.46
Strep pneumoniae urinary antigen obtained	82 (47.7)	6 (16.7)	<0.01
Positive	2 [2.4]	0 [0]	>0.99
Negative	80 [97.6]	6 [100]	>0.99
Legionella urinary antigen obtained	87 (50.5)	6 (16.7)	<0.01
MRSA nares obtained	73 (42.4)	2 (5.6)	<0.01
Positive	4 [5.5]	0 [0]	>0.99
Negative	69 [94.5]	2 [100]	>0.99

Table 3: Risk factors associated with receiving more than 1 course of antibiotics						
	Univariate Analysis			Multivariate analysis		
Variable	1 antibiotic course n= 120 (%)	> 1 antibiotic course n=52 (%)	P value	Odds ratio	95% CI	P-value
Length of stay, days (median, IQR)	8 (4-15)	25 (18.75-37)	<0.01	1.01	0.99-1.04	0.32
ICU admission	51 (42.5)	49 (94.2)	<0.01	1.96	0.38-11.10	0.42
ICU length of stay, days (median, IQR)	5 (3-9)	18 (12-26)	<0.01	1.14	1.06-1.26	<0.01
Mechanical ventilation	25 (20.8)	45 (86.5)	<0.01	1.83	0.46-7.61	0.39
More than 1 COVID-19 related admission	7 (5.8)	1 (1.9)	0.44			
Microbiologically-confirmed infection	4 (3.3)	20 (38.5)	<0.01	3.31	0.86-15.26	0.10
Duration of initial course of antibiotics, days (median, IQR)	4 (2-5.25)	5 (4-7)	<0.01			
Time from SARS-CoV-2 collect date to initial antibiotic start, days (median, IQR)	1 (0-5)	2 (0-2)	0.51			
28-day mortality	33 (27.5)	14 (38.9)	>0.99			

()= Percentages unless otherwise specified

333   **Figures**

Figure 2: Indications for Antibiotic Prescribing

CAP= community acquired pneumonia
HAP= hospital acquired pneumonia
VAP= ventilator associated pneumonia
BSI = blood stream infection

338