Prediction and control of COVID-19 infection based on a hybrid intelligent model

Gengpei Zhang 1, Xiongding Liu 2 *

1 China Three Gorges University, Yichang, Hubei, China, 2 School of Automation Science and Engineering, South China University of Technology, Guangzhou, Guangdong, China

* 201771338@yangtzeu.edu.cn

Abstract

The coronavirus (COVID-19) is a highly infectious disease that emerged in the late December 2019 in Wuhan, China, and it has caused a worldwide outbreak, which represents a major threat to global health. It is important to design prediction research and control strategies to crush its exploding. In this study, a hybrid intelligent model is proposed to simulate the spreading dynamics of COVID-19. First, considering the control measures, such as government investment, media publicity, medical treatment and law enforcement. The infection rates are optimized by genetic algorithm (GA), then a modified susceptible-infected-quarantined-recovered (SIQR) model is proposed, then the long short-term memory (LSTM) is imbedded into the SIQR model to design the hybrid intelligent model to further optimize other parameters of the system model to obtain the optimal predictive model and control measures. This study provide a reliable model to predict cases of infection and death ，and reasonable suggestion to control COVID-19.

Introduction

In the past six months, Chinese people have been on a high level of containment due to the outbreak of the coronavirus 2019 throughout China [1]. However, due to the convenient of global transportation, the virus has rapidly spread to all corners of the world. The coronavirus was first discovered in Wuhan, at the early stage, due to people lack of knowledge about the virus and the scarcity of medical resources, people were not aware of the virus. Relatively little knows about the disease, people neglect to control it in its early stages, until the earliest official announcement that the disease can be spread from person to person on January 20, 2020 [2]. Then, the government took a series of measures to prevent the spread of the disease on January 21, like Wuhan lockdown. Subsequently, some cities closured, with cities across the country sealing off.

41	Governments took many measures by closing down public places,
42	broadcasting propaganda, isolating people in their own homes, etc.,
43	which leads to the increased awareness of self-protection for people.
44	The epidemic has been successful control in China, but now the
45	world is not optimistic, with over 40 million people now cumulatively
46	diagnosed worldwide. The cumulative number of deaths is over 1.2
47	million, especially in countries such as the United States, Brazil,
48	Russia, and India, the pandemic is still quite severe today [3,4].
49	Therefore, each countries and states should adopt prevention and
50	control strategies. Currently, it is very important to establish and
51	analyze disease-spreading model to predict disease development
52	trends in order to control and prevent the spread of the COVID-19.
53	Due to the outbreak of disease, vaccines and medicine treatments
54	for the disease are still being researched, non-drug treatment
55	becomes the main strategy to slow down the spread of the disease.
56	The purpose of most of these prediction and control measures is to
57	reduce the probability of infection spreading during direct or indirect
58	contact [5]. For example, paying attention to personal hygiene and
59	wearing a mask, keeping social distancing and closing some public
60	space such as schools and workplaces, in order to decrease the
61	opportunities of propagation from person to person. Other measures
62	like cutting off the way of transmission of diseases, such as
63	disinfection in public places, improving the level of sanitation in public
64	places and so on. All of these measures can alleviate the propagation
65	of the epidemic [6-9]. How to quantify the impact of various
66	prevention and control strategies on disease spreading is of great
67	significance to guide disease control.
68	A common method utilizes mathematical modelling to describe
69	the spreading dynamics of infectious diseases, like Ebola, SARS.
70	This can accurately describe the spread of disease among individuals
71	in theoretical framework to guide the development of the prevention
72	and control measures [10-13]. There are many researchers studied
73	epidemic spreading have proposed some epidemic model and
74	obtained some meaningful results. The classical epidemic models
75	like susceptible-infected (SI), susceptible-infected-recovered (SIR),
76	susceptible-exposed-infected-recovered (SEIR),
77	susceptible-infected-quarantined-recovered-susceptible (SIQRS)
78	[14-19]. Most of these models are applicable to describe disease
79	spreading with a long incubation period, such as COVID-19. Based
80	on data-driven, a modified SEIR model was proposed to analysis and
81	prediction COVID-19 spread [20]. Considering the effect of
82	quarantine, B. K. Mishra et.al proposed three quarantine models to
83	analyze COVID-19 spreading [21]. Other recent studies, some

researchers have considered the effects of the basic reproductive
number, international conveyance, and some stochastic based
regression models to prediction and control the COVID-19 disease
[22,23]. However, traditional epidemic spreading models consider
that all infected individual have the same infection rate, and the
prediction of disease development trend has certain limitations.
Although data-driven disease spreading models can accurately
describe infection rates, the impact of government prevention and
control measures on infection rates has not been quantitatively in
detail. These measures, such as the laws, medical supplies, media
coverage and investment, can reduce the spread of the disease. It is
necessary to rationally arrange the optimal prevention and control
strategies with limited resources to minimize the death rate.

To solve this problem, the GA and ANN hybrid method is
proposed to optimize epidemic dynamics model and predict the
COVID-19 spreading [24-27]. Genetic algorithm is an adaptive global
optimization search algorithm formed by simulating the genetic and
evolutionary process of biological species in the natural environment
[28]. It uses the viewpoint of biogenetics and realizes the
improvement of individual adaptability through the mechanism of
natural selection, heredity and variation. Artificial intelligence (AI) is
considered one of the most successful achievements of computer
science, simulating the behavior of the human brain in data analysis.
One of the AI branches is the artificial neural network (ANN). This
information processing system, by a simulating strategy like
communication between brain neurons, has become a tool for
analyzing complex and real systems [29]. In recent years, ANN
models have been developed to overcome the difficulties presented
by health issues.

This article focuses on how to quantify the impact of the
government's prevention and control measures on the infection rate,
then to obtain the optimal model of disease spreading and the most
effective prevention and control strategy. Based on the proposed
SIQR model, the hybrid artificial neural network (ANN) model
embedded the genetic algorithm for predicting the COVID-19 in this
article, and it introduces the important prediction and control
strategies led by the government as well as the massive support
participation from the public into the prediction calculation process.
Furthermore, this article simulates the development of the epidemic
based on the proposed hybrid prediction model and predicts the trend
of the epidemic. Due to the fluctuation of virus detection capability,
the epidemic data of China shows a sudden change on February 12
and 13, the data consistency is reduced. As a developing country,

127	Brazil shows the epidemic data with periodicity and consistency, and
128	then this article takes the Brazil data as analytic target. The
129	simulation results based on the epidemic data of Brazil show that the
130	proposed hybrid model could provide a basis for estimating the law of
131	virus spread, and achieve accurate and robust performance.
132	Moreover, the prediction results of our hybrid ANN-GN model is
133	agree with the actual epidemic development trend, which
134	demonstrates that the openness, transparency, and efficiency of data
135	releasing are very important for establishing a modern epidemic
136	prevention system.
137	The remainder of this article is organized as follows. Section "A
138	hybrid model of COVID-19 spreading dynamics" introduces the
139	framework of the proposed hybrid epidemic spreading model. Section
140	"Methods" explains the method of GA and ANN to predict epidemic
141	spreading. Section "Simulation results and discussion" provides the
142	simulation results based on the epidemic data of Brazil and gives
143	some discussion. The conclusions are provided at last.
144	**A hybrid model of COVID-19 spreading dynamics**
145	In this section, we establish a mathematical model on COVID- 19
146	based on some assumptions. There are some literatures proposed
147	models mainly based on real clinical data, predict and control
148	measures. In this paper, we modified the previous model proposed in
149	[18], and then extend the model structure by designing different
150	infection rate. The model network diagram and the interaction
151	individual components demonstrated in Fig 1. Every individual in
152	network can only exist one of four independent states, namely,
153	susceptible, infected, quarantined and recovered. For simply, it can
154	be denoted by S, I, Q, and R, respectively. Each link represents the
155	transformation relationship between nodes. Here, infected individuals
156	include symptomatic infected individuals and asymptomatic infected
157	individuals. Susceptible individual is infected with probability m (M1,
158	M2, M3 and M4) if it is connected to an infected individual. Infective
159	individuals are quarantined with probability α . In the process of
160	isolation, the asymptomatic infected individuals turn into symptomatic
161	infected individuals with the probability ω . Quarantined individuals
162	are treated with drugs that move into the recovered individuals with
163	probability β . Some recovered individuals will relapse into infection
164	due to their weakened immunity with probability γ . The probability of
165	death during isolation is λ . Here, set a switch of city lockdown by the
166	death rate, infection probability m is different based on the city
167	situation (lockdown or not). The lockdown infection probability m is
168	much lower for the strict government regulation.

According to the epidemic spreading, the dynamic equations can be written as follows:

$$\begin{cases} \dfrac{dS_k(t)}{dt} = -mS_k(t) \\[2mm] \dfrac{dI_k(t)}{dt} = mS_k(t) + \gamma R_k(t) - \alpha I_k(t) \\[2mm] \dfrac{dQ_k(t)}{dt} = \alpha I_k(t) - \beta Q_k(t) - \lambda Q_k(t) \\[2mm] \dfrac{dR_k(t)}{dt} = \beta Q_k(t) - \gamma R_k(t) \end{cases} \tag{1}$$

Using the normalization condition, the probability of death can be obtained:

$$D = \frac{\lambda \alpha m \gamma}{\alpha \gamma (\lambda + m) + m\gamma ((\lambda + \beta) + \alpha \beta m} \tag{2}$$

In this paper, the proportion of death is a very important parameter. It not only depicts the fatality rate of the disease in the process of transmission, but also reflects the effectiveness of prevention and control measures. For example, the money invested by the government can improve the medical level and effectively cut off the transmission route of the virus. Increasing publicity and awareness of prevention and control will also reduce the risk of infection. Law enforcement can also affect the spread of diseases, such as city closures and home quarantine. This paper argues that due to the strict control and isolation measures taken by the government during the epidemic, the infected cases cannot infect susceptible people after quarantine. However, asymptomatic infected persons also have some ability to transmit, and it is difficult to detect. Therefore, there is a certain relationship between the number of newly infected cases on Day t and the number of infected cases in the past k days. Moreover, the infection rate of patients is closely related to the time of infection. Since government measures can inhibit the spread of the disease, the infection rate of newly infected cases may vary from time to time on t day over the past k days. Further analyzing this difference and assigning different weights to different measures, we quantified the contribution of different measures to the infection rate at time t in newly infected cases. Then, the weighted accumulation was used to estimate the infection rate so as to establish the relevant epidemic air defense modeling.

In addition, in order to study the relationship between the prediction and control measures with infection rate of the SIQR epidemic spreading model, the method of GA and LSTM are used to optimize the spreading model. This paper considers the relationship

between the rate of disease transmission and the measures taken by the government against the disease. The main factors include government investment, medical level, media publicity and law enforcement. Firstly, genetic algorithm is used to estimate the infection rate of the model, taking the acceptable mortality rate as fitness function and taking it as a basis for city closure parameters. At the same time, GA is further used to obtain the optimal means of government control by taking the minimum mortality rate as fitness function. The mutation law of GA is based on the interaction of four government measures. Furthermore, the infection rate bias and mortality bias were estimated through the LSTM network, the number of infected people was estimated by combining with the SIQR model, and the relevant parameters of the model were modified to obtain the best transmission model and predict the spread of the disease. By combining these two approaches, the optimal model of disease spreading and the optimal prevention and control strategies can be obtained, which can predict the number of infected and death cases based on the spreading model and development trend. The proposed framework is shown in Fig 2.

Methods

Epidemiological investigation and modeling are effective tools to study the spread of epidemic diseases and have played their due roles in the prevention and control of public health events with global impact, such as SARS, MERS and H1N1 influenza. Many researchers have also achieved some results in novel coronavirus research using epidemiology and modeling analysis. The study of J. Chan et al. [30] found the first evidence of human-machine transmission. Then, some researchers combined with the analysis of some early cases and gave the mean incubation period and mean infection cycle of novel coronavirus. Traditional models of disease transmission believe that the number of new infections is related to the number of infected and vulnerable people, but these models lack in-depth analysis of parameters in the process of disease transmission. In the process of disease transmission, the implementation of different control measures has a great impact on the prevention and control of disease and the suppression of disease transmission. For example, the government's publicity of disease prevention and control, the formulation of relevant laws and regulations, financial investment and other measures can affect the spread of disease. The study found significant differences in the rates of infection among people of different ages. The main purpose of this paper is to study the impact of government measures on the spread

248	of disease and to minimize the mortality rate, and to consider the
249	impact of different age groups on the spread of disease, finally, we
250	can obtain the optimized measures and the best performance model
251	to prediction, as showing in Fig 2.
252	
253	**Fig 2. The hybrid model for COVID-19 prediction.**
254	
255	Unlike the traditional SIQR epidemic spreading model, which
256	consists of a single infected rate to describe the probability of
257	infection. In this paper, we proposed a dynamic strategy to better
258	represent the real world and the new COVID-19 disease. It mainly
259	considers the impact of various measures taken by the government
260	on the control of disease spreading. This paper considers the impact
261	of government investment, media publicity, medical treatment and
262	law enforcement on the rate of disease spreading, and its parameters
263	are set as mi, i=1, 2, 3, 4, respectively. In addition, k_i is the weight of
264	each measure. Thus, the rate of disease spreading can be written as:

$$m = \sum_{i=1}^{4} k_i m_i \qquad (3)$$

266	The GA is mainly used to optimize parameters of the model, it can
267	be divided as two parts, and the first part describes that using the GA
268	to optimize the conditions of city closure. First initialization model, set
269	the initial value, with acceptable mortality as a condition of judgment.
270	According to data analysis and relevant government regulations, a
271	three-day mortality rate greater than 0.045 is defined here as an
272	unacceptable mortality rate. When this unacceptable mortality rate is
273	reached, will update and save the neural network parameters, as a
274	basis for the sealing city.

$$unacceptable\ mortality\ rate = \frac{\lambda \alpha m \gamma}{\alpha \gamma (\lambda + m) + m \gamma ((\lambda + \beta) + \alpha \beta m)} \geq 0.045 \quad (4)$$

276	At the same time, further to run the model, with the minimum
277	mortality as fitness function, further update neural network
278	parameters by genetic algorithm.

$$minimum\ mortality = \frac{\lambda \alpha m \gamma}{\alpha \gamma (\lambda + m) + m \gamma ((\lambda + \beta) + \alpha \beta m)} \qquad (5)$$

280	Using twice the genetic algorithm, the approximate optimized
281	neural network parameters, namely the control measures taken by
282	the government can be obtained. Further, the LSTM algorithm is
283	used to modify other parameters of the model, and the historical data
284	is compared with the data calculated by the model, such as the
285	number of infections and deaths, to determine whether the system
286	has reached the minimum error. The number of infections and deaths
287	are further predicted. The flow chart of COVID-19 prediction
288	algorithm is shown in Fig 3.

290 **Fig 3. The flow chart of COVID-19 prediction algorithm.**

292 New data on the daily increase in the number of infected cases

293 can be obtained. Using the same method, it is also possible to

294 process the number of new deaths per day.

295
$$\begin{cases} \Delta I(t) = I(t) - I(t-1) \\ \Delta D(t) = D(t) - D(t-1) \end{cases} \qquad (6)$$

Here, $I(t)$ is the cumulative number of infected cases in the

previous t-day, $I(t-1)$ is the cumulative number of infected cases in the

previous t-day, and ΔI is the new number of infected cases in the

first t-day. $D(t)$ is the cumulative number of deaths in the first t-day,

$D(t-1)$ is the cumulative number of deaths in the first t-day, and ΔD

is the additional number of deaths on the first t-day.

Fig 4. The LSTM method.

In order to get a more accurate model, the model parameters

need to be further processed. The GA has optimized the weight

parameters in the neural network, and other model parameters are

optimized through the optimized neural network model. Here, LSTM

neural network is used to optimize the model parameters. We

collected historical data from February 26 to October 13 in Brazil, to

train neural network. Here we discussed the model parameters

through two judgments. First, when the mortality rate of the model

reaches an unacceptable mortality rate, the neural network

parameters of the closed city are used to process the system model

parameters, and then the mortality rate and infection rate are further

compared with the actual data. By obtaining the minimum model error,

the optimized model parameters are finally obtained. Here, let the

actual infected rate be ΔI and the infected rate under the regression

exponential function be $\Delta I'$, and use the neural network to predict

the deviation between the actual mortality and the regressive

infection rate. Similarly, the same approach can be applied to

mortality and obtained ΔD and $\Delta D'$. Assuming that B= $\Delta I'$+ $\Delta D'$ is

the deviation characteristic of the prediction, the LSTM method can

be used to predict the model. The flow diagram of the prediction is

shown in Fig 4.

S_{R1} is the number of vulnerable infected persons on the first day,

I_{R1} is the number of real-time infected cases on the first day, I_{P1} is the

number of infected cases predicted on the first day, M_1 is the initial

model. Where S_{RX} is the number of vulnerable infected persons on

day X, I_{RX} is the number of real-time infected cases on day X, I_{PX} is

| 331 | the predicted number of infected cases on day X, and M_X is the model |
| 332 | on Day X. |

Simulation results and discussion

Based on GA in Fig 1, we computed the vulnerability coefficient k_i of four government measures. Without city lockdown, the infection probability m of government investment, media publicity, medical treatment and law enforcement are 0.174, 0.717, 0.021 and 0.853. With city lockdown, the infection probability m of government investment, media publicity, medical treatment and law enforcement are 0.085, 0.219, 0.107 and 0.349. There are two significant government measures media publicity and law enforcement, and the law enforcement is more important. Make a globally observation, each country invested huge amounts of money for COVID-19, and most medical workers tried their best to save lives. The difference of epidemic prevention results comes from the media publicity and law enforcement. Enhancing the influence of media can build a strong basis of epidemic prevention by offering the meaning of prevention measures. Considering the exception of inconsiderate and non-media audiences, the risk of disease spreading still exist. The efficiency of law enforcement is the crucial insurance to save the effort of medical workers.

We simulated the disease development and analyze the output of the model in 231 days with real epidemic data available from Brazil (February 26 to October 13). Fig 5a represents the number of predicted and confirmed per days, the model predicts infected per days of the last 17 days before October 13. Considering the lockdown from March 21 to July 31, the government realized the necessity of quarantine. From the data trend of infection confirmed per days over this period, the law enforcement shows insufficient efficiency. Supposing the government gave the best performance to control the disease, we performed the simulation about the development of infection and death per days. Fig 5b is the infected prediction with best performance of government; the possible infected number would much lower with effective government effort. With the comparison between Fig 5a and Fig 5b, it is obvious that there is a great impact on the prediction in the middle and later period due to the large changes in the data after government make best moves. Fig 5c is the daily-infected prediction error of model for last 17 days. Considering the enormousness of infection confirmed number, Fig 5c shows the accuracy of model prediction without best performance of government.

Fig 5. Brazil infected cases. a) Infection of predicted and confirmed per days, b) Infection prediction with best government performance, c) Infection prediction error.

To better understand the model performance, we simulated deaths in Brazil. Fig 6a represents the number of new death cases per day, this indicates that at the beginning of the epidemic, due to limited understanding of disease transmission and limited detection efforts, there will be omissions in disease statistics, leading to a similar increase trend with the infected cases. Meanwhile, Fig 6a represents the new predicted death cases per days of the last 17 days before October 13, which shows the difference with daily new death cases. Fig 6b is the daily death prediction with best performance of government. While the continuous improvement of detection methods and the continuous promotion of detection scope will be carried out, the government could strengthen the control of the disease, so the daily death cases would gradually decrease. Fig 6c is the daily death prediction error of model for last 17 days. Combine Fig 6a, we can see that the LSTM method have a good performance to predict COVID-19 disease. The method can predict the trend of the disease over a longer period with the performance of government.

Fig 6. Brazil death cases. a) Death of prediction and confirmed per days, b) Death prediction with best government performance, c) Death prediction error.

Conclusions

Based on the SIQR disease-spreading model, this paper seeks the best government performance from the four aspects by GA; and then proposed a hybrid prediction model with LSTM network. By analyzing Brazil data from February 26 to October 13, we analyzed new infected cases, new death cases per day. It is found that media publicity and law enforcement have more contribution to reduce transmission rate. With best government performance, the tread of COVID-19 in Brazil could make a difference. The prediction results of this model are highly consistent with the actual epidemic situation, which proves that the hybrid model proposed in this paper can efficiently analyze the transmission law and development trend of the virus.

References

1. Huang C, Wang Y, Li X, Ren L, Zhao J, et al. (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet 395: 497-506. https://doi.org/10.1016/S0140-6736(20)30183-5

415   2. Ye F, Xu S, Rong Z, Xu R, Liu X, et al. (2020) Delivery of infection
416     from asymptomatic carriers of COVID-19 in a familial cluster. Int J
417     Infect Dis 94: 133-138. https://doi.org/10.1016/j.ijid.2020.03.042

418   3. Crokidakis N (2020) Modeling the early evolution of the COVID-19
419     in   Brazil:   Results   from   a
420     Susceptible–Infectious–Quarantined–Recovered (SIQR) model. Int J
421     Mod   Phys   C:   2050135-2050144.
422     https://doi.org/10.1142/s0129183120501351

423   4. Bhatraju PK GB, Nichols M, Kim R, Jerome KR, Nalla AK,
424     Greninger AL, Pipavath S, Wurfel MM, Evans L, Kritek PA, West
425     TE, Luks A, Gerbino A, Dale CR, Goldman JD, O'Mahony S,
426     Mikacenic C. (2020) Covid-19 in Critically Ill Patients in the Seattle
427     Region - Case Series. New Engl J Med 382: 2012-2022.
428     https://doi.org/10.1056/NEJMoa2004500

429   5. Bai Y, Yao L, Wei T, Tian F, Jin D-Y, et al. (2020) Presumed
430     Asymptomatic Carrier Transmission of COVID-19. JAMA 323:
431     1406-1407. https://doi.org/10.1001/jama.2020.2565

432   6. Lai C-C, Liu YH, Wang C-Y, Wang Y-H, Hsueh S-C, et al. (2020)
433     Asymptomatic carrier state, acute respiratory disease, and
434     pneumonia due to severe acute respiratory syndrome coronavirus
435     2 (SARS-CoV-2): Facts and myths. J Microbiol Immunol 53: 404-412.
436     https://doi.org/10.1016/j.jmii.2020.02.012

437   7. Lan L XD, Ye G, Xia C, Wang S, Li Y, Xu H. (2020) Positive RT-PCR
438     Test Results in Patients Recovered From COVID-19. JAMA 323:
439     1502-1503. https://doi.org/10.1001/jama.2020.2783

440   8. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, et al. (2020) The
441     Incubation Period of Coronavirus Disease 2019 (COVID-19) From
442     Publicly Reported Confirmed Cases: Estimation and Application.
443     Ann Intern Med 172: 577-582. https://doi.org/10.7326/M20-0504

444   9. Linton N, Kobayashi T, Yang Y, Hayashi K, Akhmetzhanov A, et al.
445     (2020) Incubation Period and Other Epidemiological Characteristics
446     of 2019 Novel Coronavirus Infections with Right Truncation: A
447     Statistical Analysis of Publicly Available Case Data. J Clin Med 9:
448     538. https://doi.org/10.1101/2020.01.26.20018754

449   10. Zhou Y, Ma Z, Brauer F (2004) A discrete epidemic model for SARS
450     transmission and control in China. Math Comp Model 40:
451     1491-1506. https://doi.org/10.1016/j.mcm.2005.01.007

452   11. Poletto C, Pelat C, Levy-Bruhl D, Yazdanpanah Y, Boelle PY, et al.
453     (2014) Assessment of the Middle East respiratory syndrome
454     coronavirus (MERS-CoV) epidemic in the Middle East and risk of
455     international spread using a novel maximum likelihood analysis
456     approach.   Euro   SurvII   19:   20824.
457     https://doi.org/10.2807/1560-7917.es2014.19.23.20824

12. Zhang Q, Sun K, Chinazzi M, Pastore Y Piontti A, Dean NE, et al. (2017) Spread of Zika virus in the Americas. Proc Natl Acad Sci 114: 4334-4343. https://doi.org/10.1073/pnas.1620161114

13. Nishiura H (2011) Real-time forecasting of an epidemic using a discrete time stochastic model: a case study of pandemic influenza (H1N1-2009). Bio Med Eng OnLine 10: 15. https://doi.org/10.1186/1475-925x-10-15

14. Yang M, Chen G, Fu X (2011) A modified SIS model with an infective medium on complex networks and its global stability. Physica A 390: 2408-2413. https://doi.org/10.1016/j.physa.2011.02.007

15. Li T, Liu X, Wu J, Wan C, Guan Z-H, et al. (2016) An epidemic spreading model on adaptive scale-free networks with feedback mechanism. Physica A 450: 649-656. https://doi.org/10.1016/j.physa.2016.01.045

16. Daley DJ, Kendall DG (1964) EPIDEMICS AND RUMOURS. Nature 204: 1118. https://doi.org/10.1038/2041118a0

17. Zhu G, Fu X, Chen G (2012) Spreading dynamics and global stability of a generalized epidemic model on complex heterogeneous networks. Appl Math Model 36: 5808-5817. https://doi.org/10.1016/j.apm.2012.01.023

18. Li T, Wang Y, Guan Z-H (2014) Spreading dynamics of a SIQRS epidemic model on scale-free networks. Commun Nonlinear Sci Numer Simul 19: 686-692. https://doi.org/10.1016/j.cnsns.2013.07.010

19. Zhang J, Sun J (2014) Stability analysis of an SIS epidemic model with feedback mechanism on networks. Physica A 394: 24-32. https://doi.org/10.1016/j.physa.2013.09.058

20. Zhang Y, Yu X, Sun H, Tick GR, Wei W, et al. (2020) Applicability of time fractional derivative models for simulating the dynamics and mitigation scenarios of COVID-19. Chaos Soliton Fract 138: 109959. https://doi.org/10.1016/j.chaos.2020.109959

21. Mishra BK, Keshri AK, Rao YS, Mishra BK, Mahato B, et al. (2020) COVID-19 created chaos across the globe: Three novel quarantine epidemic models. Chaos Soliton Fract 138: 109928. https://doi.org/10.1016/j.chaos.2020.109928

22. Alkahtani BST, Alzaid SS (2020) A novel mathematics model of covid-19 with fractional derivative. Stability and numerical analysis. Chaos Soliton Fract 138: 110006. https://doi.org/10.1016/j.chaos.2020.110006

23. Mandal M, Jana S, Nandi SK, Khatua A, Adak S, et al. (2020) A model based study on the dynamics of COVID-19: Prediction and

control. Chaos, Solitons & Fractals 136: 109889. https://doi.org/10.1016/j.chaos.2020.109889

24. Car Z, Baressi Šegota S, Anđelić N, Lorencin I, Mrzljak V (2020) Modeling the Spread of COVID-19 Infection Using a Multilayer Perceptron. Comput Math Method M 2020: 5714714. https://doi.org/10.1155/2020/5714714

25. Arora P, Kumar H, Panigrahi BK (2020) Prediction and analysis of COVID-19 positive cases using deep learning models: A descriptive case study of India. Chaos Soliton Fract 139: 110017. https://doi.org/10.1016/j.chaos.2020.110017

26. Haghshenas SS, Pirouz B, Piro P, Na KS, Cho SE, et al. (2020) Prioritizing and Analyzing the Role of Climate and Urban Parameters in the Confirmed Cases of COVID-19 Based on Artificial Intelligence Applications. Int J Environ Res Public Health 17: 3730. https://doi.org/10.3390/ijerph17103730

27. Wu X, Hui H, Niu M, Li L, Wang L, et al. (2020) Deep learning-based multi-view fusion model for screening 2019 novel coronavirus pneumonia: A multicentre study. Eur J Radiol 128: 109041. https://doi.org/10.1016/j.ejrad.2020.109041

28. Miralles-Pechuán L, Jiménez F, Ponce H, Martínez-Villaseñor L (2020) A Deep Q-learning/genetic Algorithms Based Novel Methodology For Optimizing Covid-19 Pandemic Government Actions. arXiv e-prints: arXiv:2005.07656.

29. Wong Z, Zhou J, Zhang Q (2018) Artificial Intelligence for infectious disease Big Data Analytics. Infect Dis Health 24: 44-48. https://doi.org/10.1016/j.idh.2018.10.002

30. Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, et al. (2020) A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. Lancet 395: 514-523. https://doi.org/10.1016/S0140-6736(20)30154-9