Title: Large-scale population analysis of SARS-CoV-2 whole genome sequences reveals host-mediated viral evolution with emergence of mutations in the viral Spike protein associated with elevated mortality rates.

Authors

Carlos Farkas[1,2*], Andy Mella[3] and Jody J. Haigh[1,2*]

Affiliations

[1] Research Institute in Oncology and Hematology (RIOH), CancerCare Manitoba, Winnipeg, MB, Canada

[2] Department of Pharmacology and Therapeutics, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, MB, Canada

[3] Departamento de Física, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Blanco Encalada 2008, Santiago, Casilla 487-3, Chile.

*corresponding author(s) e-mail address(es)

Carlos.FarkasPool@umanitoba.ca, jody.haigh@umanitoba.ca

Abstract

Background

We aimed to further characterize and analyze in depth intra-host variation and founder variants of SARS-CoV-2 worldwide up until August 2020, by examining in excess of 94,000 SARS-CoV-2 viral sequences in order to understand SARS-CoV-2 variant evolution, how these variants arose and identify any increased mortality associated with these variants.

Methods and Findings

We combined worldwide sequencing data from GISAID and Sequence Read Archive (SRA) repositories and discovered SARS-CoV-2 hypermutation occurring in less than 2% of COVID19 patients, likely caused by host mechanisms involved APOBEC3G complexes and intra-host microdiversity. Most of this intra-host variation occurring in SARS-CoV-2 are predicted to change viral proteins with defined variant signatures, demonstrating that SARS-CoV-2 can be actively shaped by the host immune system to varying degrees. At the global population level, several SARS-CoV-2 proteins such as Nsp2, 3C-like proteinase, ORF3a and ORF8 are under active evolution, as evidenced by their increased $\pi N/\pi S$ ratios per geographical region. Importantly, two emergent variants: V1176F in co-occurrence with D614G mutation in the viral Spike protein, and S477N, located in the Receptor Binding Domain (RBD) of the Spike protein, are associated with high fatality rates and are increasingly spreading throughout the world. The S477N variant arose quickly in Australia and experimental data support that this variant increases Spike protein fitness and its binding to ACE2.

Conclusions

SARS-CoV-2 is evolving non-randomly, and human hosts shape emergent variants with positive fitness that can easily spread into the population. We propose that V1776F and S477N

44   variants occurring in the Spike protein are two novel mutations occurring in SARS-CoV-2 and

45   may pose significant public health concerns in the future.

Introduction

The novel SARS-CoV-2 coronavirus that causes COVID19 has surpassed 34 million infections worldwide within nine months of pandemic, resulting in more than one million deaths until September 2020 (https://coronavirus.jhu.edu/map.html) [1]. In-depth characterization of this virus is urgently needed to improve outbreak surveillance, vaccine development and for effective treatments now and in the immediate future. SARS-CoV-2 is a positive single-stranded RNA virus (+ssRNA) with a crown-like appearance observed by electron microscopy that is due the presence of the of spike glycoproteins on the lipid bilayer envelope [2, 3]. Another three transmembrane proteins are incorporated into the envelope: small envelope protein (E), matrix protein (M), and nucleocapsid protein (N) [4]. As seen with SARS-CoV-1, SARS-CoV-2 binds through its Spike glycoprotein to cell membrane-bound angiotensin-converting enzyme 2 (ACE2) for entry into host cells [5-8]. Advancements in COVID19 treatments have been recently developed including Remdesivir, a nucleoside analog that inhibits viral RNA-dependent RNA polymerase and is an effective treatment to reduce viral titers in rhesus macaques that is clinically approved for COVID19 treatment [9]. As well, either wild-type or catalytically inactive ACE2 has been demonstrated to block viral entry *in vitro*, and are proposed as promising treatments [10, 11]. A remaining question is how the human humoral immune response develops after SARS-CoV-2 infection. Studies in Iceland have shown that around 90% of infected patients develop antiviral antibodies that last up to four months [12], but it has also been suggested that around one third of the seropositive infections are asymptomatic and become antibody-negative early in the convalescence period [13]. Also, the unexpectedly low secondary infection risk reported for SARS-CoV-2 infection suggests innate immune responses are active in humans [14, 15]. To explore host– SARS-CoV-2 interactions at the genetic level it is useful to analyze viral sequencing results per individual and at the population level. Initiatives such as

GISAID (https://www.gisaid.org/) [16, 17] and the Sequence Read Archive (SRA,

https://www.ncbi.nlm.nih.gov/sra) have been storing SARS-CoV-2 sequencing datasets

worldwide from the beginning of the pandemic starting in January 2020, allowing researchers to

track fixed variants and follow viral evolution by geographical region. The unprecedented

amount of SARS-CoV-2 whole genome sequencing data can help to 1) characterize viral variants

that occur within a given host, 2) understand variant fixation in a given population and 3)

understand how the virus changes over time. In fact, the Spike protein mutation D614G global

transition that occurred very recently was discovered in this way and is associated with higher

viral titers and higher fatality rates [18, 19]. Thus, it is probable that more mutations are to be

discovered by tracking SARS-CoV-2 genomic changes globally. In this study we aimed to

characterize in depth intra-host variation and population-fixed variants worldwide up until the

beginning of August 2020 by using over 76,000 SARS-CoV-2 sequences and 17,500 sequencing

datasets from GISAID and SRA repositories, respectively. First, we found evidence for SARS-

CoV-2 hypermutation, occurring in less than 2% of COVID19 patients. This mechanism is

predicted to inactivate the virus and is likely caused by host mechanisms involved APOBEC3G

complexes and intra-host microdiversity, where G>T transversions and C>T transitions are

frequent signatures observed both in hypermutant and non-hypermutant samples. These results

suggest that SARS-CoV-2 is actively shaped by the host immune system to varying degrees.

From a population context, several SARS-CoV-2 proteins such as Nsp2, 3C-like proteinase,

ORF3a and ORF8 are under active evolution, evidenced by their increasing $\pi N/\pi S$ ratios.

Noteworthy, most of the population-fixed variants in SARS-CoV-2 are predicted to destabilize

viral proteins, as already reported for other RNA viruses. Of these variants, those occurring in

the ORF3a (Q57H), Nucleocapsid (I292T, RG203KR) and Spike protein (V1176F) have a

positive association with increased mortality ratios in populations from Saudi-Arabia and Brazil,

respectively. In particular, the V1176F variant co-occurs with the D614G mutation in the Spike

protein mutation in Brazil and arose independently in at least in three independent SARS-CoV-2

clades. This variant is predicted to stabilize the SARS-CoV-2 Spike trimmer complex and confer

flexibility to the stalk domain of the trimmer, potentially facilitating Spike binding properties to

ACE2. Also, this variant is associated with increased mortality ratios in Brazil and is

increasingly spreading throughout the world. Similarly, the emerging variant S477N occurring in

the Receptor Binding Domain, dramatically increase its frequency and became dominant in

Australia within two months. Experimental data support that S477N increase both fitness and

binding to ACE2 receptor, explaining its selection among other viruses in Australia. S477N also

is presently spreading across countries and is associated with higher fatalities throughout the

world. We propose that these variants are novel mutations occurring in SARS-CoV-2 and their

spread may pose serious concerns in public health in the future of the pandemic.

Methods

Data and Code Availability

76,553 FASTA genomes and associated sequencing metadata were downloaded from GISAID database from January 1, 2019 until August 3, 2020, specifying "human" as source host (https://www.gisaid.org/). The associated sequencing metadata including major variants per sample are available at Supplementary Table 1. Aggregated variants in VCF format for the latter genomes and associated consequence predictions are available here: https://usegalaxy.org/u/carlosfarkas/h/sars-cov-2-variants-gisaid-august-03-2020. 974 Brazilian FASTA sequences were downloaded from GISAID database from January 1, 2019 until September 25, 2020, specifying "human" as source host and "South America / Brazil" as location. These FASTA sequences and associated aggregated variants are available here: https://usegalaxy.org/u/carlosfarkas/h/brazil-genome-sequences-from-gisaid-sept25-2020.

FASTA sequences from GISAID genomes containing associated metadata until September 28, 2020, including the results from snpFreq program (containing Deceased-Released SNP associations) are available here: https://usegalaxy.org/u/carlosfarkas/h/gisaid-patient-metadata-sept28-2020. Acknowledgements to all laboratories/consortia involved in the generation of GISAID genomes used in this study are listed in Supplementary Table 2. 17,560 sequencing datasets were downloaded from Sequence Read Archive Repository (SRA, https://www.ncbi.nlm.nih.gov/sars-cov-2/) From December 1, 2019 until July 28, 2020. Associated sequencing run accessions, sequencing metadata and related BioProjects are listed in Supplementary Table 3. The code generated during this study to replicate most of the computational calculations performed in this manuscript is available at the following github repository: https://github.com/cfarkas/SARS-CoV-2-freebayes.

Next-generation sequencing and FASTA dataset processing

To process next generation sequencing datasets, we employed our pipeline (SARS-CoV-2_freebayes) consisting in a bash/UNIX script that pipes several programs in sequential order. Imputed list of SRA accessions is processed with sra-tools, https://github.com/ncbi/sra-tools), generating compressed FASTQ files per sequencing, automatically trimmed with fastp tool [20]. Minimap2 splice-aware aligner in preset mode -ax sr [21] align each trimmed fastq file against a provided reference genome (Wuhan-Hu-1, GenBank Accession: MN908947.3). The resulting BAM files were sorted and indexed by using Samtools [22]. Freebayes, as frequency-based pooled caller (-F 0.49) (https://github.com/ekg/freebayes) [23] perform variant calling on every sorted BAM file, obtaining major frequency viral variants per genome in VCF format,. Then, the Jacquard program (https://jacquard.readthedocs.io/en/v0.42/index.html) in python environment [24] merges every VCF file containing variants associated to each bam file into a single VCF file, containing aggregated variants from all genomes. Viral frequencies were recalculated in the merge VCF file by using several UNIX tools [25], in combination with vcflib (https://github.com/vcflib/vcflib). Variants per genome reported in the resulting file "logfile_variants_SRA_freebayes" were used to construct Figure 1B using GraphPad Prism 8 software (https://www.graphpad.com/scientific-software/prism/). GISAID FASTA genomes were processed in a similar manner. We preprocess a single GISAID genome collection with SeqKit [26] to decompose a single FASTA file into individual FASTA files, each file containing a single genome. The original genome collection from GISAID (merged.GISAID.fasta.gz) is available here: https://usegalaxy.org/u/carlosfarkas/h/sars-cov-2-variants-gisaid-august-03-2020). Then, Minimap2 aligner with preset -ax asm5 [21] align every FASTA genome against SARS-CoV-2 reference genome. Freebayes variant caller with --min-alternate-count 1 (C 1) option

(https://github.com/ekg/freebayes) perform variant calling on each BAM file, outputting variants

in VCF format. With these operations, major frequency viral variants in VCF format are obtained

from each FASTA genome. Then, variants are aggregated into a single VCF file, as described

with Jacquard. Figure 1A graph was constructed by using variants per genome, reported in the

output file "logfile_variants_GISAID_freebayes", inputted into the GraphPad Prism 8 software.

All these computational analyses are described here: https://github.com/cfarkas/SARS-CoV-2-

freebayes (case examples I and II, respectively).

Variant Visualization

The   Integrative   Genomics   Viewer   (IGV)   software

(http://software.broadinstitute.org/software/igv/home) was used to visualize next generation

sequencing alignments in bam format [27-29]. To visualize mayor viral frequency variants, the

variant frequency threshold was set at 0.49.

SnpEff annotation

Merged variants from GISAID genomes (n=76563) were annotated by using in a

repurposed version of SnpEff program, available in the Galaxy server [30-32]. The resulting

annotated VCF file was parsed by using conventional UNIX tools. Codon change chart related

from   Figure   2D   is   available   as   SnpEff   HTML   output   here:

https://usegalaxy.org/u/carlosfarkas/h/sars-cov-2-variants-gisaid-august-03-2020.   All   these

computational analyses are described here: https://github.com/cfarkas/SARS-CoV-2-freebayes

(case example III).

πN/πS calculation

We estimated nonsynonymous and synonymous nucleotide diversity (πN and πS,

respectively) in 1279, 6841, 46042, 17989, 1205 and 2924 GISAID FASTA genomes from

Africa, Asia, Europe, North America, South America, and Oceania, respectively. These genomes

were accessed and downloaded at August 3, 2020 from GISAID database and are available here:

https://usegalaxy.org/u/carlosfarkas/h/sars-cov-2-variants-gisaid-august-03-2020. The FASTA

files were processed from the alignment to the variant calling step as described in "GISAID

FASTA dataset processing section". All these computational analyses are described here:

https://github.com/cfarkas/SARS-CoV-2-freebayes (case example IV).

Microdiversity and low frequency viral variants.

We estimated nucleotide diversity in 397, 448 and 308 next generation sequencing (NGS)

samples from Australia, Spain, and USA populations, respectively by using aligned reads per

sample in BAM format against SARS-CoV-2 reference genome. These BAM files were inputted

in loop to InStrain program (https://instrain.readthedocs.io/en/latest/) [33], obtaining several

parameters such as analysis of coverage, microdiversity, SNV linkage, and sensitive SNP

detection, among others. As recommended by inStrain, we analyzed only sequencing samples

with sufficient breadth of coverage (>0.9), resulting in 397, 374 and 216 NGS samples from

Australia, Spain and from USA, respectively. The list of the NGS samples in the three

populations, including the referred calculations are detailed in the spreadsheet

inStrain_results.xlsx, available here: https://github.com/cfarkas/SARS-CoV-2-freebayes. We

correlated in each country the number of variants with viral frequency > 5% against the

nucleotide diversity (π) by using Spearman correlation. Spearman's correlation coefficients (r)

and confident p-values (P, to discard random sampling) were calculated in GraphPad Prism 8.

The significance thresholds were as follows: P<0.05 *, P<0.01 **, P<0.001 ***, P<0.0001 ****, P>0.05 ns. All these computational analyses are described here: https://github.com/cfarkas/SARS-CoV-2-freebayes (case example V).

SNP-mortality associations

We downloaded 7634 genomes with associated metadata from GISAID until September 28, 2020 and we grouped the genomes from released/deceased patients per country (India, Saudi Arabia, USA, and Brazil, respectively). Then, we parsed genomes and associated metadata by country (in particular, deceased and released cases) by using a combination of standard UNIX tools, vcflib (https://github.com/vcflib/vcflib) and BEDOPS [34]. After these steps, we uploaded to the Galaxy server (https://usegalaxy.org/) the resulting output per country (Deceased-Released.subset file) [31, 35] and we performed Fisher's exact test to identified variants with a significant difference in the viral frequencies between the groups (snpFreq program, https://rdrr.io/github/lvclark/SNPfreq/). P values from Fisher's exact test were converted with to negative logarithm in base 10 by using R version 3.6.3 (https://www.r-project.org/). The latter values per variant were used to construct graph from Figures 3E and 3H, respectively by using GraphPad Prism 8 software. All computational steps required for these analyses are available here: https://github.com/cfarkas/SARS-CoV-2-freebayes (case example VI).

Phylogenetic Tree Construction

We downloaded 393 SAR2-CoV-2 GISAID genomes containing variant V1176F until September 25, 2020 from GISAID database (https://www.gisaid.org/). We filtered countries with at least two sequences per country, leading 358 sequences encompassing five country/regions (Brazil, Scotland, USA, Australia, and Gibraltar, respectively). MAFFT multiple sequence

alignment program version 7.271 [36, 37] was used to align FASTA sequences against Wuhan-

Wu-1 reference genome (GenBank Accession: MN908947.3) by using the --auto --thread -1 --

keeplength –addfragments flags. Fasttree version 2.1 [38] was used to infer an approximately-

maximum-likelihood phylogenetic tree from the aligned sequences in fasta format, by using

heuristic neighbor-joining clustering method [38] and the Jukes-Cantor model of evolution [39].

Visualization and editing of the phylogenetic tree was performed by using Interactive Tree of

Life server (iTOL), collapsing all clades whose average branch length distance was below

0.0002 [40, 41].

Free energy estimation calculations

SARS-CoV-2 protein models for nsp2, nsp3, nsp4, 3C-Like proteinase, nsp6, nsp7, nsp8,

RNA-dependent RNA polymerase, Helicase, Spike protein trimmer, ORF3a, ORF6, ORF8 and

Nucleocapsid,   respectively   were   accessed   and   downloaded   from   I-TASSER   server

(https://zhanglab.ccmb.med.umich.edu/COVID-19/) on June 20, 2020, in Protein Data Bank

(PDB) format. These models were generated by the C-I-TASSER pipeline [42-45]. We

calculated the free energy of Gibbs upon variant changes ($\Delta\Delta G$; $\Delta G_{wild\text{-}type} - \Delta G_{variant}$, in

kcal/mol) from all missense variants listed in Figure 3A using SARS-CoV-2 protein structures as

inputs for the Foldx5 program [46, 47]. We repaired every PDB by using the following

command: foldx --command=RepairPDB --pdb=name-of-protein.pdb --ionStrength=0.05 --

pH=7 --water=CRYSTAL --vdwDesign=2 --out-pdb=true --pdbHydrogens=false. Then, we

modelled the variant and calculated the free energy upon aminoacid changes as follows: foldx --

command=BuildModel   --pdb=name-of-protein.pdb   --mutant-file=individual_list.txt   --

ionStrength=0.05   --pH=7   --water=CRYSTAL   --vdwDesign=2   --out-pdb=true   --

pdbHydrogens=false --numberOfRuns=30, where individual_list.txt contain the aminoacid

change (as example, for a serine/asparagine change occurring in the three chains of the spike

protein trimmer: SA477N,SB477N,SC477N;). We classified the energetic effects as follows (in

kcal/mol): highly stabilizing (< −1.84), stabilizing (−1.84 to −0.92), slightly stabilizing (−0.92 to

−0.46), neutral (−0.46 to +0.46), slightly destabilizing (+0.46 to +0.92), destabilizing (+0.92 to

+1.84), and highly destabilizing (> +1.84) [48]. Additionally, the effect of D614G/V1176F

variants on protein–protein interaction energy in the full Spike protein trimmer and the effect of

S477N variant in the Receptor Binding Domain (RBD) complexed with ACE2 dimer were

assessed by submitting the referred structures on mCSM-PPI2 server

(http://biosig.unimelb.edu.au/mcsm_ppi2/) [49].

Molecular dynamics simulations

We computed molecular dynamics simulations of the wild-type (Valine at position 1176)

and 1176F (Phenylalanine at position 1176) stalk domain trimmers from the spike protein

(aminoacids 1130-1273). The full Spike protein trimmer was obtained from I-TASSER and the

variant V1176F was modelled by using Foldx5, as previously described in the Free energy

estimation calculations section (--command=BuildModel, first outputted model). The wild-type

and F1176 variant trimmers were subjected to molecular dynamics by using GROMACS/2020.3

version, in gpu mode (http://manual.gromacs.org/documentation/) [50, 51] in the supercomputer

infrastructure LEFTRARU NLHPC (ECM-02), allocating one node with total 44 cores (logical)

and one compatible GPU (NVIDIA Tesla V100-PCIE-16GB). The trajectories were visualized

by using VMD 1.9.3 [52]. As example, for a given pdb (molecula_1.pdb), the commands used to

perform the molecular dynamics are the following:

srun -p general gmx pdb2gmx -f molecula_1.pdb -o molecula_2.gro -water spce

srun -p general gmx editconf -f molecula_2.gro -o molecula_3.gro -c -d 1.0 -bt cubic

srun -p general gmx solvate -cp molecula_3.gro -cs spc216.gro -o molecula_4.gro -p topol.top

srun -p general gmx grompp -f ions.mdp -c molecula_4.gro -p topol.top -o ions.tpr

srun -p general gmx genion -s ions.tpr -o molecula_5.gro -p topol.top -pname NA -nname CL -

neutral

gmx grompp -f 1.mdp -c molecula_5.gro -p topol.top -o em.tpr

gmx mdrun -nt 20 -nb gpu -deffnm em   #EM

gmx grompp -f 2.mdp -c em.gro -r em.gro -p topol.top -o nvt.tpr

gmx mdrun -nt 20 -nb gpu -deffnm nvt   # NPT

gmx grompp -f 3.mdp -c nvt.gro -r nvt.gro -t nvt.cpt -p topol.top -o npt.tpr

gmx mdrun -nt 20 -nb gpu -deffnm npt   # NVT

gmx grompp -f 4.mdp -c npt.gro -t npt.cpt -p topol.top -o md_0_1.tpr

gmx mdrun -nt 20 -nb gpu -deffnm md_0_1 # MD

Where commands starting with "srun" were executed directly in the cluster and the remaining

steps   were   submitted   via   SLURM   workload   manager

(https://slurm.schedmd.com/documentation.html). We choose forcefield OPLS-AA/L all-atom

force field (2001 aminoacid dihedrals) for step one, and SOLVENT for step five (choice 13,

SOL). Atom clashed in the system were minimized by the steepest descent method [53], until

potential energy were below 1000 kJ/(mol*nm). We considered a cutoff of 1.0 nm for non-

bonded interactions under periodic boundary conditions (PBC). NVT ensemble (constant

Number of particles, Volume, and Temperature) was performed setting no pressure coupling and

modified Berendsen thermostat at 300K, respectively. The NPT ensemble was used to keep the

constant pressure at 1 bar, using the Parrinello-Rahman barostat and temperature at 300 K,

using the modified Berendsen thermostat, respectively. Long-range electrostatic forces were

considered using the Particle Mesh Ewald for long-range electrostatics method [54]. Both

equilibrations were performed for 5000 picoseconds (5 nanoseconds). The total energy,

temperature, pressure and the of the stalk domain trimmers were used to corroborate both system

equilibrations. After these steps, production dynamics were carried out for 20 nanoseconds, by

using the leap-frog algorithm with an integration step of 2 femtoseconds, as motion setting.

Bonds were fixed using the P-LINCS method, with constrained H-bonds [55, 56]. Root mean

square deviation (RMSD) and radius of gyration (Rg) were obtained with the following

commands, respectively:

gmx rms -s md.tpr -f md_0_1.xtc -o rmsd.xvg -tu ns   # Choose twice 4 ("Backbone")

gmx gyrate -s md.tpr -f md_0_1.xtc -o gyrate.xvg   # Group 1 (Protein)

The xvg file records per picosecond were used to plot graphs from Figure 3C, on GraphPad

Prism 8 software. PDB, solvated molecules (.gro) and correspondent compressed gromacs

trajectories (with or without periodic border conditions) are available here :

https://usegalaxy.org/u/carlosfarkas/h/sars-cov-2-proteins-and-trayectories.

Protein Visualization and Rendering

The Spike protein trimmer image related to Figure 3C was rendered online with the

EzMol server (http://www.sbg.bio.ic.ac.uk/ezmol/) [57].

Statistical analysis

All statistical analyses were carried out by using GraphPad Prism 8 software

(https://www.graphpad.com/scientific-software/prism/). Mann-Whitney test was employed to

test data following non normal distribution. The significance thresholds were the following:

$P<0.05$ *, $P<0.01$ **, $P<0.001$ ***, $P>0.05$ ns. We interpreted Spearman nonparametric

correlations analyses as follows: perfect correlation (1), the two variables tend to increase or

decrease together (0 to 1), The two variables do not vary together at all (0), One variable

increases as the other decreases (-1 to 0), and perfect inverse correlations (-1). We computed an

approximate P value because in all correlations we employed more than 17 pair values. The

significance thresholds were the following: $P<0.05$ *, $P<0.01$ **, $P<0.001$ ***, $P<0.0001$ ****,

$P>0.05$ ns. We employed robust regression and outlier removal (ROUT) method [58] to remove

outliers from stacks of data, with a strict false discovery ratio (Q=1%). SNP-mortality

correlations were assessed by using the snpFreq program

(https://rdrr.io/github/lvclark/SNPfreq/), implemented in the R language (https://www.r-

project.org/). We employed the Fisher's exact test to identified variants with a significant

difference in the viral frequencies between deceased-released groups. We used as a significance

threshold a false discovery rate (q-value) of 0.005.

Results

Worldwide Intra-host variation in SARS-CoV-2 genomes reveal microdiversity and hypermutation likely elicited by APOBEC3G/ADAR complexes.

To trace intra-host viral variation worldwide, we downloaded and analyzed 76,553 SARS-CoV-2 genome sequences available in the GISAID database up until August 3, 2020 (**Supplementary Table 1,** see Acknowledgements in **Supplementary Table 2**). We also downloaded and analyzed 17,560 next-generation sequencing datasets from Sequence Read Archive (SRA) available until July 28, 2020 (**Supplementary Table 3**). SARS-CoV-2 genomes from GISAID accounted for the presence of major viral frequency variants (via a consensus calling approach) and the next-generation sequencing datasets (NGS) also allowed us to analyze intra-host microdiversity given the depth of sequencing. By analyzing the occurrence of major viral alleles per SARS-CoV-2 genome, both sources consistently demonstrate on average 7-8 viral variants with major alleles per genome (viral frequency > 0.5) (see "mean" in **Figure 1A** and **1B**, for SRA and GISAID datasets, respectively), demonstrating that our variant calling pipeline is reliable to call major viral alleles from FASTA and NGS datasets, respectively (see https://github.com/cfarkas/SARS-CoV-2-freebayes). The distribution from both sources also identified outliers with more than 16 viral sequence variants per genome including some samples harboring more than 100 variants per genome, greatly surpassing the average (2% and 0.3% in SRA and GIDAID sequencing datasets, see "outliers" in **Figure 1A and 1B**, respectively, Q=1% Grubbs's test). Integrative genomics viewer (IGV) snapshots of outlier samples from Spain, USA and Australian sequencing datasets clearly show hypermutability to varying degrees (see samples with black arrows, **Figure 1C**). Australian outlier samples represent an extreme case of

hypermutability (see **Figure 1C**, bottom). 16,307 aggregated variants from SRA datasets reflect

most recurrent single nucleotide substitutions occurring in all genomes from SRA repository are

enriched by the C>U (C>T) transitions and G>T (G>U) transversions, changes already reported

for SARS-CoV-2 and MERS-CoV genomes [59] and is likely elicited by APOBEC deaminases,

as already reported [60]. (**Figure 1D**, left). The latter observation is also consistent for genomes

containing outlier number of variants from SRA (**Figure 1D**, right). Most of the nucleotide

substitutions harboring outlier samples from Figure 1C correspond to missense/nonsense variants

rather than silent variants (**Figure 1E**) and are enriched in C>T (C>U) changes as well,

consistent with latter observations (**Figure 1F**). Also, G>T (G>U) transversions are significantly

present, implying a different mechanism than APOBEC3G editing and likely exerted by ADAR

deaminase (see discussion). We estimated intra-host nucleotide diversity occurring in 397, 374

and 215 next generation sequencing samples from Australia, Spain and USA populations,

respectively by using aligned reads per sample against the SARS-CoV-2 reference genome

(Wuhan-Hu-1, GenBank Accession: MN908947.3). This calculation has been already validated

to capture intra-host viral microdiversity, overcoming sequencing errors [61]. In the three

populations, average nucleotide diversity positively correlates with the number of Single

Nucleotide Variants (SNVs) with viral frequencies over 5% (Spearman correlation, r values from

0.32-0.66, P<0.0001). The latter supports the existence of intra-host minor variants and therefore

SARS-CoV-2 quasi-species, coexisting within the same host [62-65] (**Figure 1G**). We

hypothesized the existence of hypermutants can be explained by APOBEC3G/ADAR-mediating

RNA editing at early stages of SARS-CoV-2 infection (C>U and A>G editing) accompanied by

intra-host diversity, i.e. low frequency variance and homoplasy (different viral lineages emerged

after the infection) that probably are maintained at low frequency due the virus error correction

machinery (**Figure 1H**). Overall, we propose that human hosts substantially contribute to shape

SARS-CoV-2 genetic diversity. Although microdiversity is probably one of the main sources of

SARS-CoV-2 evolution, this is accompanied by RNA-editing at different levels, with SARS-

CoV-2 RNA hypermutation as an extreme case of the latter.

Non-neutral codon changes actively shape evolution of several SARS-CoV-2 proteins

We next analyzed all intra-host major viral alleles occurring in SARS-CoV-2 genomes

worldwide, by using the GISAID consensus called variants. 23,269 aggregated variants from

GISAID demonstrate that overall, C>U and A>G edition are the predominant nucleotide changes

(**Figure 2A**), also present in GISAID samples with outlier number of variants per genome

(**Figure 2B**). These changes are consistent with nucleotide changes occurring in SRA sequencing

datasets (see **Figure 1D**) and with the APOBEC3G-mediating RNA editing mechanisms. To

deduce aminoacid changes as consequences of these nucleotide changes, we analyzed codon

changes occurring in the aggregated GISAID variants and we predicted its consequences by

using the SnpEff program [30]. Occurrences per variant type demonstrate that missense and

synonymous variant occurrences are more frequent compared to frameshift/nonsense variant

occurrences per genome (**Figure 2C**). Overall, missense/silent ratio of GISAID aggregated

variants is 1.82, revealing a greater diversity in missense variants as well (**Supplementary**

Figure 1). Consequently, codon change analysis demonstrates frequent non-neutral changes in

the second position of the codons ACA>ATA, ACT>ATT and GCT>GTT and leads to missense

variants Thr>Ile and Ala>Val, respectively. Also, non-neutral changes in the first position of

codons CTT>TTT and GTT>TTT are frequent, leading to Leu>Phe and Val>Phe changes,

respectively (**Figure 2D**). Almost all these changes can be explained by C>T (C>U) transition,

already reported for SARS-CoV-2 protein changes. To understand these changes at the global

population level, we calculated nucleotide diversity in nonsynonymous (πN) and synonymous

(πS) sites of every SARS-CoV-2 protein across six different populations (Asia, Oceania, Europe,

Africa, North America, and South America) by using GISAID genomes per geographical region.

A given protein is evolving by natural selection if the πN/πS ratio is over one. Conversely, if

πN/πS ratio is less than one, a given protein is considered to be undergoing purifying selection,

as previously described [66-68]. An excess of shared nonsynonymous changes (implying

positive natural selection) is present in non-structural proteins nsp2, nsp7, 3C-like proteinase,

ORF3a and ORF8, respectively (**Figure 2E**). Conversely, purifying selection is present in the

RNA-dependent RNA polymerase, Membrane protein (M), 3'-to 5'-exonuclease, non-structural

proteins nsp3 and nsp4, respectively. Remarkably, ORF6 and to a lesser extent the Spike protein

(S) of SARS-CoV-2 are actively evolving in South America.

V1176F variant occurring in the Spike protein is predicted to improve fitness of Spike

protein complex and is likely a novel SARS-CoV-2 mutation associated with increased

mortality

We analyzed SARS-CoV-2 fixed alleles shared with at least a 3% variant frequency (VF)

in one of the five referred to continental populations until August 3, 2020. The merge

encompassed 51 variants. Consistent with intra-host variation, more than half of these changes

lead to missense variants. Non-structural proteins nsp2, nsp3, nsp6 and nsp7 harbors Leu>Phe

and Thr>Ile as frequent changes as seen in SARS-CoV-2 codon changes frequencies (**Figure**

3A). Also, consistent with the increased observed πN/πS ratios per population, ORF3a and ORF8

display missense variants with high viral frequencies. Some variants are shared among

populations with high viral frequencies, such as D614G variant in Spike protein, P323L variant

in RNA-dependent RNA polymerase and RG203KR variant in Nucleocapsid. Of these, D614G

variant in the Spike protein fulfill the category of mutation since this variant is positively

associated with mortality and increases SARS-CoV-2 infectivity, as previously reported [18, 19,

69]. As D614G was a rapidly emergent variant, this suggests novel emergent variants can also be

mutations. Of importance, two more novel missense variants occurring in the Spike protein

(S477N and V1176F) are exclusively from Oceania and South America (**Figure 3A,** red

variants). To gain understanding of the effects of these variants, we calculated the Gibbs' free

energy associated with the variant changes ($\Delta\Delta G$; $\Delta G_{wild-type} - \Delta G_{variant}$) from all missense

variants listed in Figure 3A using SARS-CoV-2 protein structures generated by the C-I-TASSER

pipeline as inputs for the Foldx5 program [42-47]. While most of the changes are predicted to be

neutral or slightly stabilizing/destabilizing, there are more changes predicted to be energetically

unfavorable rather than favorable, such as occurring in the RNA-polymerase (A97V), ORF3a

(G251V) and Nucleocapsid (I292T). Conversely, other changes occurring in the non-structural

proteins nsp3 (T1198K), nsp4 (F308Y) and Spike protein (D614G, V1176F) are predicted to

stabilize these proteins (**Figure 3B**). The emergent V1176F variant, containing the recurrent

signature Val>Phe (**Figure 2D**) is located at the stalk domain of Spike protein, specifically at the

beginning of the heptad repeat 2 (HR2) domain [70]. In agreement with a recent report, the

D614G variant has a mildly stabilizing effect on protein stability but also alters protein dynamics

according to mCSM-PPI2 analysis, since the predicted affinity change of the spike protein

trimmer decreases [71]. Conversely, the V1176F variant is predicted to increase affinity of the

Spike protein trimmer (**Figure 3C, upper right**). Molecular dynamics simulations of the stalk

domain trimmer demonstrate larger amplitude motions, since Root-mean-square deviations

(RMSD) from wild-type stalk domain trimmer fluctuates in ~1 nm (10 Å) over 20 nanoseconds

of simulation. The V1176F variant increases this value after the same settings, increasing

motility of the domain trimmer (~1.4 nm, **Figure 3C middle right**). Also, the V1176F induces

compactness of Stalk domain trimmer in ~0.52 nm (5 Å) (**Figure 3C low right**) [72]. Thus, the

V1176F variant confers more flexibility to the Stalk trimmer domain. These observations agree

with a recent report demonstrating extensive flexibility of this domain composed by three hinges

in the pre-fusion model of the Spike protein, potentially necessary for enhanced binding

mechanics with ligands such ACE2 [73]. Of note, up until September 25, 2020 Brazil was the

country with the highest frequency of genomes containing the V1176F variant, that was

completely linked to the D614G mutation (**Figure 3D**). We also found four additional regions

that present increasing cumulative distributions of genomes containing V1176F, including

Scotland, USA, Australia, and Gibraltar that have occurred during global travel bans/restrictions

(**Figure 3E**). Indeed, phylogenetic tree analysis containing genomes from Figure 3E demonstrate

V1176F variant arose independently in clades G (containing S: D614G mutation), clade GH

(containing ORF3a: Q57H variant) and clade GR (containing N: RG203KR variant) (**Figure**

3F). The predominant clade GR contain genomes from Brazil, Scotland, and Gibraltar while

clade GH is represented exclusively of genomes from USA. Clade G is also present in Brazil and

Scotland. Overall, this analysis suggests community transmission spread of V1176F occurred

from independent sources rather than from a single source arising from travel. To correlate if this

variant has an associated phenotype, we downloaded 7634 genomes with associated metadata

from GISAID until September 28, 2020 and we grouped the genomes from released/deceased

patients per country/region. Then, we performed Fisher's exact test and identified variants with a

significant difference in the viral frequencies between the groups (snpFreq program, available in

the Galaxy server) [31, 35]. Interestingly, a novel variant in the spike protein (QD613HG),

leading to D614G mutation and variants occurring in the ORF3a (Q57H) and Nucleocapsid

(RG203KR), are positively correlated with increased mortality ratios in Saudi-Arabia (**Figure**

3H). In Brazil, the D614G mutation along with the V1176F emergent variant in the Spike protein

are also positively correlated with increased mortality ratios, including variants occurring in the

nsp7, RNA-dependent RNA polymerase, and two emergent variants in South America occurring

in the ORF6 and the nucleocapsid proteins (I33T and I292T variants, respectively, **Figure 3A,**

Figure 3G). The RG203KR variants occurring in the Nucleocapsid was also found in Brazil.

Four out seven of these variants occurring in Brazil (nsp7: L71F, S: V1176F, ORF6: I33T and N:

I292T) are also positively correlated with increased mortality when comparing deceased patients

versus Released + Hospitalized patients, suggesting these variants are robustly correlated with

increased mortality in Brazil (**Figure 3H**). Of note, the V1176F variant arose independently and

is not ligated to the I292T variant occurring in the Nucleocapsid (**Supplementary Table 4**).

Thus, among emergent SARS-CoV-2 variants in Brazil, we prioritize the V1176F variant for

further experimental study since it likely arose independently across SARS-CoV-2 clades in

different countries, is predicted to improve fitness of the Spike protein and correlates with

increased mortality ratios in Brazil.

Variant S447N occurring in the Spike protein is a novel mutation that increases Spike-

ACE2 binding and is associated with higher worldwide fatality rates.

We next addressed whether the referred variants are associated with higher mortality

ratios in Brazil and Saudi-Arabia, as well as other emergent variants in the Spike protein, are

correlated with higher fatality ratios worldwide until September 28, 2020. Among the refereed

variants, the I292T variant occurring in the nucleocapsid is associated with higher fatality rates

across several countries ($p < 0.033$, Spearman correlation, **Figure 4A, Supplementary Table 5**).

Eleven variants occurring in the Spike protein were present in more than four countries and out

of them, variants A222V, S477N and E780Q are positively correlated with increased fatality

ratios. These findings are consistent with previous reports for the D614G mutation that is also

positively correlated with higher fatality ratios (**Figure 4B**) [19]. The V1176F variant was not

found to be correlated with increased fatality ratios across the world, probably due to highly

unbiased viral frequencies across the world. Up until August 3, 2020, the S477N spike variant

that emerged in Oceania (Australia) had a low viral population frequency (~ 3.5%, **Figure 3A**)

along with the A222V variant being absent or maintained at a very low frequency. However, a

dramatic increase of cumulative genomes was observed between the months June and July 2020

for both variants, not seen for the I292T variant occurring in the nucleocapsid (**Figure 4C**). The

rapid emergence of the S477N and A222V variants correlates with the recent second wave of

COVID19 that has occurred in Australia and the United Kingdom (**Figure 4D,** left and right,

respectively). Nevertheless, this rapid increase in the population frequencies of these variants

could be due to the founder effect usually seen in outbreaks, and not due increased fitness of

SARS-CoV-2 (see countries in Figure 4B, A222V and S477N, respectively) [69]. To discard the

former founder effect, we examined GISAID clade frequency of both variants, demonstrating

variant A222V arose independently in all major SARS-CoV-2 clades [74] and variant S477N

arose in clades G, GH and GR, respectively (**Figure 4E**, upper and lower, respectively). Variant

A222V occurred in the N-terminal domain of the Spike protein (light brown domain, **Figure 4F**,

left) and variant S477N occur in the Receptor Binding Domain (RBD) of the Spike protein

(purple domain, **Figure 4F**, left). This variant is located near the interface between ACE2 and

the RBD, the latter is expected to cause enhanced binding of the RBD to the ACE2 human

receptor (**Figure 4F**, right). We replaced serine for asparagine in position 477 in the RBD

(complexed with ACE2 dimer) with foldx [46], and we calculated the predicted binding energy

upon this change by using mCSM-PPI2 server [49]. The change is predicted to add one more

polar interaction when asparagine is present in the RBD, increasing the affinity between RBD

and ACE2 (**Figure 4G**). We examined the deep mutational scanning of amino acid changes in

the RBD performed by Starr et al [75] in a high throughput yeast-surface-display system for

measuring expression of folded RBD protein and its binding to ACE2 [76]. Among fourteen

variants occurring in the RBD, only variant S477N increased both expression of the RBD, a

parameter positively correlated with folding (**Figure 4H**) and its binding to ACE2, respectively

(**Figure 4I**). The combination of these two properties can lead to the generation of a more

infectious virus, explaining to a large extent the dramatic increase of the S477N variant in

Australia.

In conclusion, we have obtained evidence that the S477N variant is a gain of function mutation

occurring in the Spike protein, that is positively correlated with increased fatality rates and is

becoming dominant as its increases, as was also recently observed for the D614G mutation that

became dominant throughout the world [18].

Discussion

In this study we aimed to analyze over 94,000 SARS-CoV-2 sequences deposited between GISAID and SRA databases within the first eight months of this pandemic (up until the beginning of August 2020). We characterized the existence of intra-host viral hypermutation that results in an excessive number of variants per genome in less than 2% of SARS-CoV-2 sequences (**Figure 1A** and **1B**, respectively). This phenomenon was already described for HIV-1 virus *in vivo*, demonstrating that HIV-1 reverse transcriptase contributed only to 2% of mutations, and the majority was caused by host cytidine deaminases of the A3 family mediated editing [77]. Here, we present evidence that enzymatic RNA editing in combination with microdiversity contributes to SARS-CoV-2 diversity at a global level, leading to more than 23,000 major viral frequency variants within 76,000 GISAID genomes. In SARS-CoV-2 genomes, the C>T (C>U) transversion is substantially present both in hypermutants and non-hypermutant samples, suggesting APOBEC3G mediated RNA editing involvement, as previously reported in smaller sample sizes [59, 60]. Also, the A>G transversion is also present overall but not in hypermutant genomes (Figures **1D**, **2A** and **2B**, respectively), implying an active role of ADAR deaminases during SARS-CoV-2 infection [78]. We argue ADAR-mediating RNA editing is not the main enzyme involved in the hypermutation mechanism, but rather APOBEC3G deaminase complexes. Also, we have observed substantial G>T transversions in SARS-CoV-2 genomes. This transversion has been already reported for other RNA viruses such as Maize streak virus [79] and is has been linked with the formation of 8-oxoguanine, known to be the most common cause of spontaneous G>T (G>U) transversions in RNA [80]. Recently, it has been reported that tissue damage from neutrophils induces oxidative stress upon SARS-CoV-2 infection [81], implying that reactive oxidative species (ROS) mediated mutagenesis is likely the mechanism that cause this transversion. The hypermutated SARS-CoV-

2 variant signature often contains nonsense variants that are predicted to inactivate several

SARS-CoV-2 proteins, probably leading to an efficient mechanism of lethal mutagenesis to

control viral spread (**Figure 1E**). In addition, other signatures are present at the intra-host level,

implying microdiversity as another potential source of this variation (**Figure 2A** and **2B**,

respectively). It is possible that these combined forces produce quasi-species of viruses with

enough sequence diversity that may influence viral pathogenesis and drive SARS-CoV-2

evolution, sometimes also leading to viral extinction [82, 83]. Overall, we propose that human

host are major drivers of SARS-CoV-2 diversity rather than the virus itself, evidenced by the

levels of intra-host variation and hypermutation at different degrees, both fueled by enzymatic

RNA-editing mechanisms. We support the latter with the observed non-random signatures of

nucleotide changes in these mechanisms, and the presence of SARS-CoV-2 error-correction

machinery, not seen in other RNA viruses. Although we found a significant amount of intra-host

variation in SARS-CoV-2, neutral evolutionary theory predicts most of these variants as having

no or neutral effects [84]. Nevertheless, positive, or negative selection can occur within-

populations on viral variants, by conferring advantageous properties to viruses that ultimate lead

to mutations. Here we have demonstrated positive natural selection of several SARS-CoV-2

proteins per population, occurring in nsp2, nsp7, 3C-like proteinase, ORF3a and ORF8 proteins

and we highlighted mutations in the Spike protein, evolving in South America (Brazil, V1176F)

and Oceania (Australia, S477N). To begin to understand how these variations may affect their

encoded proteins, we assessed their structural consequences in protein models that demonstrate

that these variants tend to cause more unfavorable than favorable energetic changes. This

phenomenon has been observed in the spectrum of variants occurring in the RBD of the Spike

protein, proving that most of the variants occurring in the RBD constraints its folding and

binding to ACE2 [75]. In the case of SARS-CoV-2, immunological pressure from host could

lead to this type of phenomena as well and is consistent with the observed number of intra-host

missense over synonymous variants, respectively. Conversely, we found missense variant

V1176F, occurring in the Spike protein, is predicted to be energetically favorable and confers

flexibility to the Stalk domain of the viral Spike protein trimmer, previously described to be

important for Spike protein flexibility and binding to ACE2 [73]. Phylogenetic analysis

demonstrated V1176F variant likely emerged in South America and arose independently in

several countries associated with the D614G mutation, suggesting that this variant is being

positively selected among others occurring in the Spike protein (**Figure 3F**). This variant is also

correlated with higher mortality ratios (**Figure 3G** and **3H**, respectively) and it is possible that it

increases the fitness of SARS-CoV-2 infection by conferring flexibility to the stalk domain of the

spike protein. The same conclusions applied to variant S477N, occurring in the RBD of the

Spike protein: is energetically favorable in RBD-ACE2 binding and favor expression of the

RBD, with the latter conclusions being experimentally supported. This mutation also arose

independently in a noticeably short period of time and become dominant in Australia within two

months (**Figure 4D**). In addition, the S477N variant is constantly spreading across European

countries (**Figure 4B**) and correlates with higher mortality ratios. These observations provide

strong evidence that the S477N variant is a novel gain of function Spike protein mutation, as has

recently been demonstrated for the D614G mutation [18]. We argue that the constant spread of

V1176F and S477N variants over the world ultimately may lead to a further significant concern

in public health, due to their association with higher mortality rates.

A remaining question is the association of higher fatality rates of the A222V variant occurring

in the N-terminal domain of the Spike protein, and the I292T variant occurring in the

Nucleocapsid, among others. It is possible that these variants can confer antigenic escape, since

recently, it has been registered that a reinfection case containing A222V and D614G mutations

has occurred [85]. Nucleocapsid variation has also been documented in the nucleoprotein of the

influenza virus [86] [87] and nucleocapsid of the hepatitis virus [88]. Both RNA viruses escape

cellular immunity by these mechanisms and could also be the case for SARS-CoV-2.

In summary, we have presented potential molecular mechanisms that help researchers to

understand variation diversity fueled natural selection in SARS-CoV-2. It is important to

continue to track emergent viral variants with the bioinformatics tools developed and highlighted

in this manuscript since the evidence presented here lead us to propose V1776F and S477N

variants are novel SARS-CoV-2 mutations, due to their positive correlations with increased

fatality ratios, as previously evidenced with D614G mutation occurring in the Spike protein.

Further conclusions concerning the effects of these variants on viral fitness and host mortality

will be accomplished by future structure-function based studies using viral Spike protein mutants

and studying effects on viral entry and *in vivo* rodent models expressing the human ACE2

receptor.

Acknowledgments

Powered@NLHPC: This research was partially supported by the supercomputing infrastructure of the NLHPC (ECM-02). This research was partially funded by research funding from the CIHR, Research Manitoba and the CancerCare MB Research Foundation.

Author Contributions

CF conceived of this study and performed all bioinformatics analysis and wrote the manuscript. AM performed mutant Spike protein analysis and assist in biophysical studies. JH assisted study design, data interpretation and manuscript writing.

Declaration of Interests

None declared

    Figure titles and legends

    Figure 1: Intra-host variation in SARS-CoV-2 genomes worldwide reveal microdiversity

    and hypermutation likely elicited by APOBEC3G complexes.

    A) Mayor viral frequency variants (via a consensus calling approach) for 75,563 SARS-CoV-2

    GISAID genomes, separated by non-outliers (n=76,310) and outliers (n=243, Q=1%, Grubbs's

    test). Mean and outlier number of variants are depicted at left. **B**) Same as A for next generation

    sequencing (NGS) datasets downloaded from SRA (n=17,500). **C**) IGV snapshots of outliers and

    non-outlier NGS samples from B. Outliers samples are depicted with black arrows, exceeding

    number of variants from non-outliers. Single nucleotide polymorphisms are depicted in red if

    nucleotide differs from the reference sequence in greater than 50% of quality weighted reads. **D**)

    Nucleotide change frequencies from 17,560 SRA NGS aggregated variants (left) and from 360

    aggregated outlier variants (right), both annotated with SnpEff program. Frequency boxes are

    colored from white to dark red as number of changes increases. **E**) Nucleotide substitution

    frequencies in each of the eight outlier samples indicated with black arrows in C) grouped by

    silent (green), missense (dark red) and nonsense (red). **F**) Transitions and transversions occurring

    in the latter samples, denoted with blue and red, respectively. Significance of comparisons were

    assessed with Mann-Whitney test (P<0.05 *, P<0.01 **, P<0.001 ***, P>0.05 ns). **G**)

    Correlation between Average nucleotide diversity (π) provided by inStrain program and SNV

    counts (VF>5%) for Spain (n=374, left), USA (n=215, middle) and Australian NGS samples

    (n=397, right). In the three countries, the two variables tend to increase together (see r values).

    The significance thresholds were the following: P<0.05 *, P<0.01 **, P<0.001 ***, P<0.0001

657   ****, P>0.05 ns. **H**) Proposed model of how APOBEC3G/ADAR complex can lead to

658   hypermutation of SARS-CoV-2 (C>U and A>G editing). The RNA editing can be accompanied

659   by intra-host diversity (low frequency variants) and homoplasy (different viral lineages emerged

660   after the infection), maintained at low frequency due the virus error correction machinery.

Figure 2: Non-neutral codon changes actively shape evolution of several SARS-CoV-2 proteins.

A) Nucleotide change frequencies from 76,553 aggregated GISAD genome variants annotated with SnpEff program. Frequency boxes are colored from white to dark red as number of changes increases. **B**) Same as A for 243 aggregated GISAID genomes, corresponding to GISAID outlier samples. Frequencies boxes are colored from white to dark red, as the number of changes increases. **C**) Missense, nonsense, frameshift, and synonymous number of occurrences in 76553 GISAID genomes. Significance of comparisons were assessed with Mann-Whitney test ($P<0.05$ *, $P<0.01$ **, $P<0.001$ ***, $P>0.05$ ns). **D**) Plot of change frequencies across 32 codons in SARS-CoV-2. Changes are grouped in six categories and colored from light to dark red, according to the number of changes. **E**) Nonsynonymous/synonymous nucleotide diversity calculated by SNPGenie program for each SARS-CoV-2 protein across six geographical regions (Asia, Oceania, Europe, Africa, North America, and South America, respectively). Ratios are grouped in seven categories and colored from white to dark red, according to the ratio numbers.

Figure 3: V1176F variant occurring in the Spike protein is predicted to improve fitness of Spike protein complex and is likely a novel SARS-CoV-2 mutation.

A) Heatmap of viral frequencies from 51 variants shared at least 5% frequency within populations. Variants position in SARS-CoV-2 reference genome (Wuhan-Hu-1, GenBank Accession: MN908947.3) are depicted at left. At right, amino acid changes are colored by protein. **B**) Gibbs Free Energy Calculation (ΔΔG) of missense variants depicted in A, obtained by Foldx5 program. ΔΔG (kcal/mol) was colored according to the energetic effect in SARS-CoV-2 structures depicted in the Y axis, as follows (in kcal/mol): highly stabilizing (< -1.84), stabilizing (-1.84 to -0.92), slightly stabilizing (-0.92 to -0.46), neutral (-0.46 to $+0.46$), slightly destabilizing ($+0.46$ to $+0.92$), destabilizing ($+0.92$ to $+1.84$), and highly destabilizing ($> +1.84$). **C**) (left) Rendered Spike protein trimmer structure with EzMol server using rainbow palette. D614G mutation and V1176F variant are depicted in red, including their sidechains. Volume is depicted in grey. The stalk domain of the Spike protein trimmer is depicted in dark blue. (Upper right) Predicted affinity change (kcal/mol) of Spike trimmer by mCSM-PPI2 server upon D614G and V1176F amino acid Spike protein changes. (middle right) RMSD values (in nanometers, nm) of 20 nanoseconds of simulation of the wild-type stalk domain trimmer (red) or the domain containing Phenylalanine in position 1176 (blue). (lower right) Radius of gyration values (in nanometers, nm) of the latter simulation. **D**) Occurrences per genome of the D614G mutation and V1176F variant alone or in combination in 974 Brazilian GISAID genomes until September 25, 2020. Complete linkage of V1176F variant with D614G mutation was found since non a single genome contains V1176F variant alone. **E**) Cumulative distributions of 358 genomes containing V1176F variant (G25088T) from January 2020 until September 28, 2020 in five geographical regions, respectively. **F**) Phylogenetic tree of the 358 genomes from E,

constructed by Neighbor-Joining method, and visualized with the iTOL server. The genomes

were colored by SARS-CoV-2 clades as follows: G (containing S: D614G mutation, in green),

GH (containing ORF3a: Q57H variant, in red) and GR (containing N: RG203KR variant, in

blue). Brazilian genomes belonging clade GR were depicted with small blue font and the other

genomes were highlighted with higher font, for visualization purposes. **G**) Mortality correlations

associated with SARS-CoV-2 variants in Released vs Deceased patients occurring in India, USA,

Saudi Arabia, and Brazil until September 25, 2020, respectively. The corrected p-values from

fisher exact test (q-values) were obtained from the snpFreq program, available in the Galaxy

server and plotted as the negative logarithm in base 10 of each q-value (significance: q-

value>0.005). Significant variants were depicted at the right of each bar. **H**) Same as G, but with

the comparison Released + Hospitalized vs Deceased patients in Brazil.

Figure 4: Variant S447N occurring in the Spike protein is a novel mutation that increase Spike-ACE2 binding and is associated with worldwide higher fatality rates.

A) Correlation analysis of COVID-19 fatality rates among 49 countries against variant frequencies occurring in various SARS-CoV-2 proteins: ORF3a (Q57H), , RNA-dependent RNA polymerase (P323L), nsp7 (L71F), N (RG203KR) and N (I292T), respectively. Spearman's correlation coefficients (r) were calculated, including confident p-values (P) to discard random sampling. The significance thresholds were the following: $P < 0.05$ *, $P < 0.01$ **, $P < 0.001$ ***, $P < 0.0001$ ****, $P > 0.05$ ns. China were discarded from these analyses due its large population size and consistently low number of COVID19 cases since early April 2020. Fatality rate data was obtained from John Hopkins coronavirus resource center (https://coronavirus.jhu.edu/map.html), accessed on September 28, 2020. **B**) Same analysis as A, for five emergent variants occurring in the Receptor Binding Domain (RBD) portion of the Spike protein (N439K-L455F), including variants L18F, A222V, D614G, E780Q and V1176F occurring in the Spike protein. **C**) Cumulative distributions over time of genomes containing variants I292T in the nucleocapsid and emergent variants A222V, S477N and E780Q in the spike protein, respectively. Cumulated genome numbers were plotted at the end of the indicated months. **D**) Frequency plots over time of cumulated genomes containing spike protein emerging variants A222V, S477N occurring in Australia (left) and United Kingdom (right), respectively. Cumulated frequencies were plotted at the end of the indicated months. **E**) Graph chart indicating the clade frequencies of variants A222V (Upper) and S477N (lower), reported from GIDAID database, until September 28, 2020. **F**) (Left) Spike protein Trimmer indicating positions of variants N477 and V222 (red letters). Variant N477 locates in the Receptor Binding Domain (RBD, residues 331-530, depicted in purple) and Variant V222 locates in the N-terminal

domain (residues 1-316, depicted in light brown). The Spike structure was shown as cartoon with

grey color, obtained with EzMol server. (Right) Magnification of the interaction between the

RBD (purple) with ACE2 dimer (blue and red structures, respectively) showing the contact

surfaces. Variant N477 and its surface is depicted in red. The image was rendered in the EzMol

server. **G**) Polar interactions of S477 (left, wild type) and N477 (right, mutant) depicted by

discontinued lines in the RBD. The polar amino acid interactions of S477 and N477, including

the predicted $\Delta\Delta G$ (affinity, kcal/mol) between the wild-type or mutant RBD with ACE2 dimer

were obtained by using the mCSM-PPI2 server. Positive values of $\Delta\Delta G$ indicate increasing

affinity between the RBD and ACE2 dimer. **H**) Expression fitness of fourteen emergent variants

occurring in the RBD domain of the spike protein (see Supplementary Figure 2), measured by

high throughput yeast-surface-display system. Expression measurements were plotted as the

difference in log-mean fluorescence intensity (MFI) relative to wild-type ($\Delta logMFI$ =

$logMFI_{variant}$ - $logMFI_{wild-type}$). Positive values (blue) indicate higher RBD expression,

therefore higher folding fitness. Expression data was obtained from here:

https://jbloomlab.github.io/SARS-CoV-2-RBD_DMS/. I) ACE2 binding fitness of the referred

RBD variants, measured by high throughput yeast-surface-display system. Binding

measurements were plotted as the difference in log10(KD, apparent) relative to wild-type

($\Delta log10(KD, apparent) = log10(KD, apparent)_{wild-type} - log10(KD,app)_{variant}$). Positive values

(blue) indicate higher affinity between RBD and ACE2. Binding data was obtained from here:

https://jbloomlab.github.io/SARS-CoV-2-RBD_DMS/.

References

1.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time (vol 20, pg 533, 2020). Lancet Infectious Diseases. 2020;20(9):E215-E. PubMed PMID: WOS:000566754000001.

2.   Ke Z, Oton J, Qu K, Cortese M, Zila V, McKeane L, et al. Structures and distributions of SARS-CoV-2 spike proteins on intact virions. Nature. 2020. Epub 2020/08/18. doi: 10.1038/s41586-020-2665-2. PubMed PMID: 32805734.

3.   Cui J, Li F, Shi ZL. Origin and evolution of pathogenic coronaviruses. Nat Rev Microbiol. 2019;17(3):181-92. Epub 2018/12/12. doi: 10.1038/s41579-018-0118-9. PubMed PMID: 30531947; PubMed Central PMCID: PMCPMC7097006.

4.   Wu A, Peng Y, Huang B, Ding X, Wang X, Niu P, et al. Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV) Originating in China. Cell Host Microbe. 2020;27(3):325-8. Epub 2020/02/09. doi: 10.1016/j.chom.2020.02.001. PubMed PMID: 32035028; PubMed Central PMCID: PMCPMC7154514.

5.   Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell. 2020;181(2):271-80 e8. Epub 2020/03/07. doi: 10.1016/j.cell.2020.02.052. PubMed PMID: 32142651; PubMed Central PMCID: PMCPMC7102627.

6.   Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, et al. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature. 2003;426(6965):450-4. Epub 2003/12/04. doi: 10.1038/nature02145. PubMed PMID: 14647384; PubMed Central PMCID: PMCPMC7095016.

7.   Crackower MA, Sarao R, Oudit GY, Yagil C, Kozieradzki I, Scanga SE, et al. Angiotensin-converting enzyme 2 is an essential regulator of heart function. Nature. 2002;417(6891):822-8. Epub 2002/06/21. doi: 10.1038/nature00786. PubMed PMID: 12075344.

8.   Ge XY, Li JL, Yang XL, Chmura AA, Zhu G, Epstein JH, et al. Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. Nature. 2013;503(7477):535-8. Epub 2013/11/01. doi: 10.1038/nature12711. PubMed PMID: 24172901; PubMed Central PMCID: PMCPMC5389864.

9.   Williamson BN, Feldmann F, Schwarz B, Meade-White K, Porter DP, Schulz J, et al. Clinical benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. Nature. 2020;585(7824):273-6. Epub 2020/06/10. doi: 10.1038/s41586-020-2423-5. PubMed PMID: 32516797; PubMed Central PMCID: PMCPMC7486271.

10.   Lei C, Qian K, Li T, Zhang S, Fu W, Ding M, et al. Neutralization of SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig. Nat Commun. 2020;11(1):2070. Epub 2020/04/26. doi: 10.1038/s41467-020-16048-4. PubMed PMID: 32332765; PubMed Central PMCID: PMCPMC7265355.

11.   Monteil V, Kwon H, Prado P, Hagelkruys A, Wimmer RA, Stahl M, et al. Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. Cell. 2020;181(4):905-13 e7. Epub 2020/04/26. doi: 10.1016/j.cell.2020.04.004. PubMed PMID: 32333836; PubMed Central PMCID: PMCPMC7181998.

12.   Gudbjartsson DF, Norddahl GL, Melsted P, Gunnarsdottir K, Holm H, Eythorsson E, et al. Humoral Immune Response to SARS-CoV-2 in Iceland. N Engl J Med. 2020. Epub 2020/09/02. doi: 10.1056/NEJMoa2026116. PubMed PMID: 32871063.

13.   Pollan M, Perez-Gomez B, Pastor-Barriuso R, Oteo J, Hernan MA, Perez-Olmeda M, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological

study. Lancet. 2020;396(10250):535-44. Epub 2020/07/10. doi: 10.1016/S0140-6736(20)31483-5. PubMed PMID: 32645347; PubMed Central PMCID: PMCPMC7336131.

14. Li W, Zhang B, Lu J, Liu S, Chang Z, Cao P, et al. The characteristics of household transmission of COVID-19. Clin Infect Dis. 2020. Epub 2020/04/18. doi: 10.1093/cid/ciaa450. PubMed PMID: 32301964; PubMed Central PMCID: PMCPMC7184465.

15. Covid-19 National Emergency Response Center E, Case Management Team KCfDC, Prevention. Coronavirus Disease-19: Summary of 2,370 Contact Investigations of the First 30 Cases in the Republic of Korea. Osong Public Health Res Perspect. 2020;11(2):81-4. Epub 2020/04/08. doi: 10.24171/j.phrp.2020.11.2.04. PubMed PMID: 32257773; PubMed Central PMCID: PMCPMC7104686.

16. Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to global health. Glob Chall. 2017;1(1):33-46. Epub 2017/01/10. doi: 10.1002/gch2.1018. PubMed PMID: 31565258; PubMed Central PMCID: PMCPMC6607375.

17. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from vision to reality. Euro Surveill. 2017;22(13). Epub 2017/04/07. doi: 10.2807/1560-7917.ES.2017.22.13.30494. PubMed PMID: 28382917; PubMed Central PMCID: PMCPMC5388101.

18. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell. 2020;182(4):812-27 e19. Epub 2020/07/23. doi: 10.1016/j.cell.2020.06.043. PubMed PMID: 32697968; PubMed Central PMCID: PMCPMC7332439.

19. Becerra-Flores M, Cardozo T. SARS-CoV-2 viral spike G614 mutation exhibits higher case fatality rate. Int J Clin Pract. 2020:e13525. Epub 2020/05/07. doi: 10.1111/ijcp.13525. PubMed PMID: 32374903; PubMed Central PMCID: PMCPMC7267315.

20. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34(17):i884-i90. Epub 2018/11/14. doi: 10.1093/bioinformatics/bty560. PubMed PMID: 30423086; PubMed Central PMCID: PMCPMC6129281.

21. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018;34(18):3094-100. Epub 2018/05/12. doi: 10.1093/bioinformatics/bty191. PubMed PMID: 29750242; PubMed Central PMCID: PMCPMC6137996.

22. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 2009/06/10. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID: PMCPMC2723002.

23. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing2012 July 01, 2012:[arXiv:1207.3907 p.]. Available from: https://ui.adsabs.harvard.edu/abs/2012arXiv1207.3907G.

24. Sanner MF. Python: a programming language for software integration and development. J Mol Graph Model. 1999;17(1):57-61. Epub 2000/02/08. PubMed PMID: 10660911.

25. Kernighan BW, Morgan SP. The UNIX Operating System: A Model for Software Design. Science. 1982;215(4534):779-83. Epub 1982/02/12. doi: 10.1126/science.215.4534.779. PubMed PMID: 17747840.

26. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. PLoS One. 2016;11(10):e0163962. Epub 2016/10/06. doi: 10.1371/journal.pone.0163962. PubMed PMID: 27706213; PubMed Central PMCID: PMCPMC5051824.

27. Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. Nature Biotechnology. 2011;29(1):24-6. doi: 10.1038/nbt.1754. PubMed PMID: WOS:000286048900013.

28. Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Briefings in Bioinformatics. 2013;14(2):178-92. doi: 10.1093/bib/bbs017. PubMed PMID: WOS:000316694700006.

29.   Robinson JT, Thorvaldsdottir H, Wenger AM, Zehir A, Mesirov JP. Variant Review with the
Integrative Genomics Viewer. Cancer Research. 2017;77(21):E31-E4. doi: 10.1158/0008-5472.Can-17-
0337. PubMed PMID: WOS:000414248300009.

30.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, et al. A program for annotating and
predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. Fly (Austin). 2012;6(2):80-92. Epub 2012/06/26. doi:
10.4161/fly.19695. PubMed PMID: 22728672; PubMed Central PMCID: PMCPMC3679285.

31.   Giardine B, Riemer C, Hardison RC, Burhans R, Elnitski L, Shah P, et al. Galaxy: a platform for
interactive large-scale genome analysis. Genome Res. 2005;15(10):1451-5. Epub 2005/09/20. doi:
10.1101/gr.4086505. PubMed PMID: 16169926; PubMed Central PMCID: PMCPMC1240089.

32.   Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, et al. The Galaxy platform for
accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res.
2018;46(W1):W537-W44. Epub 2018/05/24. doi: 10.1093/nar/gky379. PubMed PMID: 29790989;
PubMed Central PMCID: PMCPMC6030816.

33.   Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek B, Morowitz MJ, Banfield JF. InStrain
enables population genomic analysis from metagenomic data and rigorous detection of identical
microbial strains. bioRxiv. 2020:2020.01.22.915579. doi: 10.1101/2020.01.22.915579.

34.   Neph S, Kuehn MS, Reynolds AP, Haugen E, Thurman RE, Johnson AK, et al. BEDOPS: high-
performance genomic feature operations. Bioinformatics. 2012;28(14):1919-20. Epub 2012/05/12. doi:
10.1093/bioinformatics/bts277. PubMed PMID: 22576172; PubMed Central PMCID: PMCPMC3389768.

35.   Afgan E, Baker D, van den Beek M, Blankenberg D, Bouvier D, Cech M, et al. The Galaxy platform
for accessible, reproducible and collaborative biomedical analyses: 2016 update. Nucleic Acids Res.
2016;44(W1):W3-W10. Epub 2016/05/04. doi: 10.1093/nar/gkw343. PubMed PMID: 27137889; PubMed
Central PMCID: PMCPMC4987906.

36.   Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30(14):3059-66. Epub 2002/07/24.
doi: 10.1093/nar/gkf436. PubMed PMID: 12136088; PubMed Central PMCID: PMCPMC135756.

37.   Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. Brief Bioinform. 2019;20(4):1160-6. Epub 2017/10/03.
doi: 10.1093/bib/bbx108. PubMed PMID: 28968734; PubMed Central PMCID: PMCPMC6781576.

38.   Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for large
alignments. PLoS One. 2010;5(3):e9490. Epub 2010/03/13. doi: 10.1371/journal.pone.0009490. PubMed
PMID: 20224823; PubMed Central PMCID: PMCPMC2835736.

39.   Steel MA, Fu YX. Classifying and counting linear phylogenetic invariants for the Jukes-Cantor
model. J Comput Biol. 1995;2(1):39-47. Epub 1995/01/01. doi: 10.1089/cmb.1995.2.39. PubMed PMID:
7497119.

40.   Letunic I, Bork P. Interactive Tree Of Life (iTOL): an online tool for phylogenetic tree display and
annotation. Bioinformatics. 2007;23(1):127-8. Epub 2006/10/20. doi: 10.1093/bioinformatics/btl529.
PubMed PMID: 17050570.

41.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res. 2019;47(W1):W256-W9. Epub 2019/04/02. doi: 10.1093/nar/gkz239. PubMed PMID:
30931475; PubMed Central PMCID: PMCPMC6602468.

42.   Zhang C, Mortuza SM, He B, Wang Y, Zhang Y. Template-based and free modeling of I-TASSER
and QUARK pipelines using predicted contact maps in CASP12. Proteins. 2018;86 Suppl 1:136-51. Epub
2017/10/31. doi: 10.1002/prot.25414. PubMed PMID: 29082551; PubMed Central PMCID:
PMCPMC5911180.

43.   He B, Mortuza SM, Wang Y, Shen HB, Zhang Y. NeBcon: protein contact map prediction using
neural network training coupled with naive Bayes classifiers. Bioinformatics. 2017;33(15):2296-306.

Epub 2017/04/04. doi: 10.1093/bioinformatics/btx164. PubMed PMID: 28369334; PubMed Central PMCID: PMCPMC5860114.

44.   Li Y, Hu J, Zhang C, Yu DJ, Zhang Y. ResPRE: high-accuracy protein contact prediction by coupling precision matrix with deep residual neural networks. Bioinformatics. 2019;35(22):4647-55. Epub 2019/05/10. doi: 10.1093/bioinformatics/btz291. PubMed PMID: 31070716; PubMed Central PMCID: PMCPMC6853658.

45.   Zheng W, Li Y, Zhang C, Pearce R, Mortuza SM, Zhang Y. Deep-learning contact-map guided protein structure prediction in CASP13. Proteins. 2019;87(12):1149-64. Epub 2019/08/01. doi: 10.1002/prot.25792. PubMed PMID: 31365149; PubMed Central PMCID: PMCPMC6851476.

46.   Delgado J, Radusky LG, Cianferoni D, Serrano L. FoldX 5.0: working with RNA, small molecules and a new graphical interface. Bioinformatics. 2019;35(20):4168-9. Epub 2019/03/16. doi: 10.1093/bioinformatics/btz184. PubMed PMID: 30874800; PubMed Central PMCID: PMCPMC6792092.

47.   Schymkowitz J, Borg J, Stricher F, Nys R, Rousseau F, Serrano L. The FoldX web server: an online force field. Nucleic Acids Res. 2005;33(Web Server issue):W382-8. Epub 2005/06/28. doi: 10.1093/nar/gki387. PubMed PMID: 15980494; PubMed Central PMCID: PMCPMC1160148.

48.   Studer RA, Christin PA, Williams MA, Orengo CA. Stability-activity tradeoffs constrain the adaptive evolution of RubisCO. Proc Natl Acad Sci U S A. 2014;111(6):2223-8. Epub 2014/01/29. doi: 10.1073/pnas.1310811111. PubMed PMID: 24469821; PubMed Central PMCID: PMCPMC3926066.

49.   Rodrigues CHM, Myung Y, Pires DEV, Ascher DB. mCSM-PPI2: predicting the effects of mutations on protein-protein interactions. Nucleic Acids Research. 2019;47(W1):W338-W44. doi: 10.1093/nar/gkz383. PubMed PMID: WOS:000475901600049.

50.   Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJ. GROMACS: fast, flexible, and free. J Comput Chem. 2005;26(16):1701-18. Epub 2005/10/08. doi: 10.1002/jcc.20291. PubMed PMID: 16211538.

51.   Kutzner C, Pall S, Fechner M, Esztermann A, de Groot BL, Grubmuller H. Best bang for your buck: GPU nodes for GROMACS biomolecular simulations. J Comput Chem. 2015;36(26):1990-2008. Epub 2015/08/05. doi: 10.1002/jcc.24030. PubMed PMID: 26238484; PubMed Central PMCID: PMCPMC5042102.

52.   Eargle J, Wright D, Luthey-Schulten Z. Multiple Alignment of protein structures and sequences for VMD. Bioinformatics. 2006;22(4):504-6. doi: 10.1093/bioinformatics/bti825. PubMed PMID: WOS:000235277300020.

53.   Vrejoiu C, Costescu A, Burlacu L. Generalized Calculus of Coefficients of Asymptotic Series Using Steepest Descents Method. Stud Cercet Fiz. 1978;30(4):329-45. PubMed PMID: WOS:A1978FF51800001.

54.   Deserno M, Holm C. How to mesh up Ewald sums. II. An accurate error estimate for the particle-particle-particle-mesh algorithm. Journal of Chemical Physics. 1998;109(18):7694-701. doi: Doi 10.1063/1.477415. PubMed PMID: WOS:000076663100005.

55.   Hess B. P-LINCS: A parallel linear constraint solver for molecular simulation. Journal of Chemical Theory and Computation. 2008;4(1):116-22. doi: 10.1021/ct700200b. PubMed PMID: WOS:000252198200012.

56.   Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. LINCS: A linear constraint solver for molecular simulations. Journal of Computational Chemistry. 1997;18(12):1463-72. doi: Doi 10.1002/(Sici)1096-987x(199709)18:12<1463::Aid-Jcc4>3.3.Co;2-L. PubMed PMID: WOS:A1997XT81100004.

57.   Reynolds CR, Islam SA, Sternberg MJE. EzMol: A Web Server Wizard for the Rapid Visualization and Image Production of Protein and Nucleic Acid Structures. Journal of Molecular Biology. 2018;430(15):2244-8. doi: 10.1016/j.jmb.2018.01.013. PubMed PMID: WOS:000437815400009.

58.   Motulsky HJ, Brown RE. Detecting outliers when fitting data with nonlinear regression - a new method based on robust nonlinear regression and the false discovery rate. Bmc Bioinformatics. 2006;7. doi: Artn 123

10.1186/1471-2105-7-123. PubMed PMID: WOS:000237981600001.

59.   Simmonds P. Rampant C-->U Hypermutation in the Genomes of SARS-CoV-2 and Other
Coronaviruses: Causes and Consequences for Their Short- and Long-Term Evolutionary Trajectories.
mSphere. 2020;5(3). Epub 2020/06/26. doi: 10.1128/mSphere.00408-20. PubMed PMID: 32581081;
PubMed Central PMCID: PMCPMC7316492.

60.   Di Giorgio S, Martignano F, Torcia MG, Mattiuz G, Conticello SG. Evidence for host-dependent
RNA editing in the transcriptome of SARS-CoV-2. Sci Adv. 2020;6(25):eabb5813. Epub 2020/07/01. doi:
10.1126/sciadv.abb5813. PubMed PMID: 32596474; PubMed Central PMCID: PMCPMC7299625.

61.   Nelson CW, Hughes AL. Within-host nucleotide diversity of virus populations: Insights from next-
generation sequencing. Infect Genet Evol. 2015;30:1-7. doi: 10.1016/j.meegid.2014.11.026. PubMed
PMID: WOS:000350525400001.

62.   Miralles R, Gerrish PJ, Moya A, Elena SF. Clonal interference and the evolution of RNA viruses.
Science. 1999;285(5434):1745-7. Epub 1999/09/11. doi: 10.1126/science.285.5434.1745. PubMed
PMID: 10481012.

63.   Wright CF, Morelli MJ, Thebaud G, Knowles NJ, Herzyk P, Paton DJ, et al. Beyond the consensus:
dissecting within-host viral population diversity of foot-and-mouth disease virus by using next-
generation genome sequencing. J Virol. 2011;85(5):2266-75. Epub 2010/12/17. doi: 10.1128/JVI.01396-
10. PubMed PMID: 21159860; PubMed Central PMCID: PMCPMC3067773.

64.   Domingo E, Sheldon J, Perales C. Viral quasispecies evolution. Microbiol Mol Biol Rev.
2012;76(2):159-216. Epub 2012/06/13. doi: 10.1128/MMBR.05023-11. PubMed PMID: 22688811;
PubMed Central PMCID: PMCPMC3372249.

65.   Ni M, Chen C, Qian J, Xiao HX, Shi WF, Luo Y, et al. Intra-host dynamics of Ebola virus during
2014. Nat Microbiol. 2016;1(11):16151. Epub 2016/10/27. doi: 10.1038/nmicrobiol.2016.151. PubMed
PMID: 27595345.

66.   Nelson CW, Hughes AL. Within-host nucleotide diversity of virus populations: insights from next-
generation sequencing. Infect Genet Evol. 2015;30:1-7. Epub 2014/12/08. doi:
10.1016/j.meegid.2014.11.026. PubMed PMID: 25481279; PubMed Central PMCID: PMCPMC4316684.

67.   Nelson CW, Moncla LH, Hughes AL. SNPGenie: estimating evolutionary parameters to detect
natural selection using pooled next-generation sequencing data. Bioinformatics. 2015;31(22):3709-11.
Epub 2015/08/01. doi: 10.1093/bioinformatics/btv449. PubMed PMID: 26227143; PubMed Central
PMCID: PMCPMC4757956.

68.   Nei M, Gojobori T. Simple methods for estimating the numbers of synonymous and
nonsynonymous nucleotide substitutions. Mol Biol Evol. 1986;3(5):418-26. Epub 1986/09/01. doi:
10.1093/oxfordjournals.molbev.a040410. PubMed PMID: 3444411.

69.   Day T, Gandon S, Lion S, Otto SP. On the evolutionary epidemiology of SARS-CoV-2. Curr Biol.
2020;30(15):R849-R57. Epub 2020/08/05. doi: 10.1016/j.cub.2020.06.031. PubMed PMID: 32750338;
PubMed Central PMCID: PMCPMC7287426.

70.   Woo H, Park SJ, Choi YK, Park T, Tanveer M, Cao YW, et al. Developing a Fully Glycosylated Full-
Length SARS-CoV-2 Spike Protein Model in a Viral Membrane. J Phys Chem B. 2020;124(33):7128-37.
doi: 10.1021/acs.jpcb.0c04553. PubMed PMID: WOS:000563725900004.

71.   Portelli S, Olshansky M, Rodrigues CHM, D'Souza EN, Myung Y, Silk M, et al. Exploring the
structural distribution of genetic variation in SARS-CoV-2 with the COVID-3D online resource. Nat Genet.
2020. Epub 2020/09/11. doi: 10.1038/s41588-020-0693-3. PubMed PMID: 32908256.

72.   Lobanov M, Bogatyreva NS, Galzitskaia OV. [Radius of gyration is indicator of compactness of
protein structure]. Mol Biol (Mosk). 2008;42(4):701-6. Epub 2008/10/17. PubMed PMID: 18856071.

73.   Turonova B, Sikora M, Schurmann C, Hagen WJH, Welsch S, Blanc FEC, et al. In situ structural
analysis of SARS-CoV-2 spike reveals flexibility mediated by three hinges. Science. 2020. Epub
2020/08/21. doi: 10.1126/science.abd5223. PubMed PMID: 32817270.

74.   Mercatelli D, Giorgi FM. Geographic and Genomic Distribution of SARS-CoV-2 Mutations. Front
Microbiol. 2020;11:1800. Epub 2020/08/15. doi: 10.3389/fmicb.2020.01800. PubMed PMID: 32793182;
PubMed Central PMCID: PMCPMC7387429.

75.   Starr TN, Greaney AJ, Hilton SK, Ellis D, Crawford KHD, Dingens AS, et al. Deep Mutational
Scanning of SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. Cell.
2020;182(5):1295-310 e20. Epub 2020/08/26. doi: 10.1016/j.cell.2020.08.012. PubMed PMID:
32841599; PubMed Central PMCID: PMCPMC7418704.

76.   Adams RM, Mora T, Walczak AM, Kinney JB. Measuring the sequence-affinity landscape of
antibodies with massively parallel titration curves. Elife. 2016;5. Epub 2016/12/31. doi:
10.7554/eLife.23156. PubMed PMID: 28035901; PubMed Central PMCID: PMCPMC5268739.

77.   Cuevas JM, Geller R, Garijo R, Lopez-Aldeguer J, Sanjuan R. Extremely High Mutation Rate of
HIV-1 In Vivo. PLoS Biol. 2015;13(9):e1002251. Epub 2015/09/17. doi: 10.1371/journal.pbio.1002251.
PubMed PMID: 26375597; PubMed Central PMCID: PMCPMC4574155.

78.   Nishikura K. Functions and regulation of RNA editing by ADAR deaminases. Annu Rev Biochem.
2010;79:321-49. Epub 2010/03/03. doi: 10.1146/annurev-biochem-060208-105251. PubMed PMID:
20192758; PubMed Central PMCID: PMCPMC2953425.

79.   van der Walt E, Martin DP, Varsani A, Polston JE, Rybicki EP. Experimental observations of rapid
Maize streak virus evolution reveal a strand-specific nucleotide substitution bias. Virol J. 2008;5:104.
Epub 2008/09/26. doi: 10.1186/1743-422X-5-104. PubMed PMID: 18816368; PubMed Central PMCID:
PMCPMC2572610.

80.   Li Z, Wu J, Deleo CJ. RNA damage and surveillance under oxidative stress. IUBMB Life.
2006;58(10):581-8. Epub 2006/10/20. doi: 10.1080/15216540600946456. PubMed PMID: 17050375.

81.   Laforge M, Elbim C, Frere C, Hemadi M, Massaad C, Nuss P, et al. Tissue damage from
neutrophil-induced oxidative stress in COVID-19. Nat Rev Immunol. 2020;20(9):515-6. Epub 2020/07/31.
doi: 10.1038/s41577-020-0407-1. PubMed PMID: 32728221; PubMed Central PMCID: PMCPMC7388427.

82.   Pariente N, Sierra S, Lowenstein PR, Domingo E. Efficient virus extinction by combinations of a
mutagen and antiviral inhibitors. J Virol. 2001;75(20):9723-30. Epub 2001/09/18. doi:
10.1128/JVI.75.20.9723-9730.2001. PubMed PMID: 11559805; PubMed Central PMCID:
PMCPMC114544.

83.   Elena SF, Miralles R, Cuevas JM, Turner PE, Moya A. The two faces of mutation: extinction and
adaptation in RNA viruses. IUBMB Life. 2000;49(1):5-9. Epub 2000/04/20. doi: 10.1080/713803585.
PubMed PMID: 10772334.

84.   Gojobori T, Moriyama EN, Kimura M. Molecular Clock of Viral Evolution, and the Neutral Theory.
P Natl Acad Sci USA. 1990;87(24):10015-8. doi: DOI 10.1073/pnas.87.24.10015. PubMed PMID:
WOS:A1990EN15900105.

85.   To KK, Hung IF, Ip JD, Chu AW, Chan WM, Tam AR, et al. COVID-19 re-infection by a
phylogenetically distinct SARS-coronavirus-2 strain confirmed by whole genome sequencing. Clin Infect
Dis. 2020. Epub 2020/08/26. doi: 10.1093/cid/ciaa1275. PubMed PMID: 32840608; PubMed Central
PMCID: PMCPMC7499500.

86.   Phillips AM, Ponomarenko AI, Chen K, Ashenberg O, Miao J, McHugh SM, et al. Destabilized
adaptive influenza variants critical for innate immune system escape are potentiated by host
chaperones. PLoS Biol. 2018;16(9):e3000008. Epub 2018/09/18. doi: 10.1371/journal.pbio.3000008.
PubMed PMID: 30222731; PubMed Central PMCID: PMCPMC6160216.

87.   Voeten JT, Bestebroer TM, Nieuwkoop NJ, Fouchier RA, Osterhaus AD, Rimmelzwaan GF.
Antigenic drift in the influenza A virus (H3N2) nucleoprotein and escape from recognition by cytotoxic T
lymphocytes. J Virol. 2000;74(15):6800-7. Epub 2000/07/11. doi: 10.1128/jvi.74.15.6800-6807.2000.
PubMed PMID: 10888619; PubMed Central PMCID: PMCPMC112197.

88.   Rosenberg W. Mechanisms of immune escape in viral hepatitis. Gut. 1999;44(5):759-64. Epub
1999/04/16. doi: 10.1136/gut.44.5.759. PubMed PMID: 10205220; PubMed Central PMCID:
PMCPMC1727502.

1051

1052

1053 **Supporting information**

1054

Supplementary Table 1: Sequencing metadata of 76554 GISAID genomes downloaded until

August 3, 2020.

For every GISAID genome, we provided GISAID genome name, GISAID unique

identifier (Accession ID), geographic location, host, sequencing technology, lineage, and clade

fields, among other information. The last column indicates the number of variants per genome

(Major viral variants).

1061

1062 **Supplementary Table 2: Acknowledgements from sequencing laboratories and/or consortia**

1063 **associated with GISAID genomes listed in Supplementary Table 1, plus genomes**

1064 **downloaded until September 28, 2020, containing variants V1176F, S477N and A222V**

1065 **occurring in the Spike protein.**

1066

Supplementary Table 3: Sequencing metadata of 17560 Sequencing Read Archive (SRA)

datasets downloaded until July 28, 2020.

For every SRA dataset, we provided NCBI run accession, Assay type (indicates if

amplicon, RNA-seq u other sequencing corresponds), sequencing size (bases, in nucleotides),

Biosample accession ID, Center Name (depositor), release date, SRA study accession,

BioProject and geographic location, among other information. The last column indicates the

number of variants per sample (Major viral variants, viral frequency > 0.5).

Supplementary Table 4: Sequencing metadata of 543 and 393 GISAID genomes containing

variants N: I292T and S: V1176F, respectively.

We accessed GISAID database on September 28, 2020 and we downloaded genomes

containing the variant I292T or V1176F. The associated metadata from both cohorts are

presented in this table. For every GISAID genome, we provided GISAID genome name, GISAID

unique identifier (Accession ID), collection information, geographic location, host, sequencing

technology, lineage, and clade, among other information. We used GISAID unique identifiers to

overlap both groups. No overlap was found.

1082

Supplementary Table 5: Worldwide fatality ratios per country, related to Supplementary

Figure 2.

Worldwide fatality ratios among 49 countries obtained from John Hopkins coronavirus

resource center (https://coronavirus.jhu.edu/map.html), accessed on September 28, 2020. At

right, we calculated viral allele frequencies of several SARS-CoV-2 variants per country, based

on the GISAID database, also accessed on September 28, 2020. We analyzed the following

variants: ORF3a (Q57H), N (RG203KR), RNA-dependent RNA polymerase (P323L), S

(D614G), S (V1176F), nsp7 (L71F) and N (I292T), respectively.

1091

Supplementary Figure 1: Intra Host variant effects.

23,269 aggregated variants from 76554 GISAID genomes were merged as indicated in

Figure 2A. These variants were classified by SnpEff program, available in the Galaxy server. We

plotted a heatmap with the number of changes per consequence type (see x-axis) against every

SARS-CoV-2 protein (see y-axis). We also calculated the overall missense/silent ratio occurring

in SARS-CoV-2 proteins (1.82).

1098