1 **Speech cortical activation and connectivity in typically developing children and those with**
2 **listening difficulties**
3
4 **Abbreviated title**: Cortical speech networks and listening
5

6 Hannah J. Stewart [1, 2, 3], Erin K. Cash [2], Lisa L. Hunter [2], Thomas Maloney [4], Jennifer Vannest [2,5]
7 and David R. Moore [2, 6, 7]
8
1. Division of Psychology and Language Sciences, University College London, London,
UK
2. Communication Sciences Research Center, Cincinnati Children's Hospital Medical
Center, Cincinnati, Ohio, USA
3. Department of Psychology, Lancaster University, Lancaster, UK
4. Pediatric Neuroimaging Research Consortium, Cincinnati Children's Hospital Medical
Center, Cincinnati, Ohio, USA
5. Department of Communication Sciences and Disorders, University of Cincinnati, Ohio,
USA
6. Department of Otolaryngology, College of Medicine, University of Cincinnati,
Cincinnati, Ohio, USA
7. Manchester Centre for Audiology and Deafness, University of Manchester, Manchester,
M13 9PL, UK
22
23 **Correspondence**:   Hannah J. Stewart
24   Department of Psychology
25   Lancaster University
26   Lancaster UK
27   h.stewart5@lancaster.ac.uk
28

29 **Keywords**: listening difficulties, pediatric, fMRI, resting state, connectivity, speech perception

30
31

Abstract:

Listening difficulties (LiD) in people who have normal audiometry are a widespread but poorly understood form of hearing impairment. Recent research suggests that childhood LiD are cognitive rather than auditory in origin. We examined decoding of sentences using a novel combination of behavioral testing and fMRI with 43 typically developing children and 42 age matched (6-13 years old) children with LiD, categorized by caregiver report (ECLiPS). Both groups had clinically normal hearing. For sentence listening tasks, we found no group differences in fMRI brain cortical activation by increasingly complex speech, from phonology to intelligibility to semantics. Using resting state fMRI, we examined the temporal connectivity of cortical auditory and related speech perception networks. Significant group differences were found only in cortical connections engaged by more complex speech processing. The strength of the affected connections was related to the children's performance on tests of dichotic listening, speech-in-noise, attention, memory and verbal vocabulary. Together, these results support the hypothesis that childhood LiD reflects difficulties in language processing.

Abbreviations:
CANS central auditory nervous system
DLD developmental language disorder
rs-fMRI resting state-fMRI
FOV field-of-view
FWE family wise error correction
GLM general linear model
HUSH Hemodynamics Unrelated to Sounds of Hardware
LiD listening difficulties
MEG magnetoencephalography
MNI Montreal Neurological Institute
MTG middle temporal gyrus
FDR false detection rate
ROI region-of-interest
SLI specific language impairment
STG superior temporal gyrus
STS superior temporal sulcus
TD typical developing
TE echo time
TMS transcranial magnetic stimulation
TR repetition time

INTRODUCTION

Listening is commonly defined as active attention to sound. In human communication, listening
also involves memory, language and executive function (Rudner & Signoret, 2016; Schiller et al.,
2022). Listening difficulties (LiD) are frequently reported by caregivers of children with clinically
normal hearing (Petley et al., 2021; Roeser et al., 2007). However, the mechanisms underlying
childhood LiD without hearing loss remain poorly understood. Several studies have shown that
such children typically also have a variety of academic, speech, language, attention and other
developmental learning problems (Ferguson et al., 2011; Moore et al., 2018; Sharma et al., 2009).
Some of these children receive a clinical diagnosis of auditory processing disorder (APD;
American Academy of Audiology, 2010; British Society of Audiology, 2018; Dawes & Bishop,
2009). The heterogeneous nature of their problems has led to considerable controversy surrounding
APD (Iliadou et al., 2019; Iliadou et al., 2018; McFarland & Cacace, 2009; Moore, 2018;
Neijenhuis et al., 2019; Rees, 1973). At the center of this controversy is the question of whether
the children have primarily auditory sensory problems ("bottom-up"), cognitive and language
problems ("top-down"), or some combination of the two (Dillon & Cameron, 2021; Moore et al.,
2010). In this and other recent studies (Hunter et al., 2021; Petley et al., 2021) we sidestep this
controversy by using the umbrella term LiD (Dillon & Cameron, 2021), and the quantitative
metrics of a well-validated, reliable and standardized caregiver questionnaire (ECLiPS; Barry &
Moore, 2021) to operationalize LiD.

Speech understanding requires coalescence of acoustic features, from bottom-up streaming in the
auditory pathway, with top-down linguistic processes. These latter processes derive from
widespread cortical areas involved with semantic representation and cognitive functions, notably
memory and attention in the frontal, temporal and parietal cortices (Hickok & Poeppel, 2007;
Pichora-Fuller et al., 2016; Rönnberg et al., 2019; B. G. Shinn-Cunningham, 2008). Impaired
interactions between cortical auditory and related speech perception systems, resulting in reduced
higher level processing of target speech, may hold the key to LiD in children. Poor performance
on complex auditory tasks, such as speech listening in noise, has been more closely associated
with cognitive function than with performance on simpler, non-speech based auditory tasks
(Moore et al., 2010). This current study's aim was to examine the cortical networks involved in
different levels of speech processing in children with LiD relative to TD children. To identify

cortical areas used in these processes, we adopted fMRI methods developed to examine cortical
speech processing in adults (Halai et al., 2015; Scott et al., 2000). We then used the identified
cortical areas as regions of interest (ROIs) to examine how they work together and how they relate
to behavioural assessments of speech-in-noise identification, dichotic listening and cognition.
103
Speech is a complex stimulus where multiple levels of processing competencies are required for
comprehension. First, phonology bridges bottom-up and top-down processing, where auditory
information is first classified into linguistically meaningful units (Brodbeck et al., 2018). Second,
the listener links phonemes into words to judge the intelligibility of the auditory information by
comparing the words to memory templates. Third, meaning (semantics) is attached to one or more
intelligible words. Spectral and non-spectral acoustic cues and features enable each of these
processing competencies. For example, non-spectral features of pitch, loudness, temporal order
and binary source features (i.e. voicing-nonvoicing etc.) are required to distinguish individual
phonemes. Spectral features, such as vowel quality, direction of formant transition and noise burst
cues aid in linking the phonemes into words and words into sentences.
114
Scott et al. (2000) designed a paradigm, subsequently modified by Halai et al. (2015), to investigate
cortical activity produced by different levels of speech processing competencies (Figure 1A). They
used two types of speech-like stimuli by spectrally rotating (around 2 kHz; Blesser, 1972), or
rotating and then noise vocoding (into 6 frequency bands; Shannon et al., 1995) clear speech
sentences. In each of the three stimulus types, the sequential order of phonemes was preserved,
thus retaining the prosody (i.e. rhythm) from the clear speech. In the rotated speech the non-
spectral features used to identify phonemes remained, while the spectral features were
transformed. For example, the formant transition and noise burst cues that distinguish consonants
were shifted to new frequencies and so were processed by different regions of the cochlea. This
removed the intelligibility of the stimuli while maintaining the prosodic and phonetic features.
Finally, in the rotated+vocoded speech, the intonation and phonetic features were removed, along
with intelligibility.
127
128

Analysis contrasts were created, by subtracting sentence types from one another (Figure 1A), to

explore which cortical areas were recruited for each of the three speech processing competencies.

First, phonology was assessed by contrasting rotated with rotated+vocoded sentences. Second, the

judgement of intelligibility was assessed by contrasting clear sentences with rotated sentences,

where the spectro-temporal variation was preserved but went to the wrong place on the cochlea.

Finally, the ability to attach meaning to the intelligible sentence was assessed by contrasting clear

sentences with rotated+vocoded sentences where the acoustic complexity was matched but all

meaning was removed. Speech morphs into language as we progress through the hierarchy of

speech processing competencies. For example, phonetic features must be processed to access the

intelligibility of a word. It was thus expected that the spread of cortical activation for the three

contrasts would increase from Phonology to Intelligibility to Semantics, and would include overlap

in auditory and speech areas.

Our first research question asked if the two groups of children (LiD and TD) differ in cortical

activation during speech listening, specifically in early auditory and speech processing cortical

areas. Consistent with a top-down, linguistic model, and previous research in adults (Halai et al.,

2015; Scott et al., 2000), we hypothesised that there would be no group difference in primary

auditory cortex (i.e., Heschl's gyrus) for all three contrasts of speech processing competencies.

However, Intelligibility and Semantic contrasts would show less activation of the speech

processing areas (e.g. superior temporal gyrus, Wernicke's) in children with LiD than in TD

children.

Our second question investigated how the cortical areas activated during these speech processing

competencies work together. We used the task-based cortical activation results to create regions-

of-interest (ROIs) for functional connectivity analysis of a separate rs-fMRI acquired in the same

scanning session. An rs-fMRI scan allows measurement of the temporal correlation of non-

stimulus evoked fluctuations in the BOLD signal between anatomically separated brain regions

(Biswal, 2012), capturing interactions between regions in functionally associated networks

(Damoiseaux et al., 2006). We hypothesised that children with LiD would show diminished

functional connectivity in cortical speech processing networks compared with TD children.

In this report we present data from the first wave of a longitudinal study investigating the
audiological and cognitive abilities of children (6-13 years old; Table 1) with LiD and their
typically developing peers (6-13 years old; Table 1; Petley et al., 2021). All participants had
clinically normal audiometry (≤ 20 dB HL, bilaterally; 0.25 – 8 kHz, Figure 1B), tympanometry
(Hunter et al., 2021) and afferent auditory brainstem function (Hunter et al., 2022). Participants
who scored at or below the bottom 10^{th} percentile on the total score of a caregiver checklist of
everyday listening skills, the ECLiPS (Barry et al., 2015; Barry & Moore, 2021; Roebuck & Barry,
2018), were classified as having LiD (Figure 1C).

RESULTS

Task-based cortical activation

During task-based MRI acquisition intervals (Figure 1D), children listened to clear ("man") or
distorted ("alien"; rotated and rotated+vocoded) spoken sentences (Figure 1E) that we asked them
to match to a visual cartoon representation. All children responded with high accuracy on the
sentence recognition task, and did so rapidly (Figure 1F), suggesting that they maintained attention
throughout the task. TD children were more accurate than children with LiD ($F(1, 83) = 8.77$, $p =$
.004, $\eta p2 = .096$), but there was no significant difference in reaction time between the groups ($F(1,$
$83) = .89$, $p = .35$, $\eta p2 = .011$). While there was no difference across all participants in accuracy
between sentence types ($F(1.78, 148.03) = 2.11$, $p = .13$, $\eta p2 = .025$), clear sentences were
responded to fastest ($F(2, 166) = 11.50$, $p < .001$, $\eta p2 = .12$), compared to rotated ($p < .001$, $d = -$
.41) and rotated+vocoded ($p < .001$, $d = -.46$) sentences. There were no significant interactions.

Contrasting sentence types (Figure 1E) allowed us to explore cortical activation to different levels
of speech processing competencies. The two groups did not differ significantly in cortical
activation in the Phonology, Intelligibility or Semantics contrasts after correcting for multiple
comparisons (Figure 2A-C, clustering threshold = 2.3 voxels, family wise error (FWE) = .95).
Averaged across all children (n = 85), the contrasts (Figure 2D-F) showed bilateral activation
(clustering threshold = 2.3 voxels, family wise error (FWE) = .95) for all three stages of speech
processing competencies. Coordinates for the maximum intensity of the activated regions are
shown in Supplementary Material Table 2. The Phonology contrast (rotated > rotated+vocoded;
Figure 2D) showed bilateral activation in the middle and superior temporal gyrus, including

Heschl's gyrus, and temporal pole. Activation was also found in the left hemisphere in the temporal

fusiform cortex, angular gyrus and lateral occipital cortex. The Intelligibility contrast (clear >

rotated; Figure 2E) produced similar activation to the Phonology contrast, bilaterally in the middle

and superior temporal gyrus (anterior and posterior) and left frontal orbital cortex. Activation

extended anteriorly along the left temporal gyrus and into Broca's area. The Semantics contrast

(clear > rotated+vocoded; Figure 2F) also showed bilateral activation in the auditory cortices

(middle and superior temporal gyrus including Heschl's gyrus and planum temporale) with

activation extending along the left temporal fusiform and frontal orbital cortices and right

parahippocampal gyrus.

Figure 1: All children had normal tone hearing and were able to perform the speech perception task in the scanner. (A) Stimulus contrasts, illustrating how subtracting one stimulus type from another (Fig. 1E) isolates the specific speech processing competencies of Phonology, Intelligibility and Semantics. (B) Group mean hearing thresholds. (C) Total ECLiPS score. (D) fMRI paradigm. Children were asked if the picture (alien, man) matched who spoke the sentence (stimulus type, Fig. 1E). If it was a match they pressed the right button (with their stickered hand), if it was not a match they pressed the left button. Data acquisition (grey shading) was turned off/on for the presentation of the auditory stimuli (HUSH/sparse scanning). (E) Stimulus types. Spectrograms of 'The two children are laughing'. Time is represented on the x-axis (0.0–1.60 s) and frequency on the y-axis (0.0–4.4 kHz). The shading of the trace in each time/frequency region is controlled by the amount of energy in the signal at that particular frequency and time (red = more energy, blue = less energy). Clear speech ("man") is intelligible with intonation. Rotated speech ("alien") is not intelligible, though phonetic features and original intonation are preserved. Rotated noise-vocoded speech ("alien") is completely unintelligible, but preserves the character of the envelope and some spectral detail. (F) Group mean response times and accuracy for each stimulus type. Error bars represent the standard error of the mean. Groups as in Fig. 1C.

Scan type	Group	N	Age M (SD)	Gender M, F	Maternal education*	Handedness L, both, R	History of tubes	Motion-related artifacts M (SD)
fMRI	LiD	43	10.02 (2.13)	30, 13	6, 27	3, 5, 35	12	12.16 (4.63)
	TD	42	9.78 (1.93)	25, 17	0, 42	2, 1, 39	15	11.67 (4.30)
RS	LiD	42	10.06 (2.09)	29, 13	8, 34	3, 6, 33	12	28.64 (32.17)
	TD	39	9.75 (1.93)	23, 16	0, 39	2, 1, 36	15	23.00 (29.20)

Table 1: Participant details for each scan type – fMRI speech listening task and resting state (RS).

* High school graduate or less, some college or more

Cortical functional connectivity

Networks of ROIs suitable for rs-fMRI connectivity analysis were created by parcellating activation produced by the three contrasts from the task-based MRI (Phonology, Intelligibility, and Semantics), combined across groups (Figure 2D-F). The pediatric ADHD-200 sample (Bellec et al., 2017; Craddock et al., 2012) was used as a data-driven, spatially-constrained parcellation method. ROIs smaller than 4 voxels were removed. Cortical activity from the Phonology contrast was divided into a network of 16 ROIs, Intelligibility into a network of 20 ROIs, and Semantics into a network of 24 ROIs (Figure 3A-C; Brodman's area, maximum intensity coordinates and ROI sizes are in Supplementary Material Table 3). Connectivity was compared between groups (TD = 42, LiD = 39), controlling for age, using a general linear model (GLM) for each of the three networks (Supplementary Material Table 3).

Figure 2: LiDLiD and TD groups showed similar areas of cortical activation in all three contrasts from the fMRI listening task, with no statistical difference between the two groups after correcting for multiple comparisons. Second level GLM analysis for (A-C) groups (clustering threshold = 2.3 voxels, FWE = .95) and (D-F) all participants (clustering threshold = 4.0 voxels, FWE = .95) in the fMRI task, coordinates (±60, -5, 0). Coordinates for maximum intensity voxels for (D-F) can be found in Supplementary Material Table 2. Contrasts are: (A, D) Phonology (green: rotated > rotated+vocoded), (B, E) Intelligibility (blue: clear > rotated) and (C, F) Semantics (red: clear > rotated+vocoded). We took the cortical activation across all participants and created parcellated ROIs (Figure 3) for use in the rs-fMRI analysis (figure 4). Images are in neurological orientation. MRIcroGL was used for visualization.

Figure 3 (D-F) shows resting state functional connectivity in the three speech networks, summed

across groups. In the Phonology network, each group had connectivity among regions covering

bilateral middle and superior temporal gyri, temporal pole, planum temporale, and left planum

polare and supramarginal gyrus (Figure 3D). No significant group differences (p-FDR) were found

in the Phonology network (Figure 4A, D). Each group had connectivity within the Intelligibility

network, covering bilateral middle/superior temporal gyrus and temporal pole along with left pars

opercularis, frontal orbital cortex and supramarginal gyrus (Figure 3E). After false discovery rate

(FDR) correction, TD children were found to have a significantly stronger temporal correlation

between ROIs in left temporal lobe and left middle temporal gyrus (posterior) compared to

children with LiD (connection 4-14 in Figure 4B; Table 3). In the Semantics network, both groups

of children had connectivity between bilateral middle and superior temporal gyrus, left Heschl's

gyrus, pars triangularis, frontal orbital cortex, planum temporale, temporal fusiform gyrus and

228 right parahippocampal gyrus and planum polare (Figure 3F). Group comparisons showed that,

229 compared to children with LiD, the TD children had stronger temporal correlations between ROIs

230 in the fusi (connection 7-24 in Figure 4C; Table 3).

Figure 3: ROIs used for the rs-functional connectivity and their networks across all participants. The cortical activity across all participants in the fMRI task (see Figure 2 D-F) covered large areas and so they were parcellated into smaller ROIs (A-C) for the rs-functional connectivity analysis by applying data-driven spatially constrained parcellation to the areas of activation from the fMRI sentence recognition task using the pediatric ADHD-200 sample. The red lines (D-F) indicate the ROI-to-ROI connections analyzed in each network. Maximum intensity coordinates can be found in Supplementary Material Table 3. Networks are: (A, D) Phonology (green: rotated > rotated+vocoded), (B, E) Intelligibility (blue: clear > rotated) and (C, F) Semantics (red: clear > rotated+vocoded). Images are in neurological orientation. For visualization, BrainNet software was used to display foci in (A-C) (Xia, Wang & He, 2013).

233 *Relation of cortical functional connectivity to behavioral measures*

234 We explored how the functional connectivity of anatomically separate cortical areas related to

235 behavioral tasks assessing speech in noise ability, dichotic listening and cognition. As shown by

236 Petley et al. (2021), performance on those behavioural tasks was significantly better in the TD than

237 in the LiD group.

239 The Intelligibility connection had a series of significant positive correlations with speech-in-noise

240 ability (SCAN: auditory figure ground), vocabulary and task switching, a form of executive

241 function (NIH Cognition Toolbox). In contrast, the Semantics connection, involving the left

242 posterior temporal fusiform cortex (7-24; Figure 4C), had significant positive correlations with

243 dichotic listening ability (SCAN: competing words and sentences) and vocabulary (NIH Cognition

Toolbox). Interpretation of direction of functional connectivity is ambiguous; an increase in strength of connectivity does not necessarily mean improved ability (Parente et al., 2018). However, these correlations show that, as the children's behavioural test scores improved, the connection strength within the Intelligibility and Semantic networks increased. The remaining behavioral measures did not relate significantly to connectivity (Figure 4E, Supplementary Material Table 4 and Figure 1).

Contrast	Max. intensity MNI coordinates				Brain Regions (Harvard-Oxford atlas)	Cluster size (voxels)
	BA	x	y	z		
Intelligibility normal > rotated	4	20	-46	12	-34	L temporal pole	82
	14	21	-50	-34	-6	L middle temporal gyrus, post	635
Semantics normal > rotated+vocoded	7	20	44	0	-26	R STG, ant	108
	24	37	-38	-40	-22	L temporal fusiform cortex, post	4

Table 2: Selected regions of interest (ROIs, from the fMRI task; Fig. 3B and C) used in the rs-fMRI ROI-to-ROI analysis. Clustering threshold = 4.0 voxels. Family-wise error correction = .95. The full table of all ROIs can be found in Supplementary Material Table 3.

Contrast	ROI # – ROI # (Fig. 3)	t (78)	p-FDR
Intelligibility	4 – 14	3.11	0.026
Semantics	7 - 24	3.54	< 0.001

Table 3: Network temporal connectivity. Statistical comparison between group differences (TD vs. LiD, corrected for age) for the Intelligibility and Semantics networks (Fig. 4D). No significant group difference was found in the Phonology network. False discovery rate (FDR) was used for multiple comparison correction .

Figure 4: ROI-to-ROI resting state connectivity: the difference between the groups' listening networks grew from no statistical differences in the phonology network to a minor difference in the intelligibility network to more widespread differences in the semantics network. (A-C) The group comparisons without the effect of age for (A) Phonology, (B) Intelligibility and (C) Semantics networks. Thicker, more saturated color lines represent stronger connections between cortical areas. Note that the colored bar connectivity z score scales vary slightly between connectivity wheels. (D) Details of the ROI-to-ROI connectivity values (left axis) for each group (LiD green, TD grey) and effect sizes (yellow marker, right axis) of the group comparisons without the effect of age. The connections plotted are the ones highlighted as having a significant group difference in the GLM comparing groups without the effect of age (B and C). Connections show the TD group as having significantly stronger connectivity than the LiD group. (E) Scatter plots of brain and behavioral scores demonstrate the correlated patterns of connectomic and behavioural features. Coloured dots in each panel indicate the participant group (LiD: green, TD: grey). The solid black line marks significant correlations (see Supplementary Table 4 for correlation details).

DISCUSSION

Using an active sentence listening task we found bilateral cortical activation across the primary

auditory cortex areas (e.g. Heschl's gyrus, superior temporal lobe) for all three contrasts of speech

processing competencies. The activation extended bilaterally into more advanced speech

processing areas including memory and language areas (e.g., Broca's, right parahippocampal

gyrus, left temporal fusiform cortex) as we progressed through the hierarchy of speech processing

competencies, from phonology to intelligibility to semantics. Cortical areas of activation were

unaffected by LiD for all three contrasts. This provides support for our hypothesis that the children

with LiD activate the same primary auditory areas as the TD children when listening to speech.

We did not find support for our hypothesis that children with LiD show less activation in the speech

processing areas than in TD children. However, these patterns of activation were examined using

BOLD response which does not capture the excitatory and inhibitory balance or temporal

relationships between cortical areas.

Analysis of the separate rs-fMRI data showed that temporal correlations, quantifying functional

connectivity of anatomically separated brain regions, between speech activation-defined sites

differed between TD children and those with LiD. This provides support for our functional

connectivity hypothesis, but only in networks of more complex speech processing competencies.

These results suggest that children with LiD show diminished functional connectivity in

intelligibility and semantic processing networks compared to TD children. This is consistent with

an interpretation of childhood LiD as a language disorder, with difficulties in intelligibility and

semantics, rather than as a phonological disorder.

Behavioural measures of performance were consistent with these observations. No significant

correlations were observed between functional connectivity and sensory-dominant tasks (e.g,

LiSN-S advantage scores and SCAN filtered words). However, significant correlations were

observed between functional connectivity and more cognitive-dominant tasks, specifically speech-

in-noise and task switching which are more strongly dependent on language and executive

function, respectively (Petley et al., 2021). Overall, the data are consistent with the hypothesis that

children with LiD have primarily cognitive and language processing deficits.

Lateralization of speech in 6-12 year old children

Using simple, but complete sentence stimuli, we consistently found bilateral activation in children

for all speech listening contrasts. This activation did not significantly differ between groups, after

FDR correction, suggesting that children with LiD use the same cortical areas as TD children for

increasingly complex speech processing competencies. However, it is possible that multi-voxel

pattern analysis may find finer group differences in the hierarchy for speech processing (e.g.,

Okada et al., 2010). While the same cortical areas are used by the children with LiD, it is possible

that they do so on a different time frame from the TD children. Unfortunately, fMRI does not
provide sufficient time resolution to address this possibility.

Our results support the theory of bilateral processing of speech listening in 6-12 year old
population. Lateralization of speech processing in adults has been debated across and within
imaging modalities (fMRI see Evans & McGettigan, 2017; EEG e.g., Assaneo et al., 2019). For
example, Albouy et al. (2020) discussed how acoustic structure leads to a left bias for fast
modulation (speech) and a right bias for slow modulation (music) and Rauschecker & Scott's
(2009) unilateral model suggested that the left anterior STG is the hub of successful speech
perception. In contrast, Hickok and Poeppel's (2000) bilateral model proposed a perceptual
pathway in each hemisphere processing speech sounds up to the level of semantics (Hickok &
Poeppel, 2007). A middle ground has been proposed by Peelle (2012), with right hemisphere
dominated activation for "unconnected" speech (i.e., phonemes, syllables, single words) and left
hemisphere dominated activation for "connected" speech (i.e., phrases, sentences and narratives).

Bilateral activation reported here differs from the findings of Scott et al. (2000) who reported a
left lateralized pathway for speech comprehension using PET, a silent scanning technique. Our
results extend those of Halai et al. (2015) who found bilateral activation during continuous fMRI
scanning. Both these studies tested young adults and used similar stimuli to those used here.
However, speech presentation was continuous, and passive in the sense that the participants did
not perform any task during scanning. Rather, comprehension was assessed after scanning and
outside the scanning room. In contrast, we required children to provide a behavioural response to
each short sentence presentation to encourage attention and as a metric of attention. We
additionally used a silent, "sparse" acquisition protocol (Hall et al., 1999; Vannest et al., 2009), in
contrast to the previous (Halai et al., 2015) continuous fMRI scanning protocol. Scanner noise
superimposed on speech listening has been shown to increase listening effort (Peelle et al., 2010)
and to engage additional and or different cortical areas compared with silent acquisition protocols.
For example, in a meta-analysis of 57 speech comprehension studies, Adank (2012) showed that
continuous scanning more strongly activated regions of the supplementary motor area and anterior
cingulate gyrus, while sparse scanning showed more extensive activation in the STS.

Functional connectivity: Brain and behaviour

We used temporal correlations in rs-fMRI to compare how cortical areas associated with speech processing competencies worked together in children with and without LiD. Phonology is the system of processing the smallest units of speech sounds and their linguistically appropriate combinations. We found no group difference in the connectivity of this network. As we progressed up the speech processing competency hierarchy to the Intelligibility network, we found that temporal correlation between left temporal lobe and left posterior middle temporal gyrus activity was stronger in the TD children. These areas are well known for processing auditory information (temporal lobe) and speech comprehension (left MTG, Acheson & Hagoort, 2013; Dronkers et al., 2004). Posterior MTG has been associated with lexical and semantic access in a sound-to-meaning network (Hickok & Poeppel, 2000, 2004). It is unknown whether stronger or weaker temporal correlations between brain regions is beneficial (Parente et al., 2018). However, relationships found here between the variability of brain connectivity strength and of behaviour provides evidence on this key issue. Children with stronger connectivity performed better on cognitive tasks assessing vocabulary and switching attention. Increased connectivity may thus be indicative of increased processing efficiency and/or suppression of a task-relevant network, in this case language. Note that the behavioural measures used here have been shown to be either little affected by age (audiogram: Hunter et al., 2021) or were standardized across age (ECLiPS, LiSN-S, SCAN and NIH toolbox; (Petley et al., 2021).

Group comparisons in the Semantics network highlighted that, compared to children with LiD, the TD children had stronger temporal correlations between auditory areas (right STG) and the left temporal fusiform cortex, associated with word recognition and the recovery of meaning from an impoverished acoustic signal (Davis & Johnsrude, 2003). Brain-behaviour correlations suggest that decreased connectivity between auditory and word recognition cortical areas was associated with impaired speech listening (dichotic listening and verbal vocabulary; Table 2). Recently, brain-behaviour correlations have been called into debate due to extremely large datasets being required to reach even very small correlations (Marek et al., 2022). However, we consistently found mild to moderate sized correlations with a peak of $|r| = 0.33$ (mean $|r| = 0.29$, first percentile $|r| = 0.26$; Supplementary Material Table 4).

The cortical areas highlighted in this study (Figure 4 B and C) draw focus to language production,

memory encoding and retrieval, and word recognition (Acheson & Hagoort, 2013; Davis &

Johnsrude, 2003; Dronkers et al., 2004; Hickok & Poeppel, 2000, 2004) as target areas of further

research in children with LiD. Group differences were found in networks of latter stages of speech

processing competencies, in the intelligibility and semantics networks. Specifically, between the

MTG/STG and the temporal lobe. The reduction in connectivity in these networks suggest that

childhood LiD is a semantic and intelligibility disorder. However, no group differences were found

in the phonological networks, an early stage of speech processing competencies. Suggesting that

childhood LiD is not a phonological disorder.

Further assessment where the 'break' in speech listening occurs in LiD would provide a clearer

avenue for research into effective evidence-based treatments. Future studies could utilize

paradigms with anomalous and mispronounced words with imaging techniques used in parallel to

assess if the type of speech processing errors during such tasks are affected by LiD. For example,

was it detected when presented (i.e. when judging it's intelligibility) or did the listener reach the

end of the sentence and have to 'go back' when they are unable to connect meaning to the sentence.

Complementary time-sensitive techniques (EEG, MEG) could also investigate whether this

difficulty is due to a bottleneck in processing leading to increased listening effort, cognitive effort

or fatigue (McGarrigle et al., 2014). As LiD may build up during a listening event (i.e., over several

minutes; Roebuck & Barry, 2018), it is important to assess the children's ability throughout the

task with cross-sectional time points rather than using summary values of the complete listening

event (McGarrigle et al., 2020).

Relationship to neurodevelopmental disorders and cognitive function

Children identified with LiD have difficulty in speech listening compared to TD children (Petley,

Hunter, Motlagh Zadeh, et al., 2021). These groups were further distinguished here by functional

connectivity differences between cortical areas associated with language production, memory and

word recognition, rather than by the activity or connectivity of primary auditory cortical areas.

However, at least 50% of children referred/diagnosed with auditory processing disorder (APD;

(British Society of Audiology, 2018; Dillon & Cameron, 2021; Jerger & Musiek, 2000), a clinical

label closely related to LiD (Dillon & Cameron, 2021), have also been diagnosed with

developmental language disorder (DLD), dyslexia/reading disorders, attention
deficit/hyperactivity disorder (ADHD) or more than one of these other neurodevelopmental
disorders (Dawes & Bishop, 2010; Ferguson et al., 2011; Gokula et al., 2019; Moore et al., 2018;
Sharma et al., 2009). This high level of comorbidity was echoed in the study reported here. While
LiD was the primary referral for the study, as with other neurodevelopmental disorders it does not
stand in isolation. A background caregiver questionnaire showed that half of the children with LiD
also reported a diagnosis of ADHD, 9% autism spectrum disorders and 26% had seen a speech
language pathologist. However, these data were not entered into the analysis as we did not conduct
gold standard assessments for these other developmental disorders. We are currently using a web-
based resource (Neurosynth; Yarkoni et al., 2011) that allows functional connectivity analysis of
brain areas defined by a meta-analysis of published fMRI activation coordinates to explore whether
there are shared audio-based neurological patterns in children with primary diagnoses of other
developmental disorders, such as ASD and ADHD, with LiD.

Further investigation into the neurodevelopmental basis of LiD may particularly inform the
understanding of language disorders. Our results show typical phonology but impaired non-
phonological speech connectivity in children with LiD. This differs from DLD, which presents
with both abilities impaired, and dyslexia, which presents with impaired phonological and typical
non-phonological abilities in reading (Bishop & Snowling, 2004; Delage & Durrleman, 2018). It
is possible that altered cortical language processing leads to LiD. However, it could be that altered
cortical language processing may be a consequence of LiD. Further longitudinal analysis of these
children is designed in part to address these issues.

The neuroimaging results presented here highlight the importance of non-auditory factors,
specifically language, in audiological testing at a cortical level. There is a growing recognition of
the importance of language and, specifically, speech-in-noise (SiN) intelligibility in everyday
hearing (Dillon & Cameron, 2021; Golestani et al., 2009; Killion et al., 2004; Magimairaj et al.,
2021; Smits et al., 2013). It has been proposed that such testing could supplement, or even replace
pure tone detection as an audiometric gold standard (Hewitt, 2018). However, both SiN test
instructions and test items pose a challenge to language and memory as well as auditory function.
While those cognitive aspects of auditory testing and learning have previously been dismissed as

procedural issues (Hawkey et al., 2004), they are an intimate component of a SiN test. These
neuroimaging results provide insight into mechanisms of how the advanced stages of speech
processing where auditory information is translated into language during speech perception may
be disrupted in LiD, a common form of auditory impairment in both children (Moore et al., 2018)
and adults (Edwards, 2020). They also add to a growing literature on the role of cognitive function
in hearing (Moore et al., 2014; Rönnberg et al., 2013; Sharma et al., 2019; B. Shinn-Cunningham,
2017).

Conclusions
Our results provide the first multifaceted neurological profile for children classified with LiD,
based on caregiver report and normal peripheral auditory function. Children with LiD recruited
the same cortical areas as their peers when processing increasing complexities of speech. However,
how these cortical areas work together does differ between the two groups. Differences in
functional connectivity were found at the more advanced stages of speech listening where the
intelligibility and semantics of the speech are processed, specifically in the left temporal lobe,
posterior middle temporal gyrus, posterior temporal fusiform cortex and right superior temporal
gyrus. These highlighted cortical connections related to the children's behavioural abilities in
dichotic listening, speech-in-noise, attention, memory and verbal vocabulary abilities. Overall, the
data provide support for the hypothesis that children with LID are primarily affected by cognitive
and, particularly, language processing deficits.

METHODS

Participants

Eighty-five participants aged 6-12 years completed the fMRI sentence task and 81 participants completed the rs-fMRI (see Table 1). Seventy-six of participants who completed the fMRI sentence task also completed the rs-fMRI. All participants had normal audiometric hearing with thresholds < 25 dB HL at all octave-interval frequencies from 0.25 - 8 kHz in both ears (Figure 1A). In this paper we focus on the fMRI and its relationship with behavioural responses from the baseline of our longitudinal 'SICLID' study examining correlates of LiD in children. Extensive analysis of ear function and behavioural responses of these children is reported elsewhere (Hunter et al., 2021, 2022; Petley et al., 2021)).

Caregivers of all participants completed a standardized, validated and reliable checklist of everyday listening and related skills (ECLiPS; Barry & Moore, 2021). Children recruited through advertisement all scored < 10th percentile of ECLiPS standardized scores. Additional children, who had an audiological diagnosis of auditory processing disorder (APD; n=14), were placed in the LiD group. All but one of these also scored < 10th percentile on the ECLiPS (Figure 1C). Participants scoring within the upper 90th percentile on the ECLiPS and with no history of developmental disorders or delays were classified as TD (further details in Petley et al., 2021).

Eligibility for both groups included English as the child's native language, and a reported absence of otologic, neurologic or psychiatric disease, or of intellectual insufficiency that would prevent or restrict their ability to complete testing procedures. TD participants were additionally required to have no known history of developmental delay, or attention or language disorder. While half of the children with LiD also reported a diagnosis of ADHD, 9% autism spectrum disorders and 26% had seen a speech language pathologist. Eligibility was determined based on caregiver responses on a medical and educational history 'Background' questionnaire (Supplementary Material Table 1).

This study was approved by the Institutional Review Board of Cincinnati Children's Hospital (CCH) Research Foundation. Prior to completion of study-related imaging and behavioural testing, caregivers reviewed the informed consent form with a study staff member. Children aged 11 and

above were also assented using a child-friendly version of the consent document, per institutional

policy. All participants received financial compensation for their participation.

MRI acquisition

MRI was performed via a 3T Philips Ingenia scanner with a 64-channel head coil and Avotec

audiovisual system. All participants were awake throughout the scanning. The protocol was

modified from a previous large, cross-sectional examination of brain function (Holland & Vannest,

2015) and included a high-resolution T1-weighted anatomical scan, fMRI sentence task (4.9

minutes) and rs-fMRI (5 minutes). The fMRI sentence task was acquired with a sparse scanning

protocol ('HUSH', details below); TR/TE = 2000/30 ms, voxel size = 2.5 × 2.5 × 3.5 mm, 39 axial

slices. A total of 147 volumes was acquired by alternating scanning for 6 seconds (3 volumes) and

not scanning for 6 seconds, 49 times. Cardiac and respiration signals were collected during the

fMRI sentence task using the scanner's wireless respirator bellows and Peripheral Pulse Oximeter.

The rs-fMRI acquisition was acquired with TR/TE = 2000/30 ms, voxel size = 2.5 × 2.5 × 3.5 mm,

39 axial slices in ascending slice order and 150 volumes. The high-resolution T1-weighted

anatomical scan was acquired with TR/TE=8.1/3.7 ms, FOV 25.6 x 25.6 x 16.0 cm, matrix 256 x

256 and slice thickness = 1mm.

fMRI task

With sound levels reaching 118.4 ± 1.3 dB (A) in a 3T MRI system (Price et al., 2001) special

considerations must be made when planning an auditory-based MRI study. In order to protect the

participant from the loud environment, foam ear plugs and MRI safe circumaural headphones were

worn. The scanner noise may also produce masking of the desired stimuli. Therefore, in the fMRI

task we used a 'Hemodynamics Unrelated to Sounds of Hardware' (HUSH) scanning protocol

(Deshpande et al., 2016; Edmister et al., 1999; Hall et al., 1999; Schmithorst & Holland, 2004) - a

sparse temporal sampling protocol where there was no gradient coil noise during presentation of

the auditory stimuli. Stimulus levels were elevated to produce adequate signal/noise ratios for

accurate responding. We also used a talker identification task instead of a speech recognition task.

Instead of asking the children *what* they heard, we asked them *who* had said it (Fig. 1). The children

responded with button presses throughout the task so we could ensure they maintained attention

to the task.

Sixteen linguistically simple BKB sentences (Bench et al., 1979), designed to be familiar to young
children, were recorded by a single male North American speaker, mirroring the paradigm used
by Scott et al. (2000) and Halai et al. (2015). These were the clear speech stimuli (e.g. Figure 1E).
Rotated speech stimuli were created by rotating each sentence spectrally around 2 kHz using the
Blesser (1972) technique. Rotated speech was not intelligible, though some phonetic features and
some of the original intonation was preserved. Rotated+vocoded speech stimuli were created by
applying 6 band noise-vocoding (Shannon et al., 1995) to the rotated speech stimuli. While the
rotated noise-vocoded speech was completely unintelligible, the character of the envelope and
some spectral detail was preserved.
511
Participants were told that they would be completing a matching game where they would hear a
sentence and then see a picture (of a "man" or an "alien"). If the picture matched who said the
sentence (man - clear speech, alien - rotated or rotated+vocoded speech), the participant pressed a
button with their right thumb, if the picture did not match, they pressed a second button with their
left thumb. The participants were asked to respond as fast and as accurately as possible. A sticker
was placed on the participants' right hand to provide a reminder as to which hand was correct for
matching voice and picture.
519
Before scanning, each participant was familiarized with the sentence task and completed three
practice trials with verbal feedback from the tester. If a trial was completed incorrectly, the stimuli
and instructions were reintroduced until the participant showed understanding. During scanning,
each participant completed 48 matching trials, 16 of each sentence type, with no feedback. To
maintain scanner timings the behavioural task continued regardless of whether the participant
responded. However, if the child did not press a response button on three trials in a row the tester
provided reminders/encouragement over the scanner intercom between stimulus presentations.
527
528 **fMRI data analysis**
529 First-level fMRI data were processed using FSL (FMRIB Software Library,
530 https://fsl.fmrib.ox.ac.uk/fsl/). The T1 brain data were extracted using BET and normalized and
531 resampled to the 2 mm isotropic MNI ICBM 152 non-linear 6th generation template using FLIRT.
532

For the sparse HUSH acquisition, the volumes were separated and combined into three files

according to the volume's order during the scanner-on period. Each of the three files was pre-

processed separately and first-level statistics computed. The three statistical images were then

averaged together using a one sample t-test. This was done to account for the difference in

intensity among the volumes due to T2* relaxation effects. The pre-processing steps included the

following. FSL's BET was used for brain extraction of the functional data. Outlying functional

volumes were detected with 'fsl_motion_outliers' using the RMS intensity difference metric.

AFNI's '3dretroicor' was used to regress out the cardiac and respiration signals using a

RETROICOR approach (Glover et al., 2000). Motion correction was carried out by MCFLIRT. A

GLM was used to regress motion-related artifacts from the data using 6 regressors for the motion

parameters and an additional regressor for each outlying volume. The amount of motion during

the scans (the number of outlying volumes for each participant) did not differ between groups, p

$= .62$ (Table 1). The data were spatially smoothed using a Gaussian kernel with a sigma of 3 mm

and temporally filtered with a high pass filter with a sigma of 30 seconds. The results were

interpolated to a 2 mm isotropic voxel size and aligned to the Montreal Neurological Institute

(MNI) template by first co-registering it with the participant's T1 using FSL's FLIRT.

Second-level analysis was also conducted using FSL. A GLM approach was used to create group

activation maps based on contrasts between conditions for all participants (i.e. regardless of

LiD/TD status) with age as a covariate. Group composite images were thresholded using a family-

wise error correction ($p < 0.05$) and clustering threshold of $k = 4$ voxels. Three BOLD activation

contrasts were used to search for brain loci responding to different aspects of listening to language

(Halai et al., 2015; modified from Scott, 2000). First, a 'Semantics' activation map, whereby the

signal with intelligibility, intonation, phonetics, and prosody was contrasted with one lacking all

of these attributes except prosody (clear > rotated+vocoded). Second, an 'Intelligibility' activation

map contrasted the signal with all speech attributes to one retaining intonation, phonetics and

prosody (clear > rotated). Third, a 'Phonetics' activation map contrasted a signal with intonation,

phonetics and prosody with one having only prosody (rotated > rotated+vocoded).

Behavioural responses from the fMRI task were assessed using a 2 (group: TD, LiD) $\times$ 3 (sentence

type: clear, rotated, rotated+vocoded) repeated measures analysis of variance (ANOVA) for

accuracy and again for RT (Figure 1F). Where the assumption of sphericity was violated, degrees

of freedom were corrected using Greenhouse-Geisser estimates of sphericity.

Resting state fMRI

During this second scan the participant was asked to lie still, keep their eyes open and look at the

(central) white cross on the black screen. During the 5 minute period they were not performing an

exogenous task. Eyes were monitored by the tester through CCTV and no child fell asleep during

the task.

For the rs-fMRI scan, pre-processing and analysis was performed in the CONN toolbox using

standard spatial and temporal pipelines (Whitfield-Gabrieli & Nieto-Castanon, 2012). For spatial

smoothing a FWHM of 8mm was used. The Artifact Detection Tool (ART,

https://www.nitrc.org/projects/artifact_detect) within CONN was used to regress out framewise

motion. The number of frames regressed out was compared between groups with no significant

group differences, $p = .44$ (Table 1).

Resting state ROI-to-ROI analysis

The group activation maps from the fMRI task's three contrasts (cluster threshold = 4 voxels; p-

FWE = .95) were used to define the ROIs of advancing speech processing competency networks

(Phonology, Intelligibility and Semantics). However, as these areas of activation were large (e.g.

an area of 4351 voxels in the left frontal lobe as part of the Semantics network, Supplementary

Material Table 2), we applied the parcellation from the pediatric ADHD-200 sample (Bellec et al.,

2017) to each network. This created smaller and more appropriate ROIs for connectivity analysis

of each network (Figure 3D-F, Supplementary Material Table 3). ROIs smaller than 4 voxels were

not included in the analysis.

Conn was used to test the functional relationship between each pair of ROIs identified in the fMRI

sentence listening task. The mean time course of all voxels within each ROI was used to calculate

individual pairwise Pearson correlations. The r values were normalized to z values via Fisher's z-

transformation. We then used these z values to explore the relationship between the three listening

networks and behavioural measures. Statistical thresholds were set to p < .05 (corrected) at the

single voxel level, and the resulting connections were thresholded at seed-level by intensity with

FDR correction ($p < .05$).

Caregiver questionnaire

Everyday listening skills – ECLiPS (Barry & Moore, 2021)

The ECLiPS is a standardized caregiver-report measure of listening and communication

difficulties. Caregivers rated 38 simple statements about their child on a five-point scale, ranging

from strongly disagree to strongly agree.

Behavioral measures

Resting state temporal connections with significant group differences were correlated with

behavioural measures described briefly below (see Petley et al., 2021 for further detail and data).

Study data were collected and managed using REDCap electronic data capture tools hosted at

Cincinnati Children's Hospital (Harris et al., 2009, 2019). REDCap (Research Electronic Data

Capture) is a secure, web-based software platform designed to support data capture for research

studies, providing 1) an intuitive interface for validated data capture; 2) audit trails for tracking

data manipulation and export procedures; 3) automated export procedures for seamless data

downloads to common statistical packages; and 4) procedures for data integration and

interoperability with external sources.

Listening in Spatialized Noise-Sentences (Cameron & Dillon, 2007, 2008, 2009)

LiSN-S (US version) is a standardized test assessing speech in noise ability. Binaural target (T)

sentences were presented through headphones along with two other distracting sentences (D1, D2).

The children were asked to repeat the sentences of the target voice only. Distracting sentences

remained constant at 55 db SPL. After each correct trial the target voice descended in level (4 dB),

but if the child incorrectly repeated back over 50% of the sentence the level increased (by 2 dB).

Four listening conditions are made by manipulating D1 and D2 with respect to T (same voice,

different voices; same direction, 0°, different direction, ± 90° azimuth). Three difference

("Advantage") scores are calculated to reduce the influence of language and cognitive demands of

each condition (Dillon et al., 2014): Talker Advantage (different voices - same voice); Spatial

Advantage (different directions – same direction); and Total Advantage (different voices and

directions – same voices and directions). LISN-S software calculated these difference scores for

each participant.

SCAN-3:C (Keith, 2009; 2000)

SCAN-3:C is a US-standardized test battery often used by audiologists to diagnose APD in

children (Emanuel et al., 2011). We used the task to obtain comparison measures relative to other

studies of auditory processing ability (e.g. Kelley & Littenberg, 2019), but not to group children

as LiD or TD. Subtests used in our battery were Auditory Figure Ground - assessing the ability to

repeat words presented against background multi-talker speech; Competing words - a dichotic

listening task where the child repeats different words presented simultaneously to each ear,

repeating that from a designated ear first; Filtered words - assessing ability to identify words that

are low pass filtered at 750 Hz; and Competing Sentences - a dichotic listening task where different

sentences are presented simultaneously to each ear, and the child is asked to repeat the sentence

from a designated ear. Both subtest and a standardized SCAN composite score are calculated.

Cognition - NIH toolbox (Weintraub et al., 2013)

The NIH toolbox - Cognition Battery is a collection of US-standardized tests from which we used

measures of selective attention (Flanker Inhibitory Control and Attention Test), episodic memory

(Picture Sequence Memory Test), executive function (Dimensional Change Card Sort Test) and

picture vocabulary. Each visually administered test took 5 - 15 minutes to complete on an iPad.

Age-corrected subtest and an overall 'early childhood composite' scores were calculated for each

participant.

DATA AVAILABILITY

In accordance with ethics requirements, the dataset generated and analysed during the current

study are available from the corresponding author. ROIs used in the rs-fMRI analysis are available

at GitHub (https://github.com/stewarthannahj/ROIs_SICLiD.git).

ACKNOWLEDGEMENTS

Thanks to Dr Kim Leikin and Prof. Scott K Holland for aspects of study design and to Prof. Stuart Rosen and Dr Peter Chiu for the scripts to rotate and vocode the stimuli for the fMRI sentence task. Thanks also to Audrey Perdew, Nicholette Sloat and Dr Ronan McGarrigle for assisting in MRI acquisition. Many thanks to the CCH MRI techs (Matt Lanier, Lacey Haas, Kaley Bridgewater, Kelsey Murphy, Byrnne Williams and Marty Jones) for many hours of scanning.

FUNDING

This research was supported by NIH R01DC014078, NIH 2UL1TR001425 and by Cincinnati Children's Research Foundation. DRM is supported in part by the NIHR Manchester Biomedical Research Centre.

COMPETING INTERESTS

The authors report no competing interests.

CONTRIBUTIONS

HJS - study design, MRI acquisition, MRI preprocessing and analysis, behavioural analysis, figures and tables, manuscript preparation and revisions

EC - behavioural and rs-fMRI analysis, figures, assisted with manuscript preparation and revisions

LLH - study design, manuscript preparation and revisions

TM - fMRI preprocessing, assisted with manuscript preparation

JV - study design, analysis and presentation, manuscript preparation and revisions

DRM - study design and presentation, funding, manuscript preparation and revisions

REFERENCES

Acheson, D. J., & Hagoort, P. (2013). Stimulating the brain's language network: Syntactic

ambiguity resolution after TMS to the inferior frontal gyrus and middle temporal gyrus.

Journal of Cognitive Neuroscience, *25*(10), 1664–1677.

Albouy, P., Benjamin, L., Morillon, B., & Zatorre, R. J. (2020). Distinct sensitivity to

spectrotemporal modulation supports brain asymmetry for speech and melody. *Science*,

367(6481), 1043–1047.

American Academy of Audiology. (2010). *Diagnosis , Treatment and Management of Children*

and Adults with Central Auditory Processing Disorder American Academy of Audiology

Clinical Practice Guidelines. *August*.

Barry, J. G., & Moore, D. R. (2021). *ECLIPS: Evaluation of children's listening and processing*

skills (2nd ed.). Cincinnati Children's Hospital.

Barry, J. G., Tomlin, D., Moore, D. R., & Dillon, H. (2015). *Use of Questionnaire-Based*

Measures in the Assessment of Listening Difficulties in School-Aged Children. 1–14.

Blesser, B. (1972). Speech perception under conditions of spectral transformation: I. Phonetic

characteristics. *Journal of Speech and Hearing Research*, *15*(1), 5–41.

British Society of Audiology. (2018). *Position Statement and Practice Guidance—Auditory*

Processing Disorder (APD). http://www.thebsa.org.uk/wp-

content/uploads/2018/02/Position-Statement-and-Practice-Guidance-APD-2018.pdf

Davis, M. H., & Johnsrude, I. S. (2003). Hierarchical processing in spoken language

comprehension. *The Journal of Neuroscience : The Official Journal of the Society for*

Neuroscience, *23*(8), 3423–3431. https://doi.org/23/8/3423 [pii]

Dawes, P., & Bishop, D. (2009). Auditory processing disorder in relation to developmental

disorders of language, communication and attention: A review and critique Research

Report. *International Journal of Language and Communication Disorders*, *44*(4), 440–

465. https://doi.org/10.1080/13682820902929073

Dillon, H., & Cameron, S. (2021). Separating the causes of listening difficulties in children. *Ear*

and Hearing.

Dillon, H., Cameron, S., Tomlin, D., & Glyde, H. (2014). Comments on "Factors influencing tests

of auditory processing: A perspective on current issues and relevant concerns" by Tony

Cacace and Dennis McFarland. *Journal of the American Academy of Audiology*, *25*(07),

699–703.

Dronkers, N. F., Wilkins, D. P., Van Valin, R. D., Redfern, B. B., & Jaeger, J. J. (2004). Lesion

analysis of the brain areas involved in language comprehension. *Cognition*, *92*(1–2),

145–177. https://doi.org/10.1016/j.cognition.2003.11.002

Edwards, B. (2020). *Emerging technologies, market segments, and MarkeTrak 10 insights in*

hearing health technology.

Golestani, N., Rosen, S., & Scott, S. K. (2009). Native-language benefit for understanding

speech-in-noise: The contribution of semantics. *Bilingualism: Language and Cognition*,

12(3), 385–392.

Hickok, G., & Poeppel, D. (2000). Towards a functional neuroanatomy of speech perception.

Trends in Cognitive Sciences, *4*(4), 131–138.

Hickok, G., & Poeppel, D. (2004). Dorsal and ventral streams: A framework for understanding

aspects of the functional anatomy of language. *Cognition*, *92*(1–2), 67–99.

Hickok, G., & Poeppel, D. (2007). The cortical organization of speech processing. *Nature*

Reviews Neuroscience, *8*(5), 393–402.

Holland, S., & Vannest, J. (2015). *Cincinnati MR Imaging of Neurodevelopment (C-MIND)*.

https://nda.nih.gov/edit_collection.html?id=2329

Hunter, L. L., Blankenship, C. M., Lin, L., Sloat, N. T., Perdew, A., Stewart, H., & Moore, D. R.

(2021). Peripheral auditory involvement in childhood listening difficulty. *Ear and Hearing*,

42(1), 29–41.

Hunter, L. L., Blankenship, C. M., Shinn-Cunningham, B., Hood, L., Motlagh-Zadeh, L., &

Moore, D. R. (2022). Brainstem auditory physiology in children with listening difficulties.

MedRxiv.

Iliadou, V., Chermak, G. D., Bamiou, D.-E., Rawool, V. W., Ptok, M., Purdy, S., Jutras, B.,

Moncrieff, D., Mattsson, T. S., Ferre, J. M., & others. (2018). Letter to the editor: An

affront to scientific inquiry re: Moore, DR (2018) editorial: Auditory processing disorder,

Ear Hear, 39, 617–620. *Ear and Hearing*, *39*(6), 1236–1242.

Iliadou, V., Kiese-Himmel, C., Bamiou, D.-E., Grech, H., Ptok, M., Chermak, G. D., Thai-Van,

H., Stokkereit Mattsson, T., & Musiek, F. E. (2019). Clinical expertise is core to an

evidence-based approach to auditory processing disorder: A reply to Neijenhuis et al.

2019. *Frontiers in Neurology*, *10*, 1096.

Jerger, J., & Musiek, F. E. (2000). Report of the Consensus Conference on the Diagnosis of

Auditory Processing Disorders in School-Aged Children. *Journal of the American*

Academy of Audiology, *11*(9), 467–474.

Kelley, K. S., & Littenberg, B. (2019). Comparison of Children's Double Dichotic Digits and

SCAN-3 Competing Words Free Recall Scores. *Journal of the American Academy of*

Audiology, *30*(05), 363–369.

Killion, M. C., Niquette, Patricia, A(Etymotic reseach Inc., 61 Martin lane, Elk Grove Village, I.

60007), Gudmundsen, Gail, L., Revit, L. J., & Banerjee, S. (2004). Development of a

quick speech-in-noise test for measuring signal-to-noise ratio loss in normal-hearing and

hearing impaired listeners. *J Acoust. Soc. Am*, *116*(4), 2395–2405.

https://doi.org/10.1121/1.1784440

Magimairaj, B. M., Nagaraj, N. K., Champlin, C. A., Thibodeau, L. K., Loeb, D. F., & Gillam, R.

B. (2021). Speech Perception in Noise Predicts Oral Narrative Comprehension in

Children With Developmental Language Disorder. *Frontiers in Psychology*, 4587.

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S.,

Donohue, M. R., Foran, W., Miller, R. L., Hendrickson, T. J., & others. (2022).

Reproducible brain-wide association studies require thousands of individuals. *Nature*,

603(7902), 654–660.

McFarland, D. J., & Cacace, A. T. (2009). Modality Specificity and Auditory Processing

Disorders. In D. J. Mcfarland & A. T. Cacace (Eds.), *Controversies in Central Auditory*

Processing Disorder (pp. 199–216). Plural Publishing, Inc.

McGarrigle, R., Munro, K. J., Dawes, P., Stewart, A. J., Moore, D. R., Barry, J. G., & Amitay, S.

(2014). Listening effort and fatigue: What exactly are we measuring? A British Society of

Audiology Cognition in Hearing Special Interest Group 'white paper'. *International*

Journal of Audiology, *53*(7), 433–445.

Moore, D. R. (2018). Auditory processing disorder (APD). *Ear and Hearing*, *39*(4), 617.

Moore, D. R., Edmondson-Jones, M., Dawes, P., Fortnum, H., McCormack, A., Pierzycki, R. H.,

& Munro, K. J. (2014). Relation between speech-in-noise threshold, hearing loss and

cognition from 40–69 years of age. *PloS One*, *9*(9), e107720.

Moore, D. R., Ferguson, M. A., Edmondson-Jones, A. M., Ratib, S., & Riley, A. (2010). Nature

of Auditory Processing Disorder in Children. *Pediatrics*, *126*(2), E382–E390.

https://doi.org/10.1542/peds.2009-2826

Moore, D. R., Sieswerda, S. L., Grainger, M. M., Bowling, A., Smith, N., Perdew, A., Eichert, S.,

Alston, S., Hilbert, L. W., Summers, L., & others. (2018). Referral and diagnosis of

developmental auditory processing disorder in a large, United States hospital-based

audiology service. *Journal of the American Academy of Audiology*, *29*(5), 364–377.

Neijenhuis, K., Campbell, N., Cromb, M., Luinge, M., Moore, D. R., Rosen, S., & de Wit, E.

(2019). An Evidence-based Perspective on 'Misconceptions' Regarding Pediatric

Auditory Processing Disorder. *Frontiers in Neurology*, *10*, 287.

Parente, F., Frascarelli, M., Mirigliani, A., Di Fabio, F., Biondi, M., & Colosimo, A. (2018).

Negative functional brain networks. *Brain Imaging and Behavior*, *12*(2), 467–476.

Petley, L., Hunter, L. L., Zadeh, L. M., Stewart, H. J., Sloat, N. T., Perdew, A., Lin, L., & Moore,

D. R. (2021). Listening Difficulties in Children with Normal Audiograms: Relation to

Hearing and Cognition. *Ear and Hearing*.

Pichora-Fuller, M. K., Kramer, S. E., Eckert, M. A., Edwards, B., Hornsby, B. W., Humes, L. E.,

Lemke, U., Lunner, T., Matthen, M., Mackersie, C. L., & others. (2016). Hearing

impairment and cognitive energy: The framework for understanding effortful listening

(FUEL). *Ear and Hearing*, *37*, 5S-27S.

Rees, N. S. (1973). Auditory processing factors in language disorders: The view from

Procrustes' bed. *Journal of Speech and Hearing Disorders*, *38*(3), 304–315.

Roebuck, H., & Barry, J. G. (2018). Parental perception of listening difficulties: An interaction

between weaknesses in language processing and ability to sustain attention. *Scientific*

Reports, *8*(1), 1–10.

Rönnberg, J., Holmer, E., & Rudner, M. (2019). Cognitive hearing science and ease of language

understanding. *International Journal of Audiology*, *58*(5), 247–261.

Rönnberg, J., Lunner, T., Zekveld, A., Sörqvist, P., Danielsson, H., Lyxell, B., Dahlström, Ö.,

Signoret, C., Stenfelt, S., Pichora-Fuller, M. K., & others. (2013). The Ease of Language

Understanding (ELU) model: Theoretical, empirical, and clinical advances. *Frontiers in*

Systems Neuroscience, *7*, 31.

Rudner, M., & Signoret, C. (2016). The role of working memory and executive function in

communication under adverse conditions. *Frontiers in Psychology*, *7*, 148.

Schiller, I. S., Remacle, A., Durieux, N., & Morsomme, D. (2022). Effects of Noise and a

Speaker's Impaired Voice Quality on Spoken Language Processing in School-Aged

Children: A Systematic Review and Meta-Analysis. *Journal of Speech, Language, and*

Hearing Research, *65*(1), 169–199.

Shannon, R. V., Zeng, F.-G., Kamath, V., Wygonski, J., & Ekelid, M. (1995). Speech recognition

with primarily temporal cues. *Science*, *270*(5234), 303–304.

Sharma, M., Purdy, S. C., & Humburg, P. (2019). Cluster analyses reveals subgroups of

children with suspected auditory processing disorders. *Frontiers in Psychology*, *10*,

2481.

Shinn-Cunningham, B. (2017). Cortical and Sensory Causes of Individual Differences in

Selective Attention Ability Among Listeners With Normal Hearing Thresholds. *Journal of*

Speech Language and Hearing Research, *60*(10), 2976.

https://doi.org/10.1044/2017_JSLHR-H-17-0080

Shinn-Cunningham, B. G. (2008). Object-based auditory and visual attention. *Trends in*

Cognitive Sciences, *12*(5), 182–186.

Smits, C., Goverts, S. T., & Festen, J. M. (2013). The digits-in-noise test: Assessing auditory

speech recognition abilities in noise. *The Journal of the Acoustical Society of America*,

133(3), 1693–1706.

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity Toolbox for

Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, *2*(3), 125–141.

https://doi.org/10.1089/brain.2012.0073

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-

scale automated synthesis of human functional neuroimaging data. *Nature Methods*,

8(8), 665–670.

830