Macronutrient mass intake explains ==differential== weight and fat loss in isocaloric diets

Francisco Arencibia-Albite[1]   Anssi H. Manninen[2]

[1]Universidad del Sagrado Corazón, Natural Sciences Department, PO BOX 12383, San Juan PR 00914-0383, Puerto Rico. Email: franciscom.arencibia@sagrado.edu

[2]Dominus Nutrition Oy, Ylipääntie 438, 92220 Raahe, Finland. Email: anssi@dominusnutrition.fi

Corresponding author: **Anssi H. Manninen, Director of Research & Development at Dominus Nutrition Oy.**

Summary

Currently, obesity treatment rests on the "calories-in, calories-out" rule, formally named the energy balance theory (EBT). It maintains that body weight increases as food calories are greater than expended calories but decreases when the opposite occurs; hence, weight stability is expected at energy balance meaning that over time energy-in equals energy-out.[1] It follows that dietary regimens with identical energy content should evoke similar amounts of weight and fat loss with only minor differences that emerge from diet's macronutrient composition[7], e.g., diet-induced glycogen depletion and water excretion. A vast collection of evidence shows, however, that low-carbohydrate diets typically result in much greater weight and fat loss than isocaloric low-fat diets.[7, 8, 9, 10, 11, 12] Furthermore, our recent mathematical analysis demonstrated that weight stability coincides with a persistent energy imbalance and not otherwise.[4] As an alternative, the mass balance model (MBM) was proposed that fitted weight loss data and explained the often superior weight loss evoked by low-carbohydrate diets versus low-fat diets.[4] Here, we expand on these observations by computationally contrasting the predictions of both models in two scenarios described in the literature[3, 16]: altering the diet's macronutrient composition while energy intake is kept at weight maintenance level; and the weight loss response as diet composition is changed under untreated type 1 diabetes. Our results indicate that MBM predictions are remarkably accurate while those of the EBT are clearly erroneous. These findings may represent the beginning of a paradigm shift in obesity research.

Keywords: obesity, weight loss, energy balance theory, mass balance model, macronutrients

List of abbreviations

MBM = mass balance model; MB = mass balance; EB = energy balance; EBT = energy balance theory; CIM = carbohydrate-insulin model; FFM = fat-free mass; FM = fat mass; EPM = energy-providing mass; VLCD = very-low-carbohydrate diet; KD = ketogenic diet; HCD = high-carbohydrate diet; ND = "normal diet"; BW = body weight; BF = body fat; EI = energy intake; EE = energy expenditure; T1D = type 1 diabetes; ECF = extra cellular fluid; Gly = glycogen; nEPM = non-energy-providing mass; AT = adaptive thermogenesis; PAL = physical activity level; RQ = respiratory quotient; GNG = gluconeogenesis; DNL = *de novo* lipogenesis; TG = triglycerides; HDL = high-density lipoprotein; LDL = low-density lipoprotein; HbA1c = Glycated hemoglobin; total-C = total cholesterol; MRI = magnetic resonance imaging; CT = computed tomography; NHANES = National Health and Nutrition Examination Survey; GI = glycemic index; GL = glycemic load; SGLTs = sodium-glucose transporters; GEBN = Global Energy Balance Network; MBNE = memory-based nutritional epidemiology.

Introduction

"The strongest and most evil spirits have so far advanced humanity the most: they have always rekindled the drowning passions – all ordered society puts the passion to sleep; they have always re-wakened the

sense on comparison, of contradiction, of joy in the new, the daring, and the untried; they force men to
meet the opinion with opinion, model with model [EBT *vs.* MBM]... The new is always *the evil*...".

– Friedrich Nietzsche, *The Gay Science* (1882)

Macronutrients – proteins, fats and carbohydrates – are an important part of a diet, and as such, have been
the subject of a great deal of discussion and controversy, especially among obesity researchers. According
to the energy balance theory (EBT), "[a] fundamental principle of nutrition and metabolism is that body
weight change is associated with an imbalance between the energy content of food eaten and energy
expended by the body to maintain life and to perform physical work."[1] Many current approaches to obesity
treatment are based on the EBT; however, neither behavioral prevention nor behavioral treatment is
generally successful in practice. Furthermore, this theory has major limitations for explaining obesity, as
reviewed by Wells and Siervo.[2] Weight stable obese individuals, for example, may experience substantial
changes in body weight as a reaction to isocaloric modifications in diet. In many instances these body mass
alterations are far greater than those expected from the macronutrient-induced changes in total body
water.[3] Thus, the EBT is rendered inadequate in these circumstances.

Arencibia-Albite[4] has recently used mathematical methods to investigate EBT's central claim:

As energy balance approaches zero body weight becomes stable.

The conclusion of this paper is highly polemical – to put it mildly – and possibly represents the beginning of
a paradigm shift:

Body weight stability coincides with a persistent energy imbalance which is negative under low-fat diets but
positive under low-carbohydrate diets.

This signifies, then, that the EBT is an incorrect theory, and also explains why in numerous well-controlled
studies negative energy balance measurements are frequently observed under a "normal diet" in weight
stable populations.[5, 6] These findings do not represent a violation of the First Law of Thermodynamics, as it
allows for any open system to express a positive or negative energy balance under a null mass change (**For**
further details, see Supplementary File 2).

As an alternative to the EBT, the mass balance model (MBM) of obesity was proposed. It maintains in
qualitative terms that *body weight fluctuations are ultimately dependent on the difference between daily*
nutrient mass intake and daily mass excretion (e.g., elimination of macronutrient oxidation products) and
NOT on energy imbalance. In its mathematical form the MBM fitted weight loss data from dietary
interventions of low-carbohydrate diets (LCDs) versus isocaloric low-fat diets (LFDs) leading to a simple and
convincing explanation for the often superior weight loss evoked by LCDs over LFDs (Figure 1).[4] This work
compares the MBM predictions against those derived from the EBT; and it is convincingly demonstrated
that in the experimental settings where the EBT makes erroneous forecasts the MBM results in remarkably
accurate predictions.

Figure 1. MBM simulation: LCD *vs.* LFD

A. Two hypothetical overweight 90kg adult males with identical body composition and weight maintenance energy intakes (2 750 kcal/day; 35% fat (F), 50% carbohydrate (C), 15% protein (P)) initiate two distinct isocaloric diets (2 000 kcal/day): low-fat diet (LFD; 20% F, 65% C, 15% P) *vs.* low-carbohydrate diet (LCD; 70% F, 15% C, 15% P). Consistent with published feeding data[7, 8, 9, 10, 11, 12], the MBM predicts that the LCD results in greater weight loss in contrast to the LFD. Model parameters are as follows: $BW_0 = 90$kg, $FM_0 = 22.5$kg, nEPM = 1.41kg, $F_{loss} = 0.030833$, PAL = 1.5, $VO_2 = 4.32$ L/[kg x day]. **B.** The figure depicts the body fat (BF) percentage changes that correspond to panel A. **C.** As shown, the LCD mass intake (m_{in}) is small relative to the eliminated mass (m_{out}) and so the net daily mass loss is large (i.e., $|m_{in} - m_{out}|$). In the LFD the net daily mass loss is, however, not as efficient since m_{in} cancels out a substantial fraction of m_{out} decelerating weight loss. LCDs are thus more effective in maximizing the net daily mass loss relative to LFDs and consequently the former manifest a substantially larger cumulative weight loss than the latter. **D.** Daily energy expenditure (EE) data from panel A. The horizontal line represents the daily energy intake.

112 According to the EBT, this graph suggests that the subject in the LFD should lose more weight than the one
113 in the LCD, which is not the case as illustrated in panel A. Energy imbalance is, therefore, not predicting the
114 weight loss outcome in this intervention.

Materials and Methods

117 As noted above, this study compares the MBM predictions against those derived from the EBT.

118 ***Mass Balance Model (MBM):***

119 For model derivation and details consult Arencibia-Albite[4] but briefly daily body weight (BW, in kg)
120 fluctuations are given by

$$BW(t) = M/R \; + \; \left(BW_0 \; - \; M/R\right)\left(1 \; - \; R\right)^t$$
$$M = EPM \; + \; nEPM$$
$$R = F_{loss} \; - \; \rho_{O_2} \cdot PAL \cdot VO_2$$

where t is time in days; M is the average daily mass intake defined as the sum of the energy-providing mass
(EPM; e.g., F, C and P) plus the non-energy-providing mass (nEPM; i.e., water, insoluble fiber, vitamins and
minerals); R is the average relative daily rate of mass excretion free of total daily O_2 uptake; BW_0 is the
initial body weight; F_{loss} is the average relative daily rate of mass excretion that includes of total daily O_2
uptake; $\rho_{O_2} = 1/770$ is O_2 density in kg/L at 27^0 C and 1 ATM; PAL is a dimensionless number that
represents the physical activity level computed as the ratio of total daily O_2 uptake over the total daily
resting O_2 consumption; and VO_2 is the specific daily resting O_2 uptake in L/(kg x day).

The EMP parameter is adjusted according to:

$$EPM = EI\left(x_C / \rho_C \; + \; x_F / \rho_F \; + \; x_P / \rho_P\right)$$

where EI is the energy intake; $0 \le x_i \le 1$ is the energy fraction from i = C, F , P; and is the energy density of
i with $\rho_C = 4.2\,kcal/g$, $\rho_F = 9.4\,kcal/g$, $\rho_C = 4.7\,kcal/g$. To simulate pancreatic β-cell death $x_C = 0$
as the bulk of ingested glucose is lost through urination.

Fat mass (FM) and fat free mass (FFM) alterations are given by:

$$FM(t) = 10.4W\left\{\frac{FM_0}{10.4}\exp\left[\frac{\left(M/R \; - \; BW_0\right)\left[1-\left(1-R\right)^t\right] \; + \; FM_0}{10.4}\right]\right\}$$
$$FFM(t) = BW(t) - FM(t)$$

where FM_0 is the initial fat mass and W is the lambert W function.

In MBM simulations daily energy expenditure (EE) can be estimated by

$$EE = \left(\text{kcal per } O_2L\right) \cdot PAL \cdot VO_2 \cdot BW$$

The first term at the right-hand side can approximated with the diet's macronutrient composition and the
following Weir formula[13]

$$\text{kcal per } O_2L = 5.047x_C + 4.463x_P + 4.735x_F.$$

Energy Balance Theory (EBT):

The quantitative form of the EBT use in this study is that of the United States National Institute of Health
Body Weight Planner developed by Hall et al.[14] A succinct model description is given below.

Glycogen (G), extracellular fluid (ECF), body fat (F_b), lean tissue (L) and adaptive thermogenesis (AT) are
modeled by following system of equations

148
$$\frac{dG}{dt} = \left(CI - CI_b \left(G/G_0\right)^2\right)/\rho_G$$

$$\frac{dECF}{dt} = \frac{1}{[Na]}\left(\Delta Na_{diet} - \xi_{Na}\left(ECF - ECF_0\right) - \xi_{CI}\left(1 - CI/CI_b\right)\right)$$

$$\frac{dF_b}{dt} = \left[\frac{F_b/\rho_F}{10.4\rho_L/\rho_F + F_b}\right]\left(EI - EE - CI + CI_b\left(G/G_0\right)^2\right)$$

$$\frac{dL}{dt} = \left[\frac{10.4/\rho_F}{10.4\rho_L/\rho_F + F_b}\right]\left(EI - EE - CI + CI_b\left(G/G_0\right)^2\right)$$

$$\frac{dAT}{dt} = \left(\beta_{AT}\Delta EI - AT\right)/\tau_{AT}$$

where CI_b is the carbohydrate energy intake (CI) at energy balance; G_0 is the initial body glycogen
amount; ρ_G is the glycogen energy density (4.2 kcal/g); [Na] is the extracellular Na concentration (3.22
mg/ml); ξ_{Na} is a ECF Na excretion constant (3 000 mg/[ml x day]); ξ_{CI} Cl dependent Na excretion constant
(4 000 mg/[ml x day]); ρ_L Is the energy density of lean tissue (1.815kcal/g); τ_{AT} is the AT time constant (14
days); and $\beta_{AT} = 0.14$ is the AT coefficient.

EE is defined as

155
$$EE = \frac{\left[\begin{array}{l} K + \gamma_{F_b}F_b + \gamma_L L + \left[\left(1 - \beta_{TEF}\right) \cdot PAL - 1\right]RMR + \beta_{TEF}\Delta EI + AT \\ + \left(EI - CI + CI_b\left(G/G_0\right)^2\right)\left(\dfrac{10.4\eta_L/\rho_F}{10.4\rho_L/\rho_F + F_b} + \dfrac{\eta_{F_b}F_b/\rho_F}{10.4\rho_L/\rho_F + F_b}\right) \end{array}\right]}{1 + \dfrac{10.4\eta_L/\rho_F}{10.4\rho_L/\rho_F + F_b} + \dfrac{\eta_{F_b}F_b/\rho_F}{10.4\rho_L/\rho_F + F_b}}$$

where K is arbitrary constant determined to achieved energy balance at BW_0; $\gamma_{F_b} = 3.1\,kcal/[kg \cdot day]$ is
the specific body fat metabolic rate coefficient; $\gamma_L = 21.99\,kcal/[kg \cdot day]$ is the specific lean tissue
metabolic rate coefficient; $\beta_{TEF} = 0.1$ is the thermic effect of feeding coefficient;

$RMR = 19.7(BW - F_b) + 413$ is Mifflin et al.[15] resting metabolic rate formula;

$\eta_L = 229.446\,kcal/kg$ lean tissue synthesis efficacy; and $\eta_{F_b} = 179.254\,kcal/kg$ fat tissue synthesis

efficacy.

The body weight at time t is obtained by the following sum

$BW(t) = F_b(t) + L(t) + 1.0027G(t) + ECF(t)$

To simulate pancreatic β-cell death CI = 0 as the majority of ingested glucose is lost through urination. The

system of differential equations was solved using the fourth order Runge-Kutta method with a time step of

1 day.

Results

The Figure 2 A uses the MBM to simulate the effect of exchanging a high-carbohydrate diet (HCD) for an

isocaloric very-low-carbohydrate diet (VLCD) in a weight-stable 70kg individual. Under the HCD (day 150-

400), body weight gradually increases to 73kg. However, after beginning the isocaloric VLCD (day 400),

body weight decreases towards a steady value of 67.5kg. Figure 2 B repeats the simulation in panel A, but

with Hall *et al.*[14] model of the EBT. The latter indicates, in general, that isocaloric perturbations only elicit

non-significant alterations in body weight that are mostly a consequence of changes in extracellular fluid

(ECF) and stored glycogen (Gly, Figure 2 C1). In such cases, therefore, changes in body composition are

expected to be nearly undetectable (Figure 2 C2). For further details, see the figure legend.

An interesting aspect of the MBM is its capacity to account for the degree of weight loss observed at the

onset of type 1 diabetes (T1D). Before the discovery of insulin, a common dietary treatment of T1D was a

very-low-carbohydrate diet/high-fat diet which slowed down and reduced the excessive weight loss while

alleviating the classical symptoms of polyuria, polyphagia and polydipsia [16]. The Figure 2 D1 shows that

the MBM explains the weight loss associated with the onset of T1D and also makes predictions consistent

with the diabetes treatment during the pre-insulin era. The EBT, however, predicts a substantial weight loss

at T1D onset even if the pre-onset diet is a VLCD (Figure 2 D2). For further details, see the figure legend.

FIGURE 2 Predictions under isocaloric perturbations: EBT *vs.* MBM

A. MBM response to exchanging a HCD for an isocaloric VLCD. BW is stable at 70 kg with EI = 2 000 kcal
during 150 days (control: 50% F, 35% C, 15% P). The daily carbohydrate mass intake (CMI) is 167g. At day
150 EI = 2 000 kcal but CMI has increased by 143g as the control diet is exchange for a HCD (20% F, 65% C,
15% P). On day 400, EI = 2 000 kcal but the diet is exchange again for a VLCD (75% F, 10% C, 15% P). Here

CMI decreases by 262g. Model parameters are as follows: BW_0 = 70kg, FM_0 = 14kg, nEPM = 1.5kg, F_{loss} =
0.034942857, PAL = 1.5, VO_2 = 4.464 L/[kg x day]. **B.** EBT response to the same perturbations as in panel A.
Initial conditions are as follows: K = 547.2925, BW(0) = 70kg, $F_b(0)$ = 14kg, L(0) = 36.012kg, ECF(0)=19.486kg,
AT(0) = 0, G(0) = 0.5kg, PAL =1.5 **C1.** According to the EBT, majority of the weight change under an isocaloric
perturbations result from changes in extracellular fluid (ECF) and stored glycogen (Gly) that stabilize within
few days. **C2.** MBM predicts that BF evolves in parallel to the weight change direction. In contrast, the EBT
predicts that, under isocaloric perturbations, BF evolves according to the amount of fat intake (FI). **D1.**
MBM predicts that the weight loss magnitude evoked by pancreatic β-cell death is dependent on the diet's
macronutrient composition. To simulate pancreatic β-cell death, we set CI = 0, as majority of the ingested
glucose is lost through urination. One of the mechanisms that has evolved to maintain normoglycemia is
the re-uptake (by sodium-glucose transporters [SGLTs]) of glucose that is excreted in the urine but in
untreated type I diabetes very high blood glucose concentration clearly over-power these SGLTs. Model
parameters are as in A. **D2.** The EBT predicts that changes in the diet's macronutrient distribution will elicit
measurable changes in the rate and magnitude of weight loss after β-cell death but not as potent as those
predicted by the MBM. Initial conditions are as in B.

Kong et al.[3] have recently published a well-controlled feeding study that allows to test the validity of the
predictions made by the MBM and EBT. Their young female subjects (age: 21± standard deviation [SD]: 3.7
years, weight: 65.5 ± SD: 7.7 kg, body mass index: 24.9 ± SD: 2.7 kg/m^2) were weight-stable for 4 weeks
under a normal diet (ND: 1 967kacl ± SD: 362kcal ; F 44% ± SD: 7.6%, C 39.6% ± SD: 5.8%, 15.4%± SD:
3.3%) and then switched to an isocaloric ketogenic diet (KD 1 817kacl ± SD: 285kcal; F 69% ± SD: 5.4%, C
9.2% ± SD: 4.8%, 21.9%± SD: 3.4%) for another 4 weeks leading to a substantially reduced body weight (−
2.9 kg) and body fat percentage (− 2.0%). According to the authors, "[t]he findings of this study do not
seem to support the idea that the weight-loss effect of KD is due to reduced appetite [and thus reduced
energy intake]."
216
Figure 3 A1, A2 and A3 simulate Kong et al.[3] data using the Hall et al.[14] model of the EBT. Here the KD
decreases body weight by 1.71Kg (Figure 3 A1) which is mainly a result of changes in ECF and stored
glycogen (Figure 3 A2). In such case, the EBT predicts a slight increase in body fat percentage (Figure 3 A2).
Figure 3 B1 simulates Kong et al.[3] data using the MBM. As illustrated, this model predicts a much larger
drop in body weight (−3.05kg) under KD than the EBT. Contrary to the EBT, however, the MBM predicts a
drop in body fat percentage of about 1.6%. This signifies, therefore, that at the MBM leads to more
accurate predictions in than those made by the EBT.
224
225
226

FIGURE 3. The simulation of Kong *et al*. data: EBT *vs*. MBM

A1. The figure utilizes the EBT to simulate Kong *et al*.[3] data. During the ND period (days 0-30) energy balance (EB, red curve) is zero. After day 30 EB becomes negative under the KD resulting in a cumulative weight loss of 1.71kg at day 60. Initial conditions are as follows: K = 892.7146, BW(0) = 65.5kg, F_b(0) = 22.925kg, L(0) = 25.698kg, ECF(0) = 16.375kg, AT(0) = 0, G(0) = 0.5kg, PAL =1.5. **A2.** The EBT predicts that – although the KD results in *weight* loss – BF increases since FI has been augmented by 41g. **A3.** During the ND total glycogen is 500g but after the KD this amount has dropped by 280.25g. ECF has also decreased by

1.076kg. These quantities add to 1.35525kg = 0.28025kg + 1.076kg. Consequently, of the 1.71kg weight loss
in A1, 0.35475kg = 1.71kg − 1.35525kg are from other mass sources (0.2436 kg fat mass + 0.11115 kg fat-
free mass = 0.35475kg). According to the EBT, fat mass has in fact decreased; however, only by slight
amount since FI has been augmented during the KD. Consequently, as the decline in fat-free mass is much
larger than that of fat mass, BF increases as in A2. **B1.** The figure utilizes the MBM to simulate Kong *et al.*[3]
data. During the ND period (days 0-30) mass balance (MB, red curve) is zero. After day 30 MB becomes
negative under the KD resulting in a cumulative weight loss of 3.05kg at day 60. Model parameters are as
follows: BW_0 = 65.5kg, FM_0 = 22.925kg, $nEPM_{ND}$ = 1kg, $nEPM_{KD}$ =0.96kg, F_{loss}= 0.029078, PAL = 1.5, VO_2 =
4.32 L/[kg x day] **B2.** According to the MBM, of the 3.05kg of weight loss 2.07kg came from fat and 0.98kg
from fat-free mass. Even though FI has increased the decline in fat-free mass is much smaller than that of
fat mass and thus BF decreases as shown.

Discussion

According to the EBT, at energy balance, changes in macronutrient composition elicit non-substantial
changes in body weight. These alterations are assumed to be secondary to changes in ECF and glycogen
that follow from adjustments in the diet's sodium content and carbohydrate intake.[1, 7] The EBT also claims
that body composition alterations, under these circumstances, are expected to be small but to occur in
parallel to the amount of dietary fat intake (see Figure 2 C2). Consequently, from the EBT perspective, the
MBM simulations (see Figure 2 A and C2) are unrealistic, as body weight and body fat alterations are much
greater than those predicted when energy balance is assumed to be present. If such an argument is
definitive, then the accuracy of EBT predictions should always exceed those made by the MBM. Figure 3
illustrates, however, that this is *clearly* not the case.

Moreover, the EBT implicitly asserts that blood leptin concentration should be minimally affected when
arbitrary subject switches from a weight-maintenance standard diet to an isocaloric ketogenic diet since, as
argued by this theory, under such conditions fat mass decline is insignificant (see Figure 3 A2). In contrast,
the MBM predicts a substantial drop in leptin levels since this model indicates that this type of isocaloric
exchange results in a substantial drop in fat mass. Indeed, a recent well-controlled feeding trial by Kong *et*
al.[3] showed that the interchange of a weight-preserving normal diet for an isocaloric ketogenic diet evokes
a significant reduction in leptin levels.

In addition, the MBM simulations on pancreatic β-cell death are consistent with a review of case histories
from the pre-insulin era illustrating that VLCDs could result in some weight gain after the onset of T1D.[16]
Elliot Proctor Joslin, for example, was the first United States medical doctor that specialized in diabetes
treatment during this era. Although anecdotal, one of his diabetic patients, Mary H., reportedly gained
nearly 3kg of body weight while consuming a diet containing solely protein and fat. The EBT simulations, in
contrast, do not predict such therapeutic effect.

Numerous examples in the weight management literature show that the amount of weight loss is far
greater in low-carbohydrate diets (LCDs) compared to isocaloric low-fat diets (LFDs). [7, 8, 9, 10, 11, 12] According
to the EBT, this can only occur if the energy expenditure under LCDs is larger than that in LFDs. However, in
many of these dietary interventions non-significant differences are found between the energy expenditures
of these diets.[7, 12, 17, 18, 19] Researchers that subscribe to the EBT, therefore, argue that in such cases the
most likely explanation of the observed superior weight loss is the result of energy intake underreporting

by low-fat dieters.[12] This claim has encountered minimal opposition since it is widely accepted that the
majority of the energy intake recording devices (e.g., self-reported food records) are biased toward
underestimation.[5, 6] (**For further discussion, see Supplementary File 1**). Two alternative models, however,
may explain the apparent weight loss advantage of LCDs over isocaloric LFDs: the carbohydrate-insulin
model (CIM) of obesity[20] and the MBM[4] (**See also Supplementary File 3**).

The CIM postulates that high-carbohydrate intake elevates insulin levels leading to the activation of
complex neuroendocrine responses that drive body fat deposition, increase appetite and decrease energy
expenditure[17]; which, according to the EBT, explains the persistent weight gain over time observed in obese
subjects. Conversely, LCDs, by significantly decreasing circulating insulin levels, should increase energy
expenditure by augmenting fat oxidation, which – as argued by CIM advocates – accounts for the greater
and faster weight loss observed in feeding trails of LCDs *vs.* isocaloric LFDs. The above claims are open to
discussion, however, as evidence shows that obese individuals manifest a highly elevated energy
expenditure relative to normal weight subjects[21], plus, as already mentioned, the predominant evidence
indicates that no significant energy expenditures differences exist between diets. [7, 12, 17, 18, 19] A recent meta-
analysis by Ludwig *et al.*[22] suggests that an adaptation period of at least 14 days may be required in order
to LCDs to elevate energy expenditure. If this is the case, then one would expect a slow decay of the
respiratory quotient (RQ) toward 0.71 since this would reflect the gradual metabolic dependence on fat as
the main energy fuel. In ketogenic diets, however, the RQ reaches a steady state within the first week and
continuous to be stable for at least three more weeks.[17] Taken together, it seems that the LCD-enhanced
weight loss is unlikely to be a consequence of an augmented energy expenditure as proposed by the CIM.

In contrast, the MBM describes body weight fluctuations as a ***mass imbalance problem***. Specifically, when
macronutrient mass intake exceeds the excretion of macronutrient oxidation products body weight
increases and *vice versa*. Body weight stability is, hence, expected as over time the average consumed mass
equals the average eliminated mass. Our model arrives from the following long-standing observations:

1. The physiological/biochemical activity that decreases body weight is the excretion of oxidation
products and not energy expenditure.[23] This is exemplified in the oxidation of a "general" triglyceride
molecule:

The mass entering this reaction is (in g/mol)

whereas mass exiting the reaction is only present in the reaction products

and not in the dissipated heat (i.e., Calories). Therefore, body mass decreases as the body excretes or
eliminates oxidation products but not as consequence of the heat content in the energy expenditure.

2. The food property that has the capacity to increases body weight is its macronutrient mass and not its
potential nutritional energy. For instance, the absorption and retention within body cells of 1 gram of fat,
carbohydrate or protein will increase body mass by exactly 1 gram. This observation is independent of the
potential caloric content ascribe to the macronutrient; according to the Law of Conservation of Mass, the

absorbed macronutrient mass cannot be destroyed and, thus, it will contribute to total body mass as long
as it remains within the body. Such contribution ends, however, when the macronutrient is eliminated from
the body either as products of metabolic oxidation (e.g., H_2O and CO_2) or in other forms (e.g., shedding of
dead skin cells fill with keratin protein).

3. _From points 1 and 2 it follows that body weight fluctuations can be described by the difference_
between daily mass intake and daily mass excretion. Although at first sight this statement may seem not
to have far-reaching consequences, Arencibia-Albite[4] has shown that when translated into mathematical
form it fits body weight and fat mass data from isocaloric diet trails of LCDs _vs_. LFDs, and also results in
predictions that are not evident from the qualitative inspection of points 1, 2 and 3. This work further
substantiates such predictions and demonstrates that MBM-based simulations result in highly realistic
forecasts in settings where the EBT-based simulations collapse.

The energy picture of body weight dynamics explains body fat fluctuations as the balance between fat
intake and fat oxidation. The EBT cannot foresee, therefore, that a LCD may lead to more fat loss than an
isocaloric LFD since the fat balance in the latter appears to be more negative than in the former.[7] A
multitude of feeding studies[3, 9,11,12] suggest, however, this is not the case as the contrary is a more frequent
observation. In the mass picture, in contrast, body fat fluctuations are not only dependent on fat intake and
oxidation but other routes contribute as well to fat loss. In LCDs, for instance, the elevated fatty acid
oxidation leads, in hepatocytes, to high cytoplasmic HMG-CoA levels which are rapidly reduced to
mevalonic acid for cholesterol production. The characteristic excess cholesterol of LCDs[24], in turn, is then
eliminated in in feces as bile acids. Additionally, in LCDs, carbon atoms from fatty acid breakdown circulate
in the blood stream as high acetoacetate levels which can exit the body either in the urine or through
breathing when spontaneously decarboxylated to acetone. The afore mentioned processes may thus be
responsible for the greater fat loss in LCDs _vs_. isocaloric LFD even if fat oxidation is greater in LFDs as
suggested by short-term (e.g., 6 days) studies.7 Consequently, in the light of this theoretical study, the
above argument suggests that the variable that better accounts for body fat fluctuations is mass and not
energy.

Some readers may wonder why we completely ignored almost any _epidemiological_ data as it relates to
obesity and energy intake and – thus – to the debate of relative merits of the EBT _vs. the MBM. Memory-
based nutritional epidemiology (MBNE) supposedly examines the role of nutrition in the etiology of
diseases, including obesity. Although there are certain strong defenders of this dubious branch of
epidemiology (i.e., nutritional epidemiology people at Harvard School of Public Health), we tend to agree
with Ioannidis that this branch needs "radical reform". [25] In fact, almost all central assumptions coming
from the MBNE field have been refuted in rigorously controlled feeding trials.

Conclusions

Implicit in the EBT discourse is that body fat accumulates only if fat intake exceeds fat oxidation, even
under dietary energy restriction, which explains why this theory argues in favor of LFDs as the optimal fat
loss dietary treatment.[7] Such point of view is clearly captured in Figure 2 C2 and Figure 3 A2; however,
numerous examples in experimental literature indicates the opposite. The MBM, on the other hand, is
highly consistent with these data suggesting that the relevant variable for weight fluctuation is mass and
not energy. Our model by its _razor-sharp_ yet _simple_ logic, seems to be perfectly able to account for the
deferential weight loss among isocaloric diets. The obesity community is trying to explain such

"mysterious" findings by invoking complex signaling pathways that completely ignore the *fundamental rule* in science: the Occam's razor (*novacula Occami*), i.e., the simplest explanation is usually the right one. More specifically, "entities should not be multiplied without necessity", a notion attributed to English Franciscan friar William of Ockham. Nevertheless, we are certainly not claiming that our model is definitive (or downplaying the emergent and promising therapeutic value of LCDs), but rather to stimulate researchers around the world to acknowledge that the widely accepted EBT is incorrect.

The daily intake of macronutrient mass is, obviously, influenced by the ever-present interplay between the environment and genes; thus, it is important to realize that the MBM makes no claims regarding the *behavioral* aspects of obesity. Certainly, food processing, distribution, marketing, education, etc. plays a big role in current obesity epidemics.

"We make [biological] modeling because it helps us in our attempted understanding of the intricate system. It breaks down the set of objects (e.g., macronutrients] and phenomenon too complex [e.g., obesity] to be grasped in their entirties into smaller realms that can be dealt one by one [as was done here by the MBM simulations]. There is nothing objective "true" about such models; the only proper criterion of value is their usefulness."

—Isaac Asimow, *The Human Body: It's Structure and Operation* (1963); cited in *Engineering Physiology: Bases of Human Factors Engineering/Ergonomics*, Springer Nature, 2020.

Authors contribution

Study concept and designing: Manninen, Arencibia-Albite

Acquisition of data: Manninen, Arencibia-Albite

Statistical analysis and interpretation of data: Manninen, Arencibia-Albite

Drafting the manuscript: Manninen, Arencibia-Albite

Critical revision of the manuscript for important intellectual content: Manninen, Arencibia-Albite

Administrative, technical or material support: Manninen, Arencibia-Albite.

Study supervision: N/A.

Acknowledgements

This has been a time-consuming and cliff-hanging research project with both brains working in overdrive almost around a clock. Fortunately, we are now about to close the 1st chapter of our MBM research project.

394 We would like to thank our families for support and care; our numerous colleagues for stimulating
395 discussions; and *medRxiv* for providing excellent customer service.

Funding sources

398 This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-
399 profit sectors.

Declaration of competing interests

402 The authors declare that they have no competing interests.

References

1. Hall K.D. Heymsfield S.B. Kemnitz J.W. Klein S. Schoeller D.A. Speakman J.R. Energy balance and its
components: implications for body weight regulation. Am J Clin Nutr. 2012; 95: 989-994.

2. Wells J.C.K. Siervo M. Obesity and energy balance: is the tail wagging the dog? Eur J Clin Nutr. 2011; 65:
1173–1189.

3. Kong Z, Sun S, Shi Q, Zhang H, Tong TK, Nie J. Short-Term Ketogenic Diet Improves Abdominal Obesity in
Overweight/Obese Chinese Young Females. Front Physiol. 2020; 11:856.

4. Arencibia-Albite F. Serious analytical inconsistencies challenge the validity of the energy balance theory.
Heliyon. 2020; 6(7):e04204.

5. Winkler JT. The fundamental flaw in obesity research. Obesity Reviews. 2005; 6:199–202.

6. Dhurandhar NV, Schoeller DA, Brown AW, Heymsfield SB, Thomas D, Sørensen TIA, et al. Energy balance
measurement: when something is not better than nothing. International Journal of Obesity. 2014; 39:
1109–1113.

7. Hall KD, Bemis T, Brychta TR, Chen KY, Courville A, Crayner EJ, et al. Calorie for Calorie, Dietary Fat
Restriction Results in More Body Fat Loss than Carbohydrate Restriction in People with Obesity. Cell
Metabolism. 2015; 22: 531.

8. Lean M, Han T, Prvan T, Richmond P, Avenell A. Weight loss with high and low carbohydrate 1200 kcal
diets in free living women. European Journal of Clinical Nutrition. 1997; **51**: 243–248.

9. Bazzano LA, Tian H, Reynolds K, Yao L, Bunol C, Liu Y, et al. Effects of Low-Carbohydrate and Low-Fat
Diets: a randomized trail. Annals of Internal Medicine. 2014; 161: 309.

10. Samaha FF, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, et al.

A Low-Carbohydrate as Compared with a Low-Fat Diet in Severe Obesity

New England Journal of Medicine. 2003; 348: 2074–2081.

11. Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA. A Randomized Trial Comparing a Very Low Carbohydrate
Diet and a Calorie-Restricted Low Fat Diet on Body Weight and Cardiovascular Risk Factors in Healthy
Women. The Journal of Clinical Endocrinology & Metabolism. 2003; 88: 1617–1623.

12. Brehm BJ, Spang SE, Lattin BL, Seeley RJ, Daniels RS, D'Alessio DA. The Role of Energy Expenditure in the
Differential Weight Loss in Obese Women on Low-Fat and Low-Carbohydrate Diets. The Journal of Clinical
Endocrinology & Metabolism. 2005; 90: 1475–1482.

13. WEIR JB. New methods for calculating metabolic rate with special reference to protein metabolism. J
Physiol. 1949;109(1-2):1-9.

14. Hall KD, Sacks G, Chandramohan D, Chow CC, Wang YC, Gortmaker SL, Swinburn BA. Quantification of
the effect of energy imbalance on bodyweight. Lancet. 2011;378(9793):826-37.

15. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO. A new predictive equation for resting
energy expenditure in healthy individuals. Am J Clin Nutr. 1990; 51(2):241-7.

16. Westman E.C. Yancy W.S. Humphreys M. Dietary treatment of diabetes mellitus in the pre-insulin era
(1914-1922). Perspect Biol Med. 2006; **49**: 77-83.

17. Hall KD, Chen KY, Guo J, Lam YY, Leibel RL, Mayer LE, Reitman ML, Rosenbaum M, Smith SR, Walsh BT,
Ravussin E. Energy expenditure and body composition changes after an isocaloric ketogenic diet in
overweight and obese men. Am J Clin Nutr. 2016;104(2):324-33.

18. Hall KD, Guo J. Obesity Energetics: Body Weight Regulation and the Effects of Diet Composition.

Gastroenterology. 2017; 152(7):1718-1727.e3.

19. Hall KD, Guo J, Speakman JR. Do low-carbohydrate diets increase energy expenditure? Int J Obes (Lond).
2019; 43(12):2350-2354.

20. Ludwig DS, Ebbeling CB. The Carbohydrate-Insulin Model of Obesity: Beyond "Calories In, Calories Out".

JAMA Intern Med. 2018; 178(8):1098-1103.

21. Das SK, Saltzman E, McCrory MA, Hsu LK, Shikora SA, Dolnikowski G, Kehayias JJ, Roberts SB. Energy
expenditure is very high in extremely obese women. J Nutr. 2004;134(6):1412-6.

22. Ludwig DS, Dickinson SL, Henschel B, Ebbeling CB, Allison DB. Do Lower-Carbohydrate Diets Increase
Total Energy Expenditure? An Updated and Reanalyzed Meta-Analysis of 29 Controlled-Feeding Studies. J
Nutr. 2020:nxaa350.

23. Meerman R, Brown AJ. When somebody loses weight, where does the fat go? BMJ. 2014;349:g7257.

24. Retterstøl K, Svendsen M, Narverud I, Holven KB. Effect of low carbohydrate high fat diet on LDL
cholesterol and gene expression in normal-weight, young adults: A randomized controlled study.
Atherosclerosis. 2018; 279:52-61.

25. Ioannidis JPA. The Challenge of Reforming Nutritional Epidemiologic Research. JAMA. 2018 Sep
11;320(10):969-970. doi: 10.1001/jama.2018.11025.

"Macronutrient mass intake explains deferential weight and fat loss in isocaloric diets"

SUPPLEMENTARY FILE 1

Additional discussion regarding free-living human feeding studies with special reference to Kong *et al*.

As recently discussed by Hall [1], feeding trials have their own challenges, including – but certainly not
limited to – the impracticality and difficulty of randomizing large numbers of free-living subjects to eat
different diets for a long period time while ensuring very-high levels of adherence throughout. Although so-
called domiciled feeding studies [1] can certainly provide important insights, their "artificial" environment
may seriously limit generalizability and applications to free-living individuals, not to mention the fact that
this type of studies have become prohibitively expensive to conduct in the United States since the National
Institutes of Health (NIH) ceased directly funding Clinical Research Centers [1]. Consequently, well-
controlled randomized free-living feeding studies remain the gold standard when it comes to proving
causality in the areas of nutrition and metabolism.

There are always potential problems with any kind of dietary records; although we know that these records
have relatively high error rates, they can be very meaningful – in terms of causality – if *1)* the differences in
the study arms are large enough; and *2)* there are indicators of very good adherence (e.g., the presence of
ketone bodies*). A Very-low-carbohydrate diet always win in a well-controlled face-off with an isocaloric*
high-carbohydrate diet (**see the Supplementary File 1 Figure 1**). *If such a response is not observed, then it is*
simply not a well-controlled study, as alternative results would indicate a violation of the Law of
Conversation of Mass (**see Supplementary File 2**).

Researchers guided by the erroneous EBT, however, tend to automatically assume that the subjects on the
very-low-carb arm must have overreported what they actually ate, or the people of the high-carb arm must
have underreported what they ate, or both. *How likely is that the very-low-carb arms, time after time,*
substantially overreport dietary records, whereas the high-carb arms consistently underreport dietary
records? [2]

To test the predictive power of both the MBM and the EBT, we decided to utilize the recent free-living
feeding data by Kong *et al.* (**see Figures A1, A2, A3, B1, and B2 in the main body of the manuscript**). Their
young female subjects were weight-stable for 4 weeks under a "normal diet" and then switched to an
isocaloric ketogenic diet for another 4 weeks leading to a substantially reduced body weight (− 2.9 kg) and
body fat percentage (− 2.0%), consistent with the MBM.

To assure subjects' adherence to ketogenic diet (KD), Kong *et al.* required that:

• The subjects measure urinary ketones every day (early morning or after dinner) and record 3-day
food diaries (2 weekdays and 1 weekend day) during the experimental period.

• 3-day food diaries were kept by all subjects for 8 weeks.

• All subjects were given in advance "thorough instructions" on how to estimate portion sizes and
record food/beverages intake on food composition tables.

• Subjects were asked to report to the laboratory every week to assess changes in body weight and
hand in the logbook with dietary records.

• Energy and macronutrient distribution were calculated by the same dietician using the nutrition
analysis and management system; diet compliance was evaluated based on the results of the
urinary ketones and food diaries, and subjects received follow-up dietary advice and counseling
individually from the dietician.

In summary, we believe it is safe to conclude that the energy intake values in a report by Kong et al. are
about as reliable as it gets. Of course, one can claim – and some certainly do – that such free-living feeding

studies are not well-controlled. If we follow such criteria, how many "well-controlled" studies we have on
dietary supplements, or pharmaceuticals, or other interventions lasting longer than few weeks? Zero.

Already in 1971, Young *et al.* [3] compared 3 diets that contained the same amount of calories (1800
kcal/d) and protein (115g/d) but that differed in carbohydrate content. After 9 weeks on the 30g, 60g, and
104g carbohydrate diets, weight loss was 16.2kg, 12.8kg, and 11.9kg and fat accounted for 95%, 84%, and
75% of the weight loss, respectively. Thus, the authors concluded, "[w]eight loss, fat loss, and percent of
weight loss as fat appeared to be inversely related to the level of carbohydrate in the isocaloric, isoprotein
diets", consistent with the MBM (When the energy fraction from dietary fat *increases*, while energy intake
is clamped, mass intake *decreases* due to the significantly higher energy density of fat in contrast to other
substrates). And since these authors were also guided by the erroneous EBT, "[n]o adequate explanation
can be given for weight loss differences." To our knowledge, no one has found flaws in the trial by Young *et*
al. It is worth noting that they utilized underwater weighing (i.e., hydrodensitometry) to determine body
composition, which is more accurate than other widely available methods of body composition testing.
When performed properly, underwater weighing can be accurate to 1.8 to 2.8% compared to the state-of-
the-art methods (e.g., magnetic resonance imaging [MRI], computed tomography [CT]).

Modern studies have reported similar findings. For example, a randomized, balanced, free living feeding
study by Volek *et al.* [4] compared the effects of isocaloric and energy-restricted very-low-carbohydrate
(ketogenic) and low-fat diets on weight loss and body composition in overweight and obese men and
women. Despite a significantly greater energy intake with the ketogenic diet than with the low-fat diet
(1855 kcal/d *vs.* 1562 kcal/d), both the between- and within-group comparisons revealed a quite distinct
weight and fat loss advantage of a ketogenic diet (**See Supplementary File 1 Figure**
2).

[FIGURE LEGEND STARTS] Supplementary File 1 Figure 2. A 2004 well-controlled free-living feeding trial

published by Volek *et al*. [4]. This interesting figure illustrates Individual differences between total fat loss

on a very low-carbohydrate ketogenic diet (VLCK or VLCKD) minus total fat loss on a low-fat (LF) diet for

<u>each</u> person. Positive numbers reflect greater fat loss on the VLCK, whereas negative numbers indicate

greater fat loss on the LF diet. Red circles = order of diets VLCK then LF. Blue diamonds = order of diets LF

then VLCK. *Like all such studies, the report by Volek et al. demonstrate that there is always "noise" in any*

complex biological variable such as food ingestion. Based on this data, it seems that a ketogenic diet (VLCKD

or VLCK) is more palatable to men than women, as five men showed a >10 pounds difference in weight loss.

What is clear is that every time the human body oxidizes, for example, 1 g of glucose, the heat released by

the body (~4kcals) simply can NOT change as a function of one's biological sex. Thus, the mass balance

model (MBM) applies identically to both (or all) biological sexes. There are well-established differences in

food intake regulation between biological sexes; for example, food cravings – and especially carbohydrate

cravings – may change significantly during the menstrual cycle. However, *the MBM makes no claims*

regarding the <u>behavioral</u> aspects of obesity whatsoever. Thus, further discussion of this topic is clearly

outside of scoop of this paper. Reproduced, with permission, from [4]. LF = low-fat diet. **[FIGURE LEGEND**

STARTS]

Since it is widely – and mistakenly – assumed that the fundamental cause of obesity is an energy imbalance

between calories consumed and calories expended, it was logical to recommend a low-fat diet for weight

management and obesity treatment. However, data from the National Health and Nutrition Examination

Surveys (NHANES) indicate a large increase in carbohydrate intake as a major contributor to "caloric

excess" (in reality, macronutrient mass excess) in the United States from 1974 to 2000, whereas the

absolute amount of fat decreased for men during this period and showed only a slight increase for women

(**See Supplementary File 1 Figure 3**). And it was during this very period when the incidence of obesity

raised to its current epidemic proportions, consistent with the mass balance model (MBM) predictions.

It is perhaps reasonable to look outside controlled feeding trials: where does this passion for drastic

carbohydrate restriction comes from? Over the years, numerous experts have warned that low-

carbohydrate diets are dangerous and ineffective yet a huge number of consumers nowadays believe that

the so-called "official" high-carbohydrate recommendations are – in general – useless in terms of obesity

treatment and weight management. As pointed out by Philip K. Dick, reality is that which, when you stop

believing in it, does not go away.

Finally, as discussed out by Ludwig in his well-written letter [6], a recent meta-analysis [7] indicates that

high glycemic index (GI) and high glycemic load (GL) diets increase the risk for type 2 diabetes among high-

risk populations. The authors of the meta-analysis rightly concluded that "the cumulative effect from large

quantities of high GI foods may … nullify cereal fiber's protective effect". [7]. Simply put,

"[o]verweight/obese females could shift their carbohydrate intake for higher cereal fiber to decrease T2DM

risk, but higher GL may cancel-out this effect." [7] Its must be crystal clear as Finland Vodka – from the

perspective of evolutionary biology – that staple grain consumption has no precedent in human evolution

until *very* recently [6]; see also [11].

Leroy and Garwin [8] remind us that we generally take it for granted that <u>any</u> animal will flourish best on a

diet that roughly resembles the one to which it was adapted. It is extremely unlikely that *Homo sapiens*

turns out to be the only exception to this generally accepted biological principle. As was stressed in a highly

influential 1973 essay by the evolutionary biologist Theodosius Dobzhansky, "Nothing in Biology Makes

Sense Except in the Light of Evolution". [e.g., 9,10]

References

1. Hall KD. Challenges of human nutrition research. Science 2020 Mar 20;367(6484):1298-1300. doi:

10.1126/science.aba3807.

2. Public Health Collaboration. Randomised [sic] Controlled Trials Comparing Low-Carb Diets Of Less Than

130g Carbohydrate Per Day To Low-Fat Diets Of Less Than 35% Fat Of Total Calories.

https://phcuk.org/rcts/?fbclid=IwAR0ZPiJs_Bo4dO6ipt3u1QKmxYcjzazQNtuZPc0bopBzMS9BWnENcAy6Z5A

(accessed 11/01/2021)

3. Young CM, Scanlan SS, Im HS, Lutwak L. Effect of body composition and other parameters in obese young

men of carbohydrate level of reduction diet. Am J Clin Nutr 1971;24:290–6.

4. Volek JS, Sharman MJ, Gómez AI et al. Comparison of energy-restricted very low-carbohydrate and low-

fat diets on weight loss and body composition in overweight men and women. Nutr Metab (Lond). 2004

Nov 8;1(1):13. doi: 10.1186/1743-7075-1-13.

5. Feinman RD, Pogozelski WK, Astrup A et al. Dietary carbohydrate restriction as the first approach in

diabetes management: critical review and evidence base. Nutrition 2015 Jan;31(1):1-13.

6. Ludwig DS. Reply to S Joshi. Nutr. 2020 Oct 12;150(10):2836-2837. doi: 10.1093/jn/nxaa214.

7. Hardy DS, Garvin JT, Xu H. Carbohydrate quality, glycemic index, glycemic load and cardiometabolic risks
in the US, Europe and Asia: A dose-response meta-analysis. Nutr Metab Cardiovasc Dis. 2020 Jun
9;30(6):853-871. doi: 10.1016/j.numecd.2019.12.050. Epub 2020 Jan 13.

8. Leroy F, Cofnas N. Should dietary guidelines recommend low red meat intake? Crit Rev Food Sci Nutr.
2020;60(16):2763-2772. doi: 10.1080/10408398.2019.1657063. Epub 2019 Sep 5.

9. Calabrese EJ, Agathokleous E. Theodosius Dobzhansky's view on biology and evolution v.2.0: "Nothing in
biology makes sense except in light of evolution and evolution's dependence on hormesis-mediated
acquired resilience that optimizes biological performance and numerous diverse short and longer term
protective strategies". Environ Res. 2020 Jul;186:109559. doi: 10.1016/j.envres.2020.109559. Epub 2020
Apr 21.

10. Varki A. Nothing in medicine makes sense, except in the light of evolution J Mol Med (Berl). 2012
May;90(5):481-94. doi: 10.1007/s00109-012-0900-5. Epub 2012 Apr 27.

11. Ludwig DS. The ketogenic diet: Evidence for optimism but high-quality research needed. J Nutr. 2020
Jun 1;150(6):1354-1359. doi: 10.1093/jn/nxz308.

[FIGURE LEGEND STARTS] Supplementary File 1 Figure 1. Randomized controlled trials comparing low-carbohydrate diets of less than 130g carbohydrate per day to low-fat diets of less than 35% fat of total calories. For further details, see [2]. A serious question: *How likely is that the very-low-carb arms, time after time, substantially overreport dietary records, whereas the high-carb arms consistently underreport dietary records?* Figure adapted from [2]. **[FIGURE LEGEND ENDS]**

[FIGURE LEGEND STARTS] Supplementary File 1 Figure 3. Estimated macronutrient consumption during the epidemic of obesity and type 2 diabetes. Data from the National Health and Nutrition Examination Survey (NHANES) by year, and from Centers for Disease Control and Prevention (CDC). Inset: Incidence of diabetes (millions of people with diabetes by indicated year). NHANES suggest a large increase in carbohydrate intake as a major contributor to "caloric excess" (in reality, *macronutrient mass* excess) in the United States from 1974 to 2000, whereas the absolute amount of fat decreased for men during this period and showed only a slight increase for women. It was during this *very* period when the incidence of obesity and diabetes raised to its current epidemic proportions, consistent with the mass balance model (MBM) predictions. Whatever the extent to which the correlation between dietary carbohydrate consumption and diabetes/obesity is causal, *the lack of association between the consumption of dietary fat and diabetes is of real significance*. As pointed out by Feinman *et al.* [5], a lack of association is generally considered strong evidence for a lack of causality. There is no more serious error than mistaking the effect for the cause; in fact, Nietzsche called it "the real corruption of reason (*Twilight of Idols or How One Philosophizes with Hammer*, 1888). CHO = carbohydrate; Prot = protein. Reproduced, with permission, from [5]. **[FIGURE LEGEND ENDS]**

"Macronutrient mass intake explains deferential weight and fat loss in isocaloric diets"

SUPPLEMENTARY FILE 2

Mass balance model (MBM) does not violate any laws of physics

In the "Introduction" section of the mainbody of the manuscript, it is briefly clarified that the mass balance model (MBM) does not violate the First Law of Thermodynamics. In this **SUPPLEMENTARY FILE 2**, we further discuss this topic to clear some confusion.

As explained by Arencibia-Albite [1], it is possible for an open system, such as the human body, to be at mass balance while the system experiences a persistent energy imbalance; that is, energy balance may be positive ($\Delta E > 0$) or negative ($\Delta E < 0$) yet the **mass change** that may occur during energy flux is not required by the First Law of Thermodynamics (i.e., the Law of Conservation of Energy) to mirror the energy balance **direction**.

Thus, it is clear that the MBM does not violate the First Law of Thermodynamics.

The energy balance theory (EBT), in contrast, ignores the Law of Conservation of Mass [2]. This Law dates from Antoine Lavoisier's 1789 discovery that mass is neither created nor destroyed in any chemical reaction; hence, the O_2 mass that enters cellular respiration plus the mass of macronutrients that served as energy fuel will equal the mass of the excreted oxidation products. Daily weight loss is, therefore, the result of daily the elimination of oxidation products (CO_2, H_2O, urea, etc.) and not a consequence of the heat release upon nutrient combustion (i.e., daily energy expenditure).

The **Supplementary File Figure 1** illustrates an example of how weight stability happens concomitant to a persistent and significant energy imbalance. The reader may wonder how we are certain that is not possible to obtain a set of mass-in/mass-out values that result in the expected coincidence of mass balance with energy balance? Arencibia-Albite [1], using linear algebra, analyzed under what conditions mass balance concur with energy balance and found that both ONLY coexist if the following three conditions are simultaneously satisfied:

average absorbed fat mass = average oxidized fat mass

average absorbed carbohydrate mass = average oxidized carbohydrate mass

average absorbed protein mass = average oxidized protein mass

681   For example, if

682   65g = average absorbed fat mass = average oxidized fat mass = 65g

683   250g = average absorbed carbohydrate mass = average oxidized carbohydrate mass = 250g

684   85g = average absorbed protein mass = average oxidized protein mass = 85g

685   Then,

686   Average mass in = 65g + 250g + 85g = Average mass out

And, hence

Average EI = (4.7kcal/g) x 65g + (4.2kcal/g) x 250 + (9.4kcal/g) x 85g = Average EE = 2 154.5kcal

The reason why this is the only way in which weight stability and energy balance coincide lies outside
biochemistry and physiology. In mathematics and physics any well-defined problem, such as the
coincidence of weight stability and energy balance, has one and only one possibility:

1. The problem is unsolvable.

2. The problem has one and only one solution.

3. The problem has many solutions.

Thus, not all well-defined problems can be solved and if they do it may be true that it has exactly one
solution. A classic example is the famous Königsberg bridge problem that in 1736 the Swiss mathematician
Leonard Euler was able to demonstrate, without any doubt, that the problem solution did not exist leading
to the birth of graph theory, a branch of mathematics with many important applications. Consequently, as
demonstrated by Arencibia-Albite [1], the simultaneous satisfaction of these three requirements is what
guarantees that weight stability happens together with energy balance and NO OTHER WAY IS POSSIBLE.
Consequently, body weight stability must coincide with a persistent energy imbalance, since, as exemplified
in the Supplementary File 2 Figure 1, the constitutive processes of gluconeogenesis and *de novo* lipogenesis
impede the simultaneous satisfaction of afore mentioned conditions.

[FIGURE LEGEND STARTS] Supplementary File 2 Figure 1. Mass balance can occur in energy imbalance.
Here were illustrate hypothetical macronutrient mass input *vs.* output pattern that achieves *weight*
stability without energy balance. Circles represent macronutrient (protein [P], carbohydrate [C], fat [F])
body reserves. The left dashed box contains the daily average input mass, whereas the right box encloses
the daily average output mass. Energy densities are as follows: ρF = 9.4 kcal/g, ρC = 4.2 kcal/g , ρP = 4.7
kcal/g. While mass balance is achieved, energy balance is NOT since some of the absorbed or stored protein
(amino acids) may be transformed into glucose (gluconeogenesis, GNG) or lost through energy
expenditure-independent routs DNL = *de novo* lipogenesis. [**FIGURE LEGEND ENDS**].

References

716 1 Arencibia-Albite F. Serious analytical inconsistencies challenge the validity of the energy balance theory.
717 Heliyon. 2020 Jul 10;6(7):e04204. doi: 10.1016/j.heliyon.2020.e04204. eCollection 2020 Jul.

718 2. Sterner RW, Small GE, Hood JM. (2011) The Conservation of Mass. Nature Education Knowledge
719 3(10):20.

726 **"Macronutrient mass intake explains deferential weight and fat loss in isocaloric**
727 **diets"**

SUPPLEMENTARY FILE 3
728

Energy efficiency in weight loss diets
729

Contrary what the late Dr. Atkins taught, numerous prominent obesity experts have strictly, if not
fanatically, maintained that a isocaloric very-low-carbohydrate/high-fat diet cannot lead to a greater weight
loss than a isocaloric high-carbohydrate/low-fat diet – except for the small changes in body fluids – because
it would somehow violate the laws of thermodynamics. This is, of course, *not* the case (**See Supplementary**
File 2).

Where does the greater weight loss (**See Supplementary File 1**) comes from? Some claim that it is mostly
water weight, but this is only true initially. Water loss occurs due to glycogen depletion [1]; however, in the
long term, fat loss is a substantial portion (**See the main body of the paper**). Others have claimed that a
very-low-carbohydrate diet leads to progressive loss of muscle mass, but this assumption is clearly
fallacious [e.g., 2,3,4].

Regarding the main topic of this **Supplementary File 3**, It has been suggested that "[t]he greater weight
[and fat] loss on carbohydrate restricted diets, popularly referred to as metabolic advantage can thus be
understood in terms of the principles of nonequilibrium thermodynamics and is a consequence of the
dynamic nature of bioenergetics where it is important to consider kinetic as well as thermodynamic
variables." [5] (See also [6]). In his latest paper [7], Feinman states that "[e]nergetic inefficiency, substrate
cycling and demands of gluconeogenesis support observed advantages for weight [and fat] loss."

In our humble opinion, Feinman and Fine [5,6,7] fail to recognize that efficacy is NOT a matter of
incomplete oxidation of substrates; rather, it is about the way the body uses the available fuel resources.
Let say you start to run 5km every day and you will soon realize that you can complete the same distance in
same time (let say 0.5 h) but with a smaller amount of heartbeats. Now the body has adapted to this
cardiovascular stress physiologically, biochemically and – in the long run – anatomically – and thus has
become *more efficient fuel utilizer*. That is, your body can achieve the same running average speed
(5km/0.5 h=10km/h) utilizing less fuel. However, every time the body oxidizes 1 g of glucose while running
the heat released by the body continues to be ~4kcals, which will not change as you age or as a function of
your genome, epigenome, or proteome.

Body chemistry – and thus body mass – is all about detailed mass balances, not about energy conservation.
Mass balance model (MBM) maintains that a low-carbohydrate (LCD) is more effective in minimizing the
daily macronutrient mass intake relative to a high-carbohydrate diet (HCD), and – consequently – a LCD
manifests a substantially larger daily weight and fat loss than a HCD. *According to the MBM, weight loss*

advantage of a LCD over HCD is independent of the differences in the physiological effects of the diets (e.g.,
dietary carbohydrate-induced insulin secretion). Rather, deferential weight loss simply emerges from
dissimilar macronutrient mass intakes.

Answer the question: what do you measure when you stand on the bathroom scale, your body mass or
your energy?

On the other hand, we agree with Feinman's [6] recent conclusion that the control of the glucose-insulin
axis, via drastic carbohydrate restriction, allows improvement in *all* of the markers of metabolic syndrome
and provides methods for remission and reversal of type 2 diabetes [**See Supplementary File Figure** 1].

To be fair, one of us (AHM) must admit being "guilty" of suggesting similar ideas about the "metabolic
advantage" of LCDs as Feinman and Fine [e.g., 8]. Sometimes it just happens that the whole knowledge
structure turns out to be flawed and - thus – has to be abandoned. *The whole game needs to be changed.*
However, correcting a mistake always requires sacrifice. If the mistake has been great, so must be the
sacrifice. And if the truth has been denied for a very long time, a dangerous amount of sacrificial debt may
have been accumulated [9].

The take-home message of this **Supplementary File 3** is that contrary what Coca-Cola-sponsored
researchers (e.g., Global Energy Balance Network; GEBN]) have propagated [e.g., 10, 11, 12], the most
"cost-effective" way to gain body weight/body fat is to ingest pure carbohydrate-based energy-providing
mass (EPM), i.e., sugars, syrup and/or starch. It does not really matter which one in terms of body
weight/body fat.

779 References

1. Fernández-Elías VE, Ortega JF, Nelson RK et al. Ricardo Mora-Rodriguez. Relationship between muscle
water and glycogen recovery after prolonged exercise in the heat in humans. Eur J Appl Physiol. 2015
Sep;115(9):1919-26. doi: 10.1007/s00421-015-3175-z. Epub 2015 Apr 25.

2. Manninen AH. Very-low-carbohydrate diets and preservation of muscle mass. Nutr Metab (Lond). 2006
Jan 31;3:9. doi: 10.1186/1743-7075-3-9.

3. Rondanelli M, Faliva MA 2, Gasparri C et al. Current opinion on dietary advice in order to preserve fat-
free mass during a low-calorie diet. Nutrition. 2020 Apr;72:110667. doi: 10.1016/j.nut.2019.110667. Epub
2019 Dec 5.

4. Casanueva FF, Castellana M 3 Bellido D et al. Ketogenic diets as treatment of obesity and type 2 diabetes
mellitus. Rev Endocr Metab Disord. 2020 Sep;21(3):381-397. doi: 10.1007/s11154-020-09580-7.

5. Feinman RD. Fine EJ. Nonequilibrium thermodynamics and energy efficiency in weight loss diets. Theor
Biol Med Model. 2007; 4: 27. Published online 2007 Jul 30. doi: 10.1186/1742-4682-4-27.

6. Feinman RD. Beyond "a Calorie is a Calorie": An Introduction to Thermodynamics. In: *Nutrition in Crisis:*
Flawed Studies, Misleading Advice, and the Real Science of Human Metabolism. Chelsea Green Publishing,
2019.

7. Feinman RD. The biochemistry of low-carbohydrate and ketogenic diets. Curr Opin Endocrinol Diabetes
Obes. 2020 Oct;27(5):261-268. doi: 10.1097/MED.0000000000000575.

8. Antonio J. Manninen AH. Eating to Improve Body Composition. In: *Essentials of Sports Nutrition and*
Supplements. Humana Press, 2008.

9. Peterson JB. *12 Rules for Life: An Antidote to Chaos*. Random House Canada, *2018.*

10. Barlow P, Serôdio P, Ruskin G et al.. Science organisations and Coca-Cola's 'war' with the public health
community: insights from an internal industry document J Epidemiol Community Health. 2018
Sep;72(9):761-763. doi: 10.1136/jech-2017-210375. Epub 2018 Mar 14.

11. Greenhalgh S Inside ILSI: How Coca-Cola, Working through Its Scientific Nonprofit, Created a Global
Science of Exercise for Obesity and Got It Embedded in Chinese Policy (1995-2015). Health Polit Policy Law.
2020 Sep 16;8802174. doi: 10.1215/03616878-8802174. Online ahead of print.

12. Serodio P, Ruskin G, McKee M et al. Evaluating Coca-Cola's attempts to influence public health 'in their
own words': analysis of Coca-Cola emails with public health academics leading the Global Energy Balance
Network. Public Health Nutr. 2020 Oct;23(14):2647-2653. doi: 10.1017/S1368980020002098. Epub 2020
Aug 3.

Low carbohydrate diets are better than low-GI diets or high cereal diets for
weight loss, HbA1c, triglycerides and HDL. Data from Westman, *et al* (2008)
Nutr Metab (Lond), 5 (36). and Jenkins, *et al* (2008), *JAMA* **300**: 2742-2753.

[**FIGURE LEGEND STARTS**] **Supplementary File 3 Figure 1**. Low-carbohydrate diets are more effective for
weight loss, glycated hemoglobin (HbA1c), triglycerides (TG) and high-density lipoprotein than low-glycemic
(GI) or high cereal diets, all of which are consistent with the MBM. CHO = carbohydrate; Total-C = total
cholesterol; LDL = low-density lipoprotein. Preproduced, with permission, from [6]. [**FIGURE LEGEND ENDS**]

"Macronutrient mass intake explains deferential weight and fat loss in isocaloric
diets"

821 SUPPLEMENTARY FILE 4

A re-analysis of the Hall *et al.* paper "Calorie for Calorie, Dietary Fat Restriction Results in More
Body Fat Loss than Carbohydrate Restriction in People with Obesity"

In 2015, Hall *et al.* published a highly infuential paper [1] entitled "Calorie for Calorie, Dietary Fat
Restriction Results in More Body Fat Loss than Carbohydrate Restriction in People with Obesity" in *Cell*
Metabolism. The authors restricted dietary carbohydrates *vs.* dietary fat for 6 days following a 5 day
baseline diet in 19 obese adults confined to a metabolic ward where they exercised daily. Subjects
were given both isocaloric diets, the carbohydrate restricted diet (RC) or the fat restricted diet (RF), in
random order during each of two stays in a ward. Data seems to suggest that that RF resulted in a
greater rate of body fat loss compared to isocaloric RC.

We decided to take a closer look and re-analyze the Hall *et al.* data; and we come to the firm conclusion
that their conclusion is highly, if not fatally, misleading.

833
834 **1. Hall *et al.* data demonstrate that carbohydrate restricted diet (RC) show no risk of body fat**
835 **accumulation**
836
837 A widespread postulate, defended by the energy balance theory (EBT), is that during underfeeding body fat
838 loss (FL) is given by the following difference:
839
840 FL = (Fat intake) − (Net Fat Oxidation)
841
Accordingly, RF is expected to result in a greater FL than an isocaloric RC as fat intake is substantially lower.
It should be noted, however, that this is a reasonable hypothesis only if the net fat oxidation is independent
of diet macronutrient composition. If the net fat oxidation increases as fat intake increases (and *vice versa*),
FL can be similar among isocaloric diets that vary greatly in dietary fat content. Hall *et al.* data demonstrate
that this is indeed the case. Respiratory quotient (RQ) measurements in their Figure 2C, for example,
indicate that fat oxidation increases in proportion to dietary fat intake. Consequently, the characteristic
high fat content of RC elicits high levels of fat oxidation, which counterbalances the high fat intake avoiding
excessive body fat deposition. This interpretation is valid as dual-energy X-ray absorptiometry (DXA)
measurements in Table 3 indicate that fat loss in RC was similar to that of RF even though the latter diet

contained substantially less fat (Fat intake: RC = 108g/day *vs.* RF = 17g/day; Fat loss: RC = −0.529±0.13kg *vs.*
RF = −0.588±0.14kg, p = 0.78). Therefore, it seems that the alleged risk of unwanted fat accumulation with
RC diets is highly unlikely and experimentally unjustified.
854
855 **2. The cumulative fat measurements illustrated in Table 3 (or Figure 3D) are artificial quantities**
856 **that do <u>not</u> reflect the <u>actual</u> level of body fat loss.**
857
858 DXA measurements in Hall et al. study are the only <u>direct</u> measurement of fat loss. The presented
859 cumulative fat loss illustrated in Table 3 (or in Figure 3D) is an <u>undirect</u> estimate of fat loss computed
860 according the following formulas (in grams):
861
862 (Daily fat loss) = (Daily Fat Intake) − (Daily Net Fat Oxidation)
863
864 where Daily Fat Oxidation is another estimate calculated by
865
866 (Daily Net Fat Oxidation) = 1.63xVO2 − 1.64xVCO2 − 1.84x(Urinary Nitrogen)
867
868 (see Hall *et al*. [1] methods).
869
In these formulas, the only precise numeric input is "Daily Fat Intake", whereas the "Daily Net Fat
Oxidation" is an *estimate* obtained from *estimates* which unavoidably increases the measurement
inaccuracy. Moreover, if an indirect measurement is indeed accurate, it should be very close to the most
precise direct measurements available, such as DXA. Consequently, if cumulative fat loss calculations
reported by Hall *et al*. are accurate, they should be nearly identical to values measured by DXA, which is
clearly NOT the case. *The cumulative fat measurements illustrated in Table 3 (or Figure 3D) are thus*
artificial quantities that do <u>not</u> reflect the <u>actual</u> level of fat loss.
877
878 Nevertheless, Hall *et al*. try to convince readers that the direction of accuracy is other way around, i.e.,
879 cumulative fat calculations are more accurate than direct DXA fat mass measurements. In the justification
880 of such a proposal, they argue that DXA measurements may be prone to inaccuracies in settings of
881 carbohydrate restriction as this may lead to greater water loss than fat restriction:
882
"But even high precision methods, such as DXA, may lack the sensitivity to detect small differences in body
fat change... Indeed, retrospective analysis [what analysis?] of our data suggests that the minimal
detectable difference between the diets for body fat mass using DXA was ⬚0.4 kg. Thus, we suspect that the
DXA measurements of fat mass change in the present study were insufficiently sensitive to detect
differences between the diets. Furthermore, DXA may provide inaccurate results in situations of dynamic
weight change and shifting body fluids... This could be especially important with diets differing in their level
of carbohydrate restriction since greater losses of body water are likely with lower levels of dietary
carbohydrate." [1]
891
It should be noted, however, that DXA <u>fat mass</u> measurements will be mainly unaffected since the amount
of water present in the body fat compartment is essentially non-existent. Thus, if any inaccuracy is present,
it would be substantially greater for lean mass measurements than for fat mass measurements as total

body water is basically localized in the lean mass compartment. This fact, alone, refutes the above
mentioned interpretation by Hall *et al.*

Sensitivity, nonetheless, is an academic issue with minor relevance in the treatment of obesity. Lets
assume, as Hall *et al.* claim, that DXA sensitivity is ~ 0.4kg. This means that if DXA fat mass measurement
before a dietary intervention is, let say 35kg, and a particular underfeeding procedure decreases fat mass
by exactly 200g (0.2kg) then DXA measurement will still read 35kg as the fat mass change is less than the
sensitivity of the instrument. This is, of course, immaterial during typical underfeeding interventions as the
fat loss at such procedures substantially exceeds DXA sensitivity.

3. Compared to RF, RC resulted in significantly greater weight loss <u>both</u> experimentally and
computationally

Table 3 by Hall *et al.* shows that RC resulted in substantially more weight loss than isocaloric RF (RC:−
1.85±0.15kg vs. RF: −1.3±0.16kg, p = 0.022). For instance, the simulation in Figure 3H shows that, over a six
month period, RC leads to about 4kg of extra weight loss compared to RF.

References

1. Hall KD, Bemis T, Brychta R, Chen KY, Courville A, Crayner EJ, Goodwin S, Guo J, Howard L, Knuth ND,
Miller BV 3rd, Prado CM, Siervo M, Skarulis MC, Walter M, Walter PJ, Yannai L. Calorie for Calorie, Dietary
Fat Restriction Results in More Body Fat Loss than Carbohydrate Restriction in People with Obesity. *Cell*
Metab. 2015 Sep 1;22(3):427-36. doi: 10.1016/j.cmet.2015.07.021. Epub 2015 Aug 13. PMID: 26278052;
PMCID: PMC4603544.

NOTE:

As Corresponding Author, I confirm that that all supplementary files have been read and
approved for submission by all the named authors.

Sincerely,

Anssi H. Manninen   23/03/2021

[NOTE THAT THE REFERENCES IN SUPPLEMENTARY FILES ARE SEPARATE FROM THE REFERENCES
IN THE MAINBODY OF THE MANUSCRIPT. THUS, THEY DO NOT FOLLOW THE SAME REFERENCE
NUMBERING.]

940