Re-analysis of genetic risks for Chronic Fatigue Syndrome from 23andMe data finds few remain

Felice L. Bedford[1]*, Bastian Greshake Tzovaras [2]*

[1]University of Arizona, Tucson, AZ, USA

[2]Université de Paris, Paris, France

* Correspondence:
Corresponding Authors
bedford@arizona.edu, bastian.greshake-tzovaras@cri-paris.org

Keywords: Chronic Fatigue Syndrome[1], Myalgic Encephalomyelitis[2], single nuclear polymorphism[3], 23 and Me[4], genetic predisposition[5]

Word count: 1999 Tables: 1 Supplementary Figures: 1

Abstract

It is tempting to mine the abundance of DNA data that is now available from direct-to-consumer genetic tests, but this approach has its pitfalls A recent study put forth a list of 50 single nucleotide polymorphisms (SNPs) that predispose to Chronic Fatigue Syndrome (CFS), a potentially major advance in understanding this still mysterious disease. However, only the patient cohort data came from a commercial company (23andMe) while the control was a genetic database. The extent to which 23andMe data agree with genetic reference databases is unknown. We reanalyzed the 50 purported CFS SNPs by comparing to control data from 23andMe which are available through public platform OpenSNP. In addition, large high-quality database ALFA was used as an additional control. The analysis lead to dramatic change with the top of the leaderboard for CFS risk reduced and reversed from an astronomical 129,000 times to 0.8. Errors were found both within 23andMe data and the original study-reported Kaviar database control. Only 3 of 50 SNPs survived initial study criterion of at least twice as prevalent in patients, EFCAB4B involving calcium ion channel, LINC01171, and MORN2 genes. We conclude the reported top-50 deleterious polymorphisms for Chronic Fatigue Syndrome were more likely the top-50 errors in the 23andMe and Kaviar databases. In general, however, correlation of 23andMe control with ALFA was a respectable 0.93, suggesting an overall usefulness of 23andMe results for research purposes but only if caution is taken with chips and SNPs.

1   Introduction

As part of a growing number of researchers that advocate using the plentiful genetic data from direct to consumer testing (1), we are also aware of its pitfalls. Recently, a study in Frontiers in Pediatrics (2) described an ambitious project to elucidate genetic predispositions for Chronic Fatigue Syndrome (CFS), a not-uncommon condition of debilitating fatigue, immune dysregulation, and central nervous system impairment. The study analyzed the approximate 500,000 genetic single nuclear

polymorphisms (snps) resulted by the commercial 23andMe genetic testing company in people with
CFS.

Two items from the Perez et al. results were immediate red flags. They showed the top 50 deleterious
snps with the greatest difference in frequency between CFS patients and control data. At the very top
of the leaderboard was a snp on the gene GPBAR1 that was 129,000 times more prevalent in CFS
patients. If this finding is accurate then the authors may well have discovered THE genetic cause of
CFS rather than just a predisposition. In addition, the CYP2D6 gene on their list is recognizable as
one that in the past, for a different snp, had the majority of 23andMe customers believing they had a
poor xenobiotic metabolizer phenotype that only actually affects less than one percent of the
population (from Snpedia 3).

In the Perez et al. study, the data used to compare to the 23andMe CFS participants did not come
from 23andMe. Instead, they relied on published frequencies in the Kaviar database, a compilation
from multiple projects (4). Unless the control data comes from the identical source as the
experimental data, any differences in quality or population constituency between the two datasets
may lead to inaccurate conclusions when compared. How many of the reported frequency ratios
between CFS patients and controls would remain noteworthy if both data come from the same
source? The study established a criterion that a snp should be at least twice as prevalent in CFS
patients, a ratio of 2, to be of note. In addition, how closely do 23andMe data match published
genetic reference data? Errors in direct-to-consumer genetic tests have been reported (5).

To address these questions, we accessed a publicly available control set of genomes from 23andMe
participants. We reanalyzed the frequency ratio of the highly prevalent snps in CFS from the Perez et
al. study to this more appropriate control. We also incorporated an additional new large high-quality
online control dataset for further comparison among control datasets.

2   Method

To provide control data from 23andMe for the 50 top-predisposing CFS snps, we accessed publicly
available genome files on openSNP[1], a platform co-founded and maintained by one of us (BGT).
OpenSNP allows individuals to upload their own genetic results from a variety of commercial test
companies for public sharing (6). The allele frequencies and genotypes for these SNPs were
calculated for all 23andMe data sets present in openSNP on 2020-06-19. While self-selection of
participants can skew a dataset, the platform also allows phenotypes of interest to be added by
participants. We noted the absence of CFS and Myalgic Encephalomyelitis (ME) on the list of
phenotypes which provided an initial confidence that the dataset does not contain an
overrepresentation of CFS patients compared to the general population.

For the additional control: The allele frequency aggregation project (ALFA) project was developed as
part of the National Center for Biotechnology Information (NCBI) database of genotypes and
phenotypes (dbGAP) and had their inaugural release on March 10, 2020 (7). It contains a high-quality
aggregate of over 1200 studies with a goal of one million dbGAP subjects. ALFA (build 154, release
date April 21, 2020) was accessed through NCBI dbSNP[2]. ALFA Europe was selected when

[1] https://opensnp.org

[2] https://www.ncbi.nlm.nih.gov/snp

available to match the population of 23andMe, primarily Americans of European decent. When
unavailable, ALFA Global was the second choice, followed by GnomAD – exome and then 1000
Genome project if absent from ALFA entirely. When ALFA is referred to subsequently, it is a
shortcut notation for the totality of this procedure.

To recalculate the ratio with the new controls, for the CFS data we used the frequencies provided by
Perez et al. in Table 1. Since neither the table nor supplementary materials explicitly listed the alleles,
we assumed that the frequencies always referred to the derived allele. In the event of multiple derived
alleles at a position, we further assumed the most prevalent one was used. Spot checks of their Kaviar
control supported that this was the study's intended listing. The new ratios were recalculated for the
23andMe control and for the ALFA control and subsequently compared to the original ratios. The
three control datasets at the 50 snps (Kaviar, 23andMe, ALFA) were compared to each other.

3 Results

The recalculated ratio of allele frequency in CFS patients to control subjects using 23andMe control
data, or where unavailable, ALFA frequencies, along with the elimination of 2 duplicates, found that
only 11 of the 50 polymorphisms now exceeded a ratio of 2. That is, only 22 percent of the originally
reported polymorphisms remained at the original study criterion that notable snps were at least
double the frequency in CFS patients compared to unaffected controls.

Of the 11 remaining polymorphisms with a ratio that met the criterion, the majority, 7, could be
based only on ALFA frequencies; and 1 was not reported on ALFA either. These 8 snps were only
present in the 23andMe control data set in very few samples, ranging from 17 to 0, in contrast to a
median of nearly 3000 samples in the 23andMe control data overall. These further 8 snps therefore
were also not shown to have a higher prevalence in 23andMe CFS patients than in 23andMe controls.
Dates of upload to openSNP hint that the early days of the 23andMe v5 chip could be a source of
error. All 8 of these come from the top of the original ratio leaderboard (Table 1).

Only 3 snps of 50 remained, on genes EFCAB4B, LINC01171, and MORN2, that could be shown to
meet the original criterion of at least double in CFS patients compared to a comparable 23andMe
control, all hovering at a ratio of about 2.0. None of the astronomically high ratios of patients to
controls could be shown to remain.

Table 1 presents the ratios of allele frequency in CFS patients to control subjects (original,
recalculated with the new 23andMe control, recalculated with ALFA), the frequency of allele
occurrence (original CFS patients, original Kaviar, new 23andMe control, new ALFA control), and
the number of samples (23andMe control) for each of the snps reported in the original table of the
study. Genotype frequencies found in the 23andMe control samples for each snp is in Supplementary
Figure 1.

Comparison of the control datasets (Kaviar, 23andMe, ALFA) found 2 primary patterns. Most
prevalent, 29 out of the 48 of the 23andMe control frequencies were in good agreement with ALFA
with both being substantially higher than the reported Kaviar values, suggesting a Kaviar-related
error. We call this error Type A. For 8 of the 48 snps, the 23andMe control frequencies were instead
different (higher) than both ALFA and Kaviar, which were in good agreement with each other and
point to a 23andMe error (Type B error). Like the missing 23andMe snps, Type B errors came from
the top of the leaderboard and further accounts for the original astronomic reported ratios.

Concerning the two-red flag genes mentioned at the outset, both are Type B errors. The GPBAR1
snp for derived allele A was found in 97% of the 23andMe control sample compared to practically 0
in the other control data leading to a recalculation of the ratio of prevalence between CFS patients
and controls from the reported 129,000 to 0.8, a reversal. Based on upload dates, this erroneous base
A call may be traceable specifically to the v4 chip but would require exact test dates for
confirmation. The CYP2D6 had 2 snps, one of which had more than a third erroneous 23andMe call
for base A..

The correlation between the 23andMe control and ALFA control frequencies without the genes
above. and without any 23andMe data having fewer than 50 participants, was a respectable 0.93.

Discussion

The genetic predispositions reported for Chronic Fatigue Syndrome are not supported when
reanalyzed with more appropriate control data including those drawn from the same 23andMe pool as
the CFS patients. Out of the original 50 genomic positions presented to have the most prevalent
deleterious polymorphisms among CFS sufferers, only three remained that met the original study
criterion of at least twice as frequent as healthy individuals. The top-ranked risk factor on gene
GPBAR1 with an astronomical ratio of 129,000 was reduced to the more sensible 0.8, which would,
if anything, be a protective snp against CFS.

The erroneous odds ratios were found to originate from a mixture of errors in both 23andMe and in
the reported Kaviar control dataset. The more dramatic frequencies that had been listed as dozens,
hundreds, and thousands of times higher in CFS patients were due to seeming 23andMe peculiarities
of very high frequencies for minor alleles. We found these 23andMe errors to either also be present
in high numbers in the 23andMe controls (Genes GPBAR1, CYP2D6, PLA2G4D, CYP2A6, DDX5)
or quite often were simply missing from most samples. The number of CFS subjects from Perez et al.
at each of these snps, and overall, is unknown. The majority of the errors with less-striking ratio
inflations arose from the reported Kaviar control data where we found these to be lower frequencies
than both ALFA and 23andMe control datasets for many of the snps.

The Perez et al. discussion goes through, spelling out one by one, the function of each of the genes
from the top 10 in the table by summarizing what is known and including speculation for how these
factors may tie into CFS. For example, they suggest that decreased metabolism of xenobiotics may
be relevant to multiple chemical sensitivity disorder which is in turn relevant for some CFS cases and
a gene that is downregulated by sleep deprivation which in turn is a factor in chronic fatigue states.
The present reanalysis finds at the very least that such discussion and speculation are premature as
there is no evidence that any of those genes are relevant.

The three genes with polymorphisms that remained with the original study criteria of occurring at
least twice as often in CFS patients were EFCAB4B, MORN2 and LINC01171 which involve
calcium ion channels, cell differentiation, and a long non-coding RNA transcript respectively. It is
tempting to fish for connections such as to other ion channel polymorphisms that have been found
relevant in CFS (8) but here too, it is premature to speculate; reanalysis of the full 23andMe data for
CFS patients, as is now clearly warranted, may produce an entirely different top 50 leaderboard and
functional analysis. Likewise, the criterion of a ratio of at least double may also prove too stringent
which may then put some of the snps back into consideration but this too is unknown until a full
reanalysis. It is beyond the scope of this article to raise that high CADD scores also may not always
be an appropriate filter (e.g. 9). We conclude that the top-50 table presented in the CFS study does not

reflect the top 50 deleterious differences between Chronic Fatigue Syndrome and unaffected
individuals as intended but rather the top 50 errors in the 23andMe and Kaviar databases.

The present reanalysis highlights the need to use control data from the same commercial direct-to-
consumer genetic testing company when used for research. On a positive note, quirks aside, there is
generally high agreement between 23andMe and scientific genetic database ALFA. There are 10
million direct-to-consumer genetic test results which positively dwarfs the data collected in scientific
studies. Using the abundant commercial DNA results to find genetic predispositions is very appealing
especially for disorders without known cause, like Chronic Fatigue Syndrome The promise for
successful continued mining of public data for research purposes remains but with caution over select
snps and chips.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Author Contributions

Conceived the project (FB), extraction from openSNP (BGT), data analysis (FB, BGT), manuscript
creation (FB), manuscript edits (BGT).

References

1. Bedford FL. Sephardic signature in haplogroup T mitochondrial DNA. Eur J Hum Genet. 2012
Apr;20(4):441–8.

2. Perez M, Jaundoo R, Hilton K, Del Alamo A, Gemayel K, Klimas NG, et al. Genetic
Predisposition for Immune System, Hormone, and Metabolic Dysfunction in Myalgic
Encephalomyelitis/Chronic Fatigue Syndrome: A Pilot Study. Front Pediatr [Internet]. 2019
[cited 2020 Jul 24];7. Available from:
https://www.frontiersin.org/articles/10.3389/fped.2019.00206/full

3. Cariaso M, Lennon G. SNPedia: a wiki supporting personal genome annotation, interpretation
and analysis. Nucleic Acids Res. 2012 Jan 1;40(D1):D1308–12.

4. Glusman G, Caballero J, Mauldin DE, Hood L, Roach JC. Kaviar: an accessible system for
testing SNV novelty. Bioinformatics. 2011 Nov 15;27(22):3216–7.

5. Tandy-Connor S, Guiltinan J, Krempely K, LaDuca H, Reineke P, Gutierrez S, et al. False-
positive results released by direct-to-consumer genetic tests highlight the importance of clinical
confirmation testing for appropriate patient care. Genetics in Medicine. 2018 Dec;20(12):1515–
21.

6. Greshake B, Bayer PE, Rausch H, Reda J. openSNP–A Crowdsourced Web Resource for
Personal Genomics. PLOS ONE. 2014 Mar 19;9(3):e89204.

7. Tryka KA, Hao L, Sturcke A, Jin Y, Wang ZY, Ziyabari L, et al. NCBI's Database of Genotypes
and Phenotypes: dbGaP. Nucleic Acids Res. 2014 Jan;42(Database issue):D975-979.

8.   Marshall-Gradisnik S, Smith P, Brenu E, Nilius B, Ramos S, Staines D. Examination of Single
Nucleotide Polymorphisms (SNPs) in Transient Receptor Potential (TRP) Ion Channels in
Chronic Fatigue Syndrome Patients. Immunology and Immunogenetics Insights. 2015 Apr 15;7.

9.   Mather CA, Mooney SD, Salipante SJ, Scroggins S, Wu D, Pritchard CC, et al. CADD score has
limited clinical validity for the identification of pathogenic variants in non-coding regions in a
hereditary cancer panel. Genet Med. 2016 Dec;18(12):1269–75.

Table 1 Recalculated ratios of polymorphism frequencies of CFS patients to controls

Gene	rsID	Frequency				PolyM	Ratio of CFS to Control			N	Fla
		ME/CFS	Kaviar control	23andMe control	Alfa control		Original (with Kaviar)	with 23and Me	with Alfa	23andMe control	
GPBAR1	rs199986029	77.3	0.0006	99.73	0	G to A/C	129,000	0.78	infinite	1835	
HLA-C	rs41560916	62.7	0.0013	-	0	C toA/G/T	48,200	-	infinite	0	*
BCAM	rs3810141	10.2	0.0006	-	6.3	C to A/T	17,000	-	1.62	2	*
AAAS	rs150511103	19.3	0.0013	8.33	0.01	C toA/G/T	14,900	2.32	1930.00	12	*
FGA	rs146387238	19.3	0.0013	0.00	0.03	C to A/G	14,900	infinite	643.33	17	*
SLC25A13	rs80338723	19.3	0.0013	0.00	0	C toA/G/T	14,900	infinite	infinite	17	*
MYBPC3	rs112738974	19.3	0.0019	3.13	0	C toA/G/T	10,200	6.18	infinite	16	*
PEX6	rs112298166	19.3	0.0019	-	-	C to G/T	10,200	-	-	-	*
CYP2D6	rs1135830	45.4	0.0097	35.31	0.01	G to A/T	4,680	1.29	4540.00	1892	
HLA-DRB1	rs112796209	41.5	0.0109	0.00	10.2	T to C	3,810	-	4.07	1	*
PLA2G4D	rs147516345	15.9	0.0103	18.20	0.5	T to C	1,550	0.87	31.80	1865	
CYP2A6	rs5031017	38.6	0.0264	30.50	0.1	C to A	1,460	1.27	386.00	1983	
CYP2D6	rs199535154	94.3	0.231	50.00	0.5	A to G	408	1.89	188.60	4	*
DDX51	rs201101053	15.9	0.0708	15.08	0	G to A	225	1.05	infinite	1873	
LHB	rs34349826	74.2	0.644	27.27	7	A to G	115	2.72	10.60	6	*
HLA-A	rs1137110	13.8	0.249	-	0.13	T to G	56	-	106.15	0	*
HLA-DRB1	rs1136756	43.9	1	50.00	30	T to C/G	44	-	1.46	1	*
HLA-DRB1	rs9269744	40.5	1.3	-	29.8	G to C	31	-	1.36	0	*
TPTE	rs1810540	34.5	1.16	30.29	34.8	C to A/T	30	1.14	0.99	1835	
HLA-DQA1	rs1061172	15.7	1.33	46.07	16.8	A to G	12	0.34	0.93	1922	
C6orf183	rs399561	63.2	6.46	40.68	40.7	G to A	10	1.55	1.55	3027	
C14orf37	rs3829765	81.5	9.75	51.54	54.5	G to A/T	8	1.58	1.50	3826	
EFCAB4B	**rs11062745**	**27.9**	**3.39**	**13.78**	**15.5**	**T to C**	**8**	**2.02**	**1.80**	**3783**	
PLD5	rs2810008	55.4	6.71	32.30	32.8	G toA/C/T	8	1.72	1.69	3024	
MUC19	rs11564109	24	2.95	14.18	15	G to A	8	1.69	1.60	3825	
ARHGAP42	rs17647207	14.4	1.82	8.83	9.5	G to A	8	1.63	1.52	3018	
ADAMTS19	rs30645	76.5	9.75	51.01	51.5	T to A/C	8	1.50	1.49	3791	
LINC01171	**rs11605546**	**23**	**2.97**	**10.68**	**10.4**	**G to A**	**8**	**2.15**	**2.21**	**3826**	
ANKDD1B	rs34358	83.3	10.9	62.93	63.6	G to A/T	8	1.32	1.31	2990	
ZBED5	rs2232919	12	1.61	6.56	7.4	T to C/G	7	1.83	1.62	2979	
CTC-441N14.4	rs9112	60.3	8.44	40.47	41.8	G to A/C	7	1.49	1.44	2987	

SLC35B2	rs3187	13.1	1.85	10.22	9	G to A	7	1.28	1.46	2887
PRSS41	rs61747737	11.5	1.63	7.01	7.9	T to A/G	7	1.64	1.46	1879
OTOG	rs12422210	26.4	3.76	15.09	17.3	G to A	7	1.75	1.53	2941
MTCH2	rs1064608	45.7	6.58	39.77	34.26	G to C/T	7	1.15	1.33	45
SULF1	rs6990375	51.2	7.49	30.51	30	G to A/T	7	1.68	1.71	3832
OTOG	rs11024333	29.5	4.34	16.20	16.3	G toA/C/T	7	1.82	1.81	3795
ART3	rs14773	43.3	6.41	28.20	26.7	C to A	7	1.54	1.62	2950
PPHLN1	rs12658	36.3	5.45	23.14	22.4	C to A./T	7	1.57	1.62	2991
PRICKLE1	rs12658	36.3	5.45	D	D		7	D	D	D
VARS2	rs2249464	74.7	11.4	55.74	54.4	T to C	7	1.34	1.37	3736
MORN2	**rs3099950**	**21.9**	**3.37**	**11.08**	**12.1**	**G to A**	**7**	**1.98**	**1.81**	**2990**
AC007956.1	rs2270424	36.8	5.99	21.37	20.4	G to A	6	1.72	1.80	3793
AREL1	rs2270424	36.8	5.99	D	D		6	D	D	D
PRRT4	rs359642	95	15.5	80.00	82.3	G to A	6	1.19	1.15	3751
HUS1	rs2307252	16.7	2.76	11.45	9.9	G to A	6	1.46	1.69	2990
PRSS56	rs1550094	92.2	16.2	69.83	69.2	G toA/C/T	6	1.32	1.33	3674
C5orf52	rs10051838	24	4.35	13.38	13.3	G to A	6	1.79	1.80	3024
ZNHIT1	rs17319250	40.5	7.41	24.14	23.5	T to C	5	1.68	1.72	3798
CPLX2	rs3822674	70.5	12.9	49.23	48	T to A/C	5	1.43	1.47	2903

202   D = duplicate; - = missing value; N = number of participants; PolyM = polymorphisms

203   * Ratios obtained with fewer than twenty 23andMe control subjects were considered invalid