medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

The dates of the sudden 2020 Autumn COVID-19 boost in European countries is linked to latitudes and not to temperatures.

Running title: COVID-19 second wave and latitude

Stephan Walrand, cliniques universitaires Saint-Luc, Brussels, Belgium

corresponding author:

stephan.walrand@uclouvain.be

phone : 32 495 680813

Abstract :

Purpose: Determining the triggering factor of the sudden boost of the daily reported new COVID-19 cases arising in most European countries during 2020 Autumn.

Methods: The dates of the boost were determined using an automatic fitting of the two last months reported daily new cases in 18 European countries of latitude ranging from 39° to 62°.

Results: The present study proves no correlation between the country boost date and its 2 weeks preceding temperature, but shows an impressive linear correlation with its latitude. The country boost date corresponds to the time when its sun UV daily dose drops below 30% of that of 0° latitude.

Conclusions: Previous studies compared covid-19 severity between different populations. In contrary, the date of the boost is an intrapopulation observation and can thus be only triggered by a parameter globally affecting the population, i.e. the sun UV daily dose decreases. This result supports low vitamin blood D level as a real primary cause of COVID-19 severity. Europe and US could thus sink into a long crisis, since they will get back above the October sun UV dose only end 2021 March.

Keywords: covid-19, latitude, vitamin D, second wave

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Introduction

Most European countries underwent an unexpected spectacular boost of the daily new COVID-19 cases in autumn (Fig. 1), imposing in emergency new confinement rules



Figure 1: Examples of daily new COVID-19 cases (extracted from the Google home page when searching "covid"). All curves exhibit a clear boost of the growing rates around October.

The most common reported explanation is the decreasing temperature. The aim of this data analysis study is to challenge this assumption against a pure country latitude impact.

Material and method

Data sources

The daily reported country new COVID-19 cases were obtained by digitalizing the curves from the Johns Hopkins coronavirus resource center (https://coronavirus.jhu.edu/map.html) using Digitizelt (www.digitizelt.de).

The country population weighted center (PWC) latitudes were obtained from the Baylor University population resource (http://cs.ecs.baylor.edu/~hamerly/software/europe_population_weighted_centers.txt)

The averaged 2 weeks temperatures preceding the boost dates were derived from weather.com.

The theoretical sun UV daily dose for vitamin D production, as a function of latitude and of the day of the year was derived by interpolation from [1].

Boost date determination

medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

The date of the growing rate boost was automatically determined by fitting the two last months of the daily new COVID-19 cases with:

$$N_c \, e^{(\alpha_c + (\beta_c - \alpha_c)\, l(t - t_c))\, t} \qquad (1)$$

where l is the logistic function:

$$l(t - t_c) = \frac{1}{1 + e^{-\gamma(t - t_c)}} \qquad (2)$$

t_c is considered as the boost date of the country c, i.e. the date when the exponential coefficient, coming from the initial value α_c, crosses the value $\frac{\alpha_c + \beta_c}{2}$ before tending towards β_c when $t \to \infty$. γ is the steepness of this changing. γ was assumed country independent as we search for an impact of the latitude of its own. This further allows to prevent an over-fitting of the data noise by a steepness tuned for each country.

Note that as the exponential coefficient varies with time, the doubling time around the boost date is not simply ln(2) divided by this coefficient.

Results:

Table 1 shows the fitting results (all the data and fitting process is provided in the supplementary xlsx file).

country	temp. [C]	PWC lat. [deg]	date	day	Sun UV [%]
Greece	23.4	38.7	oct 22	296	36.0
Portugal	19.9	39.7	oct 24	299	33.8
Spain	17.6	39.8	oct 28	302	31.7
Bulgaria	17.5	42.8	nov 01	306	26.3
Italy	20.6	42.9	oct 16	291	34.1
Serbia	16.0	43.8	oct 20	294	30.7
Croatia	17.4	45.3	oct 16	291	30.8
Slovenia	14.5	46.2	oct 15	289	30.1
Switzerland	14.0	47.0	oct 11	285	31.0
France	14.9	47.2	oct 22	296	25.4
Austria	15.4	47.8	oct 16	290	27.6
Belgium	18.6	50.8	oct 02	277	30.7
Germany	14.9	50.9	oct 14	289	24.5
Netherland	18.3	52.1	oct 02	277	29.0
UK	21.1	52.8	sept 27	271	30.7
Russia	15.2	54.3	sept 21	266	32.4
Sweden	17.3	59.0	sept 21	266	25.9
Finland	14.1	61.8	sept 20	264	21.9

Table 1. Temp: mean country temperature during the 2 weeks preceding its COVID-19 boost date. PWC lat: latitude of the country PWC. day: corresponding day of the year. Sun: theoretical sun UV daily dose at the boost date, expressed as a fraction at the PWC latitude versus the latitude 0°.

medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint **(which was not certified by peer review)** is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Fig. 2A clearly proves no correlation with temperatures, while fig. 2B clearly shows an impact of the country latitude.



Figure 2: COVID-19 boost date as a function of the country mean temperature during the 2 preceding weeks (A), and as a function of the country PWC latitude (B). Use of flags allows linking country positions on the two graphs.

Fig. 3 clearly shows that the boost dates set on the theoretical sun UV daily dose as a function of the latitude evidences an impact of the sun UV daily dose.



Figure 3: Solid curves: theoretical sun UV daily dose for vitamin D skin production under a clear sky corresponding to the 18 PWC country latitudes plus the 31° latitude (derived from [1]). Black circles: country boost dates positioned on their corresponding latitude curve. Black dot-dash line corresponds to the averaged countries sun UV daily doses at their boost date.

medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Fig. 4 shows that the day of the second wave boost is well predicted by the moment when the sun UV daily dose of the country becomes lower than 30% of that at latitude 0°.



Figure 4: observed day of the second wave covid-19 boost as a function of the day when the country sun UV daily dose drops lower than 30% of that at latitude 0°. Trendline forced to intercept (0,0).

Discussion:

Many studies support an impact of low vitamin D blood level on the respiratory impairment in coronavirus or virus diseases [2], and also recently for the COVID-19 pandemic [3]. Low vitamin D blood level is also more present in populations at risk: i.e. aged people [4], obese patients [5], coloured skin people living in high latitude countries [6] and diabetic patients [7].

However these studies compared the covid-19 severity between different populations. They cannot thus discriminate whether the vitamin D blood level is a primary cause of covid-19 severity or only a consequence associated to another population weakness being the primary severity cause.

In contrary the date of the boost is an intrapopulation observation and can be only triggered by a parameter globally affecting this population. As there is no temperature correlation but an impressive latitude correlation (fig. 2), the remaining common parameter affecting these populations at different times depending on the latitude is the sun UV daily dose.

The present study supports thus that lower vitamin D blood level is one of the primary cause of increased COVID-19 severity. By increasing the coronavirus load in the respiratory track, the contagiousness in the population is also increased explaining the wave boost.

The results are also in line with the low population mortality observed within ±35° latitudes [8], and also with the confinement success reported in Hubei which is located on the 31° latitude, regarding that these regions are most of the year above the sun UV daily dose threshold (fig. 3: black dot-dash line).

medRxiv preprint doi: https://doi.org/10.1101/2020.10.28.20221176; this version posted November 8, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.
It is made available under a CC-BY-NC-ND 4.0 International license .

Conclusion:

To the best of our knowledge, the present study is the first reported evidence that lower vitamin D blood level is one of the primary causes of covid-19 severity.

Europe, and US as well, are starting this autumn a long covid-19 crisis as they will get back above the October sun UV daily dose only end 2021 March.

The bet to reduce the pandemic severity during the coming winter using controlled preventive vitamin D supplementation could be considered [10].

References:

1. Grigalavicius M, Juzeniene A, Baturaite Z, Dahlback A, Moan J. Biologically efficient solar radiation: vitamin D production and induction of cutaneous malignant melanoma. Dermato-endocrinology. 2013;5:150-8.

2. Martineau Adrian R, Jolliffe David A, Richard HL et al (2017) Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. BMJ 356:i6583.

3. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, Bhattoa HP. Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. Nutrients. 2020 Apr;12(4):988.

4. Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. Aging Clinical and Experimental Research. 2020 May 6:1-4.

5. Carter SJ, Baranauskas MN, Fly AD. Considerations for Obesity, Vitamin D, and Physical Activity Amid the COVID-19 Pandemic. Obesity. 2020 May 21.

6. Boucher BJ. Vitamin D status as a predictor of Covid-19 risk in Black, Asian and other ethnic minority groups in the UK. Diabetes/Metabolism Research and Reviews. 2020 Jul 31:e3375.

7. Payne J, Ray R, Watson D, Delille C, Rimler E, Cleveland J, Lynn M, Tangpricha V, Srivastava S. Vitamin D insufficiency in diabetic retinopathy. Endocrine Practice. 2012 Mar 1;18(2):185-93.

8. Rhodes JM, Subramanian S, Laird E, Kenny RA. low population mortality from COVID-19 in countries south of latitude 35 degrees North supports vitamin D as a factor determining severity. Alimentary pharmacology & therapeutics. 2020;51:1434-7.

9. Lewiecki EM. Vitamin D and COVID-19: Is something better than nothing?. Osteoporosis and Sarcopenia. 2020. https://doi.org/10.1016/j.afos.2020.08.005