Title

U-shaped-aggressiveness of SARS-CoV-2: Period between onset of nonspecific-specific symptoms for COVID-19.

A population-based cohort study

Short Title

U-shaped aggressiveness of SARS-CoV-2

Authors

Dan Morgenstern-Kaplan[1], Bruno Buitano-Tang[1], Mercedes Martínez-Gil[1], Andrea Zaldívar-Pérez Pavón[1], Juan O. Talavera[2].

Affiliations

1. Centro de Investigación en Ciencias de la Salud Anáhuac (CICSA), Universidad Anáhuac México, CDMX, México

2. Department of Medical Education and Research, ABC Medical Center, Mexico City, Mexico.

Corresponding Author

Juan O. Talavera jotalaverap@abchospital.com,

Word Count: 2440

Abstract

Background: Early identification of different COVID-19 clinical presentations may depict distinct

pathophysiological mechanisms and guide management strategies.

Objective: To determine the aggressiveness of SARS-CoV-2 using symptom progression in COVID-19 patients.

Design: Historic cohort study of Mexican patients. Data from January-April 2020 were provided by the Health

Ministry.

Setting: Population-based. Patients registered in the Epidemiologic Surveillance System in Mexico.

Participants: Subjects who sought medical attention for suspicion of COVID-19. All patients were subjected to RT-

PCR testing for SARS-CoV-2.

Measurements: We measured the period between onset of nonspecific to specific symptoms for COVID-19

(PONSS) and compared it to the primary outcomes (mortality and pneumonia).

Results: 65,500 patients were included. Reported fatalities and pneumonia were 2176 (3.32%), and 11568

(17.66%), respectively. According to the PONSS, patients were distributed as follows: 14.89% in <24 hours,

43.25% between 1-3 days, 31.87% between 4-7 days and 9.97% >7 days. The distribution for mortality and

pneumonia was 5.2% and 22.5% in <24 hours, 2.5% and 14% between 1-3 days, 3.6% and 19.5% between 4-7 days,

4.1% and 20.6% >7 days, respectively ($p<0.001$). Adjusted-risk of mortality was (OR [95% CI], p-value): <24

hours= 1.75 [1.55-1.98], $p<0.001$; 1-3 days= 1 (reference value); 4-7 days= 1.53 [1.37-1.70], $p<0.001$; >7 days=

1.67 [1.44-1.94], $p<0.001$. For pneumonia: <24 hours= 1.49 [1.39-1.58], $p<0.001$; 1-3 days= 1; 4-7 days= 1.48

[1.41-1.56], $p<0.001$; >7 days= 1.57 [1.46-1.69], $p<0.001$.

Limitations: Using a database fed by large numbers of people carries the risk of data inaccuracy. However, this

imprecision is expected to be random and data are consistent with previous studies.

Conclusion: The PONSS shows a U-shaped SARS-CoV-2 aggressiveness pattern. Further studies are needed to

corroborate the time-related pathophysiology behind these findings.

Key Words: COVID-19, Epidemiology, Public Health, Pandemic, Mortality

Introduction

Coronaviruses are single-stranded RNA organisms capable of infecting humans and other animal species (1,2). The most recently discovered coronavirus, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), is cause of the clinical entity denominated Coronavirus Disease 2019 (COVID-19). This virus initially spread in the Wuhan province in China and later to the rest of the world, causing a pandemic (3). Reported worldwide cases are continuously growing and currently (as of July 3^{rd}, 2020) there are over 10 million infected people confirmed and over 500,000 fatalities. Global reports reveal case-fatality rate of 4.8% and more than half of the cases are in the Americas region. In Mexico, over 230,000 cases have been reported, with over 28,000 fatalities and a case-fatality rate of 12.3%, which surpasses by far the global estimate (4).

Every human in the world is susceptible to infection, for as the mean age of infected patients is 47 years, 87% of patients lie between 30 and 79 years old. COVID-19 behaves more aggressively in older patients and in patients undergoing chronic medical conditions such as obesity, diabetes, hypertension and other cardiovascular diseases, increasing the risk of mortality in these populations (5,6). Approximately 80% of cases are asymptomatic with a mild disease course, while the other 20% can be accompanied of severe complications such as pneumonia, Acute Respiratory Distress Syndrome (ARDS) and other secondary infections. Among these severe cases, 80% correspond to people over 60 years. Many of these cases can be attributed to a severe clinical entity known as "cytokine storm", which causes a rise in serum levels of many pro-inflammatory mediators and provokes massive tissue damage in several vital organs (5,7).

In patients who developed severe symptoms, dyspnea was reported between 8-12 days after onset of symptoms, and some patients deteriorate into severe disease during the first week after onset of symptoms. This accelerated worsening has been hypothesized to be caused by the cytokine storm and to thrombotic events that may be caused by infection with SARS-CoV-2 (8).

Hospitalized patients have been thoroughly described and analyzed, with an average time between onset of symptoms to intubation of 14.5 days, and a time from intubation to death ranging from 4-5 days (5,7,9). A longer period between onset of symptoms and first contact seeking medical attention has been associated with a poorer outcome in these patients. However no in-depth studies have been conducted (10).

Until now, studies have been focused on patient-centered risk factors, while SARS-CoV-2 aggressiveness has been stablished as provoking 20% of severe and critic patients, however, there are still many unanswered questions

concerning the clinical aggressiveness behavior of SARS-CoV-2. This study focuses on progression of symptoms as

a marker of such aggressiveness, using the period between onset of nonspecific to specific symptoms for COVID-19

(PONSS) to determine the risk of severe disease and mortality.

METHODS

Study Design and Data Source

A historic cohort study of Mexican patients that were classified as a suspect case of COVID-19 and sought medical

attention in either public or private health services in Mexico, was analyzed. Data was provided by the General

Directorate of Epidemiology of the Mexican Health Ministry, which is deidentified, publicly available online, and

registers all patients in the Epidemiologic Surveillance System of the 32 federal states in Mexico. This analysis was

done in all cases registered in this dataset up until April 25[th], 2020, with a total of 65,500 patients (11). The

Institutional Review Board of Anahuac University (Mexico City, Mexico) approved this study.

Variable Definition

Our dataset includes demographic characteristics such as age, gender, location, health sector, underlying medical

conditions (obesity, diabetes, COPD, asthma, immunosuppression, hypertension, cardiovascular, chronic kidney

diseases, and other comorbid diseases), pregnancy status, tobacco use, and indigenous language speaker.

Furthermore, it references the dates of the onset of symptoms and the date of medical attention, including hospital

admission, as well as the presence of pneumonia. Regarding in-hospital decisions, data include results of RT-PCR

testing for SARS-CoV-2 (reported as positive, negative or pending), admission to the Intensive Care Unit (ICU) and

requirement of mechanical ventilation. Finally, the date of death of all deceased patients is reported.

Data Analysis

Coding and substitution of variables

The state where the patient sought attention was recoded according to the socio-economic level of that particular

state into low, middle and high level, based on the Gross Domestic Product (GDP) of that state, as reported by the

National Institute of Statistics and Geography (INEGI) (12). The health sector variable was recoded in three

categories: private, with social security and without social security.

To establish the date of symptoms specific to COVID-19, we assumed that the population followed the

recommendations of the government to seek medical attention only when presenting COVID-19 specific-symptoms.

These recommendations were: Stay-at-home orders unless two of these three symptoms were present: 1) cough, 2)

fever or 3) headache, plus one or more of the following: 1) breathing difficulty, 2) sore or burning throat, 3) runny

nose, 4) red eyes, 5) pain in muscles or joints, or 6) being part of these high-risk groups: pregnancy, <5 or ≥60 years

old, or having a chronic disease such as hypertension, diabetes mellitus, cancer or HIV. These indications were

broadcast by television, radio, newspapers and internet since the beginning of the pandemic until July 1, 2020; up to

this date the government issued nationwide stay-at-home orders and only essential activities were permitted.

Therefore, the period between onset of nonspecific to specific symptoms for COVID-19 (PONSS) was calculated

between the date of onset of symptoms and the date the patients sought medical attention. Initially, PONSS was

categorized in days (<1, 1, 2, 3, etc.), but to improve comprehension and data management, adjacent days whose

frequency of death remained in similar proportions, were grouped into 4 categories (<24 hours, 1-3 days, 4-7 days

and >7 days). The primary outcomes were mortality and pneumonia. The presence of pneumonia was used as an

indicator of severe disease as reported in previous studies (5,7).

Missing data were substituted using the mode for the following categorical variables: Health Sector (338 patients,

0.5%), indigenous language speaker (1241, 2%), tobacco use (242, 0.4%), pregnancy status (166, 0.3%), diabetes

(255, 0.4%), COPD (245, 0.4%), asthma (251, 0.4%), immunosuppression (259, 0.4%), hypertension (241, 0.4%),

cardiovascular disease (252, 0.4%), obesity (220, 0.3%), chronic kidney disease (245, 0.4%), other comorbid

condition (331, 0.5%), admission to the ICU (12, <0.01%) and mechanical ventilation (12, <0.01%).

Statistical Analysis

Demographic features and comorbid conditions were initially compared between the four categories of PONSS; age

was analyzed with a one-way ANOVA and the rest of the variables with Chi squared test analysis. Afterwards, we

performed a bivariate analysis comparing the four categories of PONSS with the medical decisions made (result of

PCR testing for SARS-CoV-2, hospital admission, ICU, mechanical ventilation) and outcomes (mortality and

pneumonia) using the Chi squared test.

Finally, the four categories of PONSS against primary outcomes -mortality and pneumonia-, were compared using a

multivariable logistic regression model. The model was adjusted in five steps for the following variables: age,

gender, indigenous language speaker, state's socioeconomic status, pregnancy, tobacco use, obesity, hypertension,

diabetes, asthma, COPD, cardiovascular disease, chronic kidney disease, immunosuppression, other comorbid

conditions. This model was repeated in four groups of patients within the sample, depending on their RT-PCR

testing result for SARS-CoV-2:

a.   All patients: Every patient in the dataset regardless of their test result

b.   Positives: Only patients with a positive test result

c.   Negatives: Only patients with a negative test result

d.   Pending: Only patients with pending results of the test

The PONSS used as reference in the regression models, was 1-3 days based in the lower rate of mortality, observed

in the results of the bivariate analysis. Each logistic regression model is presented with the Odds Ratio (OR) and its

respective 95% Confidence Interval ($CI_{95\%}$). Statistical significance was set at $p < 0.05$ and performed with SPSS

version 25.0 (IBM). The full model for the group of all patients can be found in the Supplementary Material Section.

RESULTS

The study population included 65,500 patients. Among them, the average age was 41 ± 17 years, 50.2%, were

women, 55.8% belonged to a high socioeconomic level, 27.7% to a medium and 16.5% to a low one, 4.6% of

patients were treated on a private health institution, 37.7% in a facility for patients with social security and 57.7%

attended to a public hospital for patients without social security. Of all the patients, 41% had at least one

comorbidity, hypertension being the most frequent in 17%, followed by obesity in 15.6% and diabetes 12.8%. In

addition, 9.9% reported tobacco use and 2.3% of women were pregnant. Mortality was observed in 2176 patients

(3.32%), and Pneumonia in 11568 patients (17.66%).

According to PONSS patients were distributed as follows: 14.89% in <24 hours, 43.25% between 1-3 days, 31.87%

between 4-7 days and 9.97% after 7 days, with no significant difference by gender. We compared PONSS against

demographic features and comorbidities. A PONSS of <24 hours was more frequent in older patients (25.7% in

patients > 80 years old vs. 15% in <30 years old) reversing in the period of 1-3 days (41.5% vs 48.4 %,

respectively), and returning to the initial behavior in subsequent periods. This same pattern was observed when

comparing PONSS with the presence of all comorbidities, except for asthma and obesity. Demographic

Characteristics of all patients are summarized in Table 1.

Table 1. Patient demographic characteristics according to the period between onset of nonspecific to specific symptoms for COVID-19 (PONSS)										
		Period between onset of nonspecific to specific symptoms for COVID-19 (PONSS)								Global P Value
		<24 Hrs (n=9759)		1-3 Days (n=28331)		4-7 Days (n=20877)		>7 Days (n=6533)		
		N	%	N	%	N	%	N	%	
Gender	**Female**	4686	14.3%*	14371	43.7%	10562	32.1%	3260	9.9%	**<0.001**
	Male	5073	15.6%	13960	42.8%	10315	31.6%	3273	10.0%	
Age	**< 30**	2469	15.0%*	7965	48.4%	4732	28.7%**	1302	7.9%***	**<0.001**
	30-49	3745	12.8%	12813	43.6%	9785	33.3%	3016	10.3%	
	50-59	1430	14.7%	3875	39.7%	3321	34.1%	1123	11.5%	
	60-69	1095	19.8%	2006	36.2%	1785	32.2%	656	11.8%	
	70-79	615	21.9%	1018	36.3%	863	30.7%	311	11.1%	
	> 80	405	25.7%	654	41.5%	391	24.8%	125	7.9%	
	Mean (SD)	43(20)*		40 (18)		42 (17)**		44 (16)***		**<0.001**
State's Socioeconomic Status	**High**	5532	15.1%*	15745	43.1%	11352	31.1%**	3921	10.7%***	**<0.001**
	Medium	2878	15.8%	8027	44.2%	5868	32.3%	1402	7.7%	
	Low	1349	12.5%	4559	42.3%	3657	33.9%	1210	11.2%	
Health Sector	**Private**	468	15.6%*	1228	40.9%	885	29.5%**	418	13.9%***	**<0.001**
	With SS	4584	18.6%	10515	42.6%	7274	29.5%	2321	9.4%	
	Without SS	4707	12.5%	16588	43.9%	12718	33.6%	3794	10.0%	
Indigenous Language Speaker	**Yes**	98	14.4%	277	40.8%	242	35.6%	62	9.1%	0.20
	No	9661	14.9%	28054	43.4%	20635	31.8%	6471	10.0%	
Pregnancy	**Yes**	139	18.3%	383	50.5%	190	25.1%**	46	6.1%***	**<0.001**
	No	9620	14.9%	27948	43.2%	20687	32.0%	6487	10.0%	
Tobacco Use	**Yes**	946	14.5%	2786	42.8%	2122	32.6%	661	10.1%	0.49
	No	8813	14.9%	25545	43.3%	18755	31.8%	5872	10.0%	
Diabetes	**Yes**	1514	18.1%*	3307	39.5%	2684	32.1%**	867	10.4%***	**<0.001**
	NO	8245	14.4%	25024	43.8%	18193	31.8%	5666	9.9%	
COPD	**Yes**	401	22.6%*	685	38.6%	535	30.2%	153	8.6%	**<0.001**
	No	9358	14.7%	27646	43.4%	20342	31.9%	6380	10.0%	
Asthma	**Yes**	394	12.4%*	1422	44.6%	1034	32.5%	336	10.5%	**0.001**
	No	9365	15.0%	26909	43.2%	19843	31.8%	6197	9.9%	
Immunosuppression	**Yes**	455	26.1%*	697	40.0%	429	24.6%**	163	9.3%	**<0.001**
	No	9304	14.6%	27634	43.3%	20448	32.1%	6370	10.0%	
Hypertension	**Yes**	1932	17.4%*	4415	39.7%	3593	32.3%**	1194	10.7%***	**<0.001**
	No	7827	14.4%	23916	44.0%	17284	31.8%	5339	9.8%	
Cardiovascular Disease	**Yes**	470	22.0%*	820	38.3%	627	29.3%	224	10.5%***	**<0.001**
	No	9289	14.7%	27511	43.4%	20250	32.0%	6309	10.0%	
Obesity	**Yes**	1381	13.5%	4093	40.0%	3633	35.5%**	1119	10.9%***	**<0.001**
	No	8378	15.2%	24238	43.9%	17244	31.2%	5414	9.8%	
Chronic Kidney Disease	**Yes**	435	28.1%*	612	39.5%	374	24.1%**	129	8.3%	**<0.001**
	No	9324	14.6%	27719	43.3%	20503	32.1%	6404	10.0%	

Other Comorbid Condition	Yes	662	19.0%[*]	1440	41.3%	1059	30.4%	323	9.3%	<0.001
	No	9097	14.7%	26891	43.4%	19818	32.0%	6210	10.0%	

SS= Social Security. COPD=Chronic Obstructive Pulmonary Disease. SD=Standard Deviation. Statistical

*Significance p<0.05, * Significant Difference between periods of <24 hrs. and 1-3 days. ** Significant Difference*

*between periods of 4-7 days and 1-3 days.*** Significant Difference between periods of >7 days and 1-3 days.*

Table 2 shows the initial medical decisions according to their PONSS. More people were hospitalized in the first 24

hours (43.2%), with a drop towards the period of 1-3 days (19.7%), and a slight increase in subsequent days. A

similar phenomenon is observed in terms of admission to the ICU, with admission being 2.8% when the period is

<24 hours, falling to 1.8% in 1-3 days, and gradually increasing to 2.7% in 4-7 days and 3.2% in >7 days. The

proportion of patients under mechanical ventilation steadily increased over time, starting from 1.6% in the period of

<24 hours, up to 2.9% in the period of >7 days.

Table 2. Medical Decisions according period between onset of nonspecific to specific symptoms for COVID-19

(PONSS**).**

		Period between onset of nonspecific to specific symptoms for COVID-19 (PONSS)PONSS								Global P Value
		<24 Hrs (n=9759)		1-3 Days (n=28331)		4-7 Days (n=20877)		>7 Days (n=6533)		
		N	%	N	%	N	%	N	%	
Type of care	Ambulatory Care	5547	56.8%[*]	22742	80.3%	15674	75.1%[**]	4837	74.0%[***]	<0.001
	Hospital Admission	4212	43.2%	5589	19.7%	5203	24.9%	1696	26.0%	
Admission to the ICU		271	2.8%[*]	503	1.8%	563	2.7%[**]	208	3.2%[***]	<0.001
Mechanical Ventilation		155	1.6%	479	1.7%	525	2.5%[**]	189	2.9%[***]	<0.001
Result of RT-PCR Test[+]	Not Positive to SARS-CoV 2	7036	81.4%[*]	19801	80.4%	12672	69.8%[**]	3910	66.5%[***]	<0.001
	Positive to SARS-CoV-2	1601	18.6%	4818	19.6%	5462	30.2%	1961	33.5%	

ICU= Intensive Care Unit. PCR= Polymerase Chain Reaction. SD= Standard Deviation

*Statistical Significance p<0.05, *Significant Difference between periods of <24 hrs and 1-3 days. ** Significant*

*Difference between periods of 4-7 days and 1-3 days. *** Significant Difference between periods of >7 days and 1-3*

days. + Undefined were not included.

Table 3 and Figure 1 show the risks for mortality and pneumonia related to PONSS. A "U-shaped distribution" was

observed according to PONSS (<24 hrs., followed by 1-3 days, 4-7, and >7). The proportion of patients with

Mortality was 5.2%, 2.5%, 3.6%, and 4.1% (p<0.001), and for Pneumonia 22.5%, 14%, 19.5% and 20.6%

(p<0.001).The adjusted-risk of mortality for all patients evaluated for suspicion of COVID-19 according to PONSS,

was for <24 hours OR of 1.75 (95% CI, 1.55 to 1.98, p = <0.001), 1-3 days OR = 1 (reference value), 4-7 days, OR

1.53 (1.37-1.70, p = <0.001), and >7 days, OR 1.67 (1.44-1.94, p = <0.001), while for Pneumonia it was for <24

hours OR of 1.49 (95% CI, 1.39 to 1.58, p = <0.001), 1-3 days OR = 1 (reference value), 4-7 days, OR 1.48 (1.41-

1.56, p = <0.001), and >7 days, OR 1.57 (1.46-1.69, p = <0.001).

Table 3. Mortality and Pneumonia among patients with COVID-19 according to the period between onset of

nonspecific to specific symptoms (PONSS).

Variable All Patients (N=65,500)	Mortality			Pneumonia		
PONSS (N)	%*	OR (95% CI)	p-value	%*	OR (95% CI)	p-value
< 24 Hours (9759)	5.2%	1.75 (1.55-1.98)	<0.001	22.5%	1.49 (1.39-1.58)	<0.001
1-3 Days (28331)	2.5%	1	1	14%	1	1
4-7 Days (20877)	3.6%	1.53 (1.37-1.70)	<0.001	19.5	1.48 (1.41-1.56)	<0.001
> 7 Days (6533)	4.1%	1.67 (1.44-1.94)	<0.001	20.6%	1.57 (1.46-1.69)	<0.001
"p value"	<0.001			<0.001		
SUBGROUP ANALYSIS						
Positive Test (N=13,842)						
PONSS	OR (95% CI)		p-value	OR (95% CI)		p-value
< 24 Hours	2.11 (1.75-2.55)		<0.001	1.66 (1.45-1.90)		<0.001
1-3 Days	1		1	1		1
4-7 Days	1.42 (1.22-1.65)		<0.001	1.69 (1.53-1.85)		<0.001
> 7 Days	1.22 (1.00-1.49)		0.053	1.83 (1.62-2.07)		<0.001
Negative Test (N=43,419)						
PONSS	OR (95% CI)		p-value	OR (95% CI)		p-value
< 24 Hours	1.55 (1.29-1.86)		<0.001	1.58 (1.46-1.70)		<0.001
1-3 Days	1		1	1		1
4-7 Days	1.00 (0.83-1.21)		0.967	1.10 (1.01-1.16)		0.038
> 7 Days	1.39 (1.10-1.81)		0.014	1.11 (1.00-1.24)		0.061
Test Result Pending (N=8,239)						
PONSS	OR (95% CI)		p-value	OR (95% CI)		p-value
< 24 Hours	1.41 (0.68-2.90)		0.357	0.80 (0.64-0.99)		0.043
1-3 Days	1		1	1		1
4-7 Days	1.22 (0.70-2.13)		0.484	2.03 (1.77-2.34)		<0.001
> 7 Days	1.71 (0.78-3.74)		0.182	1.96 (1.57-2.44)		<0.001

This Global Multiple Logistic Regression Model is adjusted for all demographic characteristics and comorbid

conditions present in the patients. Adjustments for the group of all patients can be found in the Supplementary

Material Section.

*Bivariate analysis between PONSS vs. Mortality or Pneumonia.

Figure 1. U-Shaped Distribution of the odds ratio for the primary outcomes (mortality/pneumonia) vs. PONSS.

Caption: Association according to the period between onset of nonspecific to specific symptoms for COVID-19

(PONSS) and the primary outcomes of the study (Mortality and Pneumonia), including all patients. A U-shaped

distribution is observed, with higher OR for PONSS <24 hours and ≥4 days.

DISCUSSION

In this study, we found an association concerning the period between onset of nonspecific to specific symptoms for

COVID-19 (PONSS) with the risk of severe disease and mortality in patients with suspected COVID-19. A "U

shaped" distribution was observed, with a high risk of death and pneumonia when PONSS is <24 hours (OR 1.75,

and 1.49, respectively), with a decrease of this risk in 1-3 days (OR 1), and with an additional rise in subsequent

periods of 4-7 days (OR 1.53, and 1.48) and >7 days (OR 1.67 and 1.57).

The increased risk of mortality and pneumonia observed in patients with PONSS <24 hours, may be associated with

the presence of a cytokine storm, which has been previously described as an early factor for severity[13]. This

phenomenon is due to the uncontrolled release of pro-inflammatory mediators that lead to apoptosis of epithelial and

endothelial lung cells, causing vascular extravasation, alveolar edema and hypoxia. This inflammatory response in

conjunction with the production of reactive oxygen species triggers an acute respiratory distress syndrome (ARDS)

leading to pulmonary fibrosis and death[14]. This could support the pharmacodynamic basis for the use of

corticosteroids as adjuvant therapy in patients with COVID-19, which has been reported in other studies [15].

Chronic use of inhaled corticosteroids may be the reason why asthmatic patients manifest less severe symptoms[16],

which was consistent with our results (See Supplementary Materials).

The increased risk of mortality and pneumonia in patients with PONSS ≥4 days, could be explained by the

thrombotic events that have been reported in patients with COVID-19. These events are caused by the excessive

inflammation produced by the virus and platelet activation with accompanying endothelial damage[17,18]. This

occurs once the virus has colonized the respiratory system, impairing microvascular permeability, helping it spread

even further. Hemostatic disorders are established by the presence of thrombocytopenia, and an increase in the D-

dimer and fibrinogen, for which the use of antithrombotic therapies has been suggested[17,19].

Additionally, an increased risk of lung superinfections must be considered. So far, bacterial and fungal pneumonias

have been the most common etiologies. A study conducted in Wuhan, China reported a rate of lung superinfection

from 5-27% in adults with COVID-19(20). Historically, superinfections have been associated with increased

mortality in other viral respiratory infections, such as influenza(21).

Using a database fed by large numbers of people carries its risk, such as data inaccuracy. However, this imprecision

is expected to be random and data are consistent with results of previous studies. Furthermore, we set April 25th,

2020 as our cut-off date with the aim of including patients treated at an early stage of the pandemic in Mexico, at a

time when hospitals were not yet working at overcapacity. This increases the probability of good quality of

healthcare, decreases confounding factors for the outcomes evaluated because all required medical decisions could

be made and were not limited by medical resources available at the time (i.e. number of ventilators or ICU beds).

Plenty of studies have described the incubation period and hospital stay of affected patients(5,22,23). However,

nobody has considered the progression of symptoms in patients with COVID-19 (PONSS), as a guide for explaining

the time-specific pathophysiology associated with the U-Shaped SARS-CoV-2 aggressiveness. Further studies are

needed to corroborate the time-related pathophysiology behind these findings. Eventually, this could help identify

specific therapies aimed towards the temporal progression of the disease.

Acknowledgments

The authors report no conflicts of interest regarding the present study. The Dataset used for all statistical analyses is

available in the Mendeley Data Repository Online at: http://dx.doi.org/10.17632/k6cw45366d.1

References

1.   Tyrrell DA, Bynoe ML. Cultivation of viruses from a high proportion of patients with colds. Lancet.

244   1966;1(7428):76–7.

245   2.   Velavan TP, Meyer CG. The COVID-19 epidemic. Trop Med Int Heal. 2020;25(3):278–80.

246   3.   Battegay M, Kuehl R, Tschudin-Sutter S, Hirsch HH, Widmer AF, Neher RA. 2019-novel Coronavirus

247   (2019-nCoV): estimating the case fatality rate - a word of caution. Swiss Med Wkly.

248   2020;150(February):w20203.

249   4.   World Health Organization. WHO Coronavirus Disease (COVID-19) Dashboard. WHO. 2020.

250   5.   Wang L, He W, Yu X, Hu D, Bao M, Liu H, et al. Coronavirus disease 2019 in elderly patients:

251   Characteristics and prognostic factors based on 4-week follow-up. J Infect. 2020;(xxxx).

252   6.   Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, Evaluation and Treatment

253   Coronavirus (COVID-19) - StatPearls - NCBI Bookshelf. StatPearls. 2020. p. 1–17.

254   7.   Zhou P, Yang X Lou, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a

255   new coronavirus of probable bat origin. Nature. 2020;579(7798):270–3.

256   8.   Centers For Disease Control. Interim Clinical Guidance for Management of Patients with Confirmed

257   Coronavirus Disease (COVID-19). 2020.

258   9.   Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult

259   inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet. 2020;395(10229):1054–

260   62.

261   10.   Liang W, Guan W, Li C, Li Y, Liang H, Zhao Y, et al. Clinical characteristics and outcomes of hospitalised

262   patients with COVID-19 treated in Hubei (epicentre) and outside Hubei (non-epicentre): a nationwide

263   analysis of China. Eur Respir J. 2020 Jun;55(6):2000562.

264   11.   Direccion General de Epidemiologia, Secretaría de Salud. Datos Abiertos - Bases Históricas. 2020.

265   12.   INEGI. Comunicado De Prensa Núm . 384 / 19 31 De Julio De 2019 El Inegi Da a Conocer Los Resultados

266   De La Encuesta Nacional De Ingresos Y Gastos De Los Hogares (Enigh) 2018. Comun prensa INEGI.

267   2019;

268   13.   Henderson LA, Canna SW, Schulert GS, Volpi S, Lee PY, Kernan KF, et al. On the alert for cytokine storm:

269   Immunopathology in COVID-19. Arthritis Rheumatol. 2020;72(7):1059–63.

270   14.   Sun X, Wang T, Cai D, Hu Z, Chen J, Liao H, et al. Cytokine storm intervention in the early stages of

271   COVID-19 pneumonia. Cytokine Growth Factor Rev. 2020;53(March):38–42.

272   15.   Horby P, Lim WS, Emberson J, Mafham M, Bell J, Linsell L, et al. Effect of Dexamethasone in

273   Hospitalized Patients with COVID-19: Preliminary Report. medRxiv. 2020 Jan;2020.06.22.20137273.

274   16.   Liu S, Zhi Y, Ying S. COVID-19 and Asthma: Reflection During the Pandemic. Clin Rev Allergy Immunol.

275   2020;

276   17.   Bikdeli B, Madhavan M V., Jimenez D, Chuich T, Dreyfus I, Driggin E, et al. COVID-19 and Thrombotic

277   or Thromboembolic Disease: Implications for Prevention, Antithrombotic Therapy, and Follow-Up: JACC

278   State-of-the-Art Review. J Am Coll Cardiol. 2020;75(23):2950–73.

279   18.   Cannegieter SC, Klok FA. COVID-19 associated coagulopathy and thromboembolic disease: Commentary

280   on an interim expert guidance. Res Pract Thromb Haemost. 2020;4(4):439–45.

281   19.   Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. Clin Immunol.

282   2020;215(April).

283   20.   Clancy CJ, Nguyen MH. Coronavirus Disease 2019, Superinfections, and Antimicrobial Development:

284   What Can We Expect? Clin Infect Dis. 2020;(April).

285   21.   Rynda-Apple A, Robinson KM, Alcorn JF. Influenza and bacterial superinfection: Illuminating the

286   immunologic mechanisms of disease. Infect Immun. 2015;83(10):3764–70.

287   22.   Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The incubation period of coronavirus

288   disease 2019 (CoVID-19) from publicly reported confirmed cases: Estimation and application. Ann Intern

289   Med. 2020;172(9):577–82.

290   23.   Sun Q, Qiu H, Huang M, Yang Y. Lower mortality of COVID-19 by early recognition and intervention:

291   experience from Jiangsu Province. Ann Intensive Care. 2020;10(1):2–5.

302   **Supplementary Material Section Captions**

303   S1 Table. Stepwise analyses for binary logistic regression with Mortality as outcome for all patients in the study.

304   S2 Table. Stepwise analyses for binary logistic regression with Pneumonia as outcome for all patients in the study.