Cerebrospinal fluid total tau levels indicate aberrant neuronal plasticity in

Alzheimer's disease.

Pieter Jelle Visser[1,2,3], Lianne M. Reus[1], Johan Gobom[4,5], Iris Jansen[6], Ellen Dicks[1], Magda Tsolaki[7], Frans R.J. Verhey[2], Julius Popp[8,9], Pablo Martinez-Lage[10], Rik Vandenberghe[11,12], Alberto Lleó[13], José Luís Molinuevo[14, 15], Sebastiaan Engelborghs[16, 17], Yvonne Freund-Levi[3,18], Lutz Froelich[19], Kristel Sleegers[20,21], Valerija Dobricic[22], Shengjun Hong[22], Simon Lovestone[23], Johannes Streffer[24], Stephanie J.B. Vos[2], Isabelle Bos[1, 2], ADNI*, August B. Smit[25], Kaj Blennow[4,5], Philip Scheltens[1], Charlotte E. Teunissen[26], Lars Bertram[22,27,]Henrik Zetterberg [2,3, 28,29], and Betty M. Tijms[1].

1. Alzheimer Center Amsterdam, Department of Neurology, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, the Netherlands;
2. Alzheimer Center Limburg, School for Mental Health and Neuroscience, Maastricht University, Maastricht, the Netherlands;
3. Department of Neurobiology, Care Sciences and Society, Division of Neurogeriatrics, Karolinska Institutet, Stockholm Sweden;
4. Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden;
5. Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden;
6. Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, VU University, Amsterdam, the Netherlands;
7. 1st Department of Neurology, AHEPA University Hospital, Makedonia, Thessaloniki, Greece;
8. University Hospital Lausanne, Lausanne, Switzerland;
9. Department of Geriatric Psychiatry, University Hospita of Psychiatry and University of Zürich, Zürich, Switzerland;
10. Fundación CITA-Alzhéimer Fundazioa, San Sebastian, Spain;
11. Neurology Service, University Hospitals Leuven, Leuven, Belgium;
12. Laboratory for Cognitive Neurology, Department of Neurosciences, KU Leuven, Leuven, Belgium;
13. IIB-Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autonoma de Barcelona, Barcelona Spain;
14. Barcelonaβeta Brain Research Center (BBRC), Pasqual Maragall Foundation, Barcelona, Spain;
15. Alzheimer's Disease Unit and Other Cognitive Disorders Unit, Hospital Clinic de Barcelona, Barcelona, Spain;
16. Reference Center for Biological Markers of Dementia (BIODEM), Institute Born-Bunge, University of Antwerp, Belgium;
17. Department of Neurology, UZ Brussel and Center for Neurosciences, Vrije Universiteit Brussel, Brussels, Belgium;
18. Department of Psychiatry at School of Medical Sciences Örebro University

19. Department of Geriatric Psychiatry, Zentralinstitut für Seelische Gesundheit, University of Heidelberg, Mannheim, Germany;
20. Complex Genetics Group, VIB Center for Molecular Neurology, VIB, Antwerp, Belgium;
21. Department of Biomedical Sciences, University of Antwerp, Antwerp, Belgium;
22. Lübeck Interdisciplinary Platform for Genome Analytics, Institutes of Neurogenetics and Cardiogenetics, University of Lübeck, Lübeck, Germany;
23. University of Oxford, Oxford, United Kingdom (currently at Jansen R&D);
24. Biopharma SPRL, Brain-l'Alleud, Belgium;
25. Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands;
26. Neurochemistry laboratory, Department of Clinical Chemistry, Amsterdam University Medical Centers (AUMC), Amsterdam Neuroscience, Netherlands;
27. Lübeck Interdisciplinary Platform for Genome Analytics, Institutes of Neurogenetics and Cardiogenetics, University of Lübeck, Lübeck, Germany;
28. Center for Lifespan Changes in Brain and Cognition, Dept. of Psychology, University of Oslo, Oslo, Norway;
29. Department of Neurodegenerative Disease, UCL Institute of Neurology, London, United Kingdom;
30. UK Dementia Research Institute at UCL, London, United Kingdom.

Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Please address correspondence to:

PJ Visser, MD, PhD

Alzheimer Center and Department of Neurology, Amsterdam University

Medical Centers, Amsterdam Neuroscience, Amsterdam PO Box 7057, 1007

MB, Amsterdam, the Netherlands

Alzheimer Center Limburg, School for Mental Health and Neuroscience,

Maastricht University, Maastricht, the Netherlands

Email: pj.visser@amsterdamumc.nl; pj.visser@maastrichtuniversity.nl

Abstract

Alzheimer's disease (AD) is characterised by abnormal amyloid beta and tau processing. Previous studies reported that cerebrospinal fluid (CSF) total tau (t-tau) levels vary between patients. Here we show that CSF t-tau variability is associated with distinct impairments in neuronal plasticity mediated by gene repression factors SUZ12 and REST. AD individuals with abnormal t-tau levels have increased CSF concentrations of plasticity proteins regulated by SUZ12 and REST. AD individuals with normal t-tau, on the contrary, have decreased concentrations of these plasticity proteins and increased concentrations in proteins associated with blood-brain and blood CSF-barrier dysfunction. Genomic analyses suggested that t-tau levels in part depend on genes involved in gene expression. The distinct plasticity abnormalities in AD as signaled by t-tau urge the need for personalised treatment.

According to the amyloid cascade hypothesis, AD starts with amyloid beta (Aβ) aggregation followed by tau pathology (*1*). Between onset of Aβ aggregation and dementia, the disease has a preclinical stage of 10 years in which cognition is normal and a prodromal stage of 4 years with mild cognitive impairment (*2*). Increased CSF total tau (t-tau) levels are supposed to reflect axonal loss and disease severity, as tau stabilizes axonal microtubules (*3*). Still, CSF t-tau is abnormal in 50% of preclinical AD individuals, when neurodegeneration is limited (*4*), and normal in 25% of individuals with prodromal and dementia AD (*5,6*). An alternative explanation of variability in CSF t-tau levels may be differences in synaptic activity and neuronal plasticity (*7,8*). As tau-related mechanisms are emerging as new treatment targets for AD, it is crucial to understand drivers of CSF tau dysregulation. We hypothesized that differences in CSF t-tau levels are associated with neuronal plasticity markers, in particular in early AD stages.

We performed a proteomic, genomic and imaging study including 1380 individuals in the AD continuum (*9*), defined by abnormal CSF $A\beta_{1-42}$, in the preclinical (n=280), prodromal (n=729) and dementia stage (n=371) and 380 controls with normal cognition and normal CSF $A\beta_{1-42}$ and t-tau from the Alzheimer's Disease Neuroimaging Initiative (ADNI) and the EMIF-AD Multimodality Biomarker Discovery study (*3*). Of the individuals with AD, 788 (57%) had abnormally high CSF t-tau levels and 592 normal t-tau levels. Abnormal and normal t-tau groups showed generally similar demographics and baseline cognitive performance (sTable 1a).

We first determined stability of t-tau levels over time in a subset of 499 ADNI participants with repeated CSF sampling up to 7 years. CSF t-tau increased in each clinical stage but this increase did not differ between abnormal and normal t-tau groups, and also not from controls (sTable2b, sFigure 1a). Tau status at last measurement (on average 2.4 years after baseline), remained normal in 80% individuals with initially normal t-tau and those who became abnormal scored at baseline just below the cut-point (sTable 2b, sFigure 1b). This suggests that CSF t-tau levels reflect a trait rather than a stage.

We then studied whether the CSF proteome differed for t-tau subgroups in 559 individuals with proteomic data (sTable 1b). This included targeted proteomics of 248 proteins in ADNI and targeted and untargeted proteomic measures of 1458 proteins in EMIF-AD (sTable 2c). Compared to controls, AD individuals with abnormal CSF t-tau had increased levels of 33% (ADNI) to 52% (EMIF-AD) of the proteins measured, and decreased levels of 1 to 3% proteins (figure 1, sTable 2a,c). AD individuals with normal CSF t-tau, showed an opposite pattern with increased levels of 3 to 4% of the proteins and decreases in 28% (ADNI) to 43% (EMIF-AD) of the proteins. Around 50% of the proteins increased in individuals with abnormal t-tau were decreased in individuals with normal t-tau in both cohorts. This indicates that both t-tau groups show disruptions in the same molecular processes, but in opposite directions. Enrichment analyses using the Gene Ontology biological processes (GO-BP) for CSF proteins with increased levels in AD individuals with abnormal t-tau relative to controls, showed involvement of neuronal plasticity-related processes such as nervous system development, axongenesis, synapse assembly, myelination, gliogenesis, angiogenesis, MAPK signaling, cell-cycle, gene expression and

glycolysis (Figure 2; sTable 3a shows all biological processes; Table 3b and sFigure 2 show synapse related processes). Abnormality in each of these processes has previously been reported in AD (*10-15*), and our proteomic findings now show that they are all part of an aberrant plasticity response. Increased proteins further showed involvement of cytokine signaling, leukocyte activation, oxidative stress, mitochondrial dysfunction and apoptosis. Amyloid production was increased as well, reflected by increased levels of amyloid precursor protein (APP), $A\beta_{1-40}$, $A\beta_{1-38}$, increased levels and activity of beta-site amyloid precursor protein cleaving enzyme 1 (BACE1), and increased concentrations in substrates of BACE1, alpha secretase and gamma secretase, the main secretases involved in $A\beta$ metabolism (sTable2c). Increased $A\beta_{1-42}$ production is known to set off the amyloid cascade in autosomal dominant AD (*16*), and our data suggest that increased amyloid production may also play a role in sporadic AD. Increased $A\beta_{1-42}$ production is related to neuronal plasticity because APP and oligomeric $A\beta_{1-42}$ activate mitogenic MAPK/ERK signaling, and the APP intracellular domain (AICD) stimulates gene transcription through interaction with APBB1(*13,17,18*). To identify other potential drivers of increased protein levels we performed CHeA transcription factor-binding site enrichment analysis, which indicated SUZ12 (120 proteins, p-FDR corrected=1.62E-11) and REST (92 proteins, p-FDR corrected =1.04E-9) as most significantly enriched. SUZ12 and REST repress gene transcription through H3 histone acetylation, and play a role in neuronal development and neurodegeneration (*19,20*). SUZ12 and REST were previously associated with increased expression of plasticity genes in iPSC neuronal progenitor cells of sporadic AD individuals, of which 63 proteins were also increased in CSF in our study in individuals with abnormal $A\beta_{1-42}$ and t-tau (*15*). REST has also been associated with the increased expression of

plasticity related genes in tangle bearing neurons (*21*). Tau may increase gene expression directly or indirectly by increasing H3K9 acetylation, REST translocation or chromatin relaxation (*15,22,23*). In summary, the CSF proteome of AD individuals with abnormal t-tau reflects a disbalance of mitogenic signaling and repressive transcriptional activity leading to increased gene expression of a wide range of neuronal plasticity proteins. As this may also further increase expression of APP and t-tau, this could induce a vicious cycle.

In AD individuals with normal t-tau, the processes related to neuronal plasticity in particular were abnormally decreased with concomitant lower levels of APP, $A\beta_{1-40}$ and $A\beta_{1-38}$, and secretase substrates (Figure 2). Proteins with decreased concentrations also converged on SUZ12 and REST transcription factors, suggesting that increased gene repression activity is driving this CSF proteomic profile. Increased repressive activity of REST and SUZ12 has been reported in brain ischemia, seizures, and toxic insults supposedly reducing excitotoxicity (*19,20,24-26*). Among the 67 proteins with increased levels relative to controls, 20 proteins were associated with blood-brain barrier (BBB) dysfunction (*27*). They also included 25 proteins with high expression in choroid plexus (CP), and this CP overreprensentation was confirmed by ABAEnrichment analysis (minimum pFWER=0.004), which could suggest blood-CSF barrier (BCSFB) dysfunction. Increased proteins were mainly seen in the EMIF-AD cohort, as ADNI proteomics was targeted for brain specific proteins. Impaired BBB/BCSFB function may result from toxic effects of Aβ on cells that constitute these barriers (*28*). Barrier dysfunction reduces neuronal plasticity through impaired glucose transport, capillary perfusion, and neurogenesis (*28*), which is possibly reflected in the hypoplasticity signal we observed in CSF. When we repeated analysis

by comparing CSF protein concentrations between AD individuals with normal and abnormal t-tau we found enrichment for the same processes that differed for each group relative to controls (sTable2c, sFigure 3a). Analysis of cell-type specific proteins showed that plasticity proteins were strongly associated with neuron-specific proteins (Figure 2, sTable 2d). Stratified analyses for disease stage further showed that alternations in CSF proteome related to neuronal plasticity were already present in the preclinical stage (Figure 2, sTable 2c, e), and in both groups decreased with disease stage (sTable 2c, Figure 3, sFigure 3b). In the preclinical stage of individuals with normal t-tau levels, proteins associated with choroid plexus were increased (sTable 2e) and with increasing disease severity concentrations of BBB related proteins increased (Figure 3).

We next aimed to identify genetic factors related to increased t-tau in AD individuals in the combined ADNI and EMIF-AD cohorts (n=1067). *APOE* ε4, the major genetic AD risk factor, was more common in the abnormal t-tau group (66% vs 53% p<0.001), and in both AD groups *APOE* ε4carriership was more common than in controls (17%, p<0.001, sTable 1a). Compared to controls, both abnormal and normal t-tau groups showed higher AD polygenic risk scores (PGRS) across SNP inclusion thresholds (Figure 4, sTable4a) (*29*). In none of the clinical stages, PGRS differed between AD individuals with or without increased t-tau after correction for *APOE* ε4 carriership and age (sTable 4a). Thus, both groups have a similar presence of AD genetic risk variants. An exploratory GWAS within the AD group to identify potential genetic markers associated with abnormal t-tau in EMIF-AD and ADNI (combined by meta-analysis), showed tentative associations with SNPs in the *APBB2* gene (p-values 1.63-2.92E10-6, sTable 4b), which previously were associated with AD (*30*). *APBB2*

binds AICD and its overexpression has been associated with increased levels of $A\beta_{1-40}$ and AICD (*31*). The top 3 genes associated with abnormal t-tau from MAGMA gene-based analyses showed in addition to *ABPP2*, *TBC1D10B* (p=2.9E-04), a Rab GTPA activating protein involved in MAPK signaling, and *LRP3* (p=3.36E-04), a low-density lipoprotein receptor protein (sTable 4c). The top 3 GO-BP gene-sets associated with abnormal t-tau from MAGMA were positive regulation of cellular process (self-contained p=1.07E-04), protein K29-linked ubiquitination (p= 1.52E-04) and negative regulation of actin nucleation (p=1.57E-04, sTable 4d). Twenty-eight percent of GO-BPs associated with proteins that differed between AD individuals with abnormal and normal t-tau also differed between tau groups in MAGMA gene-set analysis (sTable 3a), suggesting that differences in CSF t-tau concentration have partly a genetic basis. Then we directly correlated CSF protein concentrations with subject level genetic risk scores (GRS) for the top 3 genes and GO-BP and 9 GO-BPs listed in figure 2 that differed between abnormal and normal t-tau groups in both CSF and MAGMA geneset analysis. Given the association of CSF profiles with REST and SUZ12 we also selected 2 histone acetylation GO-BPs with MAGMA self-contained p-value<0.05. The strongest positive correlations between genetic risk scores with CSF proteins were observed for gene expression (288 proteins), regulation of MAPK cascade (146 proteins), and histone H3 acetylation (127 proteins, sTable 5, figure 5). This further supports a genetic basis of t-tau differences between groups.

Finally, we studied effects of t-tau on disease progression using longitudinal ADNI data. Relative to controls, both preclinical AD groups showed faster cognitive decline and PET-amyloid accumulation (Figure 6, sFigure 4 and 5, sTable 6). Compared to

individuals with AD and normal t-tau, individuals with abnormal t-tau declined faster on cognitive tests, cortical thickness and glucose metabolism in prodromal AD and showed faster amyloid aggregation on PET in all stages. This faster accumulation of Aβ may result from hyperplasticity-related Aβ production.

In summary, CSF t-tau levels in AD are related to aberrant neuronal plasticity and dysfunction of the blood-brain and blood-CSF barriers. CSF proteomic analysis indicated that both AD individuals with abnormal and normal t-tau have disturbed mitogenic MAPK signaling and SUZ12/REST signaling, but in opposite ways. Furthermore, opposite CSF proteomic patterns were already observed in the preclinical stage, and persisted in prodromal and mild dementia stages, indicating that they are traits rather than a reflection of disease state.

Our exploratory GWAS, MAGMA analyses and the association between gene risk scores and CSF protein correlations suggest that t-tau levels in part depend on genes involved in gene expression. Larger samples are needed to confirm these results. This genetic pattern is different from that of AD risk genes, which are associated with lipid metabolism and immune activation (29). Possibly AD risk genes drive amyloid aggregation, and other genes determine whether an individual will follow a hypo- or hyperplastic trajectory.

Our finding that increased CSF t-tau was associated with increased levels of neuronal plasticity proteins provides further support for the role of tau in neuronal plasticity (21,23). Tau may influence gene expression itself (21-23), which may explain why knock-out of tau in AD mice models prevented hyperactivation (32,33).

A clinical implication of our results is that in the diagnostic framework for AD t-tau should be considered also a marker of dysregulated plasticity (*9*). Treatments for hyperexcitation and hyperplasticity, such as histone modification and anti-epileptic drugs, which are currently tested in clinical trials, may be only beneficial for AD individuals with abnormal t-tau before dementia onset only. AD individuals with normal t-tau and early barrier dysfunction would benefit the most from treatments focusing on prevention of barrier damage and may response differently on antibody therapy than AD individuals with increased t-tau.

2. Methods

2.1 Study participants

We selected individuals with AD pathology in CSF and a control group with normal cognition and normal AD biomarkers from the Alzheimer's Disease Neuroimaging Initiative (ADNI, n=961, adni.loni.usc.edu) and the EMIF-AD multimodality biomarker discovery (MBD) study (n=799) (*34*). ADNI started in 2003 as a public-private collaboration under the supervision of Principle Investigator Michael W. Weiner, MD. The primary goal of ADNI is to study whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological measures can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD) and has enrolled 2850 individuals so far, see www.adni-info.org for the latest information. The EMIF-AD MBD study aimed to identify novel markers for diagnosis and prognosis of predementia AD. It combined existing clinical data, samples and scans of 1218 individuals with normal cognition, MCI or mild dementia from prospective cohort studies (*34*). The study collected baseline clinical and neuropsychological data, MRI scans, and plasma, DNA and CSF samples and follow-up diagnosis and cognitive scores (*34*). The institutional review boards of all participating institutions approved the procedures for this study. Written informed consent was obtained from all participants or surrogates.

2.2 Group definition and staging

We selected individuals with AD pathology defined as abnormal CSF abeta 1-42 ($A\beta_{1-42}$), and we subdivided this group into abnormal and normal CSF total tau (t-tau) groups (see below for CSF analysis). Based on their cognitive performance, AD

individuals were classified in 3 clinical stages as preclinical AD (normal cognition), prodromal AD (mild cognitive impairment, i.e., MCI) and mild to moderate AD-type dementia according to study specific criteria (*34,35*). The control group consisted of individuals with normal cognition, as well as normal CSF $A\beta_{1-42}$ and t-tau levels.

2.3 Clinical assessment

As a measure of global cognition we used the Mini-Mental State Examination (MMSE) (*36*) and the ADAS-Cog 11 item version (*37*) (ADNI only). As a measure of memory function we used the delayed recall of the logical memory subscale II of the Wechsler Memory Scale delayed recall (ADNI) (*38*) or a center specific verbal word learning task (EMIF-AD) (*34*). We selected the Clinical Dementia Rating (CDR) scale sum of boxes score as a measure of disease severity (*39*).

2.4 CSF analysis

CSF samples were obtained as previously described (*34,40*).

<u>Abeta42 and tau measurements:</u> In EMIF-AD CSF $A\beta_{1-42}$ and t-tau were measured locally with INNOTEST ELISA or INNOBIA AlzBio3 INNOBIA AlzBio3 (Fujirebio, Ghent, Belgium) and in ADNI on the xMAP Luminex platform (Luminex Corp, Austin, TX) at the ADNI Biomarker Core laboratory at the University of Pennsylvania Medical Center. Abnormal $A\beta_{1-42}$ and t-tau were defined according to published cut-offs in ADNI (*41*). Cut-offs for $A\beta_{1-42}$ and t-tau were cohort-specific in EMIF-AD (*34*). Because EMIF-AD centers had used different approaches to determine $A\beta_{1-42}$ cut-offs, which could lead to bias, we determined for each cohort specific cut-offs using unbiased Gaussian mixture modelling (sTable 1c) (*42*). We

used longitudinal t-tau levels from ADNI (*40*), measured within the same batch within individuals, to assess changes over time.

Proteomic analysis ADNI: Details on protein measurements can be found at http://adni.loni.usc.edu/data-samples/biospecimen-data/. YKL-40, neurogranin, soluble APP beta, neurogranin, neurofilament light, alpha synuclein, phosphorylated alpha synuclein, HBB, CFH, sTREM2, VILIP-1, and BACE1 activity were measured by ELISA or related assays. 190 analytes were measured using the multiplex Human DiscoveryMAP panel from Rules Based Medicine (MAP-RBM) (*43*). This panel assesses proteins related to cytokines, chemokines, metabolic markers, hormones, growth factors, tissue remodeling proteins, angiogenesis markers, acute phase reactants, cancer markers, kidney damage markers, and CNS biomarkers. In addition, mass-spectroscopy measures of isoprostanes, $A\beta_{1-38}$, and $A\beta_{1-40}$ (*44*), and 567 peptides representing 222 proteins were performed (the "Biomarkers Consortium CSF Proteomics MRM data set") (*43,45*). Proteins and peptides were selected based upon their previous detection in CSF, relevance to AD, and results from the MAP-RBM. We used the quality checked and finalised 'Normalized Intensity' data as described in the "Data Primer" document at https://adni.loni.usc.edu/wp-content/uploads/2012/01/2011Dec28-Biomarkers-Consortium-Data-Primer-FINAL1.pdf

Proteomic analysis EMIF-AD: Neurogranin, neurofilament-light, abd YKL-40, abeta38 and abeta40 were measured by ELISA and $A\beta_{38}$ and $A\beta_{40}$, using the V-PLEX Plus Ab Peptide Panel 1 (6E10) Kit from Meso Scale Discovery (MSD, Rockville, MD) (*3*). Mass spectrometry was performed using tandem mass tag (TMT) technique with 10+1 plexing as previously described, using high-pH reverse phase HPLC for peptide prefractionation (*46*) (Tijms et al in press).The EMIF-AD mass

spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE partner repository with the dataset identifier

10.6019/PXD019910 (*47*). Normalised abundances with associated clinical data can

be requested from the EMIF-AD consortium (*3*). Peptides were mapped to 2535

proteins using the software ProteomeDiscoverer v.2.2. (Thermo Scientific), using

Mascot (MatrixScience) for protein identification (precursor Dm tolerance 15 ppm,

fragment tolerance 0.05 Da, max missed cleavage sites 2), searching the human subset

of the UniProtKB SwissProt database (www.uniprot.org). Percolator (MatrixScience)

was used for scoring peptide specific matches, and a strict 1% false discovery rate

(FDR) was set as threshold for identification. For reporter ion quantification the

following settings were used: Integration tolerance 20 ppm; Integration Method Most

Confident Centroid; normalize on the reference protein average. The median (IQR)

CV for these proteins was 5.6 (3.8, 8.0).

Protein selection: Proteins and protein fragments (ADNI) values were normalised

according the mean and standard deviation values of the control group. For ADNI, we

averaged peptides that mapped to the same protein into a composite protein score

when they correlated with r >.5, and included as a single peptide otherwise. When the

same protein was measured by different platforms in ADNI, values were averaged if

they correlated with r>.5 and otherwise we included them as separate proteins. From

the EMIF-AD MRM analysis proteins were included when observed in at least 92

individuals (30% of EMIF-AD sample). For related proteins that had identical values

(e.g. ACTA1 and ACTA2), due to fragment a-specificity we randomly selected one

protein for analysis.

2.5 Proteomic annotation

Biological process: We performed pathway enrichment analysis for Gene Ontology biological processes for proteins that differed between groups using the online Panther application (*48*) using FDR corrected Fisher exact tests.

Predominant cell-type of protein production: Based on the BRAIN RNASeq database (http://www.brainrnaseq.org) (*49*), we labelled proteins as being predominantly produced by a certain cell type when levels were higher than 40% of old individuals produced across cell types.

Transcription factor enrichment: We performed transcription factor enrichment analysis using ChIP-X ChEA through Enrichr and report on adjusted p-values (*50*).

SynGO: We mapped proteins that differed between a+t+ and a+t- to location at the synapse using SynGO 1.0 (*51*).

Other annotations:

- Proteins associated with choroid plexus expression: High expression of proteins in the choroid were based on and the Allen Brain Atlas (*52*) through Harmonizome (*53*) and ABAEnrichment analysis (*54*);

- Blood-Brain barrier associated proteins: Proteins mentioned in (*27,55,56*);

- BACE1 substrates: Proteins mentioned in one of the following articles (*57,58*);

- Alpha secretase substrates: Proteins mentioned in (*59,60*);

- Gamma secretase substrates: Proteins mentioned in (*61*).

2.6 Neuroimaging analysis

We studied neuroimaging data only from ADNI, because this study collected longitudinal scans. As a measure of brain atrophy we took cortical thickness data

from 34 cortical areas from the longitudinal processing pipeline in Freesurfer version 4.3 for 1.5T T1-weighted MRI scans, and v5.1 for 3T T1-weighted MRI scans (http://adni.loni.usc.edu). As a measure of amyloid accumulation, we used the region-specific SUVR values for florbetapir binding assessed by PET imaging in 34 cortical areas as parcellated by Freesurfer v4.5.0 (*62*). As a measure of brain metabolism we analysed fluorodeoxyglucose (FDG)-PET scans (*63*), and we determined average glucose metabolism for brain areas according to the Desikaan-Killiany atlas (*64*) matching the cortical thickness areas for 3461 preprocessed scans that were available from ADNI (see http://adni.loni.usc.edu/methods/pet-analysis-method/pet-analysis/ for preprocessing details). Briefly, the Desikan-Killiany atlas was warped from standard space to subject-space using the subject-specific normalization parameters in SPM12. Normalization failed for n=24 scans, leaving a total of n=3437 scans. Voxelwise FDG signals greater than 3×10^{-6} were averaged within each brain area, and standardized to the average uptake in the vermis and brain stem.

2.7 Genomic assessment

DNA extraction: For ADNI subjects, DNA was extracted from blood (n=1040) or cell lines (n=69). ADNI samples were genotyped using either the Illumina 2.5-M or the Illumina OmniQuad array and were retrieved online from http://adni.loni.cule.edu/ (*65*). For EMIF-AD individuals, DNA was extracted locally at the collection site (n=805) or from whole blood (n=148) using QIAamp DNA Blood Mini Kit (QIAGEN GmbH, Hilden, Germany) at the University of Lübeck. EMIF-AD MBD samples were genotyped using the Illumina Global Screening array (GSA) with shared custom content (Illumina, Incl) at the Institute of Clinical and Medical Biology

(UKSH, Campus-Kiel). A total of 936 EMIF-AD MBD samples passed post-experiment QC.

Genotype imputation and quality control: Data processing and quality control was performed using GenomeStudio Software (v2.0.04, Illumina, Inc.), PLINK software (v1.9) and bcftools (v1.9), as described previously (66). Briefly, SNPs were excluded if they had a minor allele frequency (MAF) less than 1%, deviated significantly from Hardy-Weinberg equilibrium (HWE) ($p < 5 \times 10^{-6}$) in the total sample of founder individuals, or had a call rate of less than 95%. We only used SNPs with no more than 2% genotype missingness and removed samples with excess heterozygosity rate (>5 SD). Individuals with sex mismatches were excluded from analysis. To identify ethnic outliers, a principal component analysis of ancestry (PCA) was performed based on 1000 Genomes clustering, phase 3 using PLINK) (67). Individuals of non-European descent were excluded from analysis. Relatedness was assessed through identity by descent, and family relations up to second degree were excluded. After filtering, the ADNI genotype data included 747 individuals and EMIF-AD 931 individuals. SNPs were locally imputed using Minimac 3 (68) to the Haplotype Reference Consortium (HRC) reference panel (69), leaving 4,583,773 SNPs (ADNI) and 7,778,465 SNPs (EMIF-AD) for final analyses. To account for population structure, principal components (PC1-PC20) were computed on a subset of relatively uncorrelated ($r2 < 0.2$) SNPs.

Classification of APOE genotypes: For ADNI subjects, APOE ε4 genotype was assessed with two SNPs; rs429358 the 'ε4-allele' and rs7412 the 'ε2 allele'), using DNA extracted by Cogenics from a 3mL aliquot of EDTA blood. We used imputed DNA microarray genotype data available from ADNI-1, ADNI-GO, ADNI-2 processed with GenomeStudio v2009.1. APOE ε4 genotypes in EMIF-AD were

generated as described elsewhere (*70*). Briefly, APOE ε4genotypes were determined

either directly (rs7412) or by imputation (rs429358). For 80 samples for which no

local *APOE* genotype was available, and for 45 mismatches between locally derived

and GSA–derived genotypes (4.8%), APOE ε4 genotype was determined using

TaqMan assays (ThermoFisher Scientific, Foster City, CA) on a QuantStudio-12K-

Flex system in a 384-well format.

2.8 GWAS and post-GWAS analyses

Genome-wide association study (GWAS) analyses on CSF t-tau abnormality status in

subjects with abnormal amyloid were performed separately in ADNI (a+t+=246, a+t-

=238) and EMIF-AD (a+t+=294, a+t-=155) using PLINK software (V1.90). We used

a logistic regression model including the PC1-PC3. Genome-wide significance was

defined as p≤5e-08. Meta-analysis on ADNI and EMIF-AD GWAS summary

statistics was performed using METAL (*71*). MAGMA software package was used to

compute gene scores and geneset scores for biological pathways based on p-values of

the meta-analysis summary statistics. We considered for further analysis gene scores

based on at least 6 SNPs and geneset scores with information of at least 6 genes.

2.9 Polygenic and other genetic risk score analysis

Polygenic risk scores (PGRS) for AD were computed for each subject using PRSice

(V2.3) (*72*). PGRS were calculated by adding the sum of each allele weighted by the

strength of its association with AD risk as calculated previously by the largest GWAS

on AD (71,880 cases; 383,378 controls) (*29*). AD cases were defined as patients

clinically diagnosed with AD-type dementia or individuals with a parental history of

AD (i.e., AD-by-proxy). AD-by-proxy showed strong genetic correlation with the

clinical diagnosis of AD (rg = 0.81). Clumping was performed prior to calculating

PGRS, to remove SNPs that are in LD (r2>0.1) within a slicing 1M bp window. After

clumping, PGRS were computed using various SNP inclusion thresholds (i.e., p≤1e-

30, p≤1e-8, p≤1e-5, p≤0.01, p≤0.02, p≤0.03, p≤0.04, p≤0.1, p≤0.2, p≤0.3,

p≤0.4,p≤0.5). In order to explore how specific genetic alterations were associated

with CSF proteomic profiles, we calculated for the top 3 genes and top 3 GO BP

genesets from the MAGMA analysis, a selection of 9 GO-BPs that were associated

with abnormal t-tau both in CSF enrichment analysis and MAGMA geneset analysis,

and 2 GO-BP associated H3 acetylation, as described in the text, gene-specific risk

scores (GRS) including only SNPs located on the respective genes with clumping as

described above and computation of GRS using various SNP inclusion thresholds. We

used weights from one cohort to generate GRS in the other cohort.

2.10 Statistics

Unless specified otherwise, we report on uncorrected p-values with statistical

significance set at p<0.05.

Cross-sectional analysis

The group comparisons between AD subgroups and controls were performed using

ANOVA correcting for age and gender.

Longitudinal analysis: Change in cognition and imaging markers were assessed using

ADNI data with linear mixed models including as main terms group, time and the

interaction group*time, and correcting for age and gender, and additionally for level

of education for cognitive markers.

Protein change with disease severity: We investigated within the AD normal t-tau and

AD abnormal t-tau groups whether proteins differed between stages by performing

MANOVA tests. We calculated p for trend over disease severity categories. A protein was considered to decrease or increase with disease severity if the linear trend was statically significant or if there was a difference between controls and MCI, controls and dementia, or MCI and dementia and the trend analysis supported an increase or decrease with severity.

Genetic analysis: Analysis were performed in all individuals with GWAS data and information on abeta42 and t-tau status. All (P)GRS were compared between AD subgroups using linear models. Associations between GRS scores for risk genes and CSF protein levels were tested with linear models adjusted for age and sex. In order to reduce number of tests GRS scores were selected for SNP inclusion threshold that best differentiated within AD between abnormal and normal t-tau.

Acknowledgements

This work has been supported by ZonMW Memorabel grant programme #73305056 (BMT) and #733050824 (BMT and PJV), the Swedish Research Council (#2018-02532, HZ), the European Research Council (#681712, HZ) and Swedish State Support for Clinical Research (#ALFGBG-720931, HZ), the Alzheimerfonden (Grant no. AF-930934) and Dtiftelsen Gamla tjänarinnor (JG), and the Innovative Medicines Initiative Joint Undertaking under EMIF grant agreement #115372 (PJV, HZ, SV, IB). Statistical analyses were performed at the VUmc Alzheimer Center that is part of the neurodegeneration research program of the Neuroscience Campus Amsterdam. EMIF-AD MBD proteomic analyses were performed at the Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Sweden. The VUmc Alzheimer Center is supported by Stichting Alzheimer Nederland and Stichting VUmc fonds. HZ is a Wallenberg Academy Scholar. The Leuven cohort was funded by Stichting Alzheimer Onderzoek (#11020, #15005, #13007) and the Vlaamse Impulsfinanciering voor Netwerken voor Dementie-onderzoek (IWT #135043). The Lausanne cohort was supported by the Swiss National Research Foundation SNF (#320030_141179).

We thank Drs. Andre Franke and Michael Wittig as well as Mrs. Tanja Wesse and Sanaz Sedghpour Sabet for their help with the genome-wide genotyping experiments. We thank Dr. Fabian Kilpert for his help with the processing and management of the genome-wide genotyping data.

Data was used for this project of which collection and sharing was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health

Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

Author Contributions:

Study design: BMT, PJV.

Design proteomics EMIF-AD: JG, HZ.

Design statistical analyses: BMT, PJV.

Data acquisition, processing, technical: JG, LR, IJ, ED, SH, VD, MT, FRJV, JP,

PMLA, RV, AL, JLM, SE, YFL, LF, KS, LB, SL, JS, SV, IB, DP, GS, KB, PS, CET,

HZ, PJV.

Manuscript drafting: BMT, PJV.

Manuscript revising: all authors.

References

1.	D. J. Selkoe, J. Hardy, The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol Med* **8**, 595-608 (2016).
2.	L. Vermunt *et al.*, Duration of preclinical, prodromal, and dementia stages of Alzheimer's disease in relation to age, sex, and APOE genotype. *Alzheimers Dement* **15**, 888-898 (2019).
3.	I. Bos *et al.*, Cerebrospinal fluid biomarkers of neurodegeneration, synaptic integrity, and astroglial activation across the clinical Alzheimer's disease spectrum. *Alzheimers Dement* **15**, 644-654 (2019).
4.	S. J. Vos *et al.*, Preclinical Alzheimer's disease and its outcome: a longitudinal cohort study. *Lancet Neurol* **12**, 957-965 (2013).
5.	S. J. Vos *et al.*, Prevalence and prognosis of Alzheimer's disease at the mild cognitive impairment stage. *Brain* **138**, 1327-1338 (2015).
6.	D. Ferreira *et al.*, The contribution of small vessel disease to subtypes of Alzheimer's disease: a study on cerebrospinal fluid and imaging biomarkers. *Neurobiol Aging* **70**, 18-29 (2018).
7.	A. Fuster-Matanzo, M. Llorens-Martin, J. Jurado-Arjona, J. Avila, F. Hernandez, Tau protein and adult hippocampal neurogenesis. *Front Neurosci* **6**, 104 (2012).
8.	A. M. Pooler, E. C. Phillips, D. H. Lau, W. Noble, D. P. Hanger, Physiological release of endogenous tau is stimulated by neuronal activity. *EMBO Rep* **14**, 389-394 (2013).
9.	C. R. Jack, Jr. *et al.*, NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement* **14**, 535-562 (2018).
10.	Y. Ihara, Massive somatodendritic sprouting of cortical neurons in Alzheimer's disease. *Brain Res* **459**, 138-144 (1988).
11.	M. Bulk *et al.*, Postmortem MRI and histology demonstrate differential iron accumulation and cortical myelin organization in early- and late-onset Alzheimer's disease. *Neurobiol Aging* **62**, 231-242 (2018).
12.	K. E. Biron, D. L. Dickstein, R. Gopaul, W. A. Jefferies, Amyloid triggers extensive cerebral angiogenesis causing blood brain barrier permeability and hypervascularity in Alzheimer's disease. *PLoS One* **6**, e23789 (2011).
13.	S. M. Greenberg, E. H. Koo, D. J. Selkoe, W. Q. Qiu, K. S. Kosik, Secreted beta-amyloid precursor protein stimulates mitogen-activated protein kinase and enhances tau phosphorylation. *Proc Natl Acad Sci U S A* **91**, 7104-7108 (1994).
14.	L. B. Gatta, A. Albertini, R. Ravid, D. Finazzi, Levels of beta-secretase BACE and alpha-secretase ADAM10 mRNAs in Alzheimer hippocampus. *Neuroreport* **13**, 2031-2033 (2002).
15.	K. Meyer *et al.*, REST and Neural Gene Network Dysregulation in iPSC Models of Alzheimer's Disease. *Cell Rep* **26**, 1112-1127 e1119 (2019).
16.	M. A. Israel *et al.*, Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. *Nature* **482**, 216-220 (2012).
17.	K. F. Young, S. H. Pasternak, R. J. Rylett, Oligomeric aggregates of amyloid beta peptide 1-42 activate ERK/MAPK in SH-SY5Y cells via the alpha7 nicotinic receptor. *Neurochem Int* **55**, 796-801 (2009).
18.	X. Cao, T. C. Sudhof, Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J Biol Chem* **279**, 24601-24611 (2004).

19. V. L. Kouznetsova, A. Tchekanov, X. Li, X. Yan, I. F. Tsigelny, Polycomb repressive 2 complex-Molecular mechanisms of function. *Protein Sci* **28**, 1387-1399 (2019).

20. J. Y. Hwang, R. S. Zukin, REST, a master transcriptional regulator in neurodegenerative disease. *Curr Opin Neurobiol* **48**, 193-200 (2018).

21. M. Otero-Garcia *et al.*, Single-soma transcriptomics of tangle-bearing neurons in Alzheimer's disease reveals the signatures of tau-associated synaptic dysfunction. *bioRxiv*, 2020.2005.2011.088591 (2020).

22. B. Frost, M. Hemberg, J. Lewis, M. B. Feany, Tau promotes neurodegeneration through global chromatin relaxation. *Nat Neurosci* **17**, 357-366 (2014).

23. H. U. Klein *et al.*, Epigenome-wide study uncovers large-scale changes in histone acetylation driven by tau pathology in aging and Alzheimer's human brains. *Nat Neurosci* **22**, 37-46 (2019).

24. H. Xue *et al.*, Sevoflurane post-conditioning alleviates neonatal rat hypoxic-ischemic cerebral injury via Ezh2-regulated autophagy. *Drug Des Devel Ther* **13**, 1691-1706 (2019).

25. N. Khan *et al.*, A systems approach identifies Enhancer of Zeste Homolog 2 (EZH2) as a protective factor in epilepsy. *PLoS One* **14**, e0226733 (2019).

26. E. Pajarillo, A. Rizor, D. S. Son, M. Aschner, E. Lee, The transcription factor REST up-regulates tyrosine hydroxylase and antiapoptotic genes and protects dopaminergic neurons against manganese toxicity. *J Biol Chem* **295**, 3040-3054 (2020).

27. L. Dayon *et al.*, Proteomes of Paired Human Cerebrospinal Fluid and Plasma: Relation to Blood-Brain Barrier Permeability in Older Adults. *J Proteome Res* **18**, 1162-1174 (2019).

28. F. Marques, J. C. Sousa, N. Sousa, J. A. Palha, Blood-brain-barriers in aging and in Alzheimer's disease. *Mol Neurodegener* **8**, 38 (2013).

29. I. E. Jansen *et al.*, Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. *Nat Genet* **51**, 404-413 (2019).

30. Y. Li *et al.*, Genetic association of the APP binding protein 2 gene (APBB2) with late onset Alzheimer disease. *Hum Mutat* **25**, 270-277 (2005).

31. Y. Chang *et al.*, Generation of the beta-amyloid peptide and the amyloid precursor protein C-terminal fragment gamma are potentiated by FE65L1. *J Biol Chem* **278**, 51100-51107 (2003).

32. J. J. Palop *et al.*, Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron* **55**, 697-711 (2007).

33. M. Yoshikawa, Y. Soeda, M. Michikawa, O. F. X. Almeida, A. Takashima, Tau Depletion in APP Transgenic Mice Attenuates Task-Related Hyperactivation of the Hippocampus and Differentially Influences Locomotor Activity and Spatial Memory. *Front Neurosci* **12**, 124 (2018).

34. I. Bos *et al.*, The EMIF-AD Multimodal Biomarker Discovery study: design, methods and cohort characteristics. *Alzheimers Res Ther* **10**, 64 (2018).

35. D. Bertens, D. L. Knol, P. Scheltens, P. J. Visser, I. Alzheimer's Disease Neuroimaging, Temporal evolution of biomarkers and cognitive markers in the asymptomatic, MCI, and dementia stage of Alzheimer's disease. *Alzheimers Dement* **11**, 511-522 (2015).

36.   M. F. Folstein, S. E. Folstein, P. R. McHugh, "Mini-Mental State", a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* **12**, 189-198 (1975).

37.   W. G. Rosen, R. C. Mohs, K. L. Davis, A new rating scale for Alzheimer's disease. *Am J Psychiatry* **141**, 1356-1364 (1984).

38.   D. Wechsler, *Wechsler Memory Scale- IIIrd edition. Administration and scoring manual.*, (The Psychological Corporation, San Antonio, 1997).

39.   J. Morris, The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology* **43**, 2412-2414 (1993).

40.   J. B. Toledo, S. X. Xie, J. Q. Trojanowski, L. M. Shaw, Longitudinal change in CSF Tau and Abeta biomarkers for up to 48 months in ADNI. *Acta Neuropathol* **126**, 659-670 (2013).

41.   L. M. Shaw *et al.*, Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol* **65**, 403-413 (2009).

42.   B. M. Tijms *et al.*, Unbiased Approach to Counteract Upward Drift in Cerebrospinal Fluid Amyloid-beta 1-42 Analysis Results. *Clin Chem* **64**, 576-585 (2018).

43.   Y. Y. Leung *et al.*, Identifying amyloid pathology-related cerebrospinal fluid biomarkers for Alzheimer's disease in a multicohort study. *Alzheimers Dement (Amst)* **1**, 339-348 (2015).

44.   M. Korecka *et al.*, Qualification of a surrogate matrix-based absolute quantification method for amyloid-beta(4)(2) in human cerebrospinal fluid using 2D UPLC-tandem mass spectrometry. *J Alzheimers Dis* **41**, 441-451 (2014).

45.   D. S. Spellman *et al.*, Development and evaluation of a multiplexed mass spectrometry based assay for measuring candidate peptide biomarkers in Alzheimer's Disease Neuroimaging Initiative (ADNI) CSF. *Proteomics Clin Appl* **9**, 715-731 (2015).

46.   T. S. Batth, C. Francavilla, J. V. Olsen, Off-line high-pH reversed-phase fractionation for in-depth phosphoproteomics. *J Proteome Res* **13**, 6176-6186 (2014).

47.   Y. Perez-Riverol *et al.*, The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* **47**, D442-D450 (2019).

48.   H. Mi *et al.*, Protocol Update for large-scale genome and gene function analysis with the PANTHER classification system (v.14.0). *Nat Protoc* **14**, 703-721 (2019).

49.   Y. Zhang *et al.*, An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929-11947 (2014).

50.   M. V. Kuleshov *et al.*, Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* **44**, W90-97 (2016).

51.   F. Koopmans *et al.*, SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse. *Neuron*, (2019).

52.   M. J. Hawrylycz *et al.*, An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* **489**, 391-399 (2012).

53.   A. D. Rouillard *et al.*, The harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes and proteins. *Database (Oxford)* **2016**, (2016).

54. S. Grote, K. Prufer, J. Kelso, M. Dannemann, ABAEnrichment: an R package to test for gene set expression enrichment in the adult and developing human brain. *Bioinformatics* **32**, 3201-3203 (2016).

55. S. I. Rapoport, K. D. Pettigrew, A heterogenous, pore-vesicle membrane model for protein transfer from blood to cerebrospinal fluid at the choroid plexus. *Microvasc Res* **18**, 105-119 (1979).

56. M. Neumeier *et al.*, Detection of adiponectin in cerebrospinal fluid in humans. *Am J Physiol Endocrinol Metab* **293**, E965-969 (2007).

57. B. Dislich *et al.*, Label-free Quantitative Proteomics of Mouse Cerebrospinal Fluid Detects beta-Site APP Cleaving Enzyme (BACE1) Protease Substrates In Vivo. *Mol Cell Proteomics* **14**, 2550-2563 (2015).

58. P. H. Kuhn *et al.*, Secretome protein enrichment identifies physiological BACE1 protease substrates in neurons. *EMBO J* **31**, 3157-3168 (2012).

59. P. H. Kuhn *et al.*, Systematic substrate identification indicates a central role for the metalloprotease ADAM10 in axon targeting and synapse function. *Elife* **5**, (2016).

60. T. Brummer *et al.*, NrCAM is a marker for substrate-selective activation of ADAM10 in Alzheimer's disease. *EMBO Mol Med*, (2019).

61. A. Haapasalo, D. M. Kovacs, The many substrates of presenilin/gamma-secretase. *J Alzheimers Dis* **25**, 3-28 (2011).

62. S. M. Landau *et al.*, Measurement of longitudinal beta-amyloid change with 18F-florbetapir PET and standardized uptake value ratios. *J Nucl Med* **56**, 567-574 (2015).

63. S. M. Landau *et al.*, Associations between cognitive, functional, and FDG-PET measures of decline in AD and MCI. *Neurobiol Aging* **32**, 1207-1218 (2011).

64. A. Klein, J. Tourville, 101 labeled brain images and a consistent human cortical labeling protocol. *Front Neurosci* **6**, 171 (2012).

65. A. J. Saykin *et al.*, Alzheimer's Disease Neuroimaging Initiative biomarkers as quantitative phenotypes: Genetics core aims, progress, and plans. *Alzheimers Dement* **6**, 265-273 (2010).

66. Hong S, Prokopenko D, D. V, e. al., Genome-wide association study of Alzheimer's disease CSF biomarkers in the EMIF-AD Multimodal Biomarker Discovery dataset. *BioRXiv*, (2019).

67. N. Siva, 1000 Genomes project. *Nat Biotechnol* **26**, 256 (2008).

68. S. Das *et al.*, Next-generation genotype imputation service and methods. *Nat Genet* **48**, 1284-1287 (2016).

69. S. McCarthy *et al.*, A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet* **48**, 1279-1283 (2016).

70. M. Ten Kate *et al.*, MRI predictors of amyloid pathology: results from the EMIF-AD Multimodal Biomarker Discovery study. *Alzheimers Res Ther* **10**, 100 (2018).

71. C. J. Willer, Y. Li, G. R. Abecasis, METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190-2191 (2010).

72. S. W. Choi, P. F. O'Reilly, PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* **8**, (2019).

Figure legends

Figure 1. Protein concentrations relative to control group in individuals with AD according to t-tau status.

Protein concentrations of individuals with abnormal $A\beta_{1-42}$ and abnormal total tau (a+t+, red) and abnormal $A\beta_{1-42}$ and normal t-tau (a+t-, blue) in ADNI (A) and EMIF-AD (B). Concentrations are expressed as z-score relative to control group with normal cognition, $A\beta_{1-42}$ and t-tau (z-score=0). Proteins are sorted according to change relative to control group. Shown are proteins that differed from controls in either a+t+ or a+t-. Venn diagram shows number of proteins that differed. Full data are shown in sTable2c.

Figure 2. Association of selected GO biological processes with proteins that differ from controls, MAGMA geneset score, and cell type.

Heatmap showing p-value (log p-value*-1) of selected enriched GO biological processes and SUZ12 and REST transcription factors of proteins increased (red) or decreased (blue) relative to controls in total sample (A) or in preclinical stage (B); P-value of association of GO biological processes with abnormal t-tau in GWAS-based MAGMA genset analysis (C); Proportion of proteins with cell-type specific expression (D). Data are shown for ADNI (A) and EMIF-AD (E) separately as indicated.

Data used in heatmap are shown in sTable 7. P-values of all GO biological processes are listed in sTable 3a (CSF enrichment) and sTable 4D (MAGMA analysis).

A+t+: abnormal $A\beta_{1-42}$ and t-tau; A+t-: abnormal $A\beta_{1-42}$ and normal total tau; CN A+t+: abnormal $A\beta_{1-42}$ and t-tau in preclinical stage; CN A+t-: abnormal $A\beta_{1-42}$ and

normal t-tau in preclinical stage. Oligo= oligodendrocyte; endothelial=endothelial cell.

Figure 3. Association of protein process score (PPS) with disease severity.

Data are based on cross-sectional analysis and combines data from ADNI and EMIF-AD. In grey PPS of control group with normal cognition and normal CSF $A\beta_{1-42}$ and t-tau; In red PPS of individuals with abnormal $A\beta_{1-42}$ and t-tau; In blue PPS of individuals with abnormal $A\beta_{1-42}$ and normal t-tau. CN=normal cognition; MCI=mild cognitive impairment; Dem=dementia.

Figure 4. PGRS across SNP thresholds

SNP weights were based on AD GWAS (Jansen et al 2019). Full data (including data after correction for APOE ε4 genotype and demographics, and separate for cohort and clinical stage) are shown in sTable 4a. A+t+: abnormal $A\beta_{1-42}$ and t-tau (red); A+t-: abnormal $A\beta_{1-42}$ and normal t-tau (blue).

Figure 5. Relation between genetic risk score and CSF protein correlation.

Relation between genetic risk score and CSF protein concentration. Shown are the 10 GRS from genes or GO-biological process (GO-BP) with the highest number of positive correlations with CSF proteins. The size of the arrows visually indicates how many proteins that correlate with the GRS in the coloured left outer ring belong to one of the CSF GO-BP shown in the right grey outer ring. For example, the GRS of GO-BP gene expression showed a positive correlation with 256 CSF proteins and these proteins were predominantly associated with GO-BP of nervous system development, gene expression and cell adhesion. Only arrows to CSF GO-BP are shown in case the

GRS correlated with at least 8 proteins from the CSF GO-BP. Supporting data in sTable 5. ECM= Extracellular matrix.

Figure 6. Longitudinal change in cognition and amyloid PET binding

Longitudinal change in Logical memory (A), CDR sum of boxes (B) and PET amyloid aggregation (C) according to diagnostic group and clinical stage. PET amyloid data are not shown for dementia with abnormal Aβ1-42 and normal t-tau because of low number of participants (n=6). In figure 6A and 6B change of controls are shown in grey, of individuals with a+t+ in red, and of individuals with a+t- in blue. Data are from ADNI only. Data are shown in sTable 6a (Logical memory, CDR-sum of boxes) and 6b (amyloid aggregation). CN=normal cognition; MCI=mild cognitive impairment; a+t+= abnormal $A\beta_{1-42}$ and t-tau; a+t-= abnormal $A\beta_{1-42}$ and normal t-tau.

Supplemental figures

sFigure 1. Longitudinal change in CSF total tau

A) Change in t-tau as function of group (control (grey), abnormal CSF $A\beta_{1-42}$ and abnormal t-tau (red), and abnormal $A\beta_{1-42}$ and normal t-tau (blue) and clinical stage. CN is normal cognition, MCI=mild cognitive impairment.

B) Baseline t-tau concentration of individuals with normal t-tau at baseline according to t-tau status at last follow-up. Individuals with abnormal t-tau at follow-up scored just below the cut-point of abnormal t-tau (dotted line) at baseline.

sFigure 2. Enrichment of proteins for synaptic processes from SynGO

Enrichment of SynGO synaptic biological processes in EMIF-AD (A, C) and ADNI (B, D). Enriched processes in proteins with increased levels in individuals who have

abnormal CSF Aβ_{1-42} and abnormal t-tau (a+t+) relative to control (CN, figure A, B) are similar to processes enriched in proteins with decreased level in individuals with abnormal CSF Aβ_{1-42} and normal t-tau (a+t-) relative to CN (figure C, D). Data are shown in sTable 3b.

sFigure 3. Enrichment of proteins for transcription factors and GO biological process

A) Heatmap showing p-value (log p-value*-1) of selected enriched GO biological processes and SUZ12 and REST transcription factors of proteins that were higher in in individuals with abnormal Aβ_{1-42} and t-tau (a+t+ (red)) or lower (blue) relative to individuals with abnormal Aβ_{1-42} and normal t-tau (a+t-).

B) GO-BP associated with proteins that showed a decrease in concentration with disease stage (blue) or increase with disease stage (red).

Data are shown for ADNI and EMIF-AD separately. Data used in heatmap are shown in sTable 7. P-values of all GO biological processes are listed in sTable 3a.

sFigure 4. Longitudinal change in cognitive measures according to diagnostic group and clinical stage.

MMSE (A) and ADAS-Cog (B). Change of controls are shown in grey, of individuals with abnormal Aβ_{1-42} and t-tau in red, and of individuals with abnormal Aβ_{1-42} and normal t-tau in blue. Data are shown in sTable 6a. Data are from ADNI only.

CN=normal cognition; MCI=mild cognitive impairment; a+t+= abnormal Aβ_{1-42} and t-tau; a+t-= abnormal Aβ_{1-42} and normal t-tau.

sFigure 5. Longitudinal change in imaging measures according to diagnostic group and clinical stage

(A) cortical thickness and (B) FDG-PET glucose metabolism. Data are from ADNI only. Data are shown in sTable 6b. CN=normal cognition; MCI=mild cognitive impairment; a+t+= abnormal Aβ$_{1-42}$ and t-tau; a+t-= abnormal Aβ$_{1-42}$ and normal t-tau.

Supplemental tables

Supplemental table 1. Descriptives

Supplemental Table 1a. Characteristics of individuals with information on CSF Aβ$_{1-42}$ and t-tau at first visit.

Supplemental Table 1b. Characteristics of individuals with available CSF proteomic data.

Supplemental Table 1c. Aβ$_{1-42}$ Cut-point definition.

Supplemental table 2. Protein data

Supplemental Table 2a. Number of proteins with change in concentration relative to reference or disease stage.

Supplemental Table 2b. Longitudinal t-tau data (ADNI only)

Supplemental Table 2c. Protein annotation and statistics of group comparisons.

Supplemental Table 2d. Cell-type specific proteins with largest differences between abnormal Aβ$_{1-42}$ and t-tau and abnormal Aβ$_{1-42}$ and normal t-tau.

Supplemental Table 2e. Proteins that differ in preclinical abnormal Aβ$_{1-42}$ and t-tau or preclinical abnormal Aβ$_{1-42}$ and normal t-tau from control group.

Supplemental table 3. Protein enrichment analysis

Supplemental Table 3a. Full list of GO biological processes associated with proteins that differ according to group or severity.

Supplemental Table 3b. SynGO enriched synaptic cellular components and biological processes that differ for proteins that are increased in abnormal $A\beta_{1-42}$ and abnormal t-tau versus CN or decreased in abnormal $A\beta_{1-42}$ and normal t-tau versus CN.

Supplemental Table 4 Genetic analysis

Supplemental Table 4a. Estimated marginal means (se) of AD GWAS-based polygenic risk scores in controls, abnormal $A\beta_{1-42}$ and normal t-tau and abnormal $A\beta_{1-42}$ and abnormal t-tau in total sample and according to clinical stage.

Supplemental Table 4b. Top 1000 SNPS from GWAS on abnormal $A\beta_{1-42}$ and normal t-tau versus abnormal $A\beta_{1-42}$ and abnormal t-tau in pooled ADNI and EMIF-AD cohorts.

Supplemental Table 4c. Difference in MAGMA gene score between abnormal $A\beta_{1-42}$ and abnormal t-tau and abnormal $A\beta_{1-42}$ and normal t-tau based on GWAS in pooled ADNI and EMIF-AD cohorts.

Supplemental Table 4d. Difference in GO biological process MAGMA geneset score between abnormal $A\beta_{1-42}$ and abnormal t-tau and abnormal $A\beta_{1-42}$ and normal t-tau based on GWAS in pooled ADNI and EMIF-AD cohorts.

Supplement Table 5. Correlation between genetic risk score and CSF protein concentrations

sTable 5a. Summary data on correlation of selected genetic risk scores with CSF protein correlations

Figure 1. Protein concentrations relative to control group (z-score =0) in individuals with AD according to t-tau status

Figure 2. Association of selected GO biological processes with proteins that differ from controls, MAGMA geneset score, and cell type.

Figure 3. Association of protein process score (PPS) with disease severity

Figure 4. PGRS across SNP thresholds

Figure 5. Relation between genetic risk score (GRS) and CSF protein concentration. Shown are the 10 GRS with the highest number of positive correlations with CSF proteins.The size of the arrows visually indicates how many proteins that correlate with the GRS in the coloured left outer ring belong to one of the CSF GO-BP shown in the right grey outer ring. Supporting data in sTable 5.

A)

C)

β annual change SUVr

Figure 6. Longitudinal change in cognition (A, B) and amyloid PET binding (C) according to diagnostic group and clinical stage

a)

Normal amyloid, normal tau ——— Abnormal amyloid, normal tau ——— Abnormal amyloid, abnormal tau

b)

sFigure 1. Longitudinal change in CSF total tau (data from ADNI). a) Change in total tau levels over time in each clinical stage. b) Baseline total tau concentrations in individuals with normal tau according to tau status at follow up. Supportive data are shown in sTable 2b.

sFigure 2. Enrichment of SynGO synaptic biological processes in EMIF-AD (A, C) and ADNI (B, D). Enriched processes in proteins with increased levels in individuals who have abnormal CSF Aβ1-42 and abnormal t-tau (a+t+) relative to control (CN, figure A, B) are similar to processes enriched in proteins with decreased level in individuals with abnormal CSF Aβ1-42 and normal t-tau (a+t-) relative to CN (figure C, D).
Supportive data are shown in sTable 3b.

sFigure 3. Enrichment of proteins for transcription factors and GO biological process that differ between A+T+ and A+T- (A) or with increasing disease severity in cross-sectional analysis (B). Supporting data in sTable 7

sFigure 4. Longitudinal change in MMSE (a) and ADAS-Cog (b) according to diagnostic group and clinical stage. Supporting data in sTable 6

sFigure 5 Longitudinal change in imaging measures according to diagnostic group and clinical stage. Supporting data in sTable 6

A) Cortical thickness

B) FDG-PET