A tipping point in cancer-immune dynamics leads to divergent immunotherapy responses and hampers biomarker discovery

Jeroen H.A. Creemers[1,4], Willem J. Lesterhuis[3], Niven Mehra[2], Winald R. Gerritsen[2],

Carl G. Figdor[1,4], I. Jolanda M. de Vries[1], Johannes Textor[1,5*]

1) Department of Tumor Immunology, Radboud Institute for Molecular Life Sciences; Radboudumc, Nijmegen, The Netherlands
2) Department of Medical Oncology, Radboudumc, Nijmegen, The Netherlands
3) School of Biomedical Sciences and Telethon Kids Institute, University of Western Australia, Perth, Australia
4) Oncode Institute, Nijmegen, The Netherlands
5) Data Science Department, Institute for Computing and Information Sciences, Radboud University, Nijmegen, The Netherlands

* Corresponding author:
Dr. J. Textor
Department of Tumor Immunology, Radboud Institute for Molecular Life Sciences, Radboudumc
Geert Grooteplein 26, 6500 HB Nijmegen (P.O. Box 9101), The Netherlands
E-mail: johannes.textor@radboudumc.nl

ABSTRACT

Background

 In the field of immuno-oncology, predicting treatment response or survival of cancer patients remains a challenge. Efforts to overcome these challenges focus mainly on the discovery of new biomarkers. Owing to the complexity of cancers and their tumor microenvironment, only a limited number of candidate biomarkers eventually enters clinical practice, despite advances in cellular and molecular approaches.

Methods

A computational modeling approach based on ordinary differential equations was used to simulate the fundamental mechanisms that dictate tumor-immune dynamics and show its implications on responses to immune checkpoint inhibition (ICI) and patient survival. Using *in silico* biomarker discovery trials, we extracted fundamental principles underlying the success rates of biomarker discovery programs.

Results

Our main finding is the prediction of a tipping point – a sharp state transition between immune control and immune evasion – that follows a strongly non-linear relationship between patient survival and both immunological and tumor-related parameters. In patients close to the tipping point, ICI therapy may lead to long-lasting survival benefits, whereas patients far from the tipping point may fail to benefit from these potent treatments.

Conclusion

These findings have two important implications for clinical oncology. First, the apparent conundrum that ICI induces substantial benefits in some patients yet completely fails in others could be, to a large extent, explained by the presence of a tipping point. Second, predictive biomarkers for immunotherapy should ideally combine both immunological and tumor-related markers, as the distance of a patient' status from the tipping point cannot, in general, be reliably determined from solely one of these. The notion of a tipping point in cancer-immune dynamics could help to optimize strategies in biomarker discovery to ensure accurate selection of the right patient for the right treatment.

INTRODUCTION

Immunotherapies are revolutionizing clinical care for cancer patients. The most widely used approach, immune checkpoint inhibition (ICI), can lead to long-term survival benefits in patients with advanced melanoma (1), lung cancer (2), and renal cell carcinoma (3). However, not all patients benefit from ICI therapy, and adequate predictions of treatment response have proven elusive so far (4, 5). Efforts to improve these predictions focus mainly on discovering biomarkers in aberrant molecular pathways within the tumor microenvironment that drive immunosuppression and therapeutic resistance (6, 7). These include genomic alterations in oncogenic drivers, the absence of tumor-specific antigens, and the presence of immunosuppressive molecules or cells (8, 9). Despite substantial efforts, only a limited fraction (according to one estimate, <1% (10)) of proposed cancer biomarkers find their way into the clinical practice. These apparent challenges in identifying biomarkers for immunotherapy and translating them into clinical practice could be a consequence of the inherent complexity of cancers and their interaction with the immune system.

To unravel the complexities of cancers and their treatments, researchers have adopted mathematical and computational approaches to complement laboratory research. A plethora of modeling approaches are available, ranging from simple one-variable equations to complex spatial agent-based simulation models. *In silico* modeling has contributed to fundamental insights into tumor growth and cancer progression (11-13), tumor-immune control (e.g., neoantigen prediction as targets for immunotherapy) (14), identification of tumor-associated genes (15), verification of treatment-related safety concerns such as hematological toxicity (16), prediction of treatment responses to chemo- and immunotherapy (17-19), investigation of drug-induced resistance (20), and timing of anti-cancer treatments (21-23). In the context of disease course dynamics, ordinary differential equation (ODE) models have proven useful over the years. ODE models follow the principle that a model should be "as simple as possible but not simpler". Based on plausible biological assumptions, they aim to reduce the complex reality of the modeled system to its bare essentials to enable investigation of critical underlying dynamics. For instance, Fassoni *et al.* used ODE models to predict that dose de-escalation of tyrosine kinase inhibitors targeting the oncogenic protein BCR-ABL1 in patients with chronic myeloid leukemia (chronic phase) does not lead to worse long-term outcomes (24). The recent results of the DESTINY trial support this prediction (25).

In this study, we investigate the consequences of tumor-immune dynamics on patients' responses to ICI and survival with an ODE model. Our model reveals a tipping point within tumor-immune dynamics – a critical threshold for survival culminating in an all-or-nothing principle – that has profound implications for a patient's disease course and outcome. We show how the presence of such a tipping point alone robustly induces heterogeneous immunotherapy treatment outcomes, and how this complicates the search for both prognostic and predictive biomarkers.

METHODS

Capturing core mechanisms of the tumor microenvironment in a mathematical model

We constructed a mathematical model based on a system of ordinary differential equations (ODE) to capture essential interactions between cancerous cells and lymphocytes in the tumor microenvironment. Our model represents tumorigenesis in patients, starting with the malignant transformation of a single cell.

The model describes the following processes: (1) conversion of naive T cells into antigen-specific effector T cells in lymph nodes; (2) the clonal expansion of effector T cells in the lymph nodes; (3) tumor growth; (4) subsequent attraction of tumor-infiltrating lymphocytes; and (5) formation of tumor-immune cell complexes to facilitate tumor cell killing. Below we describe the model; a mathematical formulation containing the model equations is provided as Supplementary Information. Tumor growth is represented by the generalized exponential model proposed by Mendelsohn (26), meaning that at each time interval, a fraction of tumor cells divide. The dividing fraction decreases as the tumor burden increases since substantial parts of a larger tumor mass, such as the necrotic core, are no longer able to proliferate. During tumor development, T cells are produced in the lymph nodes. This process starts with the conversion of naive T cells (i.e., not activated antigen-specific) into antigen-specific effector T cells at a rate proportional to the tumor burden. Effector T cells expand clonally and migrate into the tumor microenvironment, where they form tumor-immune complexes (i.e., an immune cell attached to a tumor cell). Within these complexes, T cells kill the tumor cells.

The killing rate expression is derived from the Michaelis-Menten kinetics for enzyme-substrate interaction with a quasi-steady-state approximation, as described by Borghans *et al.* (27). Given that complex formation and dissociation occur on a time scale several orders of magnitude smaller than the time scale for tumor growth, this approximation states that this process can be regarded as a constant process. Further simplification of the killing rate expression using the Padé approximation yields the double saturation killing rate of T cells (27). T cells follow a 'monogamous killing' strategy, meaning that one T cell interacts with one tumor cell at a time (27, 28). In addition, T cells die and exit the system at a fixed rate.

Simulation parameters

The simulation parameters are listed in Table 1.

Table 1: Simulation parameters in ODE model

Symbol	Parameter (dimension)	Default value (range[a])
ρ	Tumor growth rate (cells day^{-1})	1 (0-7)
ξ	Relative killing rate	0.001 (0-0.05)
h	Michaelis constant (cells)	571
δ	Death rate of immune cells (cells day^{-1})	0.019 (29)

α	Conversion rate of naive T cells into specific T cells	0.0025
p_S	Total production rate of effector T cells from lymph nodes (cells day^{-1})	1
m_S	Migration rate from lymph node to tumor microenvironment	1

[a]: if not fixed

The parameters were chosen to mimic realistic *in vivo* intercellular behavior. The rationale for the

choice of each parameter is explained below.

In a human adult, an estimated repertoire of approximately 10^{10} - 10^{11} naive CD8$^+$ T cells is present (30,

31). Naive CD8$^+$ T cells need to be primed to become activated effector T cells. The CD8$^+$ T cell precursor

frequency – the frequency at which any given peptide-MHC complex is recognized by naive antigen-

specific CD8$^+$ T cells – is on the order of 1 : 100.000 (30). Priming should be limited primarily to naive

CD8$^+$ T cells in one of the tumor areas draining lymph nodes. A human body contains ±600 lymph

nodes. At steady states, roughly 40% of all lymphocytes reside in lymph nodes, meaning that 40.000

naive T cells ($\approx$ 70 naive CD8$^+$ T cells per lymph node) can be primed (32, 33). We assume that priming

occurs primarily in the tumor-draining lymph node station (per station harboring around 20 lymph

nodes (34)), leading to an optimal priming rate of 1400 T cells per day. Considering that DCs might

present multiple epitopes and antigens, and that T cell priming *in vivo* might occur suboptimally, we

set a priming rate of at most 2500 cells per day. The order of magnitude of these priming rates

corresponds to priming rates found in chronic infectious diseases (35). Due to evasive mechanisms,

anti-tumor immunity is a more dormant process than an immune response to infections (36).

Therefore, we scaled the priming rate with tumor size (Supplementary Information), which translates

into a maximum production of 10^6 antigen-specific CD8$^+$ T cells per day via clonal expansion. Next, we

assume that all antigen-specific effector T cells migrate into the tumor microenvironment to interact

with tumor cells (i.e., complex formation).

Complex formation and dissociation rates are described by 'the Michaelis constant', which we derived

from literature (28). The Michaelis constant describes the ratio between complex formation and

dissociation.

The killing rate of effector T cells has been investigated mainly in the context of infectious disease. In

their review (37), Halle *et al.* discuss discrepancies between *in vitro* and *in vivo* killing rates of effector

T cells. Dependent on the context, killing rates of effector T cells vary from 1 target per 5 minutes to

0-10 targets per day (37), but tumor cells are considered difficult to kill. Extensive variation in

experimental *in vivo per capita* killing rates (i.e., the number of cells killed by an effector T cell per unit

of time) complicates the selection of a default fixed killing parameter. Therefore, we investigated T cell

dynamics over a range of overall killing rates as described using the monogamous killing regime in a

double saturation model by Gadhamsetty *et al*. (28). The double saturation model ensures that the

killing rate saturates with respect to the tumor cell and the effector cell densities. Consequently, our

model's maximum per capita killing rate is 2.5 cells^{-1} day^{-1}: one T cell can kill at most 2.5 tumor cells

per day, provided there are abundant target cells available and there is no competition with other T

cells. The default tumor growth rate is one cell day^{-1}, but we varied this parameter extensively in our

simulations. Taken together, our default parameter values led to simulations of disease courses with

realistic survival times in patients with malignancies and matched the order of magnitude of tumor

growth rates as reported by others (38).

Patient simulations

We simulated tumor development in patients up to a maximum of 5 years. Note that depending on

emergent tumor-immune dynamics, simulated patients may not reach the overall survival endpoint

during this interval. Each time step in the simulation corresponded to one day. At baseline, one tumor

cell and a pool of 10^6 naïve tumor-specific T cells are present in a patient. Activated effector T cells are

absent. We defined the time of diagnosis as the time at which the tumor burden exceeded $65 * 10^8$

cells and became clinically apparent. This cut-off corresponds to the assumption that a tumor with a

volume of 1 cm^3 contains 10^8 tumor cells (39) and that several primary tumors (e.g., lung cancer, colon

carcinoma, and renal cell carcinoma) are diagnosed as spherical structures with a median diameter of

approximately 5 cm (40-42). The 'lethal' tumor burden of patients in these simulations is estimated at

10^{12} cells, corresponding to a total tumor mass of approximately 22 * 22 * 22 cm.

Model implementation

We implemented our ODE model in C++. The Boost library 'odeint' was used to solve the system of

ordinary differential equations (43). The code is available at GitHub:

https://github.com/jeroencreemers/tipping-point-cancer-immune-dynamics. Analyses and

visualizations were performed in R.

RESULTS

Modeling tumor-immune dynamics yields realistic disease trajectories

To investigate the consequences of tumor-immune dynamics on the survival kinetics of patients, we used a computational modeling approach. We aimed to capture the interplay between tumor- and immune cells in the tumor microenvironment and simulate tumor growth in patients (see Methods). Our ODE model captured four essential processes in anti-tumor immunity: priming of naive antigen-specific CD8+ T cells, clonal expansion of effector T cells in lymph nodes, tumor growth leading to effector T cell attraction into the tumor microenvironment, and formation of tumor-immune cell complexes to enable tumor cell killing (Fig. 1A).

Figure 1: An in silico model of the tumor microenvironment generates realistic and modifiable disease courses of cancer patients. (A) The ODE model describes fundamental processes in the tumor microenvironment. Parameters: α = naive T cell priming rate, δ = effector T cell death rate, ξ = effector T cell killing rate, ρ = tumor growth rate, p_s = effector T cell proliferation rate, and m_s = effector T cell migration rate. (B) An effective anti-tumor immune response can eradicate tumor cells before the clinical manifestation of a tumor. (C) After an initial state in which the tumor outpaces the immune system, the immune system can suppress tumor growth and controls it in a subclinical state. (D) The natural course of disease for a clinically apparent tumor. An initial malignant transformation is followed by tumor growth until clinical diagnosis. Despite the activation of adaptive immunity, the tumor prevails. A stage of progressive disease follows, ultimately culminating in cancer-related death. The horizontal grey lines indicate (from bottom to top): the tumor burden at diagnosis and the tumor burden at death, respectively. Simulation parameters are added in Supplementary Table 1.

We simulated tumor development from malignant transformation of a single cell, via clinical detection

of a tumor, to advanced disease and possibly death. Depending on the tumor growth and the cytotoxic

capacity of effector T cells, the 'time to clinical manifestation' and overall survival varied. Despite this

variation, our simulations consistently showed three possible outcomes: 1) effector T cells inhibited

tumor cell outgrowth and eradicated the tumor before clinical manifestation (Figure 1B); 2) effector T

cells were initially unable to inhibit tumor cell outgrowth, but caught up and suppressed tumor growth

to a balanced subclinical state (Figure 1C); or 3) exponential tumor growth outpaced the immune

system's control and gave rise to a clinically detectable tumor (Figure 1D). These three scenarios only

led to two clinically different outcomes in patients: either a tumor became clinically evident, or the

immune system could suppress/eradicate a tumor at an early stage. A balanced equilibrium state, in

which the immune system keeps a clinically evident tumor under persistent control, does not exist in

this deterministic version of our model.

Patient survival depends on a tipping point in tumor-immune dynamics

To better characterize these dichotomous survival kinetics, we examined how tumor-immune

dynamics influenced patient survival by varying the tumor growth rate and the T cell killing rate over

a broad range of possible values.

First, we focused solely on the tumor-component by varying the tumor growth rate. An increase in

tumor growth did not gradually shorten overall survival in patients (Figure 2A). On the contrary, a

critical threshold was present. Once the threshold was exceeded, the kinetics 'flip' from a state of

immune control (Figure 2A, inset 1) to a state in which the tumor could evade immune control (Figure

2A, inset 2).

Second, we investigated the influence of the T cell killing rate on overall survival. As for the death rate,

a gradual increase in the cytotoxic capacity of effector T cells did not induce a gradual change in survival

times. Instead, a sharp state transition was again observed that differentiated short from long survival

(Figure 2B). This coincided with the phenotypes 'immune evasion' and 'immune control'.

To visualize this sudden state transition or 'tipping point' in tumor-immune dynamics as a function of

both tumor proliferation and cytotoxic killing at the same time, we visualized the joint influence of the

tumor growth rate and T cell killing rate on survival in a heatmap (Figure 2C). This 'phase diagram'

shows that the tipping point is not only present for specific parameter values but is a fundamental

property in our model. By contrast, the state of subclinical tumor control was not universally present

around the tipping point (Figure 2C, inset) but manifested itself only in a narrow range of parameters.

Nevertheless, the general presence of a tipping point indicates that small perturbations in either tumor

growth rate or T cell killing rate in the vicinity of a tipping point may result in substantial overall survival

differences in patients, whereas much larger perturbations far away from the tipping point would have far less effect.

Figure 2: A tipping point in the tumor-immune interaction determines a patient's outcome.
(A) A gradual increase in tumor growth reveals a tipping point, where long-term survival (immune control; inset 1) abruptly changes to short-term survival (immune evasion; inset 2). (B) A similar analysis reveals a tipping point along the immune axis, again differentiating short-term survival (immune evasion; inset 1) from long-term benefit (immune control; inset 2). (C) The tipping point is present across the entire range of parameters examined. Cure and progressive disease are the dominant states, whereas subclinical tumor control only occurs within a limited parameter range (inset). Simulation parameters are shown in Supplementary Table 2.

Immune checkpoint inhibitors induce a survival benefit by shifting patients over a tipping point

So far, we have described tumor-immune interactions during the natural course of malignant disease. In a clinical setting, however, therapeutic interventions are available to steer disease courses. Dependent on the treatment of choice, a specific effect is exerted on the tumor microenvironment. Treatment effects vary from constraining the proliferative capacity of tumor cells (e.g., chemotherapy or targeted therapy) to increasing the T cell pool (e.g., CAR T cells) or expanding the proliferative capacity of T cells (e.g., cancer vaccines; Figure 3A). Given the unparalleled responses of advanced malignancies to immunotherapy, we focused on the consequences of a tipping point for responses to

immee checkpoint inhibitors (ICI), but these findings could be extended to other therapies as well. In this study, we limited the treatment effect of ICI to their primary mode of action: the augmentation of the T cell killing rate (Figure 3A).

Figure 3: Tipping points induce dichotomous clinical outcomes in heterogeneous patient populations.
(A) Treatments target processes or cell populations in the tumor microenvironment. (B-C) Two criteria need to be met to induce long-term survival: (B) ICI need to augment T cell killing sufficiently, and (C) the treatment effect needs to be retained for a prolonged time. An inadequate treatment effect or limited treatment duration led at maximum to a temporary survival benefit. (D-E) In patient populations with variation in only (D) the tumor (i.e., growth rate), or (E) the immune system (i.e., T cell killing rate), the distance to a tipping point determines the clinical benefit. Without treatment, survival was limited (grey bars). In contrast, ICI induced long-term survival solely in patients close to a tipping point (green bars). See also Supplementary Table 3.

In the presence of a tipping point, ICI could induce a long-term survival benefit under two conditions: 1) the effect of treatment needs to be potent enough to shift a patient over a tipping point (Figure 3B), and 2) the treatment effect needs to be sustained long enough for a patient to benefit from the

treatment (Figure 3C). The treatment effect is defined as the multiplication factor of the T cell killing

rate. When both criteria were satisfied, ICI were able to induce a long-term survival benefit. However,

if the treatment effect (anti-PD1 effect < 12.6) or duration (less than ±5 months) proved inadequate,

any survival benefit was only temporary, and inevitable tumor progression would ultimately limit

overall survival (Figure 3B/C; insets). These survival kinetics depend not solely on therapeutic features

of ICI but rather on the interplay between patient and ICI characteristics. To illustrate this, we

simulated twenty patients with identical immune systems (i.e., identical T cell killing rates). In the

absence of ICI therapy, variation in the tumor growth rate – that is, variation in the distance to a tipping

point – led to a limited variation in survival (Figure 3D; grey bars). When these same patients were

treated with ICI, a survival benefit is induced in all patients. However, the extent of this benefit differs

and is dependent on the distance to a tipping point. Following clinical observations, long-term survival

is only induced in the subset of patients close to a tipping point (Figure 3D; green bars). Similar findings

were obtained in a population of patients with identical tumors but different immune systems.

Without treatment, hardly any survival variation is present (Figure 3E; grey bars). Again, treatment

with ICI induced dichotomous clinical outcomes: a small survival benefit in most patients, with long-

term survival in a subset (Figure 3E; green bars). Hence, the mere presence of a tipping point yields

heterogeneity in treatment outcomes.

Tipping points determine patient outcomes in dynamic patient trajectories

Thus far, our simulations considered tipping points generated in patients with fixed characteristics.

However, disease courses in patients are certainly not fixed and are, to a certain extent, subject to

(random) variation. We hypothesized that interpatient variability in clinical outcomes could be

(partially) attributable to this dynamic behavior of cancers and the interaction with the immune

system. Such variation might reflect biological processes (e.g., accumulating mutations, the expression

of checkpoint molecules, and the availability of nutrients) that alter anti-tumor immunity and promote

or hamper tumor development. We reasoned that the subsequent dynamics could drive patients

towards and ultimately over a tipping point – or move patients away from it, which would limit the

survival benefit of these treatments. To verify this hypothesis, we simulated the effect of dynamically

evolving tumors (Figure 4A) or immune systems (Figure 4B) in identical patients compared to a static

reference patient. Specifically, we varied the tumor growth rate and the T cell killing rate randomly

over time (parameter values are included in Supplementary Table 4). Upon reaching a diagnosable

tumor volume, all patients in these examples were treated with ICI. As expected, stochastic dynamics

prompted survival differences and induced a survival benefit in a subset of patients. In a

heterogeneous patient population, this led to an interesting finding: the initial distance to a tipping

point, along with the dynamics itself, determined the clinical outcome of patients treated with ICI (Figure 4C, Supplementary Figure 1). At population level, this led to a distinction between three subsets of patients: (1) patients far away from a tipping point with an unmodifiable bad prognosis (non-responders), (2) patients close to a tipping point with a favorable prognosis (responders), and, most importantly, (3) patients in between these groups (potential responders). In the last subset, tumor dynamics ultimately determined the treatment response, and thereby the clinical outcome (Figure 4C; grey box). A clinically important ramification of dynamic trajectories is that even if the subset to which a patient belongs is known at baseline, dynamics could alter the distance to a tipping point and, thereby, the prognosis of a patient. Therefore, it might be impossible to predict the prognosis solely based on characteristics measured upon diagnosis. Dynamic trajectories can significantly diversify patient outcomes, meaning that continuous variation in the tumor growth rate (Figure 4D) or T cell killing rate (Figure 4E) leads to an entire spectrum of patient outcomes.

Figure 4: Survival outcomes are strongly affected by evolving patient dynamics.
(A-B) Examples of dynamic disease courses in patients with identical tumors and immune systems at baseline, respectively. (A) Evolving tumors (i.e., random variation in tumor growth rate over time) and (B) continuous variation in the potency of the immune system (i.e., killing rate) lead to divergent survival outcomes. The grey dotted line indicates the baseline values for the growth rate and killing rate, respectively. (C) Dynamic trajectories in a heterogeneous patient population can move patients towards or away from a tipping point. The grey box indicates patients in which dynamic trajectories strongly alter survival outcomes. See also Supplementary Figure 1. In dynamic trajectories, (D) baseline tumor growth and (E) baseline T cell killing rates cannot accurately predict overall survival. Note: all patients in these examples are treated with ICI. The red and black dotted lines indicate the 25% and 75% quantiles, respectively. See also Supplementary Table 4.

Implications of tipping points for biomarker discovery studies

Biomarker discovery studies aim to improve the prediction of patient survival upon treatment. We observed that tipping points are crucial in shaping survival kinetics. Therefore, accurate survival predictions would require the consideration of tipping points. Ideally, a prognostic biomarker (or biomarker panel) would consistently distinguish long-term survivors from their counterparts. Since the non-linear survival dynamics following a tipping point weaken the correlation between a single biomarker and survival, the question is: how can we screen for biomarkers in a more efficient manner that takes this tipping point into account?

We approached this question with an *in silico* biomarker discovery study. We measured the value of two potential biomarkers at baseline in patients (n=100) that were subsequently treated with ICI. We simplified the cohort by fixing the tumor and immune characteristics of these patients over time and assumed to have access to an entirely accurate biomarker (i.e., no measurement error; Figure 5A). Within this cohort, we predicted the prognosis of patients based on either the tumor or immune marker (the first and second columns of Figure 5A, respectively). As is common in practice (though from a statistical point of view far from ideal), we dichotomized the biomarker using its median as a cut-off. Although survival differentiation based on these biomarkers alone was partially possible, it remained far from optimal. However, when we constructed a biomarker panel including both biomarkers, it highly accurately discriminated short-term from long-term survivors (third column of Figure 5A, Supplementary information). Note that despite variability in time from diagnosis, the initial plateau in the survival curves was caused by the fact that all tumors were diagnosed with identical sizes and immediately treated.

In clinical practice, the assumption of a 'fixed' patient trajectory does not hold. Therefore, we simulated this cohort again with dynamic trajectories. Due to the dynamics, a subgroup of patients did not develop clinical tumors and was excluded from the analysis. The prediction of a patient's prognosis with a single biomarker, either from the tumor or the immune system, in a dynamic cohort became increasingly challenging (the first and second columns of Figure 5B). The combination of both markers in a biomarker panel increased the predictive capacity slightly, enabling the prediction of prognosis to some extent. However, in line with the notion of personalized medicine, the accurate and individualized prediction of prognosis based on baseline characteristics was not feasible in a significant subgroup of patients due to dynamic tumor-immune interactions (third column of Figure 5B). Two important findings are derived from these observations: First, due to the non-linear tumor-immune dynamics with respect to survival, it is highly unlikely that a single biomarker can accurately predict a patients' prognosis. Since survival kinetics emerge from the interplay between a cancer and the immune system, biomarkers from both systems need to be incorporated simultaneously into a biomarker panel to improve the predictive value. Second, biomarker measurements at baseline are

361 merely a situational snapshot of the disease conditions *at a specific point in time*. Depending on the

362 magnitude of the dynamics, it might become challenging or even impossible to correctly predict the

363 prognosis of patients from these biomarkers, stressing the need for continuous monitoring.

Figure 5: Non-linear tumor-immune dynamics complicate biomarker discovery.
365
(A) An in silico biomarker discovery study in a 'fixed' patient cohort: while a single biomarker – either a tumor or an immune
366
marker – can predict survival to some extent (the first and second columns), information from both markers in a biomarker
367
panel enhances the predictive capacity greatly (third column). (B) Dynamic disease trajectories challenge survival prediction
368
with 'baseline' biomarkers and stress the need for continuous monitoring. Disease dynamics challenge the predictive value of
369
single 'baseline' biomarkers (the first and second columns; compare to Figure 5A). A biomarker panel improves survival
370
predictions in this cohort (the third column) but is still defied by evolving dynamics. See also Supplementary figure 5.
371

DISCUSSION
373

This study investigated how tumor-immune dynamics relate to ICI-induced treatment responses and

survival kinetics of patients. We predict that a tipping point is present in the tumor-immune

interaction. This finding implies that at the basis of the intricate interplay between a developing

malignancy and the immune system contrasting disease states determine disease outcome: a state

where the immune system controls tumor outgrowth and a state in which a tumor escapes immune

defense. A stable "steady state" in which tumor growth and the immune response perfectly balance

each other for extended periods seems only plausible in a subclinical setting. We show that treatment

with ICI can induce a survival benefit by shifting a patient over a tipping point, thereby tipping the

balance in tumor-immune dynamics in favor of survival. In line with clinical observations of interpatient

variability in disease courses, we found that dynamics in patient trajectories might contribute to

unpredictability in treatment responses. Moreover, we showed how the tipping point in dynamic

patient trajectories defies simple strategies for outcome prediction in biomarker discovery studies,

stressing the need for advanced mechanism-based monitoring.

Tipping points are well-known in complex systems such as financial markets and ecosystems but are

also present in medicine (44, 45). State transitions might progress gradually or abruptly. If a system

balances around a critical threshold, small perturbations might induce an abrupt transition to a

contrasting state. In oncology, phenomena as partial or complete radiologic responses during

treatment or (hyper)progression after discontinuation of treatment suggest the presence of state

transitions (46, 47). Based on these observations, a tipping point in cancer immunotherapy had been

speculated upon (48). Experimentally, tipping points are most clearly represented by early preclinical

work in the PD-1/PD-L1 axis. Consistent with our findings, dichotomous treatment responses arise in

syngeneic DBA/2 mice inoculated with P815/PD-L1 cells (49). While genetically identical with similar

tumor characteristics, anti-PD-L1 antibodies could prolong survival in only a subset of the mice, likely

due to stochastic differences in immune responses and TCR repertoire. Additional *in vivo* data

supporting the theory of tipping points in oncology is derived from studies on dynamic network

biomarkers, showing its relevance during the onset of metastasis in hepatocellular carcinoma (50) and

the development of treatment resistance in breast cancer (51). This study provides a potential

mechanistic explanation for this phenomenon in immuno-oncology and shows its implications on the

induction of long-term survival in clinical practice and biomarker discovery. From a biomechanistic

perspective, such state transitions in cancer immunotherapy arise due to fundamental differences in

proliferation kinetics between tumors and the immune system. While tumor cell proliferation is

virtually unrestricted, immune cell proliferation is much more limited and tightly controlled. Our

finding that tipping points affect not only natural disease courses but also treatment responses

underlines the importance of these kinetics.

An important implication of tipping points within tumor-immune dynamics involves biomarker

discovery. Biomarkers are developed to predict prognosis and steer clinical decision making. Disease

outcomes in cancer patients are essentially determined by the interplay between two complex

systems: the tumor and the immune system. Our model predicts that factors from both systems should

be considered to improve the predictive power of biomarkers. However, in contrast with this

seemingly straightforward prediction, current research mainly focuses on factors derived from one of

the two complex systems. Expression of programmed death ligand-1 (PD-L1) on tumor tissue illustrates

this properly: while 45% of patients with PD-L1 positive tumors show objective responses to anti-

PD(L)1 immunotherapy, 15% of patients with PD-L1 negative tumors also show objective responses

(52). Other explanations for this difference include heterogeneous intratumoral and inter-metastases

expression patterns, positivity-threshold selection, and differences in immunohistochemical staining

protocols. In that respect, tumor mutational burden (TMB) might prove to be a highly relevant

biomarker. The mutation rate is a tumor-intrinsic factor associated with the phenotypical

aggressiveness of tumors (53). Simultaneously, a high mutational burden might induce a plethora of

neoantigens, linking this tumor-intrinsic factor directly to adaptive immunity. Clinical observations of

a stronger association between TMB and response rates to anti-PDL1 immunotherapy compared to

PD-L1 expression in patients with urothelial carcinoma support this hypothesis (54). Thus, reinforcing

common calls to integrate multiple existing biomarkers for immunotherapy prediction outcomes (55,

56), our research indicates that a combination of both immunological and tumor-related parameters

should be the basis of *any* biomarker discovery effort. The strongly non-linear dynamics resulting from

the tipping point mean that a one-dimensional approach will likely be insufficient.

Our approach has to be interpreted in light of some limitations. Although the 'coarse-grained' nature

of ODE models allows focusing on the major common underlying mechanisms in many cancers, it is

also a potential pitfall. For example, metabolic processes such as hypoxia, immune-suppressive

characteristics of the tumor microenvironment such as the presence of FoxP3$^+$ regulatory T cells or

expression of transforming growth factor β, the presence of additional relevant effector cells such as

natural killer cells, and the availability of nutrients are only implicitly represented by our model in a

single killing efficacy parameter. This simplification also holds for treatments. In this study, ICI was

limited to its main mode of action: the augmentation of the T cell killing rate. While the 'true'

mechanistic effects might be more widespread, sufficient data to correctly parameterize more

complex models remains scarce. Furthermore, it should be emphasized that an ODE model contains

limited spatial information; while we distinguish between lymphatic tissue and the tumor

microenvironment, all cells within the microenvironment are identical, and all processes affect cells in

the same manner. We do not expect that explicit incorporation of these processes or translation of

the model into a spatial variant alters our findings of a tipping point; however, it might be of interest

to verify these hypotheses in future research in spatial agent-based models.

In conclusion, we used a computational modeling approach to show that the clinical outcome of cancer

patients is determined by tipping points in tumor-immune dynamics. A tipping point influences not

only response to treatment but also the prognosis of patients and has major implications for future

biomarker research.

REFERENCES

1.   Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2019;381(16):1535-46.

2.   Hellmann MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 2019;381(21):2020-31.

3.   Motzer RJ, Escudier B, McDermott DF, et al. Survival outcomes and independent response assessment with nivolumab plus ipilimumab versus sunitinib in patients with advanced renal cell carcinoma: 42-month follow-up of a randomized phase 3 clinical trial. *J Immunother Cancer*. 2020;8(2).

4.   Aly A, Mullins CD, Hussain A. Understanding heterogeneity of treatment effect in prostate cancer. *Curr Opin Oncol*. 2015;27(3):209-16.

5.   Kent DM, Alsheikh-Ali A, Hayward RA. Competing risk and heterogeneity of treatment effect in clinical trials. *Trials*. 2008;9:30.

6.   Bedard PL, Hansen AR, Ratain MJ, et al. Tumour heterogeneity in the clinic. *Nature*. 2013;501(7467):355-64.

7.   Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer therapies. *Nat Rev Clin Oncol*. 2018;15(2):81-94.

8.   Sharma P, Hu-Lieskovan S, Wargo JA, et al. Primary, Adaptive, and Acquired Resistance to Cancer Immunotherapy. *Cell*. 2017;168(4):707-23.

9.   Roskoski R, Jr. A historical overview of protein kinases and their targeted small molecule inhibitors. *Pharmacol Res*. 2015;100:1-23.

10.   Kern SE. Why your new cancer biomarker may never work: recurrent patterns and remarkable diversity in biomarker failures. *Cancer Res*. 2012;72(23):6097-101.

11.   Murphy H, Jaafari H, Dobrovolny HM. Differences in predictions of ODE models of tumor growth: a cautionary example. *BMC Cancer*. 2016;16:163.

12.   Edelman EJ, Guinney J, Chi JT, et al. Modeling cancer progression via pathway dependencies. *PLoS Comput Biol*. 2008;4(2):e28.

13.   Baratchart E, Benzekry S, Bikfalvi A, et al. Computational Modelling of Metastasis Development in Renal Cell Carcinoma. *PLoS Comput Biol*. 2015;11(11):e1004626.

14.   De Mattos-Arruda L, Vazquez M, Finotello F, et al. Neoantigen prediction and computational perspectives towards clinical benefit: recommendations from the ESMO Precision Medicine Working Group. *Ann Oncol*. 2020;31(8):978-90.

15.   Pujana MA, Han JD, Starita LM, et al. Network modeling links breast cancer susceptibility and centrosome dysfunction. *Nat Genet*. 2007;39(11):1338-49.

16.   Friberg LE, Henningsson A, Maas H, et al. Model of chemotherapy-induced myelosuppression with parameter consistency across drugs. *J Clin Oncol*. 2002;20(24):4713-21.

17.   Lee JJ, Huang J, England CG, et al. Predictive modeling of in vivo response to gemcitabine in pancreatic cancer. *PLoS Comput Biol*. 2013;9(9):e1003231.

18.   Wang HW, Milberg O, Bartelink IH, et al. In silico simulation of a clinical trial with anti-CTLA-4 and anti-PD-L1 immunotherapies in metastatic breast cancer using a systems pharmacology model. *Roy Soc Open Sci*. 2019;6(5).

19.   Valentinuzzi D, Simoncic U, Ursic K, et al. Predicting tumour response to anti-PD-1 immunotherapy with computational modelling. *Phys Med Biol*. 2019;64(2):025017.

20.   Sun X, Bao J, Shao Y. Mathematical Modeling of Therapy-induced Cancer Drug Resistance: Connecting Cancer Mechanisms to Population Survival Rates. *Sci Rep*. 2016;6:22498.

21.   Zhang J, Cunningham JJ, Brown JS, et al. Integrating evolutionary dynamics into treatment of metastatic castrate-resistant prostate cancer. *Nat Commun*. 2017;8(1):1816.

22.   Cunningham JJ, Brown JS, Gatenby RA, et al. Optimal control to develop therapeutic strategies for metastatic castrate resistant prostate cancer. *J Theor Biol*. 2018;459:67-78.

23.   Ribba B, Boetsch C, Nayak T, et al. Prediction of the Optimal Dosing Regimen Using a Mathematical Model of Tumor Uptake for Immunocytokine-Based Cancer Immunotherapy. *Clin Cancer Res*. 2018;24(14):3325-33.

24.   Fassoni AC, Baldow C, Roeder I, et al. Reduced tyrosine kinase inhibitor dose is predicted to be as effective as standard dose in chronic myeloid leukemia: a simulation study based on phase III trial data. *Haematologica*. 2018;103(11):1825-34.

25.   Clark RE, Polydoros F, Apperley JF, et al. De-escalation of tyrosine kinase inhibitor therapy before complete treatment discontinuation in patients with chronic myeloid leukaemia (DESTINY): a non-randomised, phase 2 trial. *Lancet Haematol*. 2019;6(7):e375-e83.

26.   Mendelsohn ML. Cell proliferation and tumor growth. *Oxford: Blackwell Scientific Publications*. 1963.

27.   Borghans JA, de Boer RJ, Segel LA. Extending the quasi-steady state approximation by changing variables. *Bull Math Biol*. 1996;58(1):43-63.

28.   Gadhamsetty S, Maree AF, Beltman JB, et al. A general functional response of cytotoxic T lymphocyte-mediated killing of target cells. *Biophys J*. 2014;106(8):1780-91.

29.   McDonagh M, Bell EB. The survival and turnover of mature and immature CD8 T cells. *Immunology*. 1995;84(4):514-20.

30.   Jenkins MK, Chu HH, McLachlan JB, et al. On the composition of the preimmune repertoire of T cells specific for Peptide-major histocompatibility complex ligands. *Annu Rev Immunol*. 2010;28:275-94.

31.   Coulie PG, Karanikas V, Lurquin C, et al. Cytolytic T-cell responses of cancer patients vaccinated with a MAGE antigen. *Immunol Rev*. 2002;188:33-42.

32.   Westermann J, Pabst R. Distribution of lymphocyte subsets and natural killer cells in the human body. *Clin Investig*. 1992;70(7):539-44.

33.   Ferrer R. Lymphadenopathy: differential diagnosis and evaluation. *Am Fam Physician*. 1998;58(6):1313-20.

34.   Vallini V, Ortori S, Boraschi P, et al. Staging of pelvic lymph nodes in patients with prostate cancer: Usefulness of multiple b value SE-EPI diffusion-weighted imaging on a 3.0 T MR system. *Eur J Radiol Open*. 2016;3:16-21.

35.   Linderman JJ, Riggs T, Pande M, et al. Characterizing the dynamics of CD4+ T cell priming within a lymph node. *J Immunol*. 2010;184(6):2873-85.

36.   Drake CG, Jaffee E, Pardoll DM. Mechanisms of immune evasion by tumors. *Adv Immunol*. 2006;90:51-81.

37.   Halle S, Halle O, Forster R. Mechanisms and Dynamics of T Cell-Mediated Cytotoxicity In Vivo. *Trends Immunol*. 2017;38(6):432-43.

38.   Beck RJ, Slagter M, Beltman JB. Contact-Dependent Killing by Cytotoxic T Lymphocytes Is Insufficient for EL4 Tumor Regression In Vivo. *Cancer Res*. 2019;79(13):3406-16.

39.   Del Monte U. Does the cell number 10(9) still really fit one gram of tumor tissue? *Cell Cycle*. 2009;8(3):505-6.

40.   Moreno CC, Mittal PK, Sullivan PS, et al. Colorectal Cancer Initial Diagnosis: Screening Colonoscopy, Diagnostic Colonoscopy, or Emergent Surgery, and Tumor Stage and Size at Initial Presentation. *Clin Colorectal Cancer*. 2016;15(1):67-73.

41.   Zastrow S, Phuong A, von Bar I, et al. Primary tumor size in renal cell cancer in relation to the occurrence of synchronous metastatic disease. *Urol Int*. 2014;92(4):462-7.

42.   Ball DL, Fisher RJ, Burmeister BH, et al. The complex relationship between lung tumor volume and survival in patients with non-small cell lung cancer treated by definitive radiotherapy: a prospective, observational prognostic factor study of the Trans-Tasman Radiation Oncology Group (TROG 99.05). *Radiother Oncol*. 2013;106(3):305-11.

43.   Ahnert K, Mulansky M. Odeint – Solving Ordinary Differential Equations in C++. *AIP Conference Proceedings*. 2011;1389.

44.   Scheffer M, Carpenter SR, Lenton TM, et al. Anticipating critical transitions. *Science*. 2012;338(6105):344-8.

45.   Scheffer M, Bascompte J, Brock WA, et al. Early-warning signals for critical transitions. *Nature*. 2009;461(7260):53-9.

46.   Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer*. 2009;45(2):228-47.

47.   Champiat S, Dercle L, Ammari S, et al. Hyperprogressive Disease Is a New Pattern of Progression in Cancer Patients Treated by Anti-PD-1/PD-L1. *Clin Cancer Res*. 2017;23(8):1920-8.

48.   Lesterhuis WJ, Bosco A, Millward MJ, et al. Dynamic versus static biomarkers in cancer immune checkpoint blockade: unravelling complexity. *Nat Rev Drug Discov*. 2017;16(4):264-72.

49.   Iwai Y, Ishida M, Tanaka Y, et al. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *Proc Natl Acad Sci U S A*. 2002;99(19):12293-7.

50.   Yang B, Li M, Tang W, et al. Dynamic network biomarker indicates pulmonary metastasis at the tipping point of hepatocellular carcinoma. *Nat Commun*. 2018;9(1):678.

51.   Liu R, Wang J, Ukai M, et al. Hunt for the tipping point during endocrine resistance process in breast cancer by dynamic network biomarkers. *J Mol Cell Biol*. 2019;11(8):649-64.

52.   Sunshine J, Taube JM. PD-1/PD-L1 inhibitors. *Curr Opin Pharmacol*. 2015;23:32-8.

53.   Thomas A, Routh ED, Pullikuth A, et al. Tumor mutational burden is a determinant of immune-mediated survival in breast cancer. *Oncoimmunology*. 2018;7(10):e1490854.

54.   Balar AV, Galsky MD, Rosenberg JE, et al. Atezolizumab as first-line treatment in cisplatin-ineligible patients with locally advanced and metastatic urothelial carcinoma: a single-arm, multicentre, phase 2 trial. *Lancet*. 2017;389(10064):67-76.

55.   Blank CU, Haanen JB, Ribas A, et al. CANCER IMMUNOLOGY. The "cancer immunogram". *Science*. 2016;352(6286):658-60.

56.   Galon J, Pages F, Marincola FM, et al. The immune score as a new possible approach for the classification of cancer. *J Transl Med*. 2012;10:1.

LIST OF ABBREVIATIONS

ICI: Immune Checkpoint Inhibition; ODE: Ordinary Differential Equation; PD-(L)1: Programmed Death-(Ligand) 1; TMB: Tumor Mutational Burden.

DECLARATIONS

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

The code of the ODE model is available at GitHub: https://github.com/jeroencreemers/tipping-point-cancer-immune-dynamics.

Competing interests

WJL reports consultancy activities for Douglas Pharmaceuticals and MSD; research funding from Douglas Pharmaceuticals, AstraZeneca, and ENA therapeutics; patents PCT/AU2019/050259 and

PCT/AU2015/000458 (all outside this work). NM reports personal fees from Bayer and Merck Sharp &

Dohme; grants and personal fees from Jansen-Cilag, Roche, Astellas, and Sanofi (all outside this work).

WRG reports consultancy activities for Bristol-Myers Squibb, IMS Health, Janssen-Cilag, Sanofi, and

MSD; speaker fees from ESMO and MSD; and research funding from Bayer, Astellas, Janssen-Cilag, and

Sanofi (all outside this work).

Funding

JC was funded by the Radboudumc. WJL was supported by Fellowships from the NHMRC, the Simon

Lee Foundation, and the Cancer Council Western Australia. CF received an ERC Adv Grant ARTimmune

(834618) and an NWO Spinoza grant. IV received an NWO-Vici grant (918.14.655). JT was supported

by a Young Investigator Grant (10620) from the Dutch Cancer Society and an NWO grant

(VI.Vidi.192.084).

Authors' contribution

JHAC and JT conceived this study. JHAC performed the experiments and wrote the manuscript under

the supervision of JT. All authors provided feedback on the manuscript and reviewed the manuscript

prior to submission.

Acknowledgements

Not applicable.