Title: Missing data in the medical record for oncology patients: prevalence and association with outcomes

Daniel X. Yang, MD[1]; Rohan Khera, MD, MS[2,3]; Joseph A. Miccio, MD[1]; Vikram Jairam, MD[1]; Enoch Chang, BA[1]; James B. Yu, MD, MHS[1]; Henry S. Park, MD, MPH[1]; Harlan M. Krumholz, MD, MS[2,3]; Sanjay Aneja, MD[1,3]

[1]*Department of Therapeutic Radiology, Yale School of Medicine, New Haven, CT*

[2]*Department of Internal Medicine, Yale School of Medicine, New Haven, CT*

[3]*Center for Outcomes Research and Evaluation, Yale School of Medicine, New Haven, CT*

Corresponding author:

Sanjay Aneja, MD

Assistant Professor

Department of Therapeutic Radiology, Yale School of Medicine

330 Cedar Street, CB326

New Haven, CT 06520

Email: sanjay.aneja@yale.edu

Phone: 203-200-2100

Fax: 203-737-1467

Abstract

Background: There has been increased enthusiasm regarding the use of real-world data (RWD) for clinical evidence generation. Cancer registries are important RWD sources that rely on data abstraction from the medical record, however, patients with unknown or missing data elements are under-represented in studies that use such data sources. Therefore, we sought to determine the prevalence of missing data and its associated survival outcomes among cancer patients.

Methods: All data elements within the National Cancer Database were reviewed for missing or unknown values for the three most common cancers in the United States diagnosed from 2006 to 2015. Patient records with incompletely abstracted data elements were identified. Prevalence of patients with missing data and their associated survival outcomes were determined.

Results: A total of 1,198,749 non-small cell lung cancer (NSCLC), 2,120,775 breast cancer, and 1,158,635 prostate cancer patients were included for analysis. For NSCLC, there were 851,295 (71.0%) patients with missing data in data elements identified as variables of interest; 2-year survival was 33.2% for patients with missing data and 51.6% for patients with complete data (p<0.001). For breast cancer, there were 1,161,096 (54.8%) patients with missing data; 2-year survival was 93.2% for patients with missing data and 93.9% for patients with complete data (p<0.001). For prostate cancer, there were 460,167 (39.7%) patients with missing data; 2-year survival was 91.0% for patients with missing data and 95.6% for patients with complete data (p<0.001). There were significant differences in demographics, tumor characteristics, cancer stage, and treatments received between cancer patients with complete and incomplete records.

Conclusions: The majority of patients in a large cancer registry-based RWD source have missing data that was unable to be ascertained from the medical record. Moreover, patients with

missing data show worse survival than those with more complete documentation. Increasing

documentation quality and adoption of rigorous missing data correction methods are needed to

best leverage RWD for clinical advancements.

Introduction

Real-world evidence derived from real-world data (RWD) holds substantial promise to accelerate innovation within oncology. RWD, which includes data on patient health status and/or the delivery of health care collected routinely,[1] is becoming increasingly relevant due to the high cost and slow pace of randomized clinical trials as well as the growth of near real-time access to data electronic health records (EHR) and other digital sources of comprehensive health-related data. RWD sources may represent a more flexible, cost-effective way to investigate clinical interventions and can supplement clinical trials. Within oncology, there have been investments in developing RWD sources as sources of clinical evidence both at the national level and within professional societies.[2-5] Moreover, the FDA has released guidance documents on the use of real-world evidence reflecting increased interest in using RWD for regulatory decisions.[6-8] Despite their comprehensiveness, generating meaningful insights from RWD requires strict adherence to good observational research practices, particularly how missing data are handled.

Cancer registries have long been established as important sources of RWD within oncology for generating insights spanning cancer epidemiology and comparative effectiveness of therapies.[2,9] Data quality including the completeness of data elements is a major consideration when working with these registries to generate evidence. This is particularly germane given emerging evidence suggest that treatment-associated survival outcomes using registry and similar randomized controlled trials are not concordant.[10-12] As the reliance on clinical registries for evidence generation is likely to grow, there is critical need to assess the quality of clinical

evidence generated from registry and other RWD sources, as well as their adherence to best data practices.

Of note, cancer registries rely on trained tumor registrars to abstract and record data from the patient medical record. Lack of quality documentation within the medical record can lead to incompletely abstracted data elements, and therefore lead to unknown or missing data values within cancer registries.[13-15] While there are a variety of methods to account for missing data, patients with unknown values are likely under-represented in RWD studies, as it is common practice to simply exclude patients without complete information in variables used in cohort construction.[16-19] However, because missing data within registries is a surrogate for poor quality documentation, such data may not be "missing completely at random", and the exclusion of such patients with may introduce significant bias in studies based on these data. There is a paucity of literature studying the impact of incompletely recorded data within cancer registry datasets.

In our study, we aim to characterize the impact of unknown documentation across multiple cancer types within a large national registry-based RWD source. Specifically, we examine the prevalence of incomplete data across the three most common cancer types, and whether characteristics and survival outcomes of patients with missing data are comparable to those with complete data.

Methods

We examined the prevalence of incomplete records and associated cancer patient survival in a large RWD registry commonly used for comparative effective studies in oncology for the three most common cancers in the United States. We compared survival differences between patients who have complete versus incomplete data.

Data source

The National Cancer Database (NCDB) is a cancer registry established since the 1980s and jointly sponsored by the American College of Surgeons Commission on Cancer (CoC) and the American Cancer Society.[20] All CoC-accredited facilities report eligible cases to the NCDB, and it is estimated that over 70 percent of newly diagnosed cancer cases in the United States are represented.[21] There have been a large number of observational studies using NCDB data, some of which have been cited to support national cancer treatment guidelines.[22] Data reporting to the NCDB follows national registry coding standards and the recording of individual data elements have detailed specifications from the CoC.[23] Trained tumor registrars at individual cancer programs abstract specified data elements from patient charts in accordance to CoC registry data standards.[24] If a data element is unable to be identified within the patient record, it may be recorded with a blank or unknown value following registry coding guidelines.

For each patient record within the NCDB, there are over 130 variables in the PUF capturing a range of facility and patient information, tumor characteristics, treatment information, and cancer outcomes.[19] Specific information abstracted include complex data elements such as comorbidity score, cancer stage, diagnostic procedures, treatment information

including receipt of surgery, radiation, chemotherapy, among others. Data elements in our study are categorized into facility and patient demographics (Demographics), tumor characteristics (Cancer Identification), cancer stage (Stage), cancer treatments (Treatment), and survival outcomes (Outcomes) variables based on the NCDB data dictionary.

Missing data ascertainment

We identified 96 variables which were in use for all diagnosis years and disease sites included in our analysis. From these, we identified variables containing missing data in at least one patient record. Missing data were defined as either "blank" or "unknown" for a variable included in the database. Two clinical oncologists reviewed all variables and excluded variables where blank data entry was allowed by the NCDB data dictionary and may not reflect incomplete clinical documentation. For example, days from diagnosis to chemotherapy was recorded as missing if the patient never received chemotherapy and therefore excluded from analysis. Furthermore, to be conservative, TNM pathologic staging were not considered missing if pathologic stage was not documented or a pathologic specimen was not collected. A final 63 variables were identified as "variables of interest" to compare patients with and without completely documented data (Figure 1). All variables are listed in supplemental eTable 1.

Patient selection

We chose the three most common cancers with the largest number of cancer-related deaths in the United States for analysis.[25] We identified cancer cases in the 2019 release of non-

small cell lung cancer (NSCLC), breast cancer, and prostate cancer PUFs. All cancer stages were included and the impact of incomplete data was assessed by stage as defined by the NCDB analytic stage group, which combines pathologic and clinical stage by using clinical stage where pathologic stage is missing, since a recognized limitation of the NCDB is that there is a significant portion of patients with missing stage information.[26]

We included patients with cancer diagnosis years from 2006 to 2015 for analysis. Due to changes in data coding rules which introduced new variables and lack of survival information for the most recent diagnosis year, we excluded cases diagnosed in 2016 (most recent diagnosis year available). Given changes in data reporting standards and completeness over time, we examined cancer cases diagnosed in the most recent 10 years prior to 2016.

Statistical analysis

We calculated the percentage of patients with missing values in all included variables. We used standard descriptive statistics, chi-squared test, and Wilcoxon rank-sum test to show differences in patient, tumor, and treatment characteristics between patients with incomplete versus complete data. Patient records were not used for comparison of patient, tumor, and treatment characteristics if it has a missing value in the variable being compared, which are tabulated in supplemental eTable 2. To evaluate the association of incomplete data with survival outcomes, we used Kaplan-Meier estimates to compare overall survival between patients with incomplete versus complete data. Log-rank test was used to identify statistically significant differences in survival. Sub-analysis stratifying by cancer stage and treatment were performed to

identify groups of patients for which missing data was associated with largest differences in survival.

As a sensitivity analysis, we also tested an alternative approach of identifying variables for which data was missing in 1-20% of patient records. This range was determined a priori since <1% missing is likely to have little impact on outcomes of RWD studies, and a large percentage missing is more likely to be reflective of explainable differences in coding rules rather than poor documentation quality. For example, high percentages of missing clinical stage in the NCDB have been attributable to changes in stage coding rules.[27] Different percentage thresholds of missing data were also tested (supplemental data).

Statistical analysis was performed using Stata 16 (StataCorp LLC, College Station, Texas). Our code is available at: https://github.com/Aneja-Lab-Yale

Results

Distribution of variables and incomplete data

Of the 96 data elements included for analysis, there were 22 (22.9%) demographics, 11 (11.5%) tumor characteristics, 18 (18.8%) cancer stage, 41 (42.7%) treatment, and 4 (4.2%) outcomes variables. After limiting to variables of interest, there were 14 (22.2%) demographics, 6 (9.5%) tumor characteristics, 13 (20.6%) cancer stage, and 30 (47.6%) treatment variables.

(Figure 1). The majority of patients had missing data values in at least one variable of interest across all three disease sites (Table 1).

Patient, cancer, and treatment characteristics

There were significant differences in demographics, tumor characteristics, and cancer treatments received between patients with incomplete and complete data. Patients with missing data were more likely to be non-white, not insured, have more advanced stage disease, and less likely to receive surgery for all three disease sites. Cancer cases diagnosed in earlier years were more likely to have missing data (Table 2).

Association of incomplete data with overall survival

For NSCLC, there were 851,295 patients with incomplete data and 347,454 patients with complete data in variables of interest; 2-year survival was 33.2% for patients with missing data and 51.6% for patients with complete data (p<0.001). For breast cancer, there were 1,161,096 patients with missing data and 959,679 patients with complete data; 2-year survival was 93.2% for patients with missing data and 93.9% for patients with complete data (p<0.001). For prostate cancer, there were 460,167 patients with missing data and 698,468 patients with complete data; 2-year survival was 91.0% for patients with missing data and 95.6% for patients with complete data (p<0.001). This equates to an absolute 2-year survival difference of 18.4% for NSCLC patients, 0.7% for breast cancer patients, and 4.6% for prostate cancer patients overall. Kaplan-Meier survival curves are shown in Figure 2.

Large survival differences persisted among patients with metastatic disease when stratified by cancer stage. There were also survival differences in patients with non-metastatic disease, although the absolute difference was small for breast (0.4%) and prostate (1.3%) cancer patients, as compared to larger absolute survival differences of 4.6% and 16.7% in breast and prostate cancer patients with metastatic disease, respectively (p<0.001 for both, Figure 3). For metastatic NSCLC patients, 2-year survival was 13.1% for patients with missing data and 15.0% for patients with complete data (p<0.001); whereas among non-metastatic NSCLC patients, 2-year survival was 51.5% for patients with missing data and 63.2% for patients with complete data (p<0.001). Survival stratified by stage is shown in supplemental eFigures 1-3.

While rates of cancer treatment are different between patients with missing data and patients without missing data, survival differences were also observed when stratified between receipt of surgery, radiation, or chemotherapy, although the effect was less pronounced amongst prostate and breast cancer patients (supplemental eFigure 4).

Trends in data completeness and cancer stage over time

There were temporal changes in level of incomplete data from 2006 to 2015. The percentage of patients with missing data decreased from 81.8% to 67.1% (p<0.001) for NSCLC, from 78.1% to 46.5% (p<0.001) for breast cancer, and from 50.7% to 31.8% (p<0.001) for prostate cancer (supplemental eFigure 5). The distribution of cancer stage at diagnosis within the NCDB has also changed over time, which has previously been described. For example, there is

increased proportion of metastatic NSCLC at diagnosis in recent years, possibly attributable to increased use of advanced cancer imaging.[28,29] The change in overall stage is shown in supplemental eFigures 6. Moreover, we show survival differences persisted when further stratified by year of diagnosis (supplemental eFigure 7).

Sensitivity analysis using different percentages of missing data

When repeating our analysis using variables for which data was missing in 1-20% of patient records, for NSCLC, there were 622,831 patients with missing data and 575,918 patients without missing data in variables of interest; 2-year survival was 33.9% for patients with missing data and 43.5% for patients without missing data (p<0.001). For breast cancer, there were 1,481,729 patients with missing data and 639,046 patients without missing data in variables of interest; 2-year survival was 92.4% for patients with missing data and 96.0% for patients without missing data (p<0.001). For prostate cancer, there were 700,523 patients with missing data and 458,112 patients without missing data in variables of interest; 2-year survival was 91.7% for patients with missing data and 97.0% for patients without missing data (p<0.001, supplemental eFigure 8). Survival differences also persisted when we tested different thresholds using either 1-5% or 5-30% missing data as the cutoff (supplemental eFigure 9).

Discussion

In a large national cancer registry, we showed a high prevalence of incomplete patient records in all three of the most common cancer types, with the majority of patients missing essential data. The missing data has marked implications for clinical care and research and suggests that there are major gaps in documenting and capturing data via the medical records for patients with cancer. Moreover, patients with missing data were not representative of the overall cancer population and had significantly lower survival than those with complete data. These findings were replicated across NSCLC, breast, and prostate cancer patients. While the survival differences in patients with non-metastatic breast and prostate cancer were small, when stratified by cancer stage, larger differences persisted among patients with metastatic disease, suggesting differences in survival were driven by metastatic patients with incomplete data, whose courses of oncologic care have increased complexity.

Furthermore, we showed significant differences in terms of demographics, tumor characteristics, and treatments received between patients with missing data and patients without missing data. For example, incomplete records are more prevalent among blacks and other minorities, reflecting long-standing disparities in access to healthcare and cancer treatment.[30-32] Patients with fewer comorbid conditions also appeared to more frequently have missing data, which may reflect less available documentation due to fewer medical visits. Advanced stage patients were significantly more likely to have missing data. We hypothesize this is due to increased complexity of care in advanced cancers, leading to increased difficulty in documenting and abstracting all data elements.[33] The small survival differences in early-stage breast and prostate cancer patients are reflective of this in that definitive and adjuvant therapeutic management options in these settings have relatively less complexity. Moreover, completion of

primary site surgery at the CoC facility submitting the cancer case also helps to enable more complete documentation and easier data abstraction, as primary cancer and treatment information would be available at the same facility.

Our findings have major implications for RWD studies. While incomplete documentation is ubiquitous in RWD sources, observational studies using large cancer registries often simply exclude patients with missing data, and how missing data is handled is inconsistently reported in the medical literature.[34,35] Despite an increasing number of papers describing approaches for correcting missing data in observational studies, the practice of handling incomplete data amongst RWD sources has been slow to change.[36] Recent systematic comparisons of registry studies and randomized trials do not demonstrate concordant results.[10,12] Poor quality documentation is therefore a major obstacle to modern RWD sources and can introduce significant biases when measuring survival outcomes. We demonstrate a consistent association of missing data with worse survival across multiple cancer types, particularly amongst patients with advanced cancers. Therefore, adequate correction of missing data will be crucial for accurate treatment effect estimation in RWD studies.[37,38]

The clinical explanations for survival differences associated with incomplete data are likely multi-factorial. There were significant differences in distribution of cancer stage between patients with and without missing data. Underlying demographic disparities, differences in year of diagnosis, and treatments received are also contributory factors. There are also likely uncaptured confounders inherent to the observational nature of RWD studies. Our findings are corroborated by other studies examining missing data as a potential source of bias among RWD

sources.[39-42] Our results also corroborate previous analysis showing significant under ascertainment of stage and treatment data within cancer-specific registries.[43-45] Given fragmented or multi-facility care more often occurs among advanced-stage patients with increased treatment complexity, this is another plausible explanation for differences in cancer patient survival.[46] Since registry data abstraction necessarily depends on information available within the patient record, patients receiving care from multiple facilities are significantly more likely to have incomplete information. Therefore, fragmented care may be an explanatory link between documentation quality and survival outcomes particularly affecting patients with complex disease courses.[47-49]

There are limitations to our analysis. We examined survival and cannot draw conclusions on other outcomes such as toxicity, disease recurrence, or causes of death which are also likely associated with incomplete patient records. The dataset within our study is an observational cancer registry, and there may be limitations in the data abstraction process precluding complete capture of the medical record. However, all RWD sources likely face these limitations to a varying degree, and our analysis therefore should be interpreted as an exemplification of incomplete documentation within RWD sources in oncology.[50] Our study population is also heterogenous. The patients' cancer treatment paradigms including receipt and sequence of local and systemic therapies necessarily differ and do not reflect one specific clinical scenario. Nevertheless, survival differences between patients with missing and complete data persisted despite adjusting for multiple tumor and treatment-related factors. Additionally, the proportion of patients with incomplete data also depends on the number of variables examined, since it is more difficult to have complete documentation for a larger number of data elements. Given there are a

large number of variables within the NCDB, we also undertook an alternative analysis of choosing variables with missing data in 1-20% of patient records as variables of interest to identify patients with incomplete versus complete data. We also tested this assumption in sensitivity analysis, where we show survival difference persists using either 1-5% or 5-30% missing as the cutoff.

Conclusions

In conclusion, we show that the majority of patients in a large cancer registry-based RWD source are subject to incomplete data. Patients with missing data that was unable to be ascertained from the medical record show worse survival than those with more complete documentation. Increasing documentation quality and adoption of rigorous missing data correction methods are needed to best leverage RWD for clinical advancements.

References

1.   US Food and Drug Administration. Real-World Evidence. https://www.fda.gov/science-research/science-and-research-special-topics/real-world-evidence. Accessed 10/1/2020.

2.   Booth CM, Karim S, Mackillop WJ. Real-world data: towards achieving the achievable in cancer care. *Nat Rev Clin Oncol.* 2019;16(5):312-325.

3.   Penberthy L, Rivera DR, Ward K. The contribution of cancer surveillance toward real world evidence in oncology. *Semin Radiat Oncol.* 2019;29(4):318-322.

4.   Rivera D, Rubinstein WS, Schussler NC, et al. NCI and ASCO CancerLinQ collaboration to advance quality of cancer care and surveillance. *Journal of Clinical Oncology.* 2019;37(15_suppl):e18317-e18317.

5.   Schilsky RL. Finding the evidence in real-world evidence: moving from data to information to knowledge. *J Am Coll Surg.* 2017;224(1):1-7.

6.   US Food and Drug Administration. Framework for FDA's real-world evidence program. 2018.

7.   US Food and Drug Administration. Use of Real-World Evidence to Support Regulatory Decision-Making for Medical Devices. 2017.

8.   US Food and Drug Administration. Submitting Documents Using Real-World Data and Real-World Evidence to FDA for Drugs and Biologics Guidance for Industry. 2019.

9.   Parkin DM. The evolution of the population-based cancer registry. *Nat Rev Cancer.* 2006;6(8):603-612.

10.   Soni PD, Hartman HE, Dess RT, et al. Comparison of population-based observational studies with randomized trials in oncology. *Journal of Clinical Oncology.* 2019;37(14):1209-1216.

11.   Bartlett VL, Dhruva SS, Shah ND, Ryan P, Ross JS. Feasibility of using real-world data to replicate clinical trial evidence. *JAMA Netw Open.* 2019;2(10):e1912869.

12.   Kumar A, Guss ZD, Courtney PT, et al. Evaluation of the use of cancer registry data for comparative effectiveness research. *JAMA Network Open.* 2020;3(7):e2011985-e2011985.

13.   Curtis MD, Griffith SD, Tucker M, et al. Development and validation of a high-quality composite real-world mortality endpoint. *Health Serv Res.* 2018;53(6):4460-4476.

14.   Ebben KCWJ, Sieswerda MS, Luiten EJT, et al. Impact on quality of documentation and workload of the introduction of a national information standard for tumor board reporting. *JCO Clinical Cancer Informatics.* 2020(4):346-356.

15.   Piñeros M, Parkin DM, Ward K, et al. Essential TNM: a registry tool to reduce gaps in cancer staging information. *The Lancet Oncology.* 2019;20(2):e103-e111.

16.   Boffa DJ. What's lost in what's missing: a thoughtful approach to missing data in the National Cancer Database. *Ann Surg Oncol.* 2019;26(3):709-710.

17.   Rajyaguru DJ, Borgert AJ, Smith AL, et al. Radiofrequency ablation versus stereotactic body radiotherapy for localized hepatocellular carcinoma in nonsurgically managed patients: analysis of the National Cancer Database. *Journal of Clinical Oncology.* 2018;36(6):600-608.

18. Stokes WA, Bronsert MR, Meguid RA, et al. Post-treatment mortality after surgery and stereotactic body radiotherapy for early-stage non-small-cell lung cancer. *Journal of Clinical Oncology.* 2018;36(7):642-651.

19. Merkow RP, Rademaker AW, Bilimoria KY. Practical guide to surgical data sets: National Cancer Database (NCDB). *JAMA Surgery.* 2018;153(9):850-851.

20. Winchester DP, Stewart AK, Phillips JL, Ward EE. The national cancer data base: past, present, and future. *Ann Surg Oncol.* 2010;17(1):4-7.

21. American College of Surgeons. National Cancer Database. https://www.facs.org/quality-programs/cancer/ncdb. Accessed 10/1/2020.

22. Boffa DJ, Rosen JE, Mallin K, et al. Using the National Cancer Database for outcomes research: a review. *JAMA oncology.* 2017;3(12):1722-1728.

23. American College of Surgeons. Past Facility Oncology Registry Data Standards. https://www.facs.org/quality-programs/cancer/ncdb/call-for-data/fordsolder. Accessed 10/1/2020.

24. Bilimoria KY, Stewart AK, Winchester DP, Ko CY. The National Cancer Data Base: a powerful initiative to improve cancer care in the United States. *Ann Surg Oncol.* 2008;15(3):683-690.

25. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA: a cancer journal for clinicians.* 2020;70(1):7-30.

26. Hoskin TL, Boughey JC. ASO author reflections: a statistical caution regarding missing clinical stage in the National Cancer Database. *Ann Surg Oncol.* 2019;26(Suppl 3):569-570.

27.   Hoskin TL, Boughey JC, Day CN, Habermann EB. Lessons learned regarding missing clinical stage in the National Cancer Database. *Ann Surg Oncol.* 2019;26(3):739-745.

28.   Morgensztern D, Ng SH, Gao F, Govindan R. Trends in stage distribution for patients with non-small cell lung cancer: a National Cancer Database survey. *J Thorac Oncol.* 2010;5(1):29-33.

29.   Fletcher SA, von Landenberg N, Cole AP, et al. Contemporary national trends in prostate cancer risk profile at diagnosis. *Prostate Cancer Prostatic Dis.* 2020;23(1):81-87.

30.   Shavers VL, Brown ML. Racial and ethnic disparities in the receipt of cancer treatment. *Journal of the National Cancer Institute.* 2002;94(5):334-357.

31.   Wolf A, Alpert N, Tran BV, Liu B, Flores R, Taioli E. Persistence of racial disparities in early-stage lung cancer treatment. *J Thorac Cardiovasc Surg.* 2019;157(4):1670-1679.e1674.

32.   Zavala VA, Bracci PM, Carethers JM, et al. Cancer health disparities in racial/ethnic minorities in the United States. *British Journal of Cancer.* 2020.

33.   Sumpio C, Knobf MT, Jeon S. Treatment complexity: a description of chemotherapy and supportive care treatment visits in patients with advanced-stage cancer diagnoses. *Support Care Cancer.* 2016;24(1):285-293.

34.   Karahalios A, Baglietto L, Carlin JB, English DR, Simpson JA. A review of the reporting and handling of missing data in cohort studies with repeated assessment of exposure measures. *BMC Med Res Methodol.* 2012;12:96.

35.   Eekhout I, de Boer RM, Twisk JW, de Vet HC, Heymans MW. Missing data: a systematic review of how they are reported and handled. *Epidemiology.* 2012;23(5):729-732.

36. De Silva AP, Moreno-Betancur M, De Livera AM, Lee KJ, Simpson JA. A comparison of multiple imputation methods for handling missing values in longitudinal data in the presence of a time-varying covariate with a non-linear association with time: a simulation study. *BMC Med Res Methodol.* 2017;17(1):114.

37. D'Agostino RB, Jr., D'Agostino RB, Sr. Estimating treatment effects using observational data. *JAMA.* 2007;297(3):314-316.

38. Freemantle N, Marston L, Walters K, Wood J, Reynolds MR, Petersen I. Making inferences on treatment effects from real world data: propensity scores, confounding by indication, and other perils for the unwary in observational research. *BMJ.* 2013;347:f6409.

39. Jagsi R, Bekelman JE, Chen A, et al. Considerations for observational research using large data sets in radiation oncology. *Int J Radiat Oncol Biol Phys.* 2014;90(1):11-24.

40. Egleston BL, Wong YN. Sensitivity analysis to investigate the impact of a missing covariate on survival analyses using cancer registry data. *Stat Med.* 2009;28(10):1498-1511.

41. Eisemann N, Waldmann A, Katalinic A. Imputation of missing values of tumour stage in population-based cancer registration. *BMC Med Res Methodol.* 2011;11:129.

42. Jacobs CD, Carpenter DJ, Hong JC, Havrilesky LJ, Sosa JA, Chino JP. Radiation records in the National Cancer Database: variations in coding and/or practice can significantly alter survival results. *JCO Clin Cancer Inform.* 2019;3:1-9.

43. Jagsi R, Abrahamse P, Hawley ST, Graff JJ, Hamilton AS, Katz SJ. Underascertainment of radiotherapy receipt in Surveillance, Epidemiology, and End Results registry data. *Cancer.* 2012;118(2):333-341.

44.   Walker GV, Giordano SH, Williams M, et al. Muddy water? Variation in reporting receipt of breast cancer radiation therapy by population-based tumor registries. *Int J Radiat Oncol Biol Phys.* 2013;86(4):686-693.

45.   Walker GV, Grant SR, Jagsi R, Smith BD. Reducing bias in oncology research: the end of the Radiation Variable in the Surveillance, Epidemiology, and End Results (SEER) program. *Int J Radiat Oncol Biol Phys.* 2017;99(2):302-303.

46.   Hester CA, Karbhari N, Rich NE, et al. Effect of fragmentation of cancer care on treatment use and survival in hepatocellular carcinoma. *Cancer.* 2019;125(19):3428-3436.

47.   Polnaszek B, Gilmore-Bykovskyi A, Hovanes M, et al. Overcoming the challenges of unstructured data in multisite, electronic medical record-based abstraction. *Medical care.* 2016;54(10):e65-72.

48.   Clarke CA, Glaser SL, Leung R, Davidson-Allen K, Gomez SL, Keegan TH. Prevalence and characteristics of cancer patients receiving care from single vs. multiple institutions. *Cancer Epidemiology.* 2017;46:27-33.

49.   Howlader N, Ward KC, Warren JL, Campbell DS, Coyle L, Mariotto AB. Assessment of oncology practice billing claims for supplementing chemotherapy: a pilot study in the Georgia SEER cancer registry. *JNCI Monographs.* 2020;2020(55):82-88.

50.   Jarow JP, LaVange L, Woodcock J. Multidimensional evidence generation and FDA regulatory decision making: defining and using "real-world" data. *JAMA.* 2017;318(8):703-704.

Figure legends

Figure 1. Distribution of variable types among study population.

Figure 2. Survival of non-small cell lung cancer, breast cancer, and prostate cancer patients by whether data is missing in variables of interest.

Figure 3. Survival of patients with metastatic and non-metastatic non-small cell lung, breast, and prostate cancer by whether data is missing in variables of interest.

Tables

Table 1. Percentage of patients with missing data in at least one variable and by variable category.

	NSCLC	Breast	Prostate
	N=1,198,749	N=2,120,775	N=1,158,635
Any variable	71.02%	54.75%	39.72%
Demographics	13.01%	16.25%	13.94%
Cancer Identification	46.78%	13.40%	8.04%
Stage	35.11%	29.25%	17.12%
Treatment	16.02%	19.25%	12.83%

Table 2. Patient, disease, and treatment characteristics.

A. Non-small cell lung cancer

	Patients without missing data	Patients with missing data	p-value
	N=347,454	N=851,295	
Patient and facility			

Age at Diagnosis	69 (62-76)	69 (61-77)	<0.001
Sex			<0.001
Male	177,594 (51%)	451,217 (53%)	
Female	169,860 (49%)	400,078 (47%)	
Race			<0.001
White	303,607 (87.38%)	720,765 (85.59%)	
Black	34,565 (9.95%)	95,560 (11.35%)	
Other	9,282 (2.67%)	25,802 (3.06%)	
Hispanic Ethnicity			<0.001
Non-Hispanic	338,785 (97.50%)	758,913 (96.80%)	
Hispanic	8,669 (2.50%)	25,102 (3.20%)	
Charlson-Deyo Score			<0.001
0	184,687 (53.15%)	503,684 (59.17%)	
1	108,556 (31.24%)	229,207 (26.92%)	
2	38,916 (11.20%)	83,537 (9.81%)	
>=3	15,295 (4.40%)	34,867 (4.10%)	
Insurance			<0.001
Not Insured	8,818 (2.54%)	27,945 (3.38%)	
Private	92,017 (26.48%)	226,175 (27.32%)	
Medicaid	19,886 (5.72%)	53,265 (6.43%)	
Medicare	222,107 (63.92%)	506,860 (61.22%)	
Other Government	4,626 (1.33%)	13,691 (1.65%)	
Facility Type			<0.001

Community	240,682 (69.27%)	571,663 (67.76%)	
Academic	106,772 (30.73%)	271,994 (32.24%)	

Tumor

Year of Diagnosis	2011 (2009-2013)	2010 (2008-2013)	<0.001
Overall Stage			<0.001
Stage I	145,393 (41.85%)	171,141 (22.04%)	
Stage II	44,488 (12.81%)	55,601 (7.16%)	
Stage III	74,441 (21.43%)	174,460 (22.47%)	
Stage IV	83,073 (23.91%)	375,298 (48.33%)	
Tumor Size			<0.001
<=3 cm	167,184 (48.19%)	278,361 (44.46%)	
>3 cm	179,778 (51.81%)	347,749 (55.54%)	
Lymph Nodes Involved			<0.001
No	197,933 (58.75%)	287,971 (41.59%)	
Yes	138,977 (41.25%)	404,504 (58.41%)	
Distant Metastasis			<0.001
No	263,796 (75.92%)	445,966 (54.69%)	
Yes	83,658 (24.08%)	369,411 (45.31%)	

Treatment

Surgery (Primary Site)			<0.001
No	174,754 (50.30%)	669,039 (78.92%)	

	Patients without missing data N=347,454	Patients with missing data N=840,924	
Yes	172,700 (49.70%)	178,671 (21.08%)	
Radiation			<0.001
No	223,946 (64.45%)	481,005 (57.20%)	
Yes	123,508 (35.55%)	359,919 (42.80%)	
Chemotherapy			<0.001
No	207,763 (59.80%)	434,274 (53.54%)	
Yes	139,691 (40.20%)	376,777 (46.46%)	

B. Breast cancer

	Patients without missing data N=959,679	Patients with missing data N=1,161,096	p-value
Patient and facility			
Age at Diagnosis	62 (53-72)	59 (49-70)	<0.001
Sex			0.43
Male	8,552 (1%)	10,465 (1%)	
Female	951,127 (99%)	1,150,631 (99%)	
Race			<0.001
White	814,602 (84.88%)	947,362 (83.24%)	
Black	105,594 (11.00%)	137,369 (12.07%)	
Other	39,483 (4.11%)	53,425 (4.69%)	

Hispanic Ethnicity			<0.001
Non-Hispanic	915,866 (95.43%)	982,844 (93.62%)	
Hispanic	43,813 (4.57%)	66,997 (6.38%)	
Charlson-Deyo Score			<0.001
0	786,312 (81.93%)	997,133 (85.88%)	
1	137,187 (14.30%)	131,158 (11.30%)	
2	27,511 (2.87%)	24,880 (2.14%)	
>=3	8,669 (0.90%)	7,925 (0.68%)	
Insurance			<0.001
Not Insured	17,384 (1.81%)	25,447 (2.27%)	
Private	486,495 (50.69%)	626,116 (55.78%)	
Medicaid	53,951 (5.62%)	70,871 (6.31%)	
Medicare	392,685 (40.92%)	388,308 (34.59%)	
Other Government	9,164 (0.95%)	11,747 (1.05%)	
Facility Type			<0.001
Community	684,570 (71.33%)	725,684 (67.99%)	
Academic	275,109 (28.67%)	341,691 (32.01%)	
Tumor			
Year of Diagnosis	2012 (2009-2014)	2010 (2008-2013)	<0.001
Overall Stage			<0.001
Stage 0 (DCIS)	133,409 (13.91%)	294,752 (27.05%)	
Stage I	459,031 (47.85%)	391,027 (35.89%)	

Stage II	258,213 (26.92%)	254,836 (23.39%)	
Stage III	78,254 (8.16%)	97,157 (8.92%)	
Stage IV	30,454 (3.17%)	51,889 (4.76%)	
Tumor Size			<0.001
<=2 cm	629,447 (65.80%)	610,410 (63.96%)	
>2 cm	327,146 (34.20%)	344,006 (36.04%)	
Lymph Nodes Involved			<0.001
No	731,333 (77.35%)	760,552 (76.73%)	
Yes	214,178 (22.65%)	230,644 (23.27%)	
Distant Metastasis			<0.001
No	926,319 (96.56%)	1,030,431 (95.10%)	
Yes	33,042 (3.44%)	53,149 (4.90%)	

Treatment

Surgery (Primary Site)			<0.001
No	46,302 (4.82%)	109,849 (9.50%)	
Yes	913,377 (95.18%)	1,046,754 (90.50%)	
Radiation			<0.001
No	433,200 (45.14%)	566,736 (49.75%)	
Yes	526,479 (54.86%)	572,478 (50.25%)	
Chemotherapy			<0.001
No	634,319 (66.10%)	697,497 (63.81%)	
Yes	325,360 (33.90%)	395,557 (36.19%)	

Hormonal Therapy			<0.001
No	347,616 (36.22%)	487,554 (45.72%)	
Yes	612,063 (63.78%)	578,864 (54.28%)	

C. Prostate cancer

	Patients without missing data N=698,468	Patient with missing data N=460,167	p-value
Patient and facility			
Age at Diagnosis	65 (59-71)	65 (59-72)	<0.001
Sex			
Male	698,468 (100%)	460,167 (100%)	
Race			<0.001
White	579,894 (83.02%)	361,049 (81.74%)	
Black	99,417 (14.23%)	67,160 (15.20%)	
Other	19,157 (2.74%)	13,501 (3.06%)	
Hispanic Ethnicity			<0.001
Non-Hispanic	669,071 (95.79%)	366,527 (94.79%)	
Hispanic	29,397 (4.21%)	20,141 (5.21%)	
Charlson-Deyo Score			<0.001
0	573,655 (82.13%)	379,345 (82.44%)	

1	101,891 (14.59%)	64,092 (13.93%)	
2	17,408 (2.49%)	12,523 (2.72%)	
>=3	5,514 (0.79%)	4,207 (0.91%)	
Insurance			<0.001
Not Insured	11,414 (1.63%)	9,344 (2.14%)	
Private	337,278 (48.29%)	205,477 (47.07%)	
Medicaid	17,389 (2.49%)	12,835 (2.94%)	
Medicare	318,328 (45.58%)	201,474 (46.15%)	
Other Government	14,059 (2.01%)	7,415 (1.70%)	
Facility Type			<0.001
Community	434,953 (62.27%)	278,141 (60.56%)	
Academic	263,515 (37.73%)	181,155 (39.44%)	

Tumor

Year of Diagnosis	2010 (2008-2013)	2010 (2007-2012)	<0.001
Overall Stage			<0.001
Stage I	96,492 (13.82%)	48,900 (12.12%)	
Stage II	493,798 (70.70%)	266,757 (66.10%)	
Stage III	73,637 (10.54%)	43,243 (10.72%)	
Stage IV	34,503 (4.94%)	44,650 (11.06%)	
Lymph Node Involvement			<0.001
No	650,476 (97.24%)	337,102 (94.65%)	
Yes	18,464 (2.76%)	19,071 (5.35%)	

Distant Metastasis			<0.001
No	677,567 (97.01%)	383,731 (91.83%)	
Yes	20,862 (2.99%)	34,135 (8.17%)	
Treatment			
Surgery (Primary Site)			<0.001
No	314,879 (45.08%)	207,399 (45.39%)	
Yes	383,589 (54.92%)	249,492 (54.61%)	
Radiation			<0.001
No	446,325 (63.90%)	303,962 (67.64%)	
Yes	252,143 (36.10%)	145,409 (32.36%)	
Chemotherapy			<0.001
No	694,105 (99.38%)	417,776 (98.59%)	
Yes	4,363 (0.62%)	5,967 (1.41%)	
Hormonal Therapy			<0.001
No	550,765 (78.85%)	323,474 (77.11%)	
Yes	147,703 (21.15%)	96,039 (22.89%)	

A. All variables

B. Variables with any missing

C. Variables of interest

Demographics
Cancer Identification
Stage
Treatment
Outcomes

A. Non-small cell lung cancer patients

Number at risk						
Non-missing:	347123	215941	151213	106018	75034	51750
Missing:	847871	386406	240036	161564	114698	82313

B. Breast cancer patients

Number at risk						
Non-missing:	959024	860689	730814	598337	478874	369313
Missing:	1156072	1025098	892013	763473	644230	530044

C. Prostate cancer patients

Number at risk						
Non-missing:	697552	619620	542057	464227	393985	325987
Missing:	458062	393980	346704	304642	263883	219493

A. Non-small cell lung cancer patients

B. Breast cancer patients

C. Prostate cancer patients