

# **The impact of keeping a religious beard in the COVID-19 pandemic: an online cross sectional survey study exploring experiences of male medical healthcare professionals**

## **Authors and affiliations**

Dr Amer Hamed\*

Cardiology Consultant

Stepping Hill Hospital

Stockport

Email: [aabbilal@hotmail.com](mailto:aabbilal@hotmail.com)

Asam Latif

Senior Research Fellow

School of Health Sciences

Faculty of Medicine & Health Sciences

University of Nottingham, Nottingham

E: [Asam.Latif@nottingham.ac.uk](mailto:Asam.Latif@nottingham.ac.uk)

Mohammad Haris

MBChB, MRes

Internal Medicine Trainee, Cardiology

Blackpool Victoria Hospital

Email: [mohammad.haris@doctors.org.uk](mailto:mohammad.haris@doctors.org.uk)

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Sufyan Patel

MBChB, MPH

Foundation Doctor

University Hospitals of Leicester

Email: [Sufyan.patel@doctors.org.uk](mailto:Sufyan.patel@doctors.org.uk)

Muhammad I Patel BSc

Fourth year medical student

University of Glasgow

Email: [2178029P@student.gla.ac.uk](mailto:2178029P@student.gla.ac.uk)

Dr Syed Abdur Rahman Mustafa, FRCR

Consultant Interventional Radiologist

University Hospitals of Leicester

Mr Omeair Khan BSc( Hons) MBBS MRCS PGCAP FHEA

Core Surgical Trainee & Honorary Clinical Teaching Fellow

The Princess Alexandra Hospital NHS Trust

[omeair.khan@nhs.net](mailto:omeair.khan@nhs.net)

Dr Ahmad Shoaib

MBBS, MRCP (UK)

PG dip in research training (UK)

MD Cardiology (UK)

NIHR Academic Clinical Lecturer Cardiology

Keele University & Royal Stoke University Hospital

Stoke on Trent, UK

Salman Waqar

BSc MBBS MRCGP DRCOG PGCert(MedEd)

Academic General Practitioner

University of Oxford

Department of Primary Care Health Sciences

Woodstock Road

OX2 6GG

[salman.waqar@phc.ox.ac.uk](mailto:salman.waqar@phc.ox.ac.uk)

\*= corresponding author

### **Acknowledgements**

We would like to acknowledge all the participants who generously gave their time to complete the questionnaire. We extend a special thanks to Dr Mohammed Javed Iqbal, Dr Mohammed Bilal Patel who contributed significantly to the success of the study, Sharjeel Abu Ameerah and Yahya Hassan for their contribution in the study, Fayyaz Khan for reviewing the introduction. We also give thanks to the British Islamic Medical Association (BIMA) for their support especially Dr Arshad Latif who helped in forming the group of people interested in this subject.

### **Declaration of Competing Interest**

None.

### **Funding**

This study received no funding.

**Contributors:** Amer Hamed and Mohamed Javid Iqbal conceived the study. Muhammad Patel and Sufyan patel designed the study. Ahmad Shoiab is responsible for the data & performed the statistical analysis. All the authors participated in writing the first draft of the manuscript. All authors drafted the manuscript for important intellectual content, contributed to revision of the final version of the manuscript, approved the final version submitted, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

## **Abstract**

There has been a disproportionate effect on individuals from black Asian & Minority Ethnic (BAME) in the UK in Coronavirus disease (COVID-19) pandemic, especially in the NHS staff. Many of them have been asked to remove their beard to be eligible to do the fit test which can have negative implications on their spiritual, psychological & emotional wellbeing. This paper surveyed the responses of 469 healthcare professionals (HCPs) with beards regarding the challenges they face in regard to personal protective equipment (PPE), mask fit testing and attitude of employers & colleagues.

Professional discrimination through fit testing rejection, unavailability or inadequate PPE and the pressure to shave their beard are unpleasant outcomes of this pandemic for some of the NHS staff. NHS trusts & hospitals need to adjust their practices to include staff with beard in their COVID-19 arrangements.

## **Introduction**

The COVID-19 pandemic has required unprecedented adaptation of work systems and practices globally. (1) Research indicates that there has been a disproportionate effect on individuals from BAME backgrounds, particularly affecting key healthcare workers. Indeed, the first 11 doctors who lost their lives to this pandemic belonged to BAME communities. (2)

There has been much debate on the reasons why minority groups have been adversely affected. One reason is the availability and use of PPE for HCPs. There are multiple guidelines regarding appropriate PPE to protect clinicians and patients from the spread of COVID-19. For HCPs, when dealing with aerosol-generating procedures (AGPs) (e.g. cardiopulmonary resuscitation), the guidelines are unanimous in instructing that a higher level of PPE is required. This is generally a filtering facepiece (FFP3) mask or a powered air-purifying respirator (PAPR) hood. (3-5) The FFP3 mask requires fit testing to ensure a secure seal is formed to provide adequate protection. (6) Fit testing for respiratory protective equipment (RPE) has been a requirement in the USA since 1969 and in the UK since 1999. (7-8) The purpose of fit testing is to verify that the specific RPE used has the

potential to provide adequate protection for the individual. This means it cannot guarantee adequate protection at all times as this depends on the user ensuring they always fit the mask correctly. (6)

Unfortunately, fit testing poses several challenges. (9-10) Logistically, it is difficult to fit test all clinical staff within a short period of time and urgency of testing in a pandemic makes it problematic. (11) There are also concerns about wastage of several FFP3 masks and indispensable PPE kit during this testing in the middle of a pandemic. (12) Nevertheless, many individuals are unable to pass the fit testing due to their facial anatomy limiting a perfect seal, and others fail due to the presence of facial hair (or beard). (13) The PPE protection is, however, essential in fighting this pandemic and hence making it imperative to consider the respirator hood. The hood is a worthwhile, but costly alternative that provides effective protection and is not affected by the aforementioned issues. (3)

Many clinicians have been asked to remove their beard during the COVID-19 pandemic. A beard may be kept as a fashion trend for some people, and therefore shaving it is then an understandable and duty-bound sacrifice. However, for many individuals, a beard is a sacred aspect of religious practice. (14) This includes many religions such as Islam, Judaism, Sikhism and some sects within traditional Christianity. Many followers of these religions consider their beards to be an integral part of their personal and religious identity. (14) There is anecdotal evidence to suggest that male healthcare professionals have been asked to shave their beards in order to be eligible. (15) Requesting the removal of their beards in order to be eligible for fit testing could have huge negative implications on their psychological, spiritual, and emotional wellbeing as this is considered to be a religious requirement for many faith groups. (16)

### **Primary research objectives**

- To ascertain what challenges healthcare professionals who wore religious beards faced in regards to PPE, mask fit-testing and attitudes of employers and colleagues.
- To estimate the proportion of men who kept religious beards that subsequently trimmed or shaved due to circumstances during the COVID-19 pandemic.
- To estimate the proportion of men who wore a religious beard that faced challenges in their workplace.
- To gather the views and opinions on why men who kept religious beards trimmed or shaved, or, made no changes.

### **Methods**

#### **Study design**

This cross-sectional study was conducted using online social media tools from April to May 2020. Social networking sites (Facebook, LinkedIn, Twitter and Whatsapp) were used to

distribute the survey and adhere to COVID-19 guidelines. Participants were assured that the information they provide would be kept anonymous and confidential. The survey was created using Google Forms and a link to participate was distributed.

A qualifying question was asked at the beginning of the survey and completion of the questionnaire implied informed consent. Items for the survey were derived from the literature and through thorough discussions with experts and colleagues within the profession. The instrument contained a total of 36 questions. A variety of question types were used including Likert-type scales with responses on a ten-point scale ranging from “strongly agree” to “strongly disagree”, closed-ended questions and open-ended questions. The questionnaire consisted of three sections with the final section requiring participants to provide their socio-demographic details. This study was intended to be a descriptive study thus the questionnaire used was not validated.

### **Patient and public involvement statement**

No patients were involved in this study. Any self-identifying healthcare professional who wore a religious beard was eligible to participate. For the purposes of this study we defined a healthcare professional as any individual working in either a primary or secondary healthcare setting in a public facing role.

### **Pilot Testing**

The questionnaire was piloted by both academics and lay personnel. The aim of the pilot testing was to ascertain time taken to fill out form, in what order questions were asked, whether questions were understood as intended and if the questionnaire was user friendly. The feedback alongside responses from the pilot survey were then discussed by the study team and relevant changes were made to the questionnaire. The pilot responses were then removed before the finalised questionnaire was uploaded to Google Forms.

### **Analysis Plan**

The data was downloaded from Google Forms and inputted into a Microsoft Excel spreadsheet. After cleaning and consistency checks, the data was transferred to the Statistical Package for Social Sciences (SPSS) version 25 and was again checked for consistency and correctness. Descriptive analysis using frequency distributions, confidence intervals, distribution (mean etc), was used to analyse the data.

### **Ethical consideration**

The self-complete questionnaire was optional and therefore had few ethical implications. Consent was implied upon completion of the questionnaire. Following enquiry with the Health Research Authority (HRA), approval was not deemed necessary. Rather, approval from the BIMA ethics committee was sufficient. Adherence to ethical standards are maintained by informed consent, anonymising the data & provision of study information.

## **Results:**

### **Demographic profile of participants**

A total of 469 healthcare professionals participated in this survey. The number of respondents varied for individual questions in this survey and they are clearly detailed in Tables 1 – 4.

Almost all of our cohort came from BAME backgrounds with the majority of respondents of Muslim faith. More than three quarters were doctors. The hospital workplace was the predominant place of work. More than half of the participants had permanent contracts. The majority of respondents were from Britain and London was the most common workplace (67, 15%).

The median age of respondents was 39 years (range 20 - 69). There were varying ethnicities of respondents with the vast majority coming from Black, Asian and Minority Ethnic (BAME) groups (446, 98%). Asian / British Asian ethnicity made up the bulk of the BAME group (383, 84%). Only a minority were of a white background (10, 2%) possibly accounted for by the significantly smaller proportion of white people sharing Muslim, Sikh or Jewish beliefs. The majority of respondents shared the Muslim faith (437, 94%) followed by the Sikh faith (18, 4%) then the Jewish faith (6, 1%).

The highest percentage of respondents had worked in the NHS for greater than 10 years (265, 57%). Doctors accounted for the majority of respondents (372, 80%) whilst nurses (9, 2%), allied healthcare workers (34, 7%) and other support staff (49, 11%) also responded. Places of work included hospitals (390, 83%), GP practices (36, 8%) and social care among others. The types of contracts varied with the majority having permanent posts (258, 55%) whilst trainees and locums accounted for 23% (107) and 14% (66) respectively.

Of our cohort 25% (119) of respondents by the 14th of May 2020 had confirmed positive for SARS-CoV-2 (COVID-19).

### **The beard**

From our cohort 444 (95%) respondents felt that the beard was part and parcel of their religious identity. Despite the majority of participants (332, 71%) knowing that 'religion or belief' was a protected characteristic under the Equality Act 2010, more than half (255, 54%) of respondents who have a beard for religious reasons felt moderately or highly pressured by their colleagues, departments or trusts to trim or shave their beards during the pandemic.

### **FIT testing**

The majority of respondents were not offered a FIT test for the FFP3 face mask due to wearing a beard (168, 36%) or they did not pursue getting one (85, 18%). 11% (52) of respondents failed the FIT test due to having a beard. Of those who passed the FIT test, 6%

(27) passed with their beard, 5% (21) after trimming and 11% (53) after shaving. Almost three quarters of our respondents (345, 74%) were not given the opportunity to discuss their concerns regarding PPE and their religious observance of a beard with a senior clinician or a manager.

### **Alternative PPE**

Among the alternatives to the FFP3 mask are respirator hoods which do not require fit testing for use and can accommodate clinicians that fail FIT testing or wear beards. From our respondents that wore beards for religious observance, 61% (296) were not provided any PPE or were refused any alternatives. 22% were supported by their Trusts and alternate PPE was provided. Of those who were not provided with alternative PPE, 13% (63) purchased and used their own PPE, 20% (95) considered buying their own PPE but either couldn't afford it or found an appropriate model and 5% (25) who considered buying their own PPE were not allowed to use it.

### **Amended duties**

Of those who failed FIT testing (either due to not being tested or actually failing the test) and were not offered alternative PPE; 45% (213) were not offered any alternative duties at work. In the same group, Only 9% (44) were advised to abstain from high risk procedures, and less than 5% (28) offered work in non-clinical areas or to work from home.

### **Participants who changed their beards**

From our respondents 38% (176) either trimmed or shaved their beard due to COVID-19. The majority who shaved or trimmed their beards cited lack of PPE (102, 58%) and perceived risk of becoming unwell (88, 50%) as the two main reasons. More than a quarter (48, 27%) felt pressured from their employer to remove their facial hair.

Shaving the beard affected the mental or emotional wellbeing of 42% (74) of respondents whilst 61% (107) felt guilty for doing so. Half of respondents felt their religious identity had been violated (50%, 88).

### **Participant who retained their beards**

From our respondents 62% (293) didn't trim or shave their beard due to COVID-19. Of this group, More than half (163, 56%) feel they do not have adequate PPE available to protect themselves with a beard. The majority (262, 78%) felt moderately or highly supported by their colleagues to keep a beard for religious observance. However, 22% (63) experienced little support from their colleagues. A minority of departments (19, 13%) made changes to the work pattern of their employees who retained beards for religious observance. A handful of respondents (3, 1%) were sacked for refusing to shave their beards for religious observance.

## **Discussion**

The recent Public Health England (PHE) report details the impacts of COVID on BAME individuals and explores the effect of health inequalities within these groups. It emphasises that faith “provides an important foundation for communities’ resilience” highlighting the major role faith plays in helping an individual overcome such challenges. (17) Unfortunately, there have been reports of Islamophobia within the health service which undermines efforts to support people practicing their religion. (18) This study aimed to explore and propose reasonable solutions to the challenges faced by some HCPs who observe religious practice by keeping a beard. Investigating appropriate PPE for AGPs in clinical practice, our study is committed to determine a solution that protects religious identity whilst ensuring that the priority of saving lives is not compromised.

Many HCPs who keep a beard for religious reasons are from religious minority groups and so the possible discrimination may lead to a decrease in workforce diversity. The fit test processes currently do not appear to be consistent with the Human Rights Act. (19) Others have suggested that the lack of appropriate workwear has reduced opportunities for female Muslim women. (20) A costly but viable alternative to a fitted respirator mask is a respirator hood; this is a reasonable substitute that should be considered where religious identity issues are present. However, many HCPs are unaware of such alternatives and organisations may not be prepared to consider such costly options. (16, 21)

Workforce diversity helps to improve patient care. People from ethnic minority faces racial and professional discrimination during stages of their medical careers. (22) Muslims are the highest religious group to face such discrimination. (23) This discrimination was felt by the majority of healthcare professionals in a recent survey. (18)

Recent NHS new national guidance for NHS organisations in England published by the NHS employers suggested that the staff shave off their beards and facial hair to ensure protective face masks fit properly. (24)

Maintaining a religious beard comes with its challenges for the staff of the NHS in the current pandemic. Professional discrimination through fit testing rejection, unavailability or inadequate PPE and the pressure to shave your beard are unpleasant outcomes of this pandemic for NHS staff from primarily Muslim faith as the study shows.

Discrimination against staff is linked to poorer outcomes for patients, and little progress has been made in the last two decades on this front. Our findings are consistent with previous research which demonstrated discrimination faced by Muslim women with respects to dress codes in theatre and in hospital ward environments.

Furthermore, research into PPE provision in BAME frontline staff during the Covid-19 pandemic also showed that Muslims reported a higher rate of being reprimanded for

reporting PPE issues in the workplace, reported higher rates of Covid-19, and had the most difficulty accessing PPE. (25)

NHS staff from the faith groups need to feel accommodated, supported & empowered by their colleagues and trusts. Maintaining diversity in the workplace is important to improve the NHS care as well as its benefits to the society as a whole. NHS trusts & hospitals need to adjust their practices to include staff with beard in their Covid-19 arrangements.

Staff from faith groups who are subjected to such discrimination need to approach the diversity personnel in their trusts or seek help & support from their faith group.

### **Implications:**

The findings from this study shows that not enough consideration was given to the importance of the beard to health care staff, by not accommodating those who have it through failure to supply alternative PPE or direct the staff to an alternative type of work. In spite of this, staff still have faith in the health system to accommodate their religious needs.

### **Strengths and Limitations**

For the purpose of this cross-sectional study, social media platforms were used for disseminating the survey. The anonymity of respondents in this survey would facilitate honest reflections and the ability to speak freely on their sensitivities with regards to the survey. From the dataset gathered, it is clear that the majority of respondents were Muslims and there was relatively little representation of Jewish and Sikh respondents. It is also difficult to objectively assess how representative this survey would be of the NHS given that respondents were responding to social media advertisement to participate in the survey and some respondents were working in other countries. Furthermore, the sample was self-selected thus it may not equally reflect opinions and practices across the NHS.

The cross-sectional nature of the study doesn't take into account changes in opinions of healthcare staff and the practices across different hospitals over time to adapt to the attitudes of staff, the needs of clinicians and the effect on practice as the pandemic eases. On the other hand, this descriptive study does look at how practices within the NHS were around the peak of the pandemic and highlights important issues which may have not been considered yet, which significantly affected healthcare workers in a practical and spiritual sense.

### **Further research**

Further research will be needed to expand more on these important issues and challenges facing healthcare professionals who have a beard in regards to PPE, mask fit-testing and attitudes of employers and colleagues.

## References

1. Hu Y, Sun J, Dai Z, et al. Prevalence and severity of coronavirus disease 2019 (COVID-19): A systematic review and meta-analysis. *Journal of Clinical Virology*. 2020;127:104371
2. Kirby T. Evidence mounts on the disproportionate effect of COVID-19 on ethnic minorities. *Lancet Respir Med*. 2020;8(6):547-548
3. Cook TM. Personal protective equipment during the COVID-19 pandemic – a narrative review. *Anaesthesia*. 2020;75(7):920-927.
4. Verbeek JH, Rajamaki B, Ijaz S et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database of Systematic Reviews* 2020, Issue 4
5. Lee SA, Hwang DC, Li HY, et al. Particle Size-Selective Assessment of Protection of European Standard FFP Respirators and Surgical Masks against Particles-Tested with Human Subjects. *J Healthc Eng* 2016;2016:8572493
6. Clayton M, Vaughan N. Fit for purpose? The role of fit testing in respiratory protection. *Ann Occup Hyg*. 2005;49(7):545-548
7. ANSI. (1969) American National Standard practices for respiratory protection. ANSI Z88.2. New York: American National Standards Institute.
8. HSE. (2003) HSE operational circular OC 282/28 fit testing of respiratory protective equipment. Available at: [www.hse.gov.uk](http://www.hse.gov.uk).
9. Sandaradura I, Goeman E, Pontivivo G, et al. A close shave? Performance of P2/N95 respirators in healthcare workers with facial hair: results of the BEARDS (BEnchmarking Adequate Respiratory Defences) study. *Journal of Hospital Infection*. 2020;104(4):529-533.
10. RR1029 - Review of fit test pass criteria for Filtering Facepieces Class 3 (FFP3) Respirators [Internet]. Hse.gov.uk. 2020 [cited 1 May 2020]. Available from: <https://www.hse.gov.uk/research/rrhtm/rr1029.htm>
11. De Iaco G, Puro V, Fusco FM, et al. Personal protective equipment management and policies: European Network for Highly Infectious Diseases data from 48 isolation facilities in 16 European countries. *Infect Control Hosp Epidemiol*. 2012;33(10):1008-1016
12. Rowan NJ, Laffey JG. Challenges and solutions for addressing critical shortage of supply chain for personal and protective equipment (PPE) arising from Coronavirus disease (COVID19) pandemic - Case study from the Republic of Ireland. *Sci Total Environ*. 2020;725:138532
13. Floyd EL, Henry JB, Johnson DL. Influence of facial hair length, coarseness, and areal density on seal leakage of a tight-fitting half-face respirator. *J Occup Environ Hyg*. 2018;15(4):334-340
14. Peterkin A. One thousand beards: a cultural history of facial hair. *Arsenal Pulp Press*; 2001

15. Sheridan, D. (2020) 'Hospitals inspect doctors for beards and tell those with more than a day's stubble to shave', *The Telegraph*. Available at <https://www.telegraph.co.uk/news/2020/04/01/hospitals-inspect-doctors-beards-te-ll-days-stubble-shave> [Accessed 29/09/2020]
16. Baig AS, Knapp C, Egan AE, et al. Health care workers' views about respirator use and features that should be included in the next generation of respirators. *Am J Infect Control*. 2010;38(1):18-25
17. Public Health England. Beyond the data: Understanding the impact of COVID-19 on BAME groups. 2020
18. Day A. (2020) 'Exclusive: Muslim Medics Taunted About Bacon And Alcohol – By Their Own NHS Colleagues', *Huffington Post*. Available at [https://huffingtonpost.co.uk/entry/islamophobia-nhs-muslim-doctors-institutionalis-ed\\_uk\\_5f562e80c5b62b3add43cccb?guccounter=1](https://huffingtonpost.co.uk/entry/islamophobia-nhs-muslim-doctors-institutionalis-ed_uk_5f562e80c5b62b3add43cccb?guccounter=1) [Accessed 29/09/2020]
19. Human Rights Act 1998, c. 42. Available at: <https://www.legislation.gov.uk/ukpga/1998/42/schedule/1/part/1/chapter/8> (Accessed: 01 May 2020).
20. Malik A, Qureshi H, Humayra A-R, et al. 'I decided not to go into surgery due to dress code': a cross-sectional study within the UK investigating experiences of female Muslim medical health professionals on bare below the elbows (BBE) policy and wearing headscarves (hijabs) in theatre. *BMJ Open*. 2019;9:e019954
21. Weissman BR. Enjoying the advantages of PAPRs. *Occup Health Saf*. 2008;77(11):18-22.
22. McKenzie KJ. Racial discrimination in medicine. *BMJ* 1995;310:478–9.
23. Kline R, Martin B. Discrimination by Appointment: How black and minority ethnic applicants are disadvantaged in NHS staff recruitment. *Public World*. 2013.
24. Stephenson J. (2020) 'New coronavirus guidance tackles issue of beards among NHS staff', *Nursing Times*. Available at <https://www.nursingtimes.net/news/policies-and-guidance/new-coronavirus-guidance-tackles-issue-of-beards-among-nhs-staff-24-03-2020/> [Accessed 01 Sep 2020]
25. Daga S, Jafferbhoy S, Menon G, et al. Does Gender or Religion Contribute to the Risk of COVID-19 in Hospital Doctors? *Sushruta J of Health Policy & Opin*. 13(2) epub 22.06.2020

## Supplements

**Table 1: Demographic information**

Survey Question	No of Participants	Result
<b>Q 1: Age</b> (Mean) (SD)	462	39 (10) Range: 20 -69
<b>Q 2: How long have you been working in healthcare?</b>		
Under 5 year:	465	120 (26%)
5-10 years	465	80 (17%)
Over 10 years	465	265 (57%)
<b>Q 3: Ethnicity</b>		
Asian/Asian British	456	383 (84%)
Middle Eastern	456	39 (9%)
White	456	10 (2%)
Black	456	11 (2%)
Any other ethnic group	456	11 (2%)
Mixed	456	2 (0.44%)
<b>Q 4: Country of Birth?</b>		
UK	447	169 (38%)
India/ Pakistan/Bangladesh	447	184 (41%)
Africa	447	44 (10%)
Middle east	447	18 (4%)
Other	447	32 (7%)

<b>Q 5: Country of work</b>		
UK	436	427 (94%)
Outside UK	436	27 (6%)
<b>Q 6: City of Work</b>		
London	436	67 (15%)
Manchester	436	33 (8%)
Leicester	436	30 (7%)
Birmingham	436	22 (5%)
Blackpool	436	16 (4%)
All Others	436	268 (61%)
<b>Q 7: Your role in healthcare?</b>		
Doctor	464	372 (80%)
Allied healthcare worker	464	34 (7%)
Nurse	464	9 (2%)
Others	464	49 (11%)
<b>Q 8: Place of work</b>		
Hospital	469	390 (83%)
General practice		36 (8%)
Other places		43 (9%)
<b>Q 9: What contract do you have with your board?</b>		
Permanent/substantive	467	258 (55%)
Training	467	107 (23%)
Locum/Bank	467	66 (14%)
Part time	467	16 (4%)
Other	467	20 (4%)

<b>Q 10: Religion</b>		
Muslim	467	437 (94%)
Sikh	467	18 (4%)
Jewish	467	6 (1%)
Other/Not to say	467	5 (1%)
<b>Q 11: Did you get suspected or confirmed Covid-19?</b>		
Yes	467	119 (25%)
No	467	348 (75%)

**Table 2: General Questions**

<b>Survey Question</b>	<b>No of Participants</b>	<b>Result (% , 95% CI)</b>
<b>Q 1: Do you think your beard represents your religious identity?</b>		
Yes	469	444 (95%) (92-96)
No	469	25 (5%) (4-8)
<b>Q 2: Do you know that “religion or belief” is one of nine protected characteristics under the Equality Act 2010?</b>		
Yes	469	332 (71%) (66 – 75)
No	469	137 (29%) (25-34)
<b>Q 3: To what extent do you feel pressurised by your colleagues, department or Trust to trim/shave your beard during the current pandemic (rate it from 1-10)?</b>		
1-4	469	214 (46%) (41-50)

5-7	469	116 (25%) (21 -29)
8-10	469	139 (29%) (26 – 34)
<b>Q 4: Did you get a mask fit test?</b>		
Not offered a test due to beard	469	168 (36%) (32 – 40)
Did not being bother fit test due to beard	469	85 (18%) (15 – 22)
Failed due to beard	469	52 (11%) (8 – 14)
Passed only after shaving	469	53 (11%) (9 – 15)
Passed with beard	469	27 (6%) (4 – 8)
Passed only after trimming	469	21 (5%) (3 – 7)
Not offered/Other reasons	469	63 (13%) (10 – 17)
<b>Q 5: Did your Trust provide you any alternative PPE (e.g. Hood) due to your beard?</b>		
Yes	469	103 (22%) (18 – 26)
No	469	296 (61%) (56 – 65)
Not applicable	469	80 (17%) (14 – 21)
<b>Q 6: If you failed fit test or were not provided with alternative PPE, were you offered alternate form of work?</b>		
Yes, advised not to perform high risk procedures	469	44 (9%) (7 – 12)

Yes, offered to work in non-clinical area	469	14 (3%) (2 – 5)
Yes, offered to work from home	469	14 (3%) (2 – 5)
Yes, any other alternate form of work	469	11 (2%) (1 – 4)
No alternative option of work was given	469	213 (45%) (41 – 50)
Did not fail fit test or was offered alternative PPE	469	173 (37%) (33 – 41)
<b>Q7: If you failed fit test or were not provided with alternative PPE, did you consider buying your own PPE?</b>		
Yes, I considered and used my own PPE	469	63 (13%) (11 – 17)
Yes, I considered buying my own PPE but was not allowed to use own PPE	469	25 (5%) (4 – 8)
Yes, have considered buying my own PPE but could not find or afford it	469	95 (20%) (17 – 24)
No, I didn't consider it	469	160 (34%) (30 – 39)
Did not fail fit test or was offered alternative PPE	469	126 (27%) (23 – 31)
<b>Q 8: Does your Trust have a policy on PPE and beards?</b>		
Don't know	469	246 (52%) (48 – 57)
No	469	143 (30%) (26 – 35)
Yes	469	80 (17%) (14 – 20)
<b>Q 9: Were you given the opportunity to speak to a senior clinician or manager in your Trust to discuss PPE and beard issues?</b>		

Yes	469	124 (26%) (23 – 30)
No	469	345 (74%) (69 – 78)
<b>Q 10: Were you given any written instructions to trim or shave your beard?</b>		
Given written instruction	469	46 (10%) (7 – 13)
Given verbal instruction and was denied written instruction	469	19 (4%) (3 – 6)
Given verbal instruction and I did not ask for written instruction	469	150 (32%) (28 – 36)
Was not asked to trim or shave my beard	469	254 (54%) (50 – 59)
<b>Q 11: Did you shave or trim your beard due to Covid-19?</b>		
Yes	469	176 (38%) (33 – 42)
No	469	293 (62%) (58 – 67)

**Table 3: Participant who have not changed beard**

Survey Question	No of Participants	Result (% , 95% CI)
<b>Q 1: How confident you feel that you can retain beard during this pandemic (rate form 1-10)? 1 – Not sure at all, 10 – fully confident to retain my beard</b>		
1-4	293	31 (11%) (8 – 15)
5-7	293	56 (19%) (15 – 24)

8-10	293	206 (70%) (65 – 75)
<b>Q 2: Do you feel that you have adequate PPE to protect yourself with a beard?</b>		
Yes	293	130 (44%) (39 – 50)
No	293	163 (56%) (50 – 61)
<b>Q 3: Do you feel that your colleagues are supportive for you to keep beard (rate from 1-10)?</b> 1 - Not all supportive, 10 - fully supportive		
1-4	293	63 (22%) (17 – 27)
5-7	293	117 (40%) (34 – 46)
8-10	293	113 (38%) (33 – 44)
<b>Q 4: 4. Do you feel a need for change in working pattern or place due to your beard?</b>		
Yes	293	96 (33%) (28 – 38)
No	293	197 (67%) (62 – 72)
<b>Q 5: Did your department make any changes in work pattern to accommodate your beard?</b>		
Yes	293	39 (13%) (10 – 18)
No	293	254 (87%) (82 – 90)
<b>Q 6: 6. Do you feel threatened to lose your job for keeping a beard during this pandemic?</b>		
Yes	293	48 (16%) (12 – 21)
No	293	242 (83%) (78 – 87)

Yes, already lost my job due to this issue	293	3 (1 %) (0.3 – 3)
--------------------------------------------	-----	----------------------

**Table 4: Participant who trimmed or shaved beard**

Survey Question	No of Participants	Result (% , 95% CI)
<b>Q 1: 1. The reason for trimming/shaving? (Participants can select more than one option)</b>		
Voluntary	176	67 (38%) (34 – 42)
Perceived risk of getting illness	176	88 (50%) (46 – 54)
Pressure from employer	176	48 (27%) (23 – 31)
Lack of alternative PPE	176	102 (58%) (53 – 63)
Threat to lose job	176	15 (8%) (4 – 13)
Other	176	27 (15%) (11 – 19)
<b>Q 2: Was the opinion and guidance of a clinician, manager or faith leader imposed on you?</b>		
Yes	176	37 (21%) (16 – 28)
No	176	139 (79%) (72 – 84)
<b>Q 3: Do you feel guilt for trimming/shaving your beard?</b>		
Yes	176	107 (61%) (53 – 68)
No	176	69 (39%) (32 – 47)

<b>Q 4: 5. Has trimming/shaving your beard affected your mental or emotional wellbeing?</b>		
Yes	176	74 (42%) (34 – 49)
No	176	102 (58%) (50 – 65)
<b>Q 5: 6. Do you feel your religious identity has been violated?</b>		
Yes	176	88 (50%) (43 – 57)
No	176	88 (50%) (43 – 57)
<b>Q 6: Do you feel that your civic or human rights have been violated?</b>		
Yes	176	73 (41%) (34 – 49)
No	176	103 (59%) (51 – 66)