Delirium and Post-Discharge Neuropsychological Outcomes in Critically Ill Patients with COVID-19: an Institutional Case Series

Jacqueline Ragheb, MD[1];[†] Amy McKinney, MA[1];[†] Mackenzie Zierau, BSN[1]; Joseph Brooks, BS[1]; Maria Hill-Caruthers[1]; Mina Iskander MS[2]; Yusuf Ahmed, BS[1]; Remy Lobo, MD[3]; and Phillip E. Vlisides, MD[1,4*]

[1]Department of Anesthesiology, Michigan Medicine, Ann Arbor, MI USA

[2]New York Medical College, Valhalla, NY USA

[3]Department of Radiology, Michigan Medicine, Ann Arbor, MI USA

[4]Center for Consciousness Science, University of Michigan Medical School, Ann Arbor, MI USA

[†]These authors contributed equally to the manuscript

*Correspondence:

Dr. Phillip E. Vlisides

Department of Anesthesiology

University of Michigan Medical School, 1H247 UH, SPC-5048

1500 East Medical Center Drive

Ann Arbor, MI 48109-5048.

pvliside@med.umich.edu.

Phone: 734-936-4280.

Funding Statement: Supported by the Department of Anesthesiology, University of

Michigan Medical School.

Abstract

Delirium is a serious and common complication among critically ill patients with COVID-19. The objective of this study was to characterize the clinical course of delirium for COVID-19 patients in the intensive care unit, including post-discharge cognitive outcomes. A retrospective chart review was conducted for patients diagnosed with COVID-19 (n=148) admitted to an intensive care unit at Michigan Medicine between 3/1/2020 and 5/31/2020. Delirium was identified in 107/148 (72%) patients in the study cohort, with median (interquartile range) duration lasting 10 (4 – 17) days. Sedative regimens, inflammation, deviation from delirium prevention protocols, and hypoxic-ischemic injury were likely contributing factors, and the most common disposition for delirious patients was a skilled care facility (41/148, 38%). Among patients who were delirious during hospitalization, 4/17 (24%) later tested positive for delirium at home based on caretaker assessment, 5/22 (23%) demonstrated signs of questionable cognitive impairment or cognitive impairment consistent with dementia, and 3/25 (12%) screened positive for depression within two months after discharge. Overall, patients with COVID-19 commonly experience a prolonged course of delirium in the intensive care unit, likely with multiple contributing factors. Furthermore, neuropsychological impairment may persist after discharge.

Introduction:

The outbreak of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), the virus that causes Coronavirus Disease (COVID-19), emerged as a public health threat in December 2019 and was declared a pandemic by World Health Organization in March 2020. Several studies have reported the serious nature of complications associated with this disease. Major neurological complications, such as encephalopathy, delirium, strokes, seizures, and ataxia, have all been observed.[1-5] Delirium appears to be a common complication, with previous investigations demonstrating an incidence of approximately 65-80% in the intensive care unit (ICU).[1,4] Delirium may occur due to direct coronavirus invasion into central nervous system, [6] and systemic inflammatory responses may further exacerbate neurocognitive impairment. In the ICU, synergistic factors such as sedation regimen, social isolation, and deviation from standard care protocols may further increase risk. Delirium is also associated with prolonged hospitalization, long-term cognitive and functional impairment, and increased mortality.[7-9] As such, there is a critical need to advance understanding of this syndrome in patients with COVID-19.

While a high incidence of delirium has been previously reported in COVID-19 patients, fundamental questions persist. The clinical course of delirium, including average duration and post-discharge cognitive trajectory, remains unknown. Pathophysiologic drivers of delirium (e.g., polypharmacy, inflammation, cerebrovascular events) are incompletely understood, and the extent to which standard prevention protocols are implemented is unclear. Such detailed understanding will contribute to delirium

phenotyping of COVID-19 patients and provide insight into the clinical and neurocognitive burden associated with COVID-19. In this context, the objective of this study was to determine granular details associated with delirium in ICU patients with COVID-19. Specifically, the clinical course of delirium, presence of exacerbating factors, nature of prevention strategy implementation, and post-discharge cognitive outcomes were all characterized.

Results

Baseline characteristics are presented in Table 1. The majority of patients were African-American and non-Hispanic, and the most common comorbidities were hypertension, Diabetes mellitus, and obesity. Overall, age, demographics, and comorbidity profiles were similar between patients with delirium and those who did not experience delirium during hospitalization.

Delirium and Neuropsychological Outcomes

Delirium incidence was high in the cohort (107/148, 72%), and median (interquartile range) duration was 10 (4 – 17) days (Table 2). The most common delirium prevention measure, based on the ABCDEF ICU liberation bundle,[10,11] was assessment and treatment for pain; bringing familiar objects from home and spontaneous awakening trials were the least common (Table 2). Overall, the total number ICU liberation bundle activities charted was fewer than expected for the median duration of ICU admission (see Table 2 legend for description of prevention protocol activity schedule). New antidepressant use was more common for those with delirium (26/107, 24%) compared to those without delirium (4/41, 9.8%; p=0.049). Similarly, a psychiatry consult was obtained for 21/107 (20%) delirious patients compared to 0/41 (0%) in the non-delirium group (p=0.002). Lastly, no evidence of reversal or improvement was reported for nearly 30% of patients during index hospitalization.

Hospitalization and Post-Discharge Outcomes

In terms of hospitalization (Table 3), median length of hospitalization was 25 (13 – 48) days, and median length of ICU stay 15 (7 – 31). Length of hospitalization, ICU length of stay, and duration of mechanical ventilation were all significantly prolonged in patients experiencing delirium (Table 3). Correspondingly, sedative-hypnotic use was higher in patients with delirium. Delirious patients demonstrated higher white blood cell counts and higher levels of c-reactive protein, lactate dehydrogenase, and d-dimer compared to non-delirious patients. Less than half of patients were ultimately discharged home, and the most common disposition for those with delirium was a skilled care facility (41/107, 38%) after discharge (Table 3).

Neuropsychological outcomes after discharge are reported in Table 4. Among patients who were still alive and available to complete survey materials, nearly 25% of patients (4/17) scored positive for delirium based on family assessment (FAM-CAM), and all of these patients were delirious during hospitalization. Similarly, approximately 23% of patients (5/22) demonstrated either questionable impairment or impairment consistent with dementia based on the Short Blessed Test, and these patients were also delirious during hospitalization. Lastly, 12% of patients (3/25) screened positive for depression after discharge. The three patients who screened positive were also in the delirium group.

Neuroradiological Findings

In total, 47 patients underwent neuroimaging during hospitalization. The majority of imaging results were unremarkable or demonstrated incidental findings unrelated to

COVID-19. However, some notable findings were present. A brain MRI was ordered for a 59-year-old female with COVID-19 pneumonia and recent ECMO decannulation due to severe encephalopathy (i.e., no response to commands or noxious stimulus) with preservation of brainstem reflexes. Imaging revealed abnormal fluid attenuated inversion recovery (FLAIR) hyperintensity affecting the occipital and temporal lobes (Figures 1A, 1B), microhemorrhage in the splenium of the corpus callosum (Figures 1B, 1C) and posterior leptomeningeal enhancement Figures 1C, 1D), suggestive of encephalitis. A 29-year-old female with history of ovarian malignancy presented with new-onset seizures, which for an MRI was ordered. Results revealed diffuse dural thickening and enhancement (Figure 2A) one day prior to positive COVID testing. The differential diagnosis included intracranial hypotension (recent lumbar puncture), inflammation, infection, and neoplastic processes. No definitive diagnosis was reached, though this enhancement resolved approximately one month later (Figure 2B). Lastly, one patient demonstrated diffuse parenchymal abnormalities on MRI suggestive of bilateral hypoxic-ischemic injury (Figure 3). This patient experienced two separate arrests (pulseless electrical activity) within the preceding three weeks. A non-contrast head CT two weeks later demonstrated poor sulcation bilaterally, suggesting global hypoxic-ischemic injury (Figure 4).

Discussion

In a cohort of ICU patients with COVID-19, delirium was a common complication, affecting more than 70% of patients. Furthermore, delirium was associated with prolonged hospitalization, increased length of ICU stay, discharge to skilled care facilities, and positive screens for neuropsychological impairment up to two months after discharge. Delirium occurred in the setting of multiple sedative-hypnotic agents, acute inflammatory responses, deviation from delirium prevention protocols, and cerebrovascular events, which are all factors that could have further catalyzed delirium precipitation. While ICU liberation activities were conducted, many of the interventions were infrequently implemented, particularly those that reduce delirium risk (e.g., spontaneous awakening trials). Overall, the burden of cognitive impairment was high in patients with COVID-19, as was the risk of related complications.

These results align with previous data demonstrating a high incidence of delirium in critically ill patients with COVID-19.[1-4] Cognitive dysfunction may occur as a result of direct coronavirus invasion of the central nervous system[6] or other indirect mechanisms, such as polypharmacy, systemic inflammatory responses, or cerebrovascular events. Indeed, benzodiazepine sedation was common in this patient cohort, with nearly 60% of patients receiving midazolam sedation at one point during ICU admission. Lorazepam was a common sedation agent as well, and benzodiazepine use is associated with delirium in critically ill patients.[12-14] Inflammation may have also contributed to delirium risk. Inflammatory markers (e.g., c-reactive protein, ferritin, interleukin-6, lactate dehydrogenase) were considerably elevated in this patient cohort. In fact, serum levels

observed in this study aligned with – or exceeded – previously reported values in patients with severe COVID-19,[2,5] and there was MRI evidence of neuroinflammation for at least two patients in this series. Specific inflammatory markers (e.g., c-reactive protein, lactate dehydrogenase) were elevated in delirium patients and may reflect a specific in neuroinflammatory process. Indeed, c-reactive protein increases blood-brain barrier permeability in basic science models,[15] and lactate dehydrogenase has been associated with neurocognitive disorders.[16,17] However, these were unadjusted, bivariable analyses, and further covariate adjustment may have revealed a reduced – or absent – association.[5] Lastly, cerebral ischemia may also contribute to delirium risk in patients with COVID-19. Severe hypoxic-ischemic injury occurred in a patient who experienced multiple cardiopulmonary arrests during the course of illness. Stroke has previously been reported in patients with COVID-19,[18] as thromboembolic phenomena and cerebral malperfusion may both occur during the clinical course of COVID-19. As such, multiple processes likely contribute to delirium in patients with COVID-19. Targeted case-control studies can address some of these potential risk factors, such sedative-hypnotic regimens and inflammatory profiles.

Delirium prevention and management are inherently challenging for COVID-19 patients in the ICU. While delirium prevention bundles have been consistently demonstrated to reduce risk,[19,20] unique challenges posed by COVID-19 hinder the implementation of standard prevention practices. Spontaneous awakening and breathing trials, for example, were often not possible due to illness severity and associated ventilator requirements. Indeed, the median number of total spontaneous awakening trials charted

was two across the entire cohort. Clinicians may have also been limited in terms of sedation regimen. Agitation was commonly observed, and nearly one-third of patients required antipsychotics. Agitation and hyperactive delirium likely prompted additional sedation and prolonged use of physical restraints. Early mobility was limited given illness severity, and family engagement was often not possible due to visitor policy restrictions. In-person interactions with clinicians were also limited given the intent of reducing virus transmission. As such, the culmination of disease severity, limited face-to-face time spent with patients, and visitor restriction policies likely hindered ICU liberation bundle implementation. Indeed, given that the median length of ICU stay was 15 days across the cohort, most interventions were charted less than once daily (Table 2). Limited implementation of delirium prevention bundles could have increased risk for delirium incidence, duration, and severity. Interestingly, prevention bundle activities were charted more commonly in delirious patients. This may have reflected clinician awareness of delirium and attempts to reduce risk. Novel strategies for implementing delirium prevention bundles in this patient population may help to further mitigate risk and should be tested in prospective trials.

Neuropsychological impairment after discharge was also present for some patients based on subjective reporting, caretaker assessment, and objective testing for depression and cognitive impairment. Furthermore, all patients that screened positive for possible impairment also experienced delirium in the hospital. These estimates may have been even higher, given that approximately 70% patients that were called for post-discharge cognitive assessment were still admitted to skilled care facilities, refused

participation, or were unable to be reached. Whether post-discharge cognitive impairment was related specifically to COVID-19 or critical illness more broadly is unclear. Indeed, cognitive impairment is common at discharge for patients who experienced delirium while in the ICU, and delirium is present for nearly 20% of patients newly admitted to acute care facilities.[7,21] Moreover, cognitive impairment can be present for months-to-years after acute respiratory distress syndrome and sepsis,[22-24] and symptoms of depression and post-traumatic stress disorder are commonly reported among ICU survivors.[25] Neuropsychological impairment after discharge may, in part, reflect critical illness, rather than pathophysiologic insults specific to COVID-19. Nonetheless, ICU patients with COVID-19 experience considerable neuropsychological burden, and related complications, both during and after hospitalization.

The strengths of this study include granular data with respect to delirium, potential risk factors, nature and frequency of prevention strategies, related complications, and post-discharge outcomes. Data were representative of an academic tertiary care center with nearly 150 patients. A validated chart review method was used to identify delirium,[26] and the study measures used to characterize cognitive function, such as the FAM-CAM, Short Blessed Test, and PROMIS assessments, are validated measures that increase confidence in the results. In terms of limitations, this is a this was a single center analysis, and the results are restricted to the institution studied. The study was not conducted with a matched control cohort, as the nature of this study was descriptive. Lastly, data were limited for post-discharge cognitive outcomes, as more than half of patients called were unavailable to complete assessments.

In summary, delirium is common complication of COVID-19 with multiple contributing factors. Furthermore, neuropsychological impairment may persist in some patients after discharge. Further research should aim to identify independent risk factors in this population and novel, effective prevention strategies.

Methods:

Study design and overview

This was a single-center, institutional case series from Michigan Medicine. Detailed chart review data were collected from critically ill patients with COVID-19 (3/1/2020 – 5/31/2020), and post-discharge telephone surveys were conducted to determine if cognitive impairment persisted after discharge. All study operations were conducted at Michigan Medicine, Ann Arbor MI USA, and approval was obtained from the University of Michigan Medical School Institutional Review Board (HUM00182646).

Eligibility criteria

All patients with a COVID-19 diagnosis admitted to a Michigan Medicine ICU between 03/01/2020 – 05/31/2020 were eligible for study inclusion. ICU patients admitted during this time, without a diagnosis of COVID-19, were not eligible for study inclusion.

Outcomes

The primary outcome was delirium presence (yes/no, %) at any point during admission. Delirium was evaluated via chart review method (described below). Several secondary outcomes were also collected in relation to delirium and overall clinical trajectory. These outcomes included the following: duration of delirium (days), antipsychotic administration, length of hospital stay, length of ICU stay, number of days requiring ventilator support, inflammatory laboratory values (white blood cell count, c-reactive protein, ferritin, lactate dehydrogenase, d-dimer, and interleukin-6), new psychiatry consults, new antidepressant use, and final disposition (e.g., home, long-term care

facility, death). Delirium prevention strategies, based on the ABCDEF ICU liberation bundle,[10,11] were also recorded. These included the following: assess and treat for pain, discontinue physical restraints, structured mobility exercises, removal of temporary medical lines or devices, place familiar objects from home at the bedside, family education and reassurance, promote use of visual and hearing aids, sleep promotion protocols (e.g., lights off at night, melatonin tablets), and spontaneous awakening trials. The total number of times a prevention strategy was charted was recorded for each patient. The median of these total values was then reported for each prevention strategy. Neuroimaging data were also collected and reviewed.

Lastly, a telephone survey was conducted between 30-60 days post-discharge to determine whether subjective or objective signs of cognitive impairment were present. During telephone interviews, the following tests were conducted: the Patient-Reported Outcomes Measurement Information System (PROMIS)[27] Cognitive Function Abilities 4a, Short Blessed Test (score 0-4 = normal cognition, score 5-9 = questionable impairment, score ≥10 = impairment consistent with dementia),[28] Family Confusion Assessment Method for delirium,[29] and the Patient Health Questionnaire-9 (scores ≥10 were considered positive screens for depression).[30,31]

Data collection

Screening for eligible patients was first performed for via DataDirect, a self-serve software tool from the University of Michigan Office of Research that enables research

teams to retrospectively search for patient cohorts. Charts that screened positive were then manually reviewed by study team members to confirm study eligibility.

Charts were then reviewed in further detail for outcome abstraction. Delirium was assessed via a validated chart review method.[26] Briefly, any instance of an acute confusional state was recorded in the instrument. The source of information was recorded, along with the date and time. The total number of days with acute confusion was also included in the instrument, along with any evidence of reversibility or improvement of the confusion state. Other clinical outcomes, along with laboratory values, were collected directly from the charts. Neuroimaging studies were manually reviewed by a board certified radiologist with a Certificate of Added Qualification in neuroradiology (R.L.).

Statistical Analysis

All analyses were performed using IBM SPSS version 27 (Armonk, NY USA). Exploratory data analysis techniques were used to assess the distribution of dependent measures for determining the appropriate analytical strategy. The Shapiro-Wilk test was used to assess the distribution of continuous outcomes, and Independent t-tests or Mann-Whitney U tests were used as appropriate. Mean (standard deviation) or median (interquartile range) was reported for parametric and non-parametric data, respectively. For binary outcomes and proportions, The Chi-Square Test or Fisher's Exact Test were used, as appropriate. The threshold for significance was set to $p<0.05$ across all tests. For post-discharge cognitive outcomes, descriptive statistics were reported.

References

Helms, J. *et al.* Neurologic features in severe SARS-CoV-2 infection. *N Engl J Med.* **382**, 2268-2270 (2020).

Mao, L. *et al.* Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol.* **77**, 683-690 (2020).

Varatharaj, A. *et al.* Neurological and neuropsychiatric complications of COVID-19 in 153 patients: a UK-wide surveillance study. *Lancet Psychiatry.* **7**, 875-882 (2020).

Helms, J. *et al.* Delirium and encephalopathy in severe COVID-19: a cohort analysis of ICU patients. *Crit Care.* **24**, 491 (2020).

Liotta, E. M. *et al.* Frequent neurologic manifestations and encephalopathy-associated morbidity in COVID-19 patients. *Ann Clin Transl Neurol.* 10.1002/acn3.51210 (2020).

Bergmann, C. C., Lane, T. E. & Stohlman, S. A. Coronavirus infection of the central nervous system: Host-virus stand-off. *Nat Rev Microbiol.* **4**, 121-132 (2006).

Ely, E. W. *et al.* Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA.* **291**, 1753-1762 (2004).

Girard, T. D. *et al.* Delirium as a predictor of long-term cognitive impairment in survivors of critical illness. *Crit Care Med.* **38**, 1513-1520 (2010).

Brummel, N. E. *et al.* Delirium in the ICU and subsequent long-term disability among survivors of mechanical ventilation. *Crit Care Med.* **42**, 369-377 (2014).

Marra, A., Ely, E. W., Pandharipande, P. P. & Patel, M. B. The ABCDEF bundle in critical care. *Crit Care Clin.* **33**, 225-243 (2017).

Pandharipande, P., Banerjee, A., McGrane, S. & Ely, E. W. Liberation and animation for ventilated ICU patients: The ABCDE bundle for the back-end of critical care. *Crit Care.* **14**, 157 (2010).

Pisani, M. A. *et al.* Benzodiazepine and opioid use and the duration of intensive care unit delirium in an older population. *Crit Care Med.* **37**, 177-183 (2009).

Zaal, I. J. *et al.* Benzodiazepine-associated delirium in critically ill adults. *Intensive Care Med.* **41**, 2130-2137 (2015).

Pandharipande, P. *et al.* Lorazepam is an independent risk factor for transitioning to delirium in intensive care unit patients. *Anesthesiology.* **104**, 21-26 (2006).

Hsuchou, H., Kastin, A. J., Mishra, P. K. & Pan, W. C-reactive protein increases BBB permeability: Implications for obesity and neuroinflammation. *Cell Physiol Biochem.* **30**, 1109-1119 (2012).

Schiepers, O. J. *et al.* Serum iron parameters, HFE C282Y genotype, and cognitive performance in older adults: Results from the FACIT study. *J Gerontol A Biol Sci Med Sci.* **65**, 1312-1321 (2010).

Reed, T. *et al.* Redox proteomic identification of 4-hydroxy-2-nonenal-modified brain proteins in amnestic mild cognitive impairment: Insight into the role of lipid peroxidation in the progression and pathogenesis of Alzheimer's disease. *Neurobiol Dis.* **30**, 107-120 (2008).

Ntaios, G. *et al.* Characteristics and outcomes in patients with COVID-19 and acute ischemic stroke: The global COVID-19 stroke registry. *Stroke.* **51**, e254-e258 (2020).

Ely, E. W. The abcdef bundle: Science and philosophy of how ICU liberation serves patients and families. *Crit Care Med.* **45**, 321-330 (2017).

Barnes-Daly, M. A., Phillips, G. & Ely, E. W. Improving hospital survival and reducing brain dysfunction at seven california community hospitals: Implementing PAD guidelines via the ABCDEF bundle in 6,064 patients. *Crit Care Med.* **45**, 171-178 (2017).

Kiely, D. K. *et al.* Delirium among newly admitted postacute facility patients: Prevalence, symptoms, and severity. *J Gerontol A Biol Sci Med Sci.* **58**, M441-445 (2003).

Hopkins, R. O. *et al.* Two-year cognitive, emotional, and quality-of-life outcomes in acute respiratory distress syndrome. *Am J Respir Crit Care Med.* **171**, 340-347 (2005).

Ehlenbach, W. J. *et al.* Association between acute care and critical illness hospitalization and cognitive function in older adults. *JAMA.* **303**, 763-770 (2010).

Iwashyna, T. J., Ely, E. W., Smith, D. M. & Langa, K. M. Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA.* **304**, 1787-1794 (2010).

Hatch, R. *et al.* Anxiety, depression and post traumatic stress disorder after critical illness: a UK-wide prospective cohort study. *Crit Care.* **22**, 310 (2018).

Inouye, S. K. *et al.* A chart-based method for identification of delirium: Validation compared with interviewer ratings using the confusion assessment method. *J Am Geriatr Soc.* **53**, 312-318 (2005).

Cella, D. *et al.* The patient-reported outcomes measurement information system (PROMIS): Progress of an NIH roadmap cooperative group during its first two years. *Med Care.* **45**, S3-s11 (2007).

Katzman, R. *et al.* Validation of a short orientation-memory-concentration test of cognitive impairment. *Am J Psychiatry.* **140**, 734-739 (1983).

Steis, M. R. *et al.* Screening for delirium using family caregivers: Convergent validity of the family confusion assessment method and interviewer-rated confusion assessment method. *J Am Geriatr Soc.* **60**, 2121-2126 (2012).

Kroenke, K., Spitzer, R. L. & Williams, J. B. The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med.* **16**, 606-613 (2001).

Levis, B., Benedetti, A. & Thombs, B. D. Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: Individual participant data meta-analysis. *BMJ.* **365**, l1476 (2019).

Acknowledgments:

We would like to acknowledge Dr. Michael Kenes (PharmD, BCPS, BCCCP) and Ms.

Margaret Diehl for assistance with medical chart data extraction.

Author Contributions:

The study was originally conceived by J.R., A.M., M.Z., and P.E.V. Data acquisition was

conducted by J.R., A.M., M.Z., J.B., M.H.C., Y.A., and P.E.V. Data analysis was

conducted by J.R., A.M., M.Z., J.B., M.H.C., M.I., Y.A., and P.E.V. Final statistical

analysis was conducted by P.E.V. Neuroimaging studies were acquired and analyzed

by R.L., and the final neuroimaging figures were generated by R.L. and P.E.V. All

authors contributed to the manuscript writing, critically reviewed the manuscript for

intellectual content, and approved the final manuscript.

Competing Interests Statement: The authors declare no competing interests.

Figure 1. Axial fluid-attenuated inversion recovery (FLAIR) (**a, b**) images at the level of the basal ganglia show abnormal FLAIR hyperintense signal (arrows) affecting the bilateral occipital, temporal lobes. This appears almost sulcal suggesting a higher protein component within the cerebrospinal fluid. Note the elevated FLAIR signal in the splenium of the corpus callosum (arrow) suggesting parenchymal insult. Axial susceptibility weighted imaging (SWI) (**c**) at the level of the splenium of the corpus callosum shows small areas of susceptibility (arrow) in the splenium, likely related to microhemorrhage. Axial T1 (**d**) post-contrast with fat suppression at the level of the basal ganglia shows subtle, though true, enhancement (arrows) in the posterior sulci, arachnoid pial (leptomeningeal) pattern suggesting a degree of encephalitis.

Figure 2. Axial T1 post-contrast with fat suppression images at the level of the mid lateral ventricles demonstrate smooth dural enhancement (**a**) (arrows) along the bilateral cerebral convexities. This feature resolves one month later (**b**) (arrows). The overall pattern of dura-arachnoid/pachymeningeal enhancement is non-specific, can be seen with intracranial hypotension, in the procedural setting (e.g., lumbar puncture), and other scenarios (e.g., infection, inflammation).

Figure 3. After initial non-contrast head CT (which was unremarkable), the patient had diffuse parenchymal abnormalities on MRI examination. Axial diffusion-weighted imaging (**a**) shows hyperintense signal (arrows) at the bilateral basal ganglia, thalami, and posterior cortices, regions are hypointense (arrows) on corresponding apparent diffusion coefficient map (**b**). These same locations are hyperintense on T2 (**c**) and FLAIR (**d**), especially at the basal ganglia and thalami.

Figure 4. Non-contrast head CT performed approximately two weeks after cardiopulmonary arrest shows poor sulcation (arrows) bilaterally, suggesting global insult, most likely hypoxic-ischemic in etiology.

Table 1. Baseline Characteristics

	All Patients (n=148)	Delirium (n=107)	No Delirium (n=41)	*P Value*
Age (IQR)	59 (49 – 71)	58 (47 – 70)	62 (54 – 71)	0.126
Male sex, n (%)	98 (67)	32 (30)	18 (44)	0.107
Race, n (%)				0.464
Caucasian	66 (45)	47 (44)	19 (46)	
African-American	70 (47)	53 (50)	17 (41)	
Other	1 (0.7)	1 (0.9)	0 (0)	
Not reported	11 (7.4)	6 (5.6)	5 (12)	
Ethnicity, n (%)				0.182
Non-Hispanic	137 (93)	99 (93)	38 (93)	
Hispanic	5 (3.4)	5 (4.7)	0 (0)	
Unknown/not reported	6 (4.1)	3 (2.8)	3 (7.3)	
Weight, kg (IQR)	103 (83 – 127)	105 (86 – 128)	96 (79 – 115)	0.064
BMI (IQR)	34 (28 – 40)	34 (29 – 41)	31 (28 – 39)	0.358
Comorbidities, n (%)				
Asthma	24 (16)	17 (16)	7 (17)	0.861
Atrial fibrillation	22 (15)	14 (13)	8 (20)	0.325
Cancer	25 (17)	20 (19)	5 (12)	0.345
Chronic kidney disease	40 (27)	29 (27)	11 (27)	0.973
Congestive heart failure	19 (13)	6 (15)	13 (12)	0.686
COPD	14 (9.5)	8 (7.5)	6 (14)	0.183

Coronary artery disease	27 (18)	18 (17)	9 (22)	0.470
Depression	17 (11)	11 (10)	6 (14)	0.457
Diabetes mellitus	75 (51)	57 (53)	18 (44)	0.308
Hypertension	102 (69)	73 (68)	29 (71)	0.768
Obstructive sleep apnea	31 (21)	22 (21)	9 (22)	0.825
Seizures	8 (5.4)	5 (4.7)	3 (7.3)	0.524
Stroke	9 (6.1)	5 (4.7)	4 (9.8)	0.263
Substance abuse	9 (6.1)	5 (4.7)	4 (9.8)	0.214
TIA	5 (3.4)	3 (2.8)	2 (4.9)	0.617

Median (interquartile range, IQR) data presented. *Kg* kilograms, *BMI* body mass index, *COPD* chronic obstructive pulmonary disease, *TIA* transient ischemic attack.

Table 3. Delirium and Neuropsychological Outcomes

	All Patients (n=148)	Delirium (n=107)	No Delirium (n=41)	*P* Values
Delirium measures				
Delirium, n (%)	107 (72)	107 (100)	---	---
Duration of delirium, days (IQR)	10 (4 – 17)	10 (4 – 17)	---	---
Antipsychotic use, n (%)	44 (30%)	41 (38)	3 (7.3)	<0.001
Evidence of reversal, n (%)	---	76 (71)	---	---
Prevention measures – number of instances, (IQR)				
Assess and treat for pain	32 (16 – 50)	36 (22 – 58)	16 (8 – 43)	<0.001
Discontinue restraints	7 (3 – 15)	9 (5 – 16)	3 (1 – 5)	<0.001
Mobility exercises	20 (11 – 36)	23 (14 – 41)	11 (5 – 26)	<0.001
Removal of temporary medical lines or devices	5 (2 – 10)	6 (4 – 12)	2 (1 – 5)	<0.001
Place familiar objects at bedside	2 (1 – 6)	3 (1 – 7)	1 (0 – 2)	0.001
Family education and reassurance to family	5 (2 – 10)	5 (3 – 11)	4 (1 – 8)	0.008
Promote use of visual and hearing aids	6 (2 – 11)	6 (3 – 120	5 (1 – 11)	0.237
Sleep promotion protocol	19 (9 – 20)	23 (11 – 330	13 (6 – 26)	0.003
Spontaneous awakening trial	2 (0 – 5)	3 (3 – 6)	0 (0 – 2)	<0.001
Psychiatric outcomes				
New antidepressant use, n (%)	30 (20)	26 (24)	4 (9.8)	0.049

New psychiatry consults, n (%)	21 (14)	21 (20)	0 (0)	0.002

Delirium prevention measures are based on the standard ICU liberation bundle protocols (see text for details). Per institutional protocol, clinicians conduct pain assessments every two hours (and more frequently as needed for procedures), range of motion exercises are scheduled three times daily, spontaneous awakening/breathing trials occur daily (if medically appropriate), and family updates are scheduled once daily. Given this schedule, these prevention measures were charted less frequently than expected based on a median ICU course of 15 (7 – 31) days. *IQR* interquartile range.

Table 3. Hospitalization

	All Patients (n=148)	Delirium (n=107)	No Delirium (n=41)	*P Value*
Length of hospitalization, days (IQR)	25 (13 – 48)	31 (18 – 52)	11 (8 – 27)	<0.001
Length of ICU stay, days (IQR)	15 (7 – 31)	19 (12 – 38)	4 (2 – 100	<0.001
Ventilator time, days, (IQR)	12 (3 – 28)	18 (10 – 29)	0 (0 – 7)	<0.001
Required hemodialysis, n (%)	45 (30)	39 (36)	6 (14)	0.010
Sedative-hypnotics, n (%)				
Propofol	113 (76)	97 (91)	16 (39)	<0.001
Midazolam	87 (59)	74 (69)	13 (32)	<0.001
Dexmedetomidine	97 (66)	88 (82)	9 (22)	<0.001
Lorazepam	58 (39)	52 (49)	6 (15)	<0.001
Laboratories				
WBC (n=148) 4.0 – 10.0 (K/µL)	10.1 (7.8 – 13.9)	11.4 (8.4 – 15.2)	8.8 (6.4 – 10.9)	0.002
C-reactive protein (n=145) 1.0 – 3.0 (mg/L)	10.2 (5.1 – 18.0)	11.5 (5.33 – 20.7)	9.7 (6.5)	0.038
Ferritin (n=147)	1,208 (591 – 1,786)	1266 (710 – 1992)	1065 (490 – 1466)	0.124

18.0 – 320.0 (ng/mL)				
Lactate dehydrogenase (n=135) 120 – 240 (IU/L)	455 (328 – 572)	461 (351 – 636)	392 (248 – 507)	0.027
IL-6 (n=52) < 17.4 pg/mL	69 (28 – 201)	73 (30 – 194)	50 (22 – 232)	0.655
D-dimer (n=142) < 0.59 mg/L	3.1 (1.5 – 6.8)	3.63 (1.82 – 7.75)	1.71 (1.27 – 4.82)	0.003
Disposition, n (%)				0.021
Home (unassisted)	62 (42)	40 (37)	22 (54)	
Skilled care facility	47 (32)	41 (38)	6 (15)	
Death	39 (26)	13 (32)	26 (24)	

Institutional reference ranges are reported for laboratory values. *ICU* intensive care unit, *IQR* interquartile range, *WBC* white blood cell count, *IL-6* interleukin-6.

Table 4. Post-Discharge Neuropsychological Outcomes[†]

	All Patients (n=148)	Delirium (n=107)	No Delirium (n=41)
Positive FAM-CAM, n (%) (n=17)	4 (24)	4 (31)	0 (0)
PROMIS 4A Cognitive Abilities Score, median (IQR) (n=25)	16 (10 – 20)	17 (9 – 20)	14 (6)
Short Blessed Test – Normal, n (%) (n=22)	17 (74)	10 (67)	7 (100)
Short Blessed Test – questionable cognitive impairment, n (%) (n=22)	2 (8.7)	2 (13)	0 (0)
Short Blessed Test – cognitive impairment, n (%) (n=22)	3 (13)	3 (20)	0 (0)
PHQ-9 screen positive, n, (%) (n=25)	3 (12)	3 (17)	0 (0)

FAM-CAM Family-based Confusion Assessment Method for delirium, *PROMIS* Patient-Reported Outcomes Measurement Information System, *PHQ* Patient Health Questionnaire. †For each post-discharge survey, proportions are calculated based on the total numbers of surveys completed. In total, 25 surveys were completed for the PROMIS 4A test, 25 surveys were completed for the PHQ-9, 17 surveys were completed for the FAM-CAM, and 22 surveys were completed for the Short Blessed Test.

Reasons for not completing a test included the following: patient deceased, patient remains admitted to an inpatient facility, patient (or family member) refused survey participation, or patient was unable to be reached.

Figure 1

Figure 2

Figure 3

Figure 4