Using Capture-Recapture Methods to Estimate Influenza Hospitalization

Incidence Rates

Running title: Estimating Influenza Hospitalization Burden

Authors:

GK Balasubramani, PhD,[1] Mary Patricia Nowalk, PhD, RDN,[2] Lloyd G. Clarke, BS

Hons,[3] Klancie Dauer,[1] Fernanda Silveira, MD,[4] Donald B. Middleton, MD,[5] Mohamed

Yassin, MD,[6] Richard K. Zimmerman, MD, MPH, MA[2]

Affiliations:

[1]University of Pittsburgh Department of Epidemiology, Suite 600 Schenley Place, 4420

Bayard St., Pittsburgh, PA 15260 USA

[2]University of Pittsburgh Department of Family Medicine, Suite 520 Schenley Place,

4420 Bayard St., Pittsburgh, PA 15260 USA

[3]UPMC Health System Department of Pharmacy, Division of Infectious Diseases/

Pharmacy Department – AMP 5[TH] Floor Falk Medical Building, 3601 Fifth Ave; Pittsburgh

PA USA

[4]University of Pittsburgh Department of Medicine, 3601 Fifth Avenue, Suite 5B,

Pittsburgh, PA 15213 USA

[5]Department of Medical Education, UPMC St. Margaret, 815 Freeport Rd., Pittsburgh

PA 15215

[6]Infection Control Department UPMC Mercy, 1400 Locust Street Suite 10550 B Building

Pittsburgh, PA, USA 15219

Corresponding author:

Mary Patricia Nowalk, PhD, RD

Department of Family Medicine

4420 Bayard St., Suite 520

Pittsburgh, PA 15260

tnowalk@pitt.edu

412-383-2355

Word counts:

Abstract – 231; Text – 2722; Tables – 3; Figures – 1; Supplemental Tables – 2

Funding:

This work was supported by the Centers for Disease Control and Prevention (CDC)

[5U01IP001035-02] and by National Institutes of Health (NIH) [UL1TR001857]. This

work represents the views of the authors and not the CDC or NIH. It is subject to CDC's

public access policy.

Conflict of Interest:

Drs. Nowalk and Balasubramani, and Mr. Lyons have grant funding from Merck & Co.,

Inc. for an unrelated project. Drs. Zimmerman, Nowalk and Balasubramani have grant

funding from Sanofi Pasteur. Dr. Silveira has grant funding from Shire, Ansun, Novartis

for unrelated projects. Dr. Middleton reports personal fees from Seqirus, grants and

personal fees from Pfizer, personal fees from Sanofi Pasteur. Dr. Yassin and Mr.

Clarke have no conflicts to report.

Abstract

Background: Accurate population estimates of disease incidence and burden are needed to set appropriate public health policy. The capture-recapture (C-R) method combines data from multiple sources to provide better estimates than is possible using single sources.

Methods: Data were derived from clinical virology test results and from an influenza vaccine effectiveness study from seasons 2016-2017 to 2018-2019. The Petersen C-R method was used to estimate the population size of influenza cases; these estimates were then used to calculate adult influenza hospitalization burden using a Centers for Disease Control and Prevention (CDC) multiplier method.

Results: Over all seasons, 343 influenza cases were reported in the clinical database and 313 in the research database. Fifty-nine cases (17%) reported in the clinical database were not captured in the research database, and 29 (9%) cases in the research database were not captured in the clinical database. Influenza hospitalizations were higher among vaccinated (58%) than the unvaccinated (35%) in the current season and were similar among unvaccinated (51%) and vaccinated (49%) in the previous year. Completeness of the influenza hospitalization capture was estimated to be 76%. The incidence rates for influenza hospitalizations varied by age and season and averaged 307-309 cases/100,000 adult population annually.

Conclusion: Using Capture-Recapture methods with more than one database, along with a multiplier method with adjustments improves the population estimates of influenza disease burden compared with relying on a single data source.

Introduction

Introduction

Policy makers and planners need accurate estimates of the incidence or

prevalence of diseases and health conditions to anticipate, prevent, and mitigate the

effects of those diseases. The decentralized nature of U.S. health care makes overall

population burden estimates difficult to calculate. Thus, policy makers must rely on

population sampling for estimates, leaving true population burden unknown. The

accuracy of population estimates depends largely upon the quality of sampling, which in

turn, is dependent upon many factors including consistency of the population being

sampled across all capture occasions [1], and affected individuals having contact with

the healthcare system to allow for enumeration. Cases of unreported disease are more

difficult to account for.

Statistical methods to improve population disease incidence and prevalence

detection include the capture-recapture (C-R) method. The Lincoln-Petersen (Petersen)

method was the earliest C-R method for estimating population size. It was developed

for field studies of animals in which only a sample of the population can be caught and

marked (captured). Frequently, animals are re-caught (recaptured) and this method

allows for recaptured animals to improve the population estimate [2]. In health-related

research, C-R uses the overlap of subjects from two or more data sources and log-

linear methods to more accurately estimate true population disease burden [3]. C-R has

the advantage of using both research and clinical databases to measure the same data,

thereby creating a fuller perspective; however, the best method to adjust for complex

denominators in urban areas with competing health systems is not straightforward

within C-R.

96    The Centers for Disease Control and Prevention (CDC) has developed methods

97 to estimate population influenza burden (L Kim, personal communication, 2020) that

98 account for some of the complexities of multi-center study design and the incomplete

99 nature of surveillance data of the Hospitalized Adult Influenza Vaccine Effectiveness

100 Network (HAIVEN) study. The current study used available data from a single health

101 system, HAIVEN methods and adjustments, and C-R calculations, to estimate adult

102 influenza hospitalization burden in Allegheny County in Southwestern Pennsylvania.

Methods

The study was approved by the University of Pittsburgh IRB and consists of a three-phase analytic plan. The phases are: (1) C-R to more accurately estimate influenza cases; (2) statistical analyses for population burden based on the CDC HAIVEN network methods (Lindsay Kim, MD, personal communication); and (3) an adjustment to this resultant population burden using C-R incidence estimates. The HAIVEN population burden methods may not account for the richness of the clinical virology data available in our particular locale. Our adjustment to the HAIVEN methods was intended to capitalize on both the robust nature of the HAIVEN methods and the richness of the local virology data.

Phase 1: Statistical analyses for C-R. Data used for this analysis were collected from two sources: 1) the local health system's clinical surveillance software system (Theradoc®), which extracts virology test results from the electronic medical record (EMR); and 2) research data from selected hospitals participating in the HAIVEN study.

An IRB-approved honest broker extracted a data list from Theradoc® of Allegheny County residents who received an inpatient clinical respiratory viral panel (RVP) test at two to five (depending upon the season) general acute care hospitals in the health system during the study period that included the 2015-2016 through 2018-2019 influenza seasons. This list also contained basic demographic data of race, sex and age and is henceforth called the "clinical" database. The clinical database was prepared for analysis by limiting it to data from hospitals in which research enrollments were taking place for each influenza season. For example, in 2015-2016, two hospitals were enrolling participants, whereas in 2018-2019 there were five participating hospitals. Secondly, data were limited to the periods during which research enrollments

were taking place. Thirdly, patients <18 years of age were eliminated. Finally, patients

were separated into influenza cases and non-cases. The "research" database was

derived from adults who were recruited from the hospitals during the 2015-2016 through

2018-2019 influenza seasons for the HAIVEN study that only included inpatients ≥18

years of age. Detailed study methods for the HAIVEN study have been described

elsewhere [4]. Briefly, patients aged ≥18 years admitted with an acute respiratory

infection (ARI) including cough or worsening symptoms of a respiratory illness

beginning within 10 days were enrolled. Patients who had been enrolled in the prior 14

days were ineligible. Following informed consent, study staff collected respiratory

specimens (nasal and throat swabs from patients) for influenza virus testing (including

virus type and subtype) by reverse-transcription polymerase chain reaction (RT-PCR),

or used results from a clinical RVP test, if available. Demographic data were obtained

from interview. Vaccination status was based on documented receipt of each year's

influenza vaccine from the local electronic health record and/or the Pennsylvania

Statewide Immunization Information System (PA-SIIS).

The following variables were used in the calculations:

M = number cases identified in the clinical database;

n = number of cases identified in the research database;

m = number of cases identified in both databases (matched);

N_1 = number of cases reported *only* in the clinical database;

N_2 = number of cases reported *only* in the research database;

X = number of cases missing/not captured in either database;

Summary statistics of the demographic and clinical characteristics were

determined for the patients found in the matched database. The number of observed

influenza cases in the two databases and the Petersen's C-R method were used to

estimate influenza incidence ($\widehat{N}$) [5].

Petersen estimate $(\widehat{N}) = \dfrac{(Observed\ \ clinical\ cases\ (M)) * (Observed\ research\ cases\ (n))}{(Observed\ cases\ from\ matched\ database\ (m))}$ Eq. (1)

The variance and 95% confidence intervals (CIs) were calculated for the C-R estimates

using the formulae:

Variance $(\widehat{N}) = \dfrac{M * n * N_1 * N_2}{m^3}$ Eq. (2)

95% CI $= \widehat{N} \pm 1.96 * \sqrt{\text{Variance}(\widehat{N})}$ Eq. (3)

The C-R calculations were made assuming that: 1) the population is closed, i.e., there

was no outmigration or loss to follow-up because the capture and recapture would have

usually occurred during the same hospitalization. Calculation of completeness of

reporting by the two sources of the C-R method is determined by calculating the number

of missing cases, X

$X = (N_1 * N_2)/m$ Eq. (4)

An example of a C-R estimate is shown in Supplemental Table S1.

Secondly, it is assumed that the populations are homogeneous i.e., each

hospitalized patient has the same and constant probability of being captured by any

combination of the databases. Thirdly, it is assumed that the clinical database and the

HAIVEN research database are independent of each other. That is, the population

estimate assumes that the probability of being captured by one source does not affect

the probability of being captured by the other source [12]. Independence can be tested

by calculating the probability of influenza positives being captured by both databases. If

that probability is equal to the product of the marginal probabilities of being influenza

positive captured by clinical and research databases, then the samples are

independent.

Independence equality: $\frac{m}{N} = \frac{M}{N} * \frac{n}{N}$ Eq. (5)

Independence was tested for the 3-year total samples and the 15 subpopulations

derived by stratifying on demographic factors (age, sex, race), influenza season,

vaccination status, and prior vaccination status (Supplemental Table S2).

Phase 2: Statistical Analyses for Population Burden based on HAIVEN Methods

The Pennsylvania Health Care Cost Containment Council (PHC4) provided data

for ARI-specific hospitalizations based on CDC ARI ICD codes in all county hospitals for

all four years of the study.

The HAIVEN methods to calculate disease burden estimates were used as

follows:

Influenza burden estimate $= \frac{Adjusted\ number\ of\ cases}{Adult\ population\ in\ Allegheny\ County}$ Eq. (6)

Proportion of all ARI cases in all county hospitals represented by the study-

specific hospitals was determined using quarterly data from PHC4.

r = cases identified through research enrollment (research cases) among Allegheny

County residents.

V_1 = number of ARI hospitalizations from county residents who are enrolled in the

research database

V_2 = number of ARI hospitalizations in study-specific hospitals during influenza months

among Allegheny County residents in PHC4 database.

V_3 = number of ARI hospitalizations in all county hospitals during the same time period

among Allegheny County residents in PHC4 database; rationale is that both V_2 and

V_3 should come from the same database.

V_4 = number of influenza cases in study-specific hospitals during research enrollment

period from clinical database

V_5 = number of influenza cases in study-specific hospitals over the entire year from

clinical database; rationale is that both V_4 and V_5 should come from the same

database.

Adjusted number of cases $=$

206
$$research\ cases\ (r) * \left(\frac{1}{Proportion\ of\ all\ ARI\ enrolled\ in\ research\ database\ \left(\frac{V1}{V2}\right)} \right) *$$

207
$$\left(\frac{1}{Proportion\ of\ all\ ARI\ in\ study-specific\ hospitals\ \left(\frac{V2}{V3}\right)} \right) *$$

208
$$\left(\frac{1}{Proportion\ of\ all\ influenza\ cases\ detected\ during\ research\ enrollment\ period\ \left(\frac{V4}{V5}\right)} \right) \quad \text{Eq. 7}$$

210 *Phase 3: Combination of C-R and HAIVEN methods for adjusted population burden*

211 To incorporate the C-R method into the HAIVEN methods to account for cases

212 estimated by C-R but not due to the enrollment fraction, the following modification of

213 equation 7 was used:

214 Adjusted number of cases $=$

215
$$research\ cases\ (r) * \left(\frac{C-R\ estimated\ cases\ (\widehat{N})}{Total\ Observed\ cases\ (m+N1+N2)} \right) *$$

$$\left(\frac{1}{Proportion\ of\ all\ ARI\ enrolled\ in\ research\ database\ \ \left(\frac{V1}{V2}\right)} \right) *$$

$$\left(\frac{1}{Proportion\ of\ all\ ARI\ in\ study-specific\ hospitals\ \left(\frac{V2}{V3}\right)} \right) *$$

$$\left(\frac{1}{Proportion\ of\ all\ influenza\ cases\ detected\ during\ research\ enrollment\ period\ \left(\frac{V4}{V5}\right)} \right) \quad \text{Eq. 8}$$

For influenza burden calculations by race, the county population used as the

denominator was the total adult population of the county multiplied by 0.78 for Whites

and 0.13 for Blacks, representing their relative proportions of the population. Data were

analyzed using SAS version 9.4 (SAS Institute, Cary, NC, USA).

Results

     The viral test result analytic databases are shown in Figure 1. The clinical

database consisted of 8,994 patients of whom 7,684 patients were unmatched; the

research database consisted of 2,154 patients of whom 844 were unmatched patients;

and 1,310 patients were found in both databases (matched). Demographic

characteristics of the patients found in the matched database are shown in Table 1.

The highest proportion of the group was 50-64 years old (34.3%) with less than one

quarter each among patients who were 65-74 and 75+ years old and one fifth who were

18-49 years old. The patients were predominantly white (64.7%), female (62.9%) and

vaccinated ≥14 days prior to illness onset (58.5%). Half of them had been vaccinated in

the previous season and 25.7% were influenza cases.

     Table 2 shows the observed cases among persons hospitalized with a cough

illness in the clinical, research and matched databases, and C-R estimated influenza

hospitalizations over all seasons, by season and by other factors.

     The HAIVEN population influenza burden estimates (using Equation 7) and the

HAIVEN + C-R (using Equation 8) population influenza burden estimates were

calculated using the values shown in Table 2. Over all three influenza seasons, the

average incidence rates for hospitalized influenza in the research hospitals were 307-

309/100,000 (HAIVEN and HAIVEN + C-R, respectively). The lowest rates over all

seasons were 17/100,000 (for both HAIVEN and HAIVEN + C-R) among 18-49-year-

olds. The highest seasonal rates for the entire adult population were 494-497/100,000

(HAIVEN and HAIVEN + C-R, respectively), in 2017-2018, an especially severe

influenza season. Although there were two to three times as many cases among

246   Whites as Blacks, the influenza burden per 100,000 population were 10%-30% higher

247   among Whites than Blacks.

Discussion

This study used capture-recapture methods to calculate influenza incidence

among hospitalized patients and then used two methods for population burden: CDC's

HAIVEN burden estimate methods and a combination of HAIVEN and C-R estimates.

C-R has been adapted from ecological studies and used in a wide array of health-

related studies including Alzheimer's disease, heart attack, HIV infection, gun injury,

pediatric disease surveillance, gastric cancer, and norovirus infections [6-11]. The

Petersen C-R method using two sources of data in this study is a special case of the

generalized C-R method for estimating burden using multiple data sources. In this

study, estimates were nearly identical between observed and estimated incidence. By

comparison, in a C-R study of norovirus cases, the combined databases yielded

incidence at a rate 2.5 times the level of the rate of the highest individual database [11].

The assumptions of the Petersen estimator allow for any amount of overlap in the

databases. We have shown in Supplemental Table S2 that the independence

assumption is met for the 3-year total and all sub-populations. With few exceptions,

each database's probability of being captured by the other is high, resulting in a

considerable overlap and high probability of being captured by both databases.

This study met or partially met four of the assumptions needed for confidence in

the reliability of the outcomes. 1) The populations could be considered closed, that is,

there was little chance of loss of cases due to outmigration or death. In a disease of

short incubation and duration such as influenza, loss to outmigration is minimal. 2) High

overlap of cases improves reliability of estimates because of low missed cases. In this

study, there was a large overlap of cases between the databases allowing for matching

of pairs of cases (same person, both databases). 3) Databases should be

homogeneous. The two databases in this study are believed to be homogeneous, in

that cases in both have a nearly equal probability of being identified. 4) The data

sources should be independent to prevent over- or underestimation of missing cases.

We have demonstrated their independence. Although they were identified using

different methods, the cases in the research database were all hospitalized in the same

hospitals from which the clinical database was drawn. This fact does not affect the

independence assumptions.

The question about cases not captured by each of the databases deserves

comment. The research databases were limited by funding for intensity and duration of

recruitment, by volunteer participation, by obtaining specimens later in the course of

illness, and by specimen collection and testing techniques. The research specimen,

obtained by trained research assistants with a mid-turbinate collection, used the CDC's

quantitative PCR. This test is slightly more sensitive than the clinical qualitative PCR,

which used a nasopharyngeal specimen obtained by clinical staff. Thus, a small

amount of discrepancy between the two databases was expected.

Because of our confidence in the hospitalization estimates, county-wide

incidence rates were calculated. Influenza hospitalization incidence was highest in

2017-2018, an influenza A/H3N2-dominated year and lowest in 2018-2019 a season in

which influenza A/H1N1 and A/H3N2 cases were approximately equal [12]. This finding

is not unexpected, given influenza A/H3N2's higher severity than influenza A/H1N1.

Using C-R methods, influenza hospitalization incidence estimates among children have

ranged from 240/100,000 in 2003-2004 [13] and 860/100,000 in 2004-2005 [14] to

89/100,000 for the 2009 A/H1N1 influenza pandemic [15].   Among adults, during the

2009 A/H1N1 pandemic, influenza related hospitalizations were 178/100,000 for adults

18-49 years and 76/100,000 for those ≥50 years of age [15].

For comparison, CDC estimates of influenza hospitalizations[16] and an average

U.S. adult population of 250 million were used to estimate influenza hospitalization

burden per 100,000 adult population. For 2016-2017 influenza burden was

≈200/100,000 compared with 286/100,000 for Allegheny County; for 2017-2018

influenza burden was ≈324/100,000 compared with 494-497/100,000 for Allegheny

County and for 2018-2019 influenza burden was ≈196/100,000 compared with 172-

174/100,000 for Allegheny County. These similarities support the strength of our

methods for calculating burden.

Strengths and Limitations

Some inaccuracy of burden estimates based on surveillance data would be

expected given its inherent weaknesses, compared with true population-based data.

Hence, adjustments were made to account for some of those weaknesses. The

methods were further enhanced by using C-R to estimate hospitalized cases. This study

is one of only a few capture-recapture papers for influenza hospitalization rates among

adults and the only one of which we are aware for these seasons. It was conducted in a

complex urban/suburban county with detailed PCR-based clinical virology and in a

health system with the majority of the market share of hospitalizations. It has the

advantages of comparing and combining the robust HAIVEN and the C-R methods.

Limitations include conducting the study in only one county, although county-

specific population burden is a key outcome. The sources of the positive or negative

dependence may lead to underestimation or over estimation of the population size.

Finally, funding and time constraints limited the duration of active surveillance during the

influenza season for the research database.

Conclusions

Influenza illness is associated with significant costs that include lost productivity

due to absenteeism and presenteeism, lost wages, and costs of medical care.

Understanding the burden of influenza hospitalization is important for policy makers to

allocate resources for the prevention and treatment of influenza. Petersen's Capture-

Recapture method, in combination with the CDC HAIVEN burden method improved

population estimates.

Acknowledgement

References

1. Hwang W-H, Huggins R: **An examination of the effect of heterogeneity on the estimation of population size using capture-recapture data. .** *Biometrika* 2005, **92:**229–233.

2. Nichols JD: **Capture-Recapture Models: Using marked animals to study population dynamics.** *BioScience* 1992, **42:**94-102.

3. Hook EB, Regal RR: **Recommendations for presentation and evaluation of capture-recapture estimates in epidemiology.** *Journal of clinical epidemiology* 1999, **52:**917-926; discussion 929-933.

4. Ferdinands JM, Gaglani M, Martin ET, Middleton D, Monto AS, Murthy K, Silveira FP, Talbot HK, Zimmerman R, Alyanak E: **Prevention of influenza hospitalization among adults in the United States, 2015–2016: results from the US Hospitalized Adult Influenza Vaccine Effectiveness Network (HAIVEN).** *The Journal of infectious diseases* 2018.

5. Ackman DM, Birkhead G, Flynn M: **Assessment of surveillance for meningococcal disease in New York State, 1991.** *American journal of epidemiology* 1996, **144:**78-82.

6. Waller M, Mishra GD, Dobson AJ: **Estimating the prevalence of dementia using multiple linked administrative health records and capture–recapture methodology.** *Emerging themes in epidemiology* 2017, **14:**3.

7. Razzak JA, Mawani M, Azam I, Robinson C, Talib U, Kadir MM: **Burden of out-of-hospital cardiac arrest in Karachi, Pakistan: Estimation through the capture-recapture method.** *J Pak Med Assoc* 2018, **68:**990-993.

8. Poorolajal J, Mohammadi Y, Farzinara F: **Using the capture-recapture method**

to estimate the human immunodeficiency virus-positive population.

Epidemiology and Health 2017, **39**.

9. Post LA, Balsen Z, Spano R, Vaca FE: **Bolstering gun injury surveillance**

accuracy using capture–recapture methods. *Journal of behavioral medicine*

2019, **42:**674-680.

10. Knowles RL, Smith A, Lynn R, Rahi JS: **Using multiple sources to improve**

and measure case ascertainment in surveillance studies: 20 years of the

British Paediatric Surveillance Unit. *Journal of public health* 2006, **28:**157-165.

11. Hardstaff J, Clough H, Harris J, Lowther J, Lees D, O'Brien S: **The use of**

capture-recapture methods to provide better estimates of the burden of

norovirus outbreaks from seafood in England, 2004–2011. *Epidemiology &*

Infection 2019, **147**.

12. **National and Regional Level Outpatient Illness and Viral Surveillance**

[http://gis.cdc.gov/grasp/fluview/fluportaldashboard.html]

13. Grijalva CG, Weinberg GA, Bennett NM, Staat MA, Craig AS, Dupont WD, Iwane

MK, Postema AS, Schaffner W, Edwards KM, Griffin MR: **Estimating the**

undetected burden of influenza hospitalizations in children. *Epidemiol Infect*

2007, **135:**951-958.

14. Grijalva CG, Craig AS, Dupont WD, Bridges CB, Schrag SJ, Iwane MK,

Schaffner W, Edwards KM, Griffin MR: **Estimating influenza hospitalizations**

among children. *Emerging infectious diseases* 2006, **12:**103-109.

15.   Jules A, Grijalva CG, Zhu Y, Talbot KH, Williams JV, Dupont WD, Edwards KM,

Schaffner W, Shay DK, Griffin MR: **Estimating age-specific influenza-related**

hospitalization rates during the pandemic (H1N1) 2009 in Davidson Co, TN.

Influenza Other Respir Viruses 2012, **6:**e63-71.

16.   **Disease burden of influenza** [https://www.cdc.gov/flu/about/burden/index.html.]

Table 1. Demographic characteristics of patients identified in the matched database (N=1,310)	
Variable	**n (%)**
Age group	
18-49 years	249 (19.0)
50-64 years	449 (34.3)
65-74 years	311 (23.7)
75+ years	301 (23.0)
Race	
White	848 (64.7)
Black	421 (32.1)
Other, unknown	41 (3.2)
Sex	
Female	824 (62.9)
Male	486 (37.1)
Season	
2016-2017	356 (27.2)
2017-2018	449 (34.3)
2018-2019	505 (38.5)
Vaccination Status	
Unvaccinated	447 (34.1)
Vaccinated ≥14 days prior to illness onset	766 (58.5)
Vaccinated < 14 days prior to illness onset	97 (7.4)

Prior year vaccination (total for all seasons)	
No	645 (49.2)
Yes	665 (50.8)
Influenza Status	
No	973 (74.3)
Yes	337 (25.7)

Table 2. Estimated population influenza hospitalizations using the capture-recapture method

| | Observed influenza cases | | | | C-R estimated influenza cases |
	Clinical (M)	Research (n)	Matched (m)	Total ($m+N_1+N_2$)	$\hat{N}\ (\widehat{95\%\ CI})$ *
3 year total	308	313	284	337	339 (336, 342)
Age group					
18-49 years	55	54	50	59	59 (58, 60)
50-64 years	94	95	85	104	105 (103, 107)
65-74 years	82	82	75	89	90 (88, 92)
75+	77	82	74	85	85 (84, 86)
Race[†]					
White	191	199	179	211	212 (210, 214)
Black	103	100	92	111	112 (110, 114)
Sex					
Female	191	195	175	211	213 (210, 216)
Male	117	118	109	126	127 (125, 129)
Season					
2016-2017	84	82	77	89	89 (88, 90)
2017-2018	139	138	127	150	151 (149, 153)
2018-2019	85	93	80	98	99 (97,101)
Vaccination Status[†]					
Unvaccinated	124	124	111	137	139 (136, 142)
Vaccinated	165	169	155	179	180 (178, 182)
Prior vaccination					
No	167	167	153	181	182 (180, 184)
Yes	141	146	131	156	157 (155, 159)

Note: N_1 = number of cases reported *only* in the clinical database; N_2 = number of cases reported *only* in the research database;

$^*\widehat{N} = (M * n)/m$

†Sum ≠ 337 due to missing data.

Table 3. Influenza hospitalization estimates per 100,000 Allegheny County adult population using CDC HAIVEN and capture-recapture methods

		Adjustment factors					Influenza Burden	
Variable	Influenza cases in study-specific hospitals among Allegheny County residents (source=research database)	ARI* cases in study-specific hospitals among Allegheny County residents (source=research database)	ARI* cases in study-specific hospitals during influenza months among Allegheny County residents (source=PHC4)	ARI* cases in all county hospitals during influenza months among Allegheny County residents (source=PHC4)	Influenza cases in study-specific hospitals during research enrollment period (source= clinical database)	Influenza cases in study-specific hospitals over the entire year (source= clinical database)	Cases/100K Allegheny County adult population (based on Eq. 7)	Cases/100K Allegheny County adult population (based on Eq.8 and using C-R estimates)
	(r)	(V_1)	(V_2)	(V_3)	(V_4)	(V_5)		
3-year total	385	2,154	13,863	46,450	1,345	1,487	921	927
Annual average							307	309

2016-2017								
Overall	107	665	4,439	16,171	289	316	286	286
18-49 years	18	125	656	1,950	63	65	29	29
50-64 years	27	235	1,154	3,851	58	66	51	51
65-74 years	31	153	1,016	3,458	59	68	81	81
75+ years	31	152	1,613	6,912	109	117	152	152
White	77	481	3,323	14,071	215	236	318	318
Black	26	166	1,082	2,066	68	75	276	276
2017-2018								
Overall	168	656	4,135	16,353	539	633	494	497
18-49 years	35	142	612	1,955	126	143	55	55
50-64 years	48	213	1,039	3,838	121	153	110	110
65-74 years	42	157	997	3,644	108	124	112	110

75+ years	43	144	1,487	6,916	184	213	240	240
White	103	398	3,051	13,899	361	426	546	552
Black	57	241	1,084	2,324	159	189	504	504
2018-2019								
Overall	110	833	5,289	13,926	517	538	192	194
18-49 years	17	175	828	1,725	159	164	17	17
50-64 years	47	335	1,481	3,360	154	161	49	49
65-74 years	26	188	1,233	3,299	104	109	48	48
75+ years	20	135	1,747	5,542	100	104	86	86
White	72	498	3,985	11,679	316	331	227	227
Black	34	299	1,294	1,921	181	188	175	175

*ARI=acute respiratory illness

Figure 1 Legend.   Flow chart for clinical and research databases in study-specific hospitals resulting in the final analytic database, including influenza status.