COVID-19's U.S. Temperature Response Profile

Richard T. Carson*
Department of Economics
University of California, San Diego

Samuel L. Carson
Laserfiche

Thayne Keegan Dye
James Hardie Building Products

Samuel A. Mayfield
Department of Economics
University of California, San Diego

Daniel C. Moyer
Computer Science & Artificial Intelligence Lab
Massachusetts Institute of Technology

Chu A. (Alex) Yu
Department of Economics
Wake Forest University

November 3, 2020

Significance Statement: COVID-19 transmission, in the form of daily state-level death counts and positive cases, ramps up and down with changes in maximum daily temperature. This relationship, known as COVID-19's temperature response profile, is reliably estimated by assembling a new dataset that largely repairs the sizable divergence between when COVID-19 deaths occur and when they are reported. Over time this effect is can be quite large and, as such, important to efforts to forecast and manage COVID-19.

Abstract: We estimate the U.S. temperature response curve for COVID-19 and show transmission is quite sensitive to temperature variation. This is despite summer outbreaks widely assumed to show otherwise. By largely replacing the death counts states report daily, with counts based on death certificate date, we build a week-ahead statistical forecasting model that explains most of the daily variation ($R^2 = 0.97$) and isolates COVID-19's temperature response profile ($p < 0.001$). These counts normalized at 31°C (U.S. mid-summer average) scale up nearly 160% at 5°C. Positive cases are more temperature sensitive; scaling up by almost 400% between 31°C and 5°C. Dynamic feedback amplifies these effects. There is a short window to get COVID-19 under control before cooler weather makes the task substantially more challenging.

Key Words: COVID-19, data alignment, forecasting, environmental epidemiology, temperature

*Correspondence to: rcarson@ucsd.edu

Funding: The University of California, San Diego provided partial financial support for the work reported on in this paper. No outside support was received.

Competing interests: The authors declare no competing interests. **Author contributions:**

Richard T. Carson: Conceptualization, Methodology, Software, Validation, Formal Analysis, Investigation, Data Curation, Writing – Original Draft, Visualization, Supervision, Project Administration. **Samuel L. Carson:** Investigation, Data Curation, Writing – Review and Editing. **Thayne Keegan Dye:** Software, Investigation, Data Curation, Writing – Review and Editing, Visualization. **Samuel A. Mayfield:** Software, Investigation, Data Curation, Writing – Review and Editing, Visualization. **Daniel C. Moyer:** Methodology, Investigation, Writing – Review and Editing. **Chu A. (Alex) Yu:** Methodology, Formal Analysis, Investigation, Writing – Review and Editing, Visualization.

Introduction

The question of whether COVID-19 exhibits a pronounced temperature response profile has garnered attention since the early days of the pandemic *(1,2,3)*. Kissler et al.'s examination of medium and long-term management of the pandemic assigns a prominent role to understanding how the temperature response profile (hereafter "TRP") of COVID-19 might influence the pandemic's progression in the United States *(3,4)*. This follows conventional wisdom regarding the strong seasonal weather-driven pattern of influenza *(5,6)*, which helped mask COVID-19's early U.S. ascent *(7)*. The recent rise in positive COVID-19 cases and related deaths across much of the United States has called into question whether COVID-19 transmission is adversely sensitive to summer weather. Nevertheless, some modeling groups such as the University of Washington's Institute for Health Metrics (IHME) have assumed COVID-19 activity will increase as temperatures decline and are now using information from the U.S. influenza monitoring network *(8)*, which was previously shown to be predictive of the pandemic's path last spring *(7)*, to help incorporate that effect.

Daily state reports often bear little resemblance to the actual number of COVID-19-related deaths that occurred on that day. The resulting temporal data misalignment creates a substantial impediment to recovering any relationships with a crucial dependence on event timing. By reconstructing the set of death counts states report daily, largely by substituting in retroactive corrections based on death certificate dates, we are able to reliably estimate the TRP for COVID-19 deaths.

We assemble over 2,500 state-level daily observations from April 16–July 15, 2020, after COVID-19 became well-established across the United States. We largely follow the literature in environmental economics on estimating pollution and temperature related impacts on a range of health and other outcomes (9). SARS-CoV-2 (the virus which causes COVID-19) being a novel virus makes full implementation of the standard approach infeasible, because multiple years of both spatially and temporally delineated data are not available. The specific issue that we cannot currently resolved is whether the TRP over a given temperature range is the same in both directions: the cooler to warmer direction reflected in our data set, and the warmer to cooler direction, occurring as the U.S. enters its fall and winter seasons. In what follows, it is also important to recognize that we estimate the joint, not separate, effect of any biological response by the virus and any behavioral response by the public.

Prior efforts

Early attempts to pin down COVID-19's temperature response profile (TRP) have proven elusive. The now well-accepted approach for estimating influenza's TRP is epitomized by Barreca and Shimshack *(6)*, which draws heavily on the modelling of climate impacts on human populations *(9,10,11)*. Under this approach, a panel data set of political entities, such as countries or their political subregions (states, counties, etc.), is assembled and the outcome of interest observed across a long time horizon (e.g., a 20-year period). The ability to employ fixed-effect indicator variables to correct for time-invariant differences between political jurisdictions, coupled with the ability to use short-run weather variability, provides statistical identification of a variety of impact response functions.

Prior research has three important limitations. First, cross-sectional data cannot statistically identify the desired function without making the implausible assumption that all possible confounding variables are adequately controlled for. Routinely updated time-series models slowly incorporate environmental conditions into their forecasts without ever isolating it. Short panel datasets, where the stimulus of interest has limited range (temperature, humidity, UV light, air pollution in each location), often lack the statistical power to pin down such response functions. Consequently, these estimated response functions

are often fragile in the sense that statistical significance is lost when time trends or demographic variables are added to models *(12,13,14,15)*.

Second, early work, often using Chinese or cross-country data, focused on the speed at which the pandemic ramped up in different locations, which used variants of derivative statistics like growth rates or R0 as the dependent variable *(16,17)*. These works point to temperatures in the 0-10°C range as being most conducive to spreading COVID-19, with a possible humidity effect *(17,18,19)*. Current interest is now focused on situations where COVID-19 is spatially well-seeded and its effective R0 can move up or down with actions like state reopening plans.

Third, the quality of reported COVID-19 statistics is often suspect, particularly from the early phase of the pandemic. We move past this period to an observational window where reporting has stabilized. However, this reveals a different problem: temporal mismatches between when an event (e.g., a COVID-19-related death) was reported and the weather variables potentially influencing that event *(11)*. When reported event dates significantly differ from actual event dates, the resulting measurement error can overwhelm standard sources of biological variation such as individual differences in incubation periods.

Correcting state-level COVID-19 statistics and why it matters

Our ability to isolate the TRP for COVID-19 related deaths stems largely from our reconstruction of state-level COVID-19 data. The most important of these is replacing the daily death counts initially reported by states with the actual daily counts based on death certificate date where possible. Second, if death certificate data is not available, we use the retroactively corrected data series that a number of states have produced. When available, this type of data generally rectifies many initial reporting errors. Third, we correct other implausible data reports such as implicit negative daily death counts and zero counts on one day followed by a clear double-count the following day using a consistent protocol. This effort is described in detail in Supplementary Information (SI) section on Data Preparation.

Figure 1 displays why repairing daily state-level COVID-19 death counts matters. Panel (A) shows the originally reported (COVID Tracking Project [CPT]) $CPTDailyDead_{it}$ daily death counts for (Florida) in blue with the "Actual" death counts by death certificate date overlaid in red. Actual curves follow the general shape predicted by epidemic models; curves based on the originally reported counts have clear day of the week patterns and large spikes, neither of which are predicted by biological models. Panel (B) shows the two implications. First, the confidence interval from fitting a simple quadratic trend model is dramatically larger for the $CPTDailyDead_{it}$ (i.e., the Reported data) than that fitting the same model using the Actuals. Second, the model based on the $CPTDailyDead_{it}$ is slow to pick up the sharp rise in deaths in Florida because the reported data is temporally misaligned, specifically originally reported counts temporally lag the actual counts. Correct temporal alignment of the death count data is also critical to being able to isolate COVID's temperature response profile. Figure 1, Panels (C) and (D) display results for Georgia which show how use of $CTPDailyDead_{it}$ can lead to both missing a downturn and an upturn, while supporting an incorrect slow steady progress story. Figure S1 provides similar graphs for Arizona and Texas.

Fig. 1. COVID-19 Deaths in Florida and Georgia. April 16-July 15. **(A)** Florida Actual vs. Reported. **(B)** Florida quadratic time trend forecasts and confidence intervals. **(C)** Georgia Actual vs. Reported. **(D)** Georgia quadratic time trend forecasts and confidence intervals.

Figure 2(A) shows the week ahead forecast from the University of Washington's Institute of Health Metrics and Evaluation (UW-IHME) plotted against our model's DailyDead$_{it}$ at the state-level for the entire United States over the three-month period we examine. This model explains a reasonable amount of the variance in the data reflected in the R^2 of .66 obtained by regressing the Actuals on their forecasts. Substituting in the week-ahead forecasts from one of the other heavily used forecasting models results in a similar overall impression.

Figure 2(B) plots the predicted DailyDead$_{it}$ using Eq. 1 which only uses information available a week earlier. This regression's R^2 of .97 is reflected in the very tight scatter plot.

The difference between Figure 2 (A) and (B) is striking. Notably it is not driven by our superior modelling ability, since our model is intentionally designed to be simple and to transparently isolate the TRP. Rather, the difference is driven by our use of dramatically higher-quality, almost textbook-like temporally aligned death counts. It suggests this pandemic's short run daily trajectory at the state-level is (now) predictable with high accuracy, if resources were devoted to ensuring that actual daily death counts were reported in

a timely enough manner for use by the modeling community, rather than trickling in over a period of months as they do now.

Fig. 2. DailyDead$_{it}$ vs. predicted values using information available 7 days earlier. U.S. States April 16-July 15. **(A)** UW-IHME forecasts (R^2 = .66). **(B)** Eq. 1 base model predictions (R^2 = .97).

Modelling approach

Our objective is to isolate COVID-19's TRPs with respect to its two visible indicators, daily death counts (DailyDead$_{it}$) and new positive test counts (NewPositives$_{it}$) at the state level. Any TRP specification should allow for the possibility of non-responsiveness (i.e., zero temperature dependence). TRPs should not incorporate fixed factors like state demographics, nor other factors associated with calendar date or a clear temporal profile. Rather, daily exogenous variation in temperature on any specific day should be the source of statistical information for identifying the TRP of interest. Intuitively, the TRP is being statistically identified by having days where the lagged count of the COVID-statistic of interest is approximately the same magnitude but a range of different temperatures is observed. The number of such comparable days is substantially increased by the introduction of controls for conditions that remain fixed across states and a flexible time trend. Statistical modeling issues revolve around functional forms and specification of relevant lag structure. The slow-moving systematic changes in temperature over time imply that, over short time horizons, temperature will not be the driving force behind DailyDead$_{it}$ and NewPositives$_{it}$. However, over longer time horizons, a virus's TRP can be a major factor in its long-run trajectory.

We focus on two directly observable COVID-19 statistics: daily deaths (DailyDead$_{it}$) and new (test-diagnosed) positive cases (NewPositives$_{it}$), indexed by state (i) and day (t). For each we build a simple week-ahead forecasting model, controlling for state-level fixed effects and including a quadratic time trend. We then examine how lagged (t-k) maximum daily temperature (MaxTemp$_{it-k}$) scales the model's predicted DailyDead$_{it}$ or NewPositives$_{it}$. Attention is restricted to conditions where MaxTemp$_{it}$ ≥ 5°C, since data below this level is sparse during our sample period and concentrated in a few sparsely populated states like Alaska and Montana. Our empirical estimates of the TRPs are normalized to 100 at 31°C (~88°F), the U.S. population-weighted average for the last week of our sample to aid interpretability.

The pandemic modelling community has largely concentrated on DailyDead$_{it}$, believing it to be a more reliable indicator of the COVID-19 infection pool than NewPositives$_{it}$ due to the large differences in testing regimes across states and time. We proceed in a similar manner, but also produce the TRP for NewPositives$_{it}$, conditioning on available testing information, since it is positive cases that are potentially directly influenced by temperature.

Model components

Our base DailyDead$_{it}$ statistical model (Eq. 1) is comprised of two multiplicative components. The first produces expected current period deaths as a function of past observed deaths at a fixed temperature. The second allows expected DailyDead$_{it}$ to (potentially) vary with past values of MaxTemp$_{it-k}$.

The terms inside the first component are the infection pool proxy, DailyDead$_{it-7}$, a set of state-level fixed effect indicators, StateIndicator$_i$ and a quadratic time trend in Days$_t$ (t=1, ..., 137; initialized to March 1 to aid interpretability). The StateIndicator$_i$ captures the influence of a wide range of variables which remain constant over the period examined, such as demographic composition, geographic links between locales that influence infections, and public health infrastructure. The time trend variables pick up the decline in the case fatality rate and the average effect over time of initial lockdowns, social distancing, and state re-openings. The first component is exponentiated to incorporate the restriction that expected DailyDead$_{it}$ should be positive if COVID-19 transmission is active anywhere in the set of connected units being examined. Commensurately, we use LogDailyDead$_{it-k}$ and LogDays$_t$ as regressors so this component can be interpreted as a log-log regression model with state-level fixed effects.

The second component is a logistic function scaling predicted DailyDead$_{it}$ up or down with MaxTemp$_{it-k}$. Deaths on any specific day are the result of infections propagated over an earlier period. We use LogMaxTemp$_{it-7}$ and LogMaxTemp$_{it-14}$ to roughly encompass the relevant period for temperature influencing current period deaths. Our panel data model specification uses an additive error term, which necessitates using nonlinear least squares to solve the model (*20,21*), but decouples conditional mean estimates from the estimated error component.

Formally, our base model for U.S. state *i* on day *t* is given by:

$$DailyDead_{it} = EXP(\Sigma_i StateIndicator_i + \alpha_1 LogDays_t + \alpha_2 LogDays_t^2 + \beta LogDailyDead_{it-7}) \ X$$

$$(1/(1 + EXP(LogMaxTemp_{it-7})^{\Upsilon 1}))*(1/(1 + EXP(LogMaxTemp_{it-14})^{\Upsilon 2})) + \varepsilon_{it}, \qquad (1)$$

where estimated coefficients for the set of StateIndicator$_i$, and the Greek letter parameters minimize the sum of the squared error term of the estimated ε_{it}. The NewPositives$_{it}$ model is similarly structured.

We investigate (Table S6) the sensitivity of our results to range of alternative specifications (e.g., different infection pool indicators, different temperature scaling functions, adding absolute humidity, relative humidity, ultraviolet radiation, the inclusion of shelter-in-place/reopening orders, lagged cumulative death counts, and use of the CTP death counts) and provide further discussion of modelling issues in the Supplementary Information (SI) section on Modeling Approach.

Data

Our analysis uses three main types of data:

1. COVID-19 statistics for state-level death counts, positive cases, and tests, using The COVID Tracking Project (CTP; covidtracking.com) as our base information source.
2. Temperature data from the U.S. National Weather Service Integrated Surface Database.
3. State-level indicators and time variables.

We undertook extensive repair of the COVID-19 data reported daily by states, particularly those involving death counts. A dominant feature of these data are the substantial lags between when many of these events occurred and when they are reported. It is not uncommon to see states include deaths that occurred several weeks prior in any given day's count.

Over half the U.S. states have made the number of COVID-related deaths publicly available by their death certificate dates in some manner (SI Data Preparation). A simple OLS regression of death counts by death certificate date on originally reported death counts for many of these states yields an R^2 of less than 0.5. Other states have made corrections to originally reported COVID-19 statistics. These updated counts tend to retroactively correct a myriad of reporting errors by states contained in the COVID Tracking Project's daily data snapshot. These issues range from failing to report any information on specific days, decisions on "probable" COVID-related deaths, and the resolution of duplicated death certificates. We use states' self-corrected counts when available.

In instances where neither of these two sources of information were available, we undertook a consistent set of data repair operations. These include: averaging across days with no reporting, prorating backwards in time large batches of deaths (or other statistics) reported on an arbitrary date when it is known those deaths occurred over a much longer period (typically from nursing homes), and making the minimum set of changes to effectively correct logical violations such as negative counts of new deaths, positive cases,

and tests. Details on our data repairs are contained in the SI Data Preparation section; we make this dataset available for use by other researchers, along with a line-by-line account of the corrections made.

For each state, weather variables are taken from the airport with the highest volume of commercial traffic. We focus on maximum daily temperature. Results with mean temperature are quite similar. State-level aggregation requires taking weather data from a single station. The measurement error induced by this compromise is likely to be less than one might think. Many states are small spatially, or have a single concentrated metropolitan area (e.g., Illinois). In spatially large states with large populations, most of the population often lives within reasonable proximity to the largest airport. Even in Texas, most people live along the corridor between Dallas (DFW is our representative airport) and Houston. As a result, over 60% of the American population lives within 300km of the representative airport for their state. Similarly, positive cases are also concentrated in major metropolitan areas near those airports. More generally, a single source of classical measurement error attenuates parameter estimates toward finding no effect.

Daily dead model results

Eq. 1 is estimated using non-linear least squares and has an R^2 of 0.97. All parameter estimates (provided in Table S1) using robust standard errors clustered at the state-level are significant at $p < 0.001$. Figure 2(B) displays the actual (corrected) $DailyDead_{it}$ versus the model's in-sample predicted values. States where death certificate date were available have considerably smaller prediction errors ($p < 0.001$, Model 2 in Table S6).

The model's quadratic time trend suggests $DailyDead_{it}$ has fallen over time; at a declining rate from mid-April through the end of May, and then almost flat (rising at a slow rate). $LogDailyDead_{it-7}$ is the dominant predictor and its coefficient estimate of 0.8686 (t=35.32) has a standard elasticity interpretation.

Figure 3 shows the estimated TRP implied by the parameter estimates for the two $MaxTemp_{it}$ lags in the logistic scaling function. The vertical axis represents expected $DailyDead_{it}$ at each temperature value when both lags are set to the same temperature and the TRP is normalized to 100 at 31°C. (Figure S9 show the corresponding TRP using the original CTP death counts.) Figure S2 provides the contour plot which allows the values of the two $MaxTemp_{it-k}$ lags to independently vary.

Fig. 3. COVID-19 daily dead temperature response profile. U.S. states: April 16-July 15. Temperatures for 7th & 14th lags set equal.

Differentiating Eq. 1 with respect to DailyDead$_{it-7}$ produces a measure of how the expected DailyDead$_{it}$ increases from a one death increase in DailyDead$_{it-7}$. Setting t and MaxTemp$_{it-k}$ to chosen values, yields [TRP*β*EXP(StateBase$_{it}$)]/(DailyDead$_{it-7}$)$^{(1-β)}$, where the TRP directly scales the elasticity parameter, β, on LogDailyDead$_{it-7}$ and StateBase$_{it}$ is the temporally vary sum of the fixed StateIndicator$_i$ and the quadratic time trend. As an example, for Georgia on July 15 (StateBase$_{it}$ = 3.9184, MaxTemp$_{it-7}$ = 29.4, MaxTemp$_{it-14}$ = 27.8 and the corresponding non-normalized TRP = 0.0368), changing DailyDead$_{it-7}$ from 27 to 28, is predicted to increase the expected DailyDead$_{it}$ in Georgia on July 15 by 1.0457. Figure S3(B) displays a variant of this calculation for individual states as DailyDead$_{it-7}$ increases from 0 to 1.

We simulate series of DailyDead$_{it}$ with both static and dynamic variants of our estimated base model (Eq 1). To do so, we take the DailyDead$_{it-7}$ from the last week of our sample period and MaxTemp$_{it}$ = 31°C as the initial values to propagate the simulation. We fix MaxTemp$_{it-7}$ and MaxTemp$_{it-14}$ at 31°C from day 1 to day 45 to mimic the rest of the summer period, then progressively decrease them by 0.2°C each day until 5°C is hit, which occurs on day 175, after which temperature is held constant.

The brown dashed line in Fig. 4 provides a stylized static [left vertical axis] representation of information contained in our base model (Eq. 1) using Georgia as an example since it starts out at just below our 31C° normalization point and in a cold year hits our 5C° endpoint. The two MaxTemp$_{it-k}$ change in tandem, with

all other variables fixed at their initial values. In this static response mode, expected DailyDead$_{it}$ increases through only one channel – the direct impact of lowering temperature.

The blue dashed curve in Figure 4 shows the dynamic counterpart [right vertical axis] that allows MaxTemp$_{it-k}$ to influence DailyDead$_{it}$, which is then used as the lagged model input for subsequent projections. In this dynamic mode, temperature affects expected DailyDead$_{it}$ through two channels: first, the direct effect of lagged temperature on daily death count, as shown in the static model; second, the indirect compounding effect of these temperature-driven daily death counts when the initial death count of each cycle of the simulation is set to a lag of these outputs. The indirect effect of maximum temperature is the dominant mechanism, accounting for more than 90% of the increase in the expected death count as maximum temperature falls from 31°C to 5°C. The inset in Fig. 4 plots the two curves under the same scale and conveys the magnitude of the differences between the pure static and dynamic responses, both of which are hard to achieve in practice. The static response requires continual reductions in effective contact rates that exactly offset the increase in transmission potential, in the precise sense that the current death count is always held to be equal to last week's death count. Observing the full dynamic effect would require both the absence of offsetting government actions and endogenous actions by the public such as increased social distancing and face mask adherence as COVID-19 activity quickly increased. A possible example of the dynamic feedback our model allows when this is not done is the early path of the pandemic through Northern Italy in late February and March 2020.

Fig. 4. Stylized static and dynamic TRPs for Georgia derived from Eq. 1 parameter estimates. The brown solid curve [left vertical axis] is stylized static TRP where temperature influences DailyDead$_{it}$ through changes to MaxTemp$_{it-k}$, with DailyDead$_{it-7}$ held constant. The blue dashed curve [right vertical axis] is stylized dynamic TRP under the same assumptions, except dynamic feedback is allowed in the form of temperature influencing subsequent DailyDead$_{it-7}$, so that temperature affects expected death count both directly and through lagged death counts.

StateIndicator$_i$ values for some geographically isolated states with small populations like Hawaii are small enough to suggest their COVID-19 death counts would be unsustainable in warm enough weather. This is not true, though, of most states, which is consistent with Baker et al.'s finding from examining earlier emergent viruses that warming weather is not enough by itself to stop their spread *(22)*.

The SI section on Alternative Specifications for Base Death Count Model describes additional analyses that (a) look at alternative temperature scaling functions, (b) substitute DailyDead$_{it-14}$ or NewPositives$_{it-7}$ for DailyDead$_{it-7}$, and (c) add the dates of state actions such as shelter-in-place orders as control variables. This work shows that our finding of a strong TRP for DailyDead$_{it}$ is quite robust. Some specifications suggest the TRP is flatter in the 10-20°C range but steeper in the 5-10°C range (Fig. S8).

New positive case model results

NewPositives$_{it}$ are modeled similarly to Eq. 1, substituting LogNewPositives$_{it-7}$ as the infection pool regressor. Testing information is required to interpret reported state-level positives cases. We use

LogNewTest$_{it}$, to control for current testing intensity, LogNewTest$_{it-7}$ (which allows for a past positivity rate interpretation), total tests administered per thousand lagged by one week (PerCapitaTests$_{it-7}$) to help control for prior testing intensity and an indicator variable for systematically lower reporting on Monday. We chose LogMaxTemp$_{it-7}$ for consistency with the death count model. For the other temperature lag, the 2nd lag fits best.

The resulting model's (Table S2) R^2 is 0.95. All regressors are significant at $p < 0.001$, except for some of the test related variables: LogNewTest$_{it-7}$ ($p=0.005$), PerCapitaTests$_{it-7}$ ($p=0.002$) and Monday ($p=0.015$). Estimated parameters for the two lagged temperature variables are considerably larger than their DailyDead$_{it}$ counterparts. The implied TRP for NewPositives$_{it}$ is displayed in Fig. 5, with the contour plot that allows the two temperature lags to vary independently is provided in Fig. S4.

Fig. 5. COVID-19: new positives temperature response profile. U.S. states: April 16-July 15. Temperatures for 2nd and 7th lags set equal.

Discussion

We show that DailyDead$_{it}$ predictably varies with changes in maximum daily temperature (Fig. 3). This relationship is considerably more pronounced for new positive cases (Fig. 5). Our TRPs are normalized to 100 at 31°C, which is near the maximum of U.S. summer temperatures. These two TRPs suggest current COVID-19 infection pools in many states need to be brought under control while high temperatures are helping to reduce the virus's transmission. Cooler temperatures with the progression of fall and winter will dramatically ramp up the number of new positive cases and the deaths that follow unless current

infection pools are dramatically reduced. Dynamic feedback between rising infection pool indicators and cooling temperatures (Fig. 4) suggests delay in responding to increased virus activity signs will result in rapid escalation. This is already being seen in the current spatial pattern of outbreaks and the ever-increasing medium-run death count forecasts (e.g., UW-IHME *(8)*). Investment in providing the pandemic modeling community with timely counts based on death certificate dates would allow them to deliver substantially more accurate and timely warnings of impending upturns.

Warming temperatures during the spring and summer actively aided efforts to reduce the spread of COVID-19 and may have contributed to a false sense of how effective those efforts were. Cooling temperatures going into the fall and winter will present a very different challenge. Figure 6 shows the average date over the past 30-years when each U.S. county enters the particularly dangerous 10°C to 5°C range for COVID-19. Effects of temperature amplification show up with a lag.

There has long been fear of facing COVID-19 during influenza's October to March season *(23)*. Our results suggest, that in addition to this concern, COVID-19 transmission will become increasingly more efficient at transmission than it was during the summer. Further, if the TRPs for deaths and positive cases behave like influenza *(6)*, both will continue to increase from 5°C until a few degrees below freezing.

Fig. 6. Expected date by U.S. county for entering 5-10°C range (30-year average).

References

1. National Academies of Sciences, Engineering, Medicine. Rapid expert consultation on SARS-CoV-2 survival in relation to temperature and humidity and potential for seasonality for the COVID-19 pandemic. (2020).

2. M. Kanzawa, H. Spindler, A. Anglemyer, and G. W. Rutherford, Will coronavirus disease 2019 become seasonal? *J. Infect. Dis* **222**, 719-721 (2020).

3. Trump, D.J. February 11, 2020 statement. https://www.factcheck.org/2020/02/will-the-new-coronavirus-go-away-in-april/, retrieved 30 May 2020.

3. S.M. Kissler, C. Tedijanto, E. Goldstein, Y. H. Grad, M. Lipsitch, Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science* **368**, 860-868 (2020).

4. C. J. Carlson, A. C. R. Gomez, S. Bansal, S. J. Ryan, Misconceptions about weather and seasonality must not misguide COVID-19 response. *Nat. Commun.* **11**, 1-4 (2020).

5. J. Shaman, E. Goldstein, M. Lipsitch, Absolute humidity and pandemic versus epidemic influenza. *Am. J. Epidemiol.* **173**, 127-135 (2011).

6. A.I. Barreca, J.P. Shimshack, Absolute humidity, temperature, and influenza mortality: 30 years of county-level evidence from the United States. *Am. J. Epidemiol.* **176**, S114-S122 (2012).

7. J.D. Silverman, N. Hupert, A. D. Washburne, Using influenza surveillance networks to estimate state-specific prevalence of SARS-CoV-2 in the United States. *Sci. Transl. Med.* **12**, (2020).

8. Institute for Health Metrics, University of Washington, http://www.healthdata.org/news-release/ ihme-models-show-second-wave-covid-19-beginning-september-15-us

9. Graff Zivin, J., M. Neidell, Environment, health, and human capital. *J. Econ. Lit.* **51**, 689-730 (2013).

10. M. Auffhammer, S. M. Hsiang, W. Schlenker, A. Sobel, Using weather data and climate model output in economic analyses of climate change. *Rev. Environ. Econ. Policy.* **7**, 181-198 (2013).

11. Hsiang, S. Climate econometrics. *Annu. Rev. Resour. Econ.* **8**, 43-75 (2020).

12. R. Xu, H. Rahmandad, M. Gupta, C. DiGennaro, N. Ghaffarzadegan, H. Amini, M. S. Jalali. The modest impact of weather and air Pollution on COVID-19 transmission. doi: http://dx.doi.org/10.2139/ssrn.3593879 (5 May 2020).

13. Á. Briz-Redón, Á. Serrano-Aroca, The effect of climate on the spread of the COVID-19 pandemic: A review of findings, and statistical and modelling techniques. *Progress in Physical Geography: Earth and Environment*, 0309133320946302 (2020).

14. P. Jüni, M. Rothenbühler, P. Bobos, K. E. Thorpe, B. R. da Costa, D. N. Fisman, A. S. Slutsky, D. Gesink, Impact of climate and public health interventions on the COVID-19 pandemic: a prospective cohort study. *Can. Med. Assoc. J.* **192**, E566-E573 (2020).

15. R.H.L. Pedrosa, The dynamics of COVID-19: weather, demographics and infection timeline. doi: https://doi.org/10.1101/2020.04.21.20074450 (10 May 2020).

16. T. Carleton, J. Cornetet, P. Huybers, K. Meng, J. Proctor, Evidence for Ultraviolet Radiation Decreasing COVID-19 Growth Rates: Global Estimates and Seasonal Implications. doi: http://dx.doi.org/10.2139/ssrn.3588601 (28 April 2020).

17. J. Wang, K. Tang, K. Feng, W. Lv, High Temperature and High Humidity Reduce the Transmission of COVID-19. doi: http://dx.doi.org/10.2139/ssrn.3551767 (9 March 2020).

18. M.M. Sajadi, P. Habibzadeh, A. Vintzileos, S. Shokouhi, F. Miralles-Wilhelm, A. Amoroso, Temperature, Humidity and Latitude Analysis to Predict Potential Spread and Seasonality for COVID-19. doi: http://dx.doi.org/10.2139/ssrn.3550308 (5 March 2020).

19. G.F. Ficetola, D. Rubolini, Climate affects global patterns of COVID-19 early outbreak dynamics. doi: https://doi.org/10.1101/2020.03.23.20040501 (27 March 2020).

20. S.M. Goldfeld, R. E. Quandt, The estimation of Cobb-Douglas type functions with multiplicative and additive errors. *Int. Econ. Rev.* **11**, 251-257 (1970).

21. J.M. Wooldridge, *Econometric analysis of cross section and panel data*. (MIT Press, Cambridge, MA, 2010).

22. R.E. Baker, W. Yang, G. A. Vecchi, C. J. E. Metcalf, B. T. Grenfell, Susceptible supply limits the role of climate in the early SARS-CoV-2 pandemic. *Science* **369**, 315-319 (2020).

23. U.S. Centers of Disease Control and Prevention. When is flu season? https://www.cdc.gov/flu/about/season/flu-season.htm, retrieved 30 May 2020.

Supporting Information For

COVID-19's U.S. Temperature Response Profile

Richard T. Carson*, Samuel L. Carson, Thayne Keegan Dye, Samuel A. Mayfield, Daniel C. Moyer, Chu A. (Alex) Yu

*Correspondence to: rcarson@ucsd.edu

Includes:

Data preparation

Construction of temperature, humidity and ultraviolet radiation data

Modeling approach

Interpretation of State-level fixed effects and time-related effects

Alternative specifications for base death count model

Constructing temperature response profiles (TRPs)

Further DailyDead$_{it}$ specifications involving weather, positives and cumulative dead

Univariate DailyDead$_{it}$ and MaxTemp$_{it-7}$ relationship

Death Count TRPs based on DailyDead vs. CTP originally reported death counts

Structure of Table S6

Data and code availability

Figs. S1-S9

Tables S1-S6

Additional references

Data Preparation

The data set used in this paper starts with the COVID-19 statistics reported by individual states as aggregated daily by The COVID Tracking Project (covidtracking.com). In preliminary work, we (and other modelling groups) found there was no plausible epidemiological model that would produce the wild up and down jumps in the state-level COVID-19 statistics reported daily. As a result, there is a major fork in the path of any modeling effort. Do you build a predictive model for the reported COVID-19 daily death count that policymakers see, where success is judged by minimizing the forecast error around those observable quantities? Or do you attempt to model the actual underlying process with an eye toward understanding key aspects of it? When the reported dependent variable systematically diverges from that being generated by the underlying process, these two approaches fundamentally diverge. For the first path, success, to some degree, comes in uncovering the administrative procedures influencing the divergence. The second path led us to undertake a major effort to rectify and repair reported COVID-19 statistics focused on recovering the temporal structure needed to provide TRP estimates.

The most important correction was replacing originally reported death counts with death counts by date of death as reflected on death certificates. Figures 1 and S1 illustrate the nature of the differences for between the original $CTPDailyDead_{it}$ and $DailyDead_{it}$ for four important states: Arizona, Florida, Georgia and Texas. The originally reported data results in substantially larger confidence intervals in predictive models and because it lags the actual death counts substantially reduce the ability to detect and respond changes in COVID-19 activity. We were able to make this correction for 26 states where deaths by death certificate information could be located. Our cutoff date of July 15 allowed for two and half months for states to make these death counts available, which should subsume most of the COVID-19 deaths in these states over our sample period. The states where we have been unable to obtain deaths by death certificate dates tend to be smaller and to have had relatively fewer COVID deaths. Typically, they did not face large peaks, which appear to be associated with increased testing delays. However, we have been unable to obtain this data from three large states – California, Illinois, and New York – and of the 20 observations (out of 4,567) with residuals from Eq. 1 whose absolute value is 50 or more, over half are from these three states.

The methods for collecting data backed by death certificates varied based on how the individual states chose to present them. Some states publish their counts based on death certificates in a downloadable format on their official COVID-19 website. Others publish them inside of longer reports in such a way that they can be hard to find, or present tables or graphs that needed to be extracted by hand into spreadsheets. Because there are no official rules for how to publish this information, even the charts themselves vary in structure and presentation. Commonality across states is largely dependent on the specific software vendor they are using for their public-facing COVID-19 dashboards. Data collection procedures sometimes required hunting through source code to find full datasets within, or by hovering cursors over each bar in a bar chart individually over the collection window to make them display the precise value represented. In each case we were able to verify via labeling or official statements that the data was compiled using verified death certificate dates. It's worth noting here that, while the CDC collects deaths by death certificate date (https://www.cdc.gov/nchs/nvss/vsrr/COVID19/index.htm), they only do so at the weekly level, which makes it unusable for the sort of modeling effort we have undertaken and, relative to the data we have obtained from individual states, suffers from even longer reporting lags.

For states where deaths by death certificate date were not available, we first sought to determine if an individual state, either on their COVID-19 reporting "dashboard" or in a downloadable file, had deaths by reported date. These datasets often contain substantial corrections to what a state originally reported on

a specific day. These include reports that missed covidtracking.com's daily reporting deadline, more accurate end-of-day tallies (e.g., late reporting counties/hospitals), correction of testing dumps that resulted in the appearance of testing spikes on certain days, the removal of duplicate death certificates, and the resolution of probable cases. When these differed from the COVID-19 death counts originally reported (COVIDTracking.com has a set of "snapshots" of the originally reported information) we used the revised state data. In some smaller states, we believe, but have been unable to fully verify, that these corrected datasets are "close" to deaths by death certificate date.

Two of these types of corrections turn out to be particularly important. First, when a state misses a reporting deadline, this is typically recorded as there having been no COVID-19 events (new deaths positives, or tests) on that date. Because covidtracking.com (and similar aggregator sites) operate off cumulative counts, the next day contains the events that happened during the previous day. Second, "probable" cases have often been treated differently across states and time. State-level resolution of this issue removes an extraneous source of variation. As with the data on death certificate dates, these corrections by states of their originally reported COVID-19 statistics were obtained through a variety of sources ranging from reading counts off interactive bar charts to downloadable *.csv files.

Next, we corrected two obvious problems with the data. First, occasional, exceptionally large spikes accompanied by auxiliary information (e.g., a news/twitter release by the state's department of health) noting that this spike was due to an accumulation of deaths over an extended time period, typically from one or more congregate living facilities, e.g., nursing homes and prisons. For these, the initial correction involved proportionately increasing death counts over the relevant period, with days where this would result in negative death counts not included in the reallocation. Second, we corrected reports of zero deaths on days surrounded by death counts that were sufficiently large that a zero-death count was highly unlikely. Such days are generally also characterized by a failure to report some other COVID-19 statistics (like new positive cases) and by an abnormally high death count on the following day (or two if it is a weekend). Our approach for these was to assume no reporting followed by double reporting so that the indicated correction is to average counts across the two and, in some instances, three days.

We also corrected data in situations where a state had "corrected" an earlier report, but where an entire rectified data series from the state was not available. A typical example is a state that initially reported all deaths except for those in the state's largest county, with the corrected version containing the death count for the whole state contained in a press/twitter release. A similar situation was when a state failed a logical consistency check by having the difference between the reported cumulative death counts on two consecutive days generate a negative daily death count. This typically occurs in a state with a small population and few COVID-19 deaths, which, without comment, reduces their cumulative death count by one or two. Here we "rollback" that correction to the closest date that no longer produces a negative daily death count.

Similar corrections have been made to daily positive case counts and new daily test counts. The major difference is that very few states have made available positive case counts by day of test administration (rather than the day the test result is reported). This means that few states publish datasets that can readily subsume the positive case count data from the COVID Tracking project. This is likely due to these types of information having different reporting standards. Specifically, a state eventually knows the actual date of COVID-related death, but this information is subject to reporting delays due to waiting for confirming test results or autopsies. Because the date of death is on the death certificate, obtaining actual death counts for all states is eventually feasible. However, while the lab doing the test knows the date of test administration, this information is often not shared with the state. States could require more

complete reporting, but it is unclear whether the past is capturable. Total test counts are even messier due to the common practice – particularly in the earlier part of our sample period – of reporting all newly returned positive test results daily but reporting negative results inconsistently or in batches. Some negative tests results (often by the state's lab) were reported along with the positives, while other negative tests (often those by private labs) were reported once a week. We performed rollbacks of antibody tests that for a while were mixed with diagnostic tests, but these are messy because information on the period over antibody tests were induced has rarely been disclosed. We have endeavored to average and prorate testing data where the nature of this practice could be reasonably inferred.

The influence of our data correct effort can quickly be gleaned from Table S5, which describes four simple autoregressive models with a constant term and 7^{th} death count lag. The first uses the original "Reported" dataset (covidtracking.com) as both the source of the dependent variable, $CTPDailyDead_{it}$ and the lagged death count. The second uses our corrected version as the "Actuals" for the dependent variable and uses the originally Reported for the lagged regressor. The third uses $CTPDailyDead_{it}$ as the dependent variable and the Actuals for the lagged regressor. The fourth uses Actuals for both. The parameter estimates for lagged deaths are similar in versions using the same lagged variable and substantively larger in the two versions using lagged actual counts. The R^2 starts at 0.69 for the Reported/Reported model, stays roughly the same (.68) if Actuals are predicted from $CTPDailyDead_{it-7}$, and increases somewhat further to 0.74 for the Reported/Actual combination. The Actual/Actual combination, though, has an R^2 of 0.91, which clearly illustrates that the large gain in explanatory power in the base Eq. (1) model in Table S1 comes mainly from our data repair and rectification effort. Note that the models in Table S5 do set the massive NJ (June 25) reported death count outlier of 1877 (Actual death count is 16) to missing, since it is so large many modeling groups have either dropped it or prorate it over earlier time periods. The R^2 of this Reported/Reported model using these observations falls to 0.42.

Temporal misalignment of $NewPositives_{it}$ has received more attention than $DailyDead_{it}$ because of the often-large gap in time between when a diagnostic test is administered and when the result is returned *(24)*. Indeed, this gap, coupled with different state testing regimes, has led most modeling groups to concentrate on predicting $DailyDead_{it}$. We have made substantial repairs to $NewPositives_{it}$ and $NewTests_{it}$ data by reference to corrected state reports, and prorated rollbacks of initially reported antibody tests. We have been much less successful in locating information on $NewPositives_{it}$ by date of test administration. However, temporal misalignment of the $NewPositives_{it-k}$ is somewhat less important than for deaths than it might first appear because there is a shorter window for positive-to-positive transmission than the positive to death transition and because test results for hospital patients, the persistent high positivity pool, are typically returned quickly. Further, even very noisy testing information can be helpful in serving as controls for variation in state-level testing behavior over time.

Construction of temperature, humidity and ultraviolet radiation data

Weather data for our main analysis are drawn from the National Centers for Environmental Information (NCEI) Integrated Surface Database (ISD), which report hourly temperature and humidity data for most airports in the world. For each state, weather variables are taken from the airport with the highest volume of commercial traffic, where the volume information is found in the Federal Aviation Administration's 2018 Commercial Service Enplanements report. Our key variable of interest is daily maximum temperature (MaxTemp). We also look at measures of humidity and ultraviolet radiation. Hourly relative humidity is calculated as a function of hourly observed temperature and dewpoint temperature. Under the assumption of ideal gas behavior, we calculate hourly absolute humidity as well (details can be found at https://www.hatchability.com/Vaisala.pdf). We then pick the highest readings within each 24-hour period as daily MaxTemp, MaxRelativeHumidity and MaxAbsoluteHumidity. Minimum daily temperature

is obtained by picking the lowest reading and the mean by averaging the hourly readings. Our measure of ultraviolet radiation is UV index, which provides a forecast of the expected risk of overexposure to UV radiation from the sun. UV index data at our representative airports is obtained from OpenWeather Ltd., which publishes daily UV index forecast calculated by National Weather Service.

To calculate the expected date by U.S. county for entering the 5-10°C range (Fig. 6), we use reanalysis data provided by PRISM Climate group (https://prism.oregonstate.edu/) which provide reliable weather data at a high spatial resolution of 4km by 4km for the contiguous U.S. We extract daily maximum temperature from 1990 to 2019 and count the average number of days it takes since Oct 1 for the daily maximum temperature to fall below 10°C.

Modelling approach

There are two competing modelling approaches to predicting future COVID-19 deaths and positive cases. The first builds on a standard SEIR model; the other, a production function approach. The first has a strong epidemiological conceptualization and is clearly better in the early phase of a pandemic when data is scarce. The second is largely agnostic as to the underlying structure of transmission but requires dramatically more data to offset this flexibility. We follow the second, with (Eq. 1) using a simple production function approach. Conceptually, there is an infection pool, the seeds planted, and various inputs ranging from a state's health care system to temperature, that influence the output, the number of deaths observed to today. We primarily use past death counts as the infection pool proxy, take into account state-level fixed effects (amalgamating effects due to constant factors such as demographic characteristics, fixed resources like health care and transportation networks, and average differences in factors including climate variables and mobility), and a simple quadratic time trend. We exponentiate the right-hand-side variables to effectively impose the restriction that under the conditions we observe, expected death counts must be positive.

The use of a multiplicative scaling function incorporates the logic that temperature by itself cannot generate changes in $DailyDead_{it}$. Because those dying at t became infected not on one day but rather over an extended period, some means of representing temperature in this setting is required. The two options, due to the strong correlation between closely adjacent $MaxTemp_{it-k}$, are a distributed lag structure that imposes structure on a set of $MaxTemp_{it-k}$, or parameterizing the scaling function as the product of individual scaling functions, each with a different $MaxTemp_{it-k}$. We find two lags – the 7^{th} and 14^{th} – are sufficient and reasonably consistent with what is known about the biology of the virus and its methods of attack.

Model results are reasonably robust to small shifts in temperature lag positions, with the following two caveats. First, the 7^{th} lag is a natural one to use; many people's lives follow a typical weekly pattern of contacts and activities, and administrative reporting procedures often have a day-of-the week pattern. Second, lags too far back are insignificant. Notably we use the 18th lag of $MaxTemp_{it}$ in the weekly model rather than its naturally shifted counterpart $LogMaxTemp_{it-21}$.

The most commonly used scaling function for our purposes is the logistic function $1/(1 + EXP(X)^Y)$, where X is the variable of interest, and Y is the single estimated parameter. This function converges to a constant as Y goes to zero. Use of $MaxTemp_{it}$ rather than $LogMaxTemp_{it-k}$ as the stimulus variable provides a function with a different curvature. Statistically, it provides a similar fit, largely because the corresponding shifts in the estimate of Y makes the two scaling functions reasonably similar after normalization.

We also consider another commonly used scaling function, $X/(X + \psi)$, where an estimate of $\psi > 0$ results in smaller values of X being scaled up more than large values of X and an estimate of $\psi < 0$ results in larger

values of X being scaled up more than smaller values of X, and an estimate of $\psi = 0$ results in a function exhibiting no temperature responsiveness. This function can be used with either $LogMaxTemp_{it-k}$ or $MaxTemp_{it-k}$, and both will approximate a reasonable range of weakly monotonic scaling functions. It is well behaved as long as the estimate of $-\psi$ is bounded away from $MIN(X)$, which appears not to be an issue in the situations we examine. Table S4 provides estimates for a set of competing specifications and their TRPs are displayed in Fig. S8.

Our first major empirical decision was to use the three-month period: April 16-July 15. This time window allows for the seeding of the virus (to different degrees) across the i=1, …, 51 U.S. states (including DC) over a three-month period. We effectively start tracking the observable outcomes, deaths, positive cases, and tests on April 9th, because we generally use a one-week lag of the COVID-19 statistics of interest. Our time variable, t=1, …, 137 denoted in days starts with 1 on March 1st, the approximate date individual states first started reporting COVID-19 statistics.

Our second major decision involved how to implement the core epidemiological concept of a pool of infected individuals who can potentially infect other individuals. This pool is dynamic; existing positives transition to being no longer infectious, while newly infected individuals enter the pool. The totality of the currently infected individuals is unobservable without universal administration at each point in time of a 100% accurate diagnostic test. The question, thus, is whether to use a lagged variant of the test diagnosed positives, $NewPositives_{it}$, which logically are part of the infection pool but not necessarily representative of it due to differential testing, or a lagged variant of $DailyDead_{it}$.

The lagged $DailyDead_{it}$ measure is one step removed but was previously assumed to always be observed *(25,26)*. This assumption is demonstrably false. In the early run of the pandemic many deaths failed to be classified as COVID-related, in part because some instances were thought to be influenza-related, while COVID's role in inducing cardiac and kidney failure was not widely recognized. We avoid many of these problems by only using $DailyDead_{it}$ data generated after the pandemic was well established.

Other problems with using lagged $DailyDead_{it}$ (or $NewPositives_{it}$) as the infection pool indicator (such as a state having a proportionately larger elderly or black population) are readily addressed using the standard statistical approach of including time-invariant state-level fixed effects.

One way the use of a lagged version $DailyDead_{it}$ vs. $NewPositives_{it}$ will vary is with respect to timing. Current positives are temporally closer to future positives or deaths than current deaths and can potentially pick up the virus spreading among healthy young adults. The shorter link between a lagged $NewPositives_{it}$ is enhanced by the short period over which a current positive can influence future COVID outcomes, which makes TRP estimation less sensitive to the choice of temperature lags.

Subject to the same measurement error, using $NewPositives_{it-7}$ should be preferred to $DailyDead_{it-7}$ as the infection pool indicator. These results are provided in Table S6 (Model 16). The main lesson is that the model using $LogNewPositives_{it-7}$ is a good predictor of $DailyDead_{it}$ ($R^2 = 0.94$), but not as good as $LogDailyDead_{it-7}$. This suggests that either the positives have more measurement error than our corrected version of death counts or that the $DailyDead_{it-7}$ is a better reflection of the current risk-adjusted infection pool than $NewPositives_{it-7}$. The other result worth noting is that with the infection pool indicator temporally advanced, the coefficient on $LogMaxTemp_{it-14}$, as expected, is no longer significant.

The parameter estimates for the base model (Table S1) can be used to provide an estimate of the minimum infection pool each state faces by setting $DailyDead_{it-7} = 0$ (from a statistical perspective), obtaining expected $DailyDead_{it}$, and dividing it by the CDC's point estimate of the infection fatality rate (0.0065) *(27)*. Setting the value of $Days_t$ equal to the end of the sample period, across states, the sum of

these minimum infection pools is 9853 cases. Fig. S3(A) provides a visual display of this information for the continental U.S. states.

Interpretation of State-level fixed effects and time-related effects

We calculate a variant of the state-level fixed effects from our main regression model (Table S1) by setting the time trend equal to the last day of our sample period, the lagged number of deaths equal to zero, and calculating the expected number of deaths for each state. Small isolated states like Hawaii generate estimates close to zero while the most populous states tend to generate estimates above 2. These estimates suggest the minimum infection pool in each state varies by a factor of approximately 20 (Fig. S3(A)). A regression (Table S3) of the minimum expected death count on a small set of state-level demographics taken from the U.S. Census Bureau (LogPopulation, %Black, %Hispanic, and %Age80+) explains 79% of the variation in these counts.

We do not want our TRP estimates to be confounded by other factors changing over time (ranging from state and locally mandated shelter-in-place orders, reopening plans, endogenous social distancing, propensity to wear face masks and changing fatality rates). Since we are agnostic as to the mechanism, the straightforward specification is a polynomial time trend, where we find a quadratic term justified. Higher order terms add little insight or predictive power. The quadratic trend for predicting DailyDead$_{it}$ falls sharply from mid-April to the end of May, after which it very slowly starts to turn up.

A model which adds (a) the number of days since a state issued a mandatory shelter-in-place order and (b) the number of days since a state started to formally reopen its economy can be found in Table S6 (Model 8). The coefficients on these two variables are small and insignificant. The two LogMaxTemp$_{it-k}$ parameters fell on average by less than 1%.

This might seem puzzling until it is recognized that the two sets of time-related variables are highly correlated. Dropping the quadratic trend (Model 9 in Table S6) provides a different picture because, while still statistically significant at conventional levels, it was substantially diminished. The earlier a state shelter in-place order was issued, the lower the predicted death count (p = 0.028), and the earlier a state started to reopen the higher the predicted death count (p < 0.001). We are reluctant to provide any substantive interpretation to this result. Papers that have tried to determine the role of state actions and the behavior they are intended to influence show the need to (a) extensively model both state and local orders, and (b) incorporate spatially disaggregated mobility data in order to identify the effects of these government mandates from endogenous social distancing by the public that often occurs before these actions *(28,29,30)*. The two temperature coefficients in this model fall on average by less than 20% and remain highly significant.

Alternative specifications for base death count model

Table S4 compares our base model to alternative specifications. These alternative specifications were chosen to look at the sensitivity of the implied TRP because they all have reasonably similar fits relative to our base model. This allows us to observe how robust the TRP is to a substantial shift in various modelling decisions that we made. The first specification replaces the LogMaxTemp$_{it-k}$ with their linear counterparts. On the surface, this seems like a decision about which of two different scales fits better, but because the models have estimated parameters in the scale function, it may be possible for the two different specifications to provide reasonably similar TRPs. The second uses a popular ratio scaling function LogMaxTemp$_{it-k}$/(LogMaxTemp$_{it-k}$ + α), where α is an estimated parameter. The third replaces LogMaxTemp$_{it-k}$ with MaxTemp$_{it-k}$ in this ratio scaling function. A potential issue with Eq. 1, is the possibility that LogMaxTemp$_{it-14}$ directly influences our main infection pool indicator LogDailyDead$_{it-7}$. Our next specification replaces our infection pool indicator with an alternative, LogDailyDead$_{it-14}$, so both

temperature variables are now clearly exogenous from the temporal perspective of the infection pool indicator.

The last is a weekly variant of the model Eq. 2 where the dependent variable is the sum of the daily death count of the next seven days with corresponding pushbacks of the lagged variables. This model averages out much of the daily variation and many types of administrative reporting practices. In forecasting the pandemic's progression, it has become common to use data aggregated to a weekly level in an effort to average out many of the administratively-induced reporting issues that our extensive reconstruction and repair of the daily death count data sought to alleviate. It is possible to estimate a variant of Eq. (1) that uses death count data aggregated into seven-day periods, $WeeklyDead_{it} = \Sigma_t DailyDead_{it}$, where the summation is over t=1 to t=7. This makes $WeeklyDead_{it-7}$ the sum of the 7th through 13th lags of $DailyDead_{it}$. Importantly, this aggregation does not reduce the number of observations because on each day, the weekly aggregation at t=1 adds a new observation $DailyDead_{i1}$ and drops $DailyDead_{i8}$. Lags of the $WeeklyDead_{it}$ variable can then be used in the standard way. One implication of this specification is that the temperature variables also need to shift backwards. For conceptual consistency, we use $LogMaxTemp_{it-14}$ in place of $LogMaxTemp_{it-7}$. Empirically, the model fits best with the second temperature lag being $LogMaxTemp_{t-18}$. The model fit is reasonably similar using the 13th lag and 19th lags, but beyond the 19th lag, the temperature variable becomes insignificant, suggesting temperature information farther back in time than this is not useful. The model we report is thus:

$$WeeklyDead_{it} = EXP(\Sigma_i StateIndicator_i + \alpha_1 LogDays_t + \alpha_2 LogDays_t^2 + \beta LogWeeklyDead_{it-7})*$$

$$(1/(1 + EXP(LogMaxTemp_{t-14})^{\Upsilon 1}))*(1/(1 + EXP(LogMaxTemp_{t-18})^{\Upsilon 2})) + \varepsilon_{it}. \quad (2)$$

The overall impression from Table S4 is the general stability of most of the common parameters for time and the infection pool. Some of the specifications offer insights into the latter variable. It is little influenced by whether the logistic or ratio scaling function was used, nor whether $LogMaxTemp_{it}$ or $MaxTemp_{it}$ was the stimulus variable. Using $LogDailyDead_{it-14}$ in place of $LogDailyDead_{it-7}$ results in an estimated TRP being similar to the base model (Fig. S6), although it does rise more sharply between 10°C and 5°C suggesting, while there may be a endogeneity effect, it is not large and that our base TRP is conservative. In this model, the coefficient on the infection pool indicator shifts from the .86 of the base model to .80; in the weekly version model it jumps to .92, The R^2 for the weekly model increases to 0.99 (from the base model's 0.97) and falls to 0.94 in the model using $LogDailyDead_{it-14}$, which is less informative than $LogDailyDead_{it-7}$ in term of the infection pool influencing current death counts .

Constructing temperature response profiles (TRPs)

To compare the TRP's implied by the models in Table S4, we plot the functions using two independent random uniform variables RTemp and RTemp2, defined over the range 5°C and 40°C. The logistic scaling function with two temperature variables, $LogMaxTemp_{it-7}$ and $LogMaxTemp_{it-14}$, and corresponding Eq. 1 estimated parameters $\Upsilon 1$ and $\Upsilon 2$ (Table S4), results in the following scaling function in the base model:

$$(1/(1 + exp(LogRTemp^{\Upsilon 1})))*(1/(1 + exp(LogRTemp2^{\Upsilon 2}))). \quad (3)$$

There is a fundamental indeterminacy in such a scaling function, in that multiplying the production function part of Eq. 1 by a constant will result in an offsetting change in the scaling function which maintains the same expected value for the dependent variable. Note that each of the two multiplicative components of Eq. 3 converges to .5 irrespective of temperature values as $\Upsilon 1$ and $\Upsilon 2$ become increasingly negative. Often logistic functions are normalized to lie between 0 and 1 by changing the "1" in the numerator to "2", but this is not needed with our normalization to 31°C, which solves the indeterminacy

from the perspective of comparing curves. This is done by calculating the value of the estimated scale function at 31°C:

$$(1 / (1 + \exp(\log(31)^{\gamma 1}))) * (1 / (1 + \exp(\log(31)^{\gamma 2}))). \tag{4}$$

Dividing Eq. 3 by Eq. 4 produces a function which equals 1 at 31°C. Multiplying this quantity by 100 produces a function which has a natural percentage interpretation and equals 100 at 31°C. Note that for forecasting purposes, the original scaling function parameters need to be used.

The ratio scaling function with estimated parameters α_1 and α_2 using RTemp and RTemp2 is:

$$(\text{LogRTemp}/ (\text{LogRTemp} + \alpha_1)) * (\text{LogRTemp2}/(\text{LogRTemp2} + \alpha_2)). \tag{5}$$

As α_1 and α_2 converge to zero, both multiplicative components of SI.1c converge to 1 irrespective of temperature values. The value of this function at 31°C can be calculated in a manner similar to that described for the logistic. Dividing Eq. 5 with the value of that function at 31°C and multiplying by 100 produces the desired TRP.

Figure S8 graphs the daily death TRPs from the model specifications in Table S4. The alternative specifications produce TRPs remarkably similar to that of our base model and tend to bracket it. The one systematic difference is that TRPs based on the ratio scaling functions tend to be flatter over 10°C to 20°C but rise more sharply in the 5°C to 10°C range.

Further DailyDead$_{it}$ specifications involving weather, positives and cumulative dead

The other weather variables which have received considerable attention are absolute humidity, relative humidity, and ultraviolet (UV) radiation. Details of construction are provided below in Data Preparation Section. The limitation with all these variables is that they are correlated with MaxTemp$_{it}$ (i.e., absolute humidity: 0.58, relative humidity: -0.21, UV: 0.79). Maximum absolute humidity has a reasonable size effect ($p < 0.001$) in the model where it replaces the corresponding MaxTemp$_{it}$ variables. However, the LogAbsoluteHumidity$_{it-k}$ parameter estimates are close to zero and are no longer significant when the two parallel logistic functions comprising the temperature scaling function are added (Model 10 in Table S6). Relative humidity has a marginally significant relationship with DailyDead$_{it}$ when MaxTemp$_{it}$ is not in the model and its effect is close to zero in a model with MaxTemp$_{it}$ (Model 11 in Table S6).

The situation with UV$_{it}$ for which (15) found support is more complex (Table S4 &Table S6, Models 12 and 13). The MaxTemp$_{it}$ lags are marginally better predictors in a head-to-head comparison. These two variables are strongly correlated. Adding UV$_{it}$ marginally decreases prediction errors, but the signs of high multicollinearity are obvious; the first MaxTemp$_{it}$ lag is still highly significant while the second is insignificant and one of the UV$_{it}$ lags is only marginally significant (Table S4). It is effectively impossible to disentangle the influence of LogMaxTemp$_{it}$ and UV$_{it}$. Their relationship is displayed in Fig. S5, which plots LogMaxTemp$_{it}$ and LogUV$_{it}$ for two states over our sample period: Georgia (Atlanta) and New York (New York City). We cast our results in terms of MaxTemp$_{it}$ rather than UV$_{it}$ because it is more widely reported and understood, without making any claim our work supports a joint versus singular causal mechanism.

The specification in Eq. (1) is cast in terms of DailyDead$_{it}$. If MaxTemp$_{it-k}$ influences COVID transmission via the link between positive cases, then we should be able to replace DailyDead$_{it-7}$ with NewPositives$_{it-7}$. Estimates for this model (Table S4) show LogNewPositives$_{it-7}$ being significant at $p < 0.001$ and the R^2 measure falling a bit. The coefficient on MaxTemp$_{it-14}$ is insignificant, which would be expected if

NewPositives$_{it-7}$ incorporates that information, while the coefficient on LogMaxTemp$_{it-7}$ is larger than in the base specification.

A different aspect of the COVID-19 death statistics that has not been incorporated into the model is lagged cumulative death count, TotalDead$_{it-k}$. If DailyDead$_{it-k}$ can be seen as a proxy for the infection pool influencing DailyDead$_{it}$, then TotalDead$_{it-k}$ (normalized by population) is proxy for the fraction of the population that no longer at risk in the sense of being either removed by death or recovered. Adding LogPCTotalDead$_{it-7}$ effectively makes the StateBase$_{it}$ dynamic in a potentially different way than the quadratic time trend by letting each state evolve according to its own pattern of deaths. Table S6 (Models 14 and 15) displays the results of this model with LogPCTotalDead$_{it-7}$ enter as (a) a second order polynomial and (b) a fourth order polynomial.

Three results are worth noting. First, the increase in explanatory power is small with the most noticeable changes being as expected in the StateIndicator$_i$ and a substantial reduction in importance of the overall quadratic time trend. Second, the LogMaxTemp$_{it-k}$ parameter estimates are similar to that of Eq. (1) suggesting that our TRP is robust to a substantial dynamic reparameterization of the model. Third, in the quadratic specification, DailyDead$_{it}$ is declining with as LogPCTotalDead$_{it-7}$ at a declining rate. In the fourth order specification, all the LogPCTotalDead$_{it-7}$ terms are insignificant (although jointly significant). Figure S6 displays, starting at .05, the two response functions for LogPCTotalDead$_{it-7}$. Like the quadratic time trend, they suggest a sharp drop in how DailyDead$_{it}$ is influence by DailyDead$_{it-7}$ as LogPCTotalDead$_{it-7}$ increases from low levels with the fourth order polynomial being flatter at high levels of LogPCTotalDead$_{it-7}$ than the quadratic. The influence of this factor is reasonably small by the time a state hits 10 deaths per 100,000, a condition that characterizes 80% of the states at the end of our sample period. Earlier, we noted one interpretation of the quadratic time trend was that medical care (and hence death rates) had improved sharply at first and then at a declining rate. This specification has a similar interpretation but suggests that some of that learning is state specific and related to its prior COVID-19 caseload. There is no indication that this rate of decline is accelerating even in the hard-hit Northeastern states where deaths per 100,000 can be as high as 175 (NJ), looking at the fourth order polynomial which should allow this feature to emerge if the data supports it. This suggests the magnitude of the fraction of the population previously infection is still too small to be a substantial factor in slowing transmission of the virus.

Univariate DailyDead$_{it}$ and MaxTemp$_{it-7}$ relationship

If we have succeeded in isolating the TRP through use of the set of StateIndicator$_i$ and a quadratic time trend, the simple regression of DailyDead$_{it}$ on lagged MaxTemp$_{it-7}$ without the state fixed effects and quadratic time trend should reveal a substantially different curve than our estimated TRP displayed in Fig. 4. Figure S7 displays this relationship using a robust LOWESS smoother (bandwidth 0.2) on MaxTemp$_{it-7}$. This curve is dramatically more sensitive to temperature between 10°C and 30°C. Below 10°C it drops, which was expected since states with temperatures near 5°C tend to be more isolated and smaller population-wise. The curve bends up near 35°C and beyond where most of the observations come from a few states with (relatively) high June and July death counts.

Pinning down the TRP further will require: (a) obtaining more data over a longer time horizon with more temperature variation and, in particular, the important -5°C to 5°C range, where there has been little U.S. experience since the virus became widely dispersed, (b) obtaining death certificate data information from the few remaining large states (California, Illinois and New York) where it is not yet available, since states with death certificate date reporting have prediction errors that are substantially smaller (p < 0.001) than those that don't (Table S6, Model 2), or (c) having high quality temporally aligned COVID-19 statistics at the county level, which would provide a better temperature match and dramatically increase sample size.

Death Count TRPs based on DailyDead vs. The COVID Tracking Project's originally reported death counts

What does a TRP based on The COVID Tracking Project's (CTP) originally reported death counts look like compared to our base (Eq. 1) Model 1 which uses DailyDead? To examine this issue, we estimate Model 26 (Table S6) a direct analogue of Model 1 that substitutes $CTPDailyDead_{it}$ and $CTPDailyDead_{it-7}$ for their DailyDead counterparts. Consistent with Model 18 (Table S6) and other models employing CTPDailyDead, we exclude the two observations that contain the large New Jersey outlier as either the dependent variable or regressor and further excluded twenty observations where $LogCTPDailyDead_{it-7}$ is undefined because $CTPDailyDead_{it-7}$ is negative.

There are some clear differences between Model 1 and Model 26. First, in Model 26 there is a large drop in the R^2 from 0.97 to 0.81 and the RMSE measure more than doubles. Second, the elasticity parameter on lagged LogCTPDailyDead is just over 60% as large as it is in the DailyDead version. This is the expected result from introducing substantial measurement error, but one that also has large implications for drawing inferences about the size of the infection pool or in undertaking any dynamic forecasting exercise. Third, the quadratic time trend, while still sizeable is substantially diminished in both magnitude and statistical significance. Fourth, there are differences between the TRPs implied by their temperature response parameters in Eq. (1).

The TRPs for Model 1 and Model 26 are plotted in Fig. S9. The TRPs for the CTPDailyDead variant effectively lies above its DailyDead counterpart. At 5°C, it predicts almost 80% more deaths than our base Model 1. This result may seem counterintuitive to the usual belief that measurement error induces the parameter estimate to be attenuated toward zero That, intuition, generally correct, helps explain the fall in coefficient when CTPDailyDeadit-7 rather than DailyDeadit-7 is used as the infection pool indicator. However, it does not say anything about the implication of introducing measurement error into another covariate in the model. It is easy to show that as the scale of the measurement error in the lagged death count increases, the magnitude of the temperature responsiveness parameters adjusts to incorporate covariance with $DailyDead_{it-k}$. All else held constant, this would increase the magnitude of the estimated temperature effect under classical measurement error, since the parameters on infection pool indicator and temperature variables should have the same sign. The situation here is more complicated because measurement error from using CTDailyDead also substantially reduces the ability to pin down the quadratic time trend. Thus, neither the sign nor magnitude of the difference between the temperature response parameters across comparable specifications is therefore known a priori. Our parameter estimates (Table S6) suggest an upward bias is likely in temperature effect estimates obtained from models similar to ours that (a) use reported death counts and (b) have specifications where the infection pool and temperature response variables are expected to share the same sign. Note though that in a simple OLS regression model, the infection pool and temperature variables should have opposite signs, resulting in an estimate of the temperature parameter that is biased in the direction of finding no effect.

Structure of Table S6

The parameter estimates and standard summary statistics for models discussed is this paper are provided in Table S6. It contains three columns for each model. The first column contains the variables included in the model, the second the parameter estimates, and the third the standard errors. After the parameter estimates, the model's R^2, root mean square error (RMSE), and the number of observations on which the model was fit are provided.

The order in which the models appear are:

1. The base model represented by Eq. 1 using $LogDays_t$, $LogDays_t^2$, and $LogDailyDead_{it-7}$ as the predictors along with a set of state-level indicator variables and using $LogMaxTemp_{it-7}$ and $LogMaxTemp_{it-14}$ in the temperature scaling function.

2. A model which regresses the squared residuals from (1) on the log of state population and an indicator variable, GOOD_DATE for $DailyDead_{it}$ representing death counts by death certificate date.

3. Base model using linear versions of the two temperature variables.

4. Base model using alternative ratio scaling function Temp/(Temp + α), where Temp is temperature variable and α is the estimated parameter.

5. Base model using the alternative ratio scaling function and linear versions of the $MaxTemp_{it-k}$.

6. Base model with $LogDailyDead_{it-14}$ substituted for $LogDailyDead_{it-7}$.

7. A weekly version of the base model (see Eq. 1) substituting $LogDailyDead_{it-14}$ for $LogDailyDead_{it-7}$ and $LogDailyDead_{it-18}$ for $LogDailyDead_{it-14}$.

8. A version of the base model that adds two state-level government policy variables, the log of the number of days since a shelter-in-place order was first issued ($LogDaysShelterInPlace_{it}$) and the log of the number of days since a state began to formally reopen its economy ($LogDaysReopen_{it}$).

9. A version of (8) that drops the quadratic time trend of the base model.

10. The base model adding parallel scaling functions using the $LogMaxAbsoluteHumidity_{it-7}$ and $LogMaxAbsoluteHumidity_{it-14}$.

11. The base model adding parallel scaling functions $LogMaxRelativeHumidity_{it-7}$ and $LogMaxRelativeHumidity_{it-7}$.

12. Base model adding additional parallel scaling functions for $LogUV_{it-7}$ and $LogUV_{it-14}$.

13. Base model substituting $LogUV_{it-k}$ for $LogMaxTemp_{it-k}$.

14. Base modeling adding a quadratic in terms of $LogTotalDead_{it-7}$.

15. Base modeling adding a 4th order polynomial $LogTotalDead_{it-7}$.

16. Base model substituting $LogNewPositives_{it-7}$ and testing variables in place of $LogDailyDead_{it-7}$.

17. The model used in the paper to predict $NewPositives_{it}$.

18. The AR(7) regression of the originally reported death counts from the COVID Tracking Project ($CPTDailyDead_{it}$) on the 7th lag of itself ($CPTDailyDead_{it-7}$) for Table S5.

19. The same as (18) but with $DailyDead_{it}$ regressed on the 7th lag of the $CTPDailyDead_{it}$.

20. The same as (18) but $CTPDailyDead_{it}$ regressed on the 7th lag of $DailyDead_{it}$.

21. The same as (18) but $DailyDead_{it}$ regressed on its 7th lag.

22. The $StateBase_{it}$ calculation for each state.

23. The model reported in Table S3 which predicts the state-level fixed-effect estimates obtained in (1) as a function of a small set of demographic variables.

24. State-level estimates of the minimum infection pool.

25. State-level change in expected $DailyDead_{it}$ when $DailyDead_{it-7}$ shifts from 0 to 1 on July 15 at 31°C.

26. Base model (1) using the uncorrected death count data.

Data and materials availability: The data used in this study are archived at https://github.com/xxx in the form of a Stata ".dta" file. An Excel version of this file was created using StatTransfer. The Stata "do" file creating the data set contains a line-by-line set of the changes made to the original CovidTracking.com data set, and the providence of those changes. Three additional Stata "do" files are available in this archive. The first contains the code (Stata 16.1) for the regression models reported in this paper. The second provides an example of how to estimate the static and dynamic temperature response profiles for an individual state using Georgia as an example. The third contains Stata code for creating the basic versions (fine labeling was done using Stata's graph editor) of the figures in this paper. We also provide a further Stata dataset and corresponding Excel files at the state level (using our representative airport) with maximum daily temperature readings from the last thirty years.

Fig. S1. COVID-19 deaths in Arizona and Texas. April 16-July 15. **(A)** Arizona Actual vs. Reported. **(B)** Arizona quadratic time trend forecasts and confidence intervals. **(C)** Texas Actual vs. Reported. and **(D)** Texas quadratic time trend forecasts and confidence intervals.

Fig. S2. DailyDead TRP contour plot representation.

(A) | Implied number of daily new infections when DailyDead$_{it-7}$ is zero.

Individuals Infected
- (250, 450]
- (200, 250]
- (150, 200]
- (100, 150]
- [20, 100]

(B) | Change in E[DailyDead$_{it}$] with DailyDead$_{it-7}$ changing from 0 to 1.

Change in Expected Deily Dead
- (2.0, 2.5]
- (1.5, 2.0]
- (1.0, 1.5]
- (0.5, 1.0]
- [0.0, 0.5]

Fig. S3. Visual representation of information contained in the parameter estimates for base model (Eq. 1). (A) displays the infection pool for each state in the continental U.S. implied if observed DailyDead$_{it-7}$ is set to zero on July 15 and the two lagged MaxTemp$_{it-k}$ are set to 31°C. **(B)** shows, under the same conditions, how the expected DailyDead$_{it}$ changes if DailyDead$_{it-7}$ moves from 0 to 1.

Fig. S4. NewPositives TRP contour plot representation.

Fig. S5. LogMaxTemp vs. LogUV. (A) displays Georgia. **(B)** displays New York.

Fig. S6. DailyDead$_{it}$ responsiveness to lagged per capita total dead.

Fig. S7. Bivariate relationship DailyDead & MaxTemp. U.S. states: April 16-July 15. Lowess smoother: .2 bandwidth.

Fig. S8. Alternative DailyDead TRPs. U.S. states: April 16-July 15.

Fig. S9. Death count TRPs based on DailyDead vs. CTP originally reported. U.S. states: April 16-July 15.

Table S1. Eq. (1) base model predicting DailyDead$_{it}$.

Variable	Coefficient	Robust S.E.	t-statistic	p-value
Alaska	24.2468	3.2668	7.42	0.000
Alabama	26.2830	3.2504	8.09	0.000
Arkansas	25.6412	3.2496	7.89	0.000
Arizona	26.6339	3.2277	8.25	0.000
California	26.5480	3.2434	8.19	0.000
Colorado	26.1724	3.2608	8.03	0.000
Connecticut	25.3270	3.2578	8.08	0.000
District of Columbia	26.0314	3.2699	7.96	0.000
Delaware	26.0525	3.2719	7.96	0.000
Florida	26.7896	3.2194	8.32	0.000
Georgia	26.4536	3.2545	8.13	0.000
Hawaii	23.8933	3.2676	7.31	0.000
Iowa	26.1666	3.2669	8.01	0.000
Idaho	25.3286	3.2630	7.76	0.000
Illinois	26.5186	3.2569	8.14	0.000
Indiana	26.3159	3.2619	8.07	0.000
Kansas	25.6968	3.2630	7.88	0.000
Kentucky	25.8271	3.2613	7.92	0.000
Louisiana	26.3165	3.2548	8.09	0.000
Massachusetts	26.4245	3.2557	8.12	0.000
Maryland	26.4151	3.2614	8.10	0.000
Maine	25.0056	3.2684	7.65	0.000
Michigan	26.2274	3.2515	8.07	0.000
Minnesota	26.3019	3.2628	8.06	0.000
Missouri	25.9973	3.2607	7.97	0.000
Mississippi	26.2978	3.2530	8.08	0.000
Montana	24.7679	3.2679	7.58	0.000
North Carolina	26.3227	3.2556	8.09	0.000
North Dakota	25.2256	3.2686	7.89	0.000
Nebraska	25.7297	3.2615	7.92	0.000
New Hampshire	25.8414	3.2635	7.92	0.000
New Jersey	26.3998	3.2477	8.13	0.000
New Mexico	26.1024	3.2650	7.99	0.000
Nevada	26.1382	3.2566	8.03	0.000
New York	26.2963	3.2417	8.11	0.000
Ohio	26.3468	3.2608	8.08	0.000
Oklahoma	25.8892	3.2644	7.93	0.000
Oregon	25.4513	3.2582	7.81	0.000
Pennsylvania	26.4789	3.2573	8.13	0.000
Rhode Island	25.1521	3.2683	8.00	0.000
South Carolina	26.4216	3.2433	8.15	0.000
South Dakota	25.4075	3.2665	7.78	0.000
Tennessee	26.0621	3.2686	8.02	0.000

Texas	26.8796	3.2213	8.34	0.000
Utah	25.7632	3.2536	7.92	0.000
Virginia	26.3402	3.2635	8.07	0.000
Vermont	24.3453	3.2683	7.45	0.000
Washington	26.0738	3.2581	8.00	0.000
Wisconsin	25.9419	3.2617	7.95	0.000
West Virginia	25.2368	3.2718	7.71	0.000
Wyoming	23.9943	3.2677	7.34	0.000
$LogDays_t$	-10.4430	1.5361	-6.80	0.000
$LogDays_t*LogDays_t$	1.1916	0.1799	6.30	0.000
$LogDailyDead_{it-7}$	0.8697	0.0246	35.32	0.000
$LogMaxTemp_{it-7}$	0.2988	0.0506	5.91	0.000
$LogMaxTemp_{it-14}$	0.3011	0.0545	5.52	0.000
R^2	0.9667			
Root MSE	9.3198			
Observations	4567			

Table S2. Model predicting NewPositives$_{it}$.

Variable	Coefficient	Robust S.E.	t-statistic	p-value
Alaska	44.3754	7.8220	5.67	0.000
Alabama	45.0719	7.7439	5.82	0.000
Arkansas	45.0296	7.7613	5.80	0.000
Arizona	45.4897	7.7481	5.87	0.000
California	45.2471	7.8135	5.79	0.000
Colorado	44.7237	7.7036	5.81	0.000
Connecticut	44.9710	7.8123	5.76	0.000
District of Columbia	44.8480	7.7886	5.76	0.000
Delaware	44.8834	7.7766	5.77	0.000
Florida	45.7236	7.8081	5.86	0.000
Georgia	45.2116	7.7504	5.83	0.000
Hawaii	43.5360	7.5748	5.75	0.000
Iowa	44.9210	7.7604	5.79	0.000
Idaho	44.4862	7.6268	5.83	0.000
Illinois	45.2043	7.8320	5.777	0.000
Indiana	44.7918	7.7378	5.79	0.000
Kansas	44.8676	7.6937	5.83	0.000
Kentucky	44.4355	7.7336	5.75	0.000
Louisiana	46.0003	7.9379	5.80	0.000
Massachusetts	45.0429	7.8516	5.74	0.000
Maryland	45.1639	7.8065	5.79	0.000
Maine	43.7347	7.6623	5.71	0.000
Michigan	44.6836	7.7889	5.74	0.000
Minnesota	44.8802	7.7795	5.77	0.000
Missouri	44.9420	7.7386	5.81	0.000
Mississippi	45.1696	7.7728	5.81	0.000
Montana	44.2548	7.6490	5.79	0.000
North Carolina	44.0362	7.549	5.81	0.000
North Dakota	44.4879	7.7784	5.72	0.000
Nebraska	44.8758	7.7590	5.78	0.000
New Hampshire	44.2714	7.7374	5.72	0.000
New Jersey	45.2459	7.8555	5.76	0.000
New Mexico	44.0833	7.8327	5.76	0.000
Nevada	45.1552	7.7471	5.83	0.000
New York	45.2907	7.8897	5.74	0.000
Ohio	44.6637	7.7159	5.79	0.000
Oklahoma	44.7409	7.7293	5.79	0.000
Oregon	43.9776	7.6495	5.75	0.000
Pennsylvania	44.8583	7.7437	5.79	0.000
Rhode Island	44.8480	7.8233	5.73	0.000
South Carolina	45.0818	7.7231	5.84	0.000
South Dakota	44.4405	7.7294	5.75	0.000
Tennessee	45.3233	7.8145	5.80	0.000

Texas	45.3508	7.7418	5.86	0.000
Utah	44.9916	7.7777	5.78	0.000
Virginia	45.0741	7.7745	5.80	0.000
Vermont	43.1094	7.6813	5.61	0.000
Washington	44.5409	7.7176	5.77	0.000
Wisconsin	44.7343	7.7540	5.77	0.000
West Virginia	44.1364	7.7090	5.73	0.000
Wyoming	43.7190	7.6626	5.71	0.000
$LogNewTests_{it}$	0.3855	0.0833	4.63	0.000
MONDAY	-0.0801	0.0329	-2.44	0.015
$LogDays_t$	-19.5476	3.7152	-5.62	0.000
$LogDays_t*LogDays_t$	2.3839	0.4551	5.24	0.000
$LogNewPositives_{it-7}$	0.5856	0.0469	12.48	0.000
$LogNewTests_{it-7}$	-0.1755	0.0584	-2.93	0.005
$PerCapitaTests_{it-7}$	-0.0096	0.0031	-3.13	0.002
$LogMaxTemp_{it-2}$	0.5435	0.1082	4.94	0.000
$LogMaxTemp_{it-7}$	0.6873	0.0460	14.94	0.000
R^2	0.9507			
Root MSE	303.503			
Observations	4592			

Table S3. Model predicting StateBase$_i$ taken from DailyDead$_{it}$ model.

Variable	Coefficient	Robust S.E.	t-statistic	p-value
Constant	-2.6082	0.2842	-9.18	0.000
LogPopulation	0.3652	0.0430	8.48	0.000
%Black	0.0197	0.0030	6.52	0.000
%Hispanic	0.0172	0.0049	3.54	0.001
%Age 80+	0.1163	0.0627	1.86	0.070
R^2	0.7938			
Root MSE	0.2845			
Observations	51			

Table S4. Alternative specifications for DailyDead$_{it}$ model. StateIndicator$_i$ for models provided in Table S6. (Robust standard errors clustered at the state level).

Variable	Eq. 1 Base Model	Eq. 3 Linear MaxTemp	Eq. 4 Ratio Scaler	Eq. 5 Ratio Scaler Linear MaxTemp	Eq. 6 14th Lag DailyDead	Eq. 7 Rolling Weekly	Eq. 12 Add UV
LogDays$_t$	-10.4430	-10.5842	-10.2210	-10.1283	-30.2125	-13.9640	-8.7000
	(1.5361)	(1.5407)	(1.5499)	(1.5545)	(2.3673)	(1.2216)	(1.3241)
LogDays$_t$*LogDays$_t$	1.1916	1.2093	1.1627	1.1524	3.4253	1.5895	1.0093
	(0.1799)	(0.1806)	(0.1810)	(0.1815)	(0.2850)	(0.1453)	(0.1565)
LogDailyDead$_{it-7}$	0.8697	0.8681	0.8738	0.8752			0.8585
	(0.0246)	(0.0248)	(0.0243)	(0.0243)			(0.0256)
LogDailyDead$_{it-14}$					0.8199		
					(0.0478)		
LogWeeklyDead$_{it-7}$						0.9280	
						(0.0373)	
LogMaxTemp$_{it-7}$	0.2988		-0.5907		0.1843		0.1364
	(0.0506)		(0.1019)		(0.1651)		(0.1326)
LogMaxTemp$_{it-14}$	0.3011		-0.5790		0.4681		0.2487
	(0.0545)		(0.0887)		(0.0726)		(0.0578)
MaxTemp$_{it-7}$		0.1131		-1.3414			
		(0.0190)		(0.2704)			
MaxTemp$_{it-14}$		0.1160		-1.3016			
		(0.0222)		(0.2224)			
LogMaxTemp$_{it-14}$						0.2215	
						(0.0454)	
LogMaxTemp$_{it-18}$						0.3292	
						(0.0523)	
							0.5213
							(0.2100)
							0.2962
							(0.1318)
R^2	0.9667	0.9667	0.9666	0.9666	0.9425	0.9903	0.9674
Root MSE	9.3198	9.3151	9.3322	9.3394	12.2483	37.8452	9.2235
Observations	4567	4567	4567	4567	4576	4552	4567

Table S5. AR(7) DailyDead$_{it}$ models using different Reported/Actual combinations. U.S. states: April 16-July 15. June 25 NJ death count (1877) set to missing in reported. (Robust standard errors clustered at state level).

	Eq. 18 Reported/Reported	Eq. 19 Actual/Reported	Eq. 20 Reported/Actual	Eq. 21 Actual/Actual
DailyDead$_{it-7}$	0.6986	0.6681	0.7360	0.7854
	(0.0069)	(0.0067)	(0.0064)	(0.0035)
Constant	5.0046	6.6079	3.4596	3.1725
	(0.4210)	(0.4096)	(0.3862)	(0.2138)
R^2	0.6893	0.6820	0.7428	0.9148
RMSE	26.545	25.828	24.154	13.371
Observations	4639	4639	4639	4639

Table S6. Parameter estimates including state-level fixed effects for all models.

Model 1

Variable	Coefficient	S.E.
State_AK	24.2468	3.2668
State_AL	26.2830	3.2504
State_AR	25.6412	3.2496
State_AZ	26.6399	3.2277
State_CA	26.5480	3.2434
State_CO	26.1724	3.2608
State_CT	26.3270	3.2578
State_DC	26.0314	3.2699
State_DE	26.0525	3.2719
State_FL	26.7896	3.2194
State_GA	26.4536	3.2545
State_HI	23.8933	3.2676
State_IA	26.1666	3.2669
State_ID	25.3286	3.2630
State_IL	26.5186	3.2569
State_IN	26.3159	3.2619
State_KS	25.6968	3.2630
State_KY	25.8271	3.2613
State_LA	26.3165	3.2548
State_MA	26.4245	3.2557
State_MD	26.4151	3.2614
State_ME	25.0056	3.2684
State_MI	26.2274	3.2515
State_MN	26.3019	3.2628
State_MO	25.9973	3.2607
State_MS	26.2978	3.2530
State_MT	24.7679	3.2679
State_NC	26.3227	3.2556
State_ND	25.2256	3.2686
State_NE	25.7297	3.2615
State_NH	25.8414	3.2635
State_NJ	26.3998	3.2477
State_NM	26.1024	3.2650
State_NV	26.1382	3.2566
State_NY	26.2963	3.2417
State_OH	26.3468	3.2608
State_OK	25.8892	3.2644
State_OR	25.4513	3.2582
State_PA	26.4789	3.2573
State_RI	26.1521	3.2683
State_SC	26.4216	3.2433
State_SD	25.4075	3.2665
State_TN	26.0621	3.2486
State_TX	26.8796	3.2213
State_UT	25.7632	3.2536
State_VA	26.3402	3.2635
State_VT	24.3453	3.2683
State_WA	26.0738	3.2581
State_WI	25.9419	3.2617
State_WV	25.2368	3.2718
State_WY	23.9943	3.2677
t0	-10.4430	1.5361
t2	1.1916	0.1799
b1	0.8697	0.0246
w1	0.2988	0.0506
w2	0.3011	0.0545
R-square	0.9667	
Root MSE	9.3198	
Observations	4567	

Model 2

Variable	Coefficient	S.E.
LogPopulation	110.6656	5.0218
Good_Dates	-58.9392	10.3053
_cons	-799.2542	40.5126
R-square	0.0965	
Root MSE	330.1528	
Observations	4567	

Model 3

Variable	Coefficient	S.E.
State_AK	24.5529	3.2749
State_AL	26.6085	3.2602
State_AR	25.9674	3.2590
State_AZ	26.9905	3.2380
State_CA	26.8647	3.2529
State_CO	26.4937	3.2708
State_CT	26.6431	3.2674
State_DC	26.3474	3.2795
State_DE	26.3659	3.2813
State_FL	27.1275	3.2294
State_GA	26.7781	3.2646
State_HI	24.2219	3.2771
State_IA	26.4856	3.2764
State_ID	25.6418	3.2723
State_IL	26.8400	3.2668
State_IN	26.6345	3.2716
State_KS	26.0204	3.2729
State_KY	26.1429	3.2709
State_LA	26.6510	3.2655
State_MA	26.7435	3.2653
State_MD	26.7333	3.2713
State_ME	25.3192	3.2773
State_MI	26.5469	3.2613
State_MN	26.6185	3.2723
State_MO	26.3193	3.2706
State_MS	26.6264	3.2628
State_MT	25.0807	3.2770
State_NC	26.6467	3.2654
State_ND	25.5440	3.2777
State_NE	26.0525	3.2711
State_NH	26.1593	3.2730
State_NJ	26.7177	3.2579
State_NM	26.4317	3.2749
State_NV	26.4738	3.2667
State_NY	26.6136	3.2519
State_OH	26.6642	3.2703
State_OK	26.2148	3.2743
State_OR	25.7649	3.2675
State_PA	26.7961	3.2673
State_RI	26.4658	3.2776
State_SC	26.7494	3.2530
State_SD	25.7246	3.2758
State_TN	26.3870	3.2582
State_TX	27.2139	3.2313
State_UT	26.0825	3.2629
State_VA	26.6573	3.2733
State_VT	24.6667	3.2772
State_WA	26.3857	3.2676
State_WI	26.2584	3.2708
State_WV	25.5517	3.2811
State_WY	24.3020	3.2761
t0	-10.5842	1.5407
t2	1.2093	0.1806
b1	0.8681	0.0248
k1	0.1131	0.0190
k2	0.1160	0.0222
R-square	0.9667	
Root MSE	9.3152	
Observations	4567	

Model 4		
Variable	Coefficient	S.E.
State_AK	20.1337	3.3174
State_AL	22.1346	3.2978
State_AR	21.4935	3.2976
State_AZ	22.4619	3.2750
State_CA	22.4087	3.2909
State_CO	22.0283	3.3069
State_CT	22.1941	3.3051
State_DC	21.8982	3.3169
State_DE	21.9243	3.3194
State_FL	22.6230	3.2668
State_GA	22.3049	3.3007
State_HI	19.7449	3.3147
State_IA	22.0278	3.3149
State_ID	21.2024	3.3106
State_IL	22.3739	3.3042
State_IN	22.1782	3.3090
State_KS	21.5547	3.3097
State_KY	21.6948	3.3089
State_LA	22.1540	3.2993
State_MA	22.2860	3.3037
State_MD	22.2768	3.3081
State_ME	20.8797	3.3174
State_MI	22.0856	3.2989
State_MN	22.1667	3.3107
State_MO	21.8541	3.3075
State_MS	22.1459	3.3003
State_MT	20.6421	3.3164
State_NC	22.1764	3.3026
State_ND	21.0901	3.3174
State_NE	21.5877	3.3092
State_NH	21.7067	3.3115
State_NJ	22.2613	3.2935
State_NM	21.9509	3.3120
State_NV	21.9790	3.3031
State_NY	22.1590	3.2876
State_OH	22.2103	3.3087
State_OK	21.7425	3.3109
State_OR	21.3224	3.3057
State_PA	22.3421	3.3038
State_RI	22.0239	3.3165
State_SC	22.2712	3.2909
State_SD	21.2743	3.3149
State_TN	21.9162	3.2961
State_TX	22.7186	3.2689
State_UT	21.6275	3.3019
State_VA	22.2043	3.3103
State_VT	20.2029	3.3185
State_WA	21.9464	3.3051
State_WI	21.8095	3.3107
State_WV	21.1075	3.3194
State_WY	19.8756	3.3181
t0	-10.2110	1.5499
t2	1.1627	0.1810
b1	0.8738	0.0243
a1	-0.5907	0.1019
a2	-0.5790	0.0887
R-square	0.9666	
Root MSE	9.3322	
Observations	4567	

Model 5		
Variable	Coefficient	S.E.
State_AK	20.2699	3.3231
State_AL	22.2581	3.3039
State_AR	21.6173	3.3036
State_AZ	22.5749	3.2812
State_CA	22.5354	3.2970
State_CO	22.1544	3.3129
State_CT	22.3229	3.3110
State_DC	22.0268	3.3228
State_DE	22.0545	3.3253
State_FL	22.7402	3.2728
State_GA	22.4284	3.3067
State_HI	19.8683	3.3209
State_IA	22.1551	3.3209
State_ID	21.3334	3.3165
State_IL	22.4993	3.3102
State_IN	22.3055	3.3149
State_KS	21.6802	3.3157
State_KY	21.8238	3.3147
State_LA	22.2725	3.3055
State_MA	22.4036	3.3096
State_MD	22.4036	3.3140
State_ME	21.0117	3.3232
State_MI	22.2125	3.3050
State_MN	22.2947	3.3166
State_MO	21.9793	3.3134
State_MS	22.2682	3.3064
State_MT	20.7733	3.3223
State_NC	22.3006	3.3086
State_ND	21.2190	3.3233
State_NE	21.7133	3.3151
State_NH	21.8348	3.3174
State_NJ	22.3879	3.2995
State_NM	22.0732	3.3181
State_NV	22.0985	3.3093
State_NY	22.2860	3.2936
State_OH	22.3383	3.3147
State_OK	21.8670	3.3169
State_OR	21.4524	3.3116
State_PA	22.4694	3.3097
State_RI	22.1545	3.3224
State_SC	22.3941	3.2969
State_SD	21.4031	3.3208
State_TN	22.0405	3.3021
State_TX	22.8377	3.2749
State_UT	21.7553	3.3078
State_VA	22.3319	3.3163
State_VT	20.3324	3.3245
State_WA	22.0770	3.3110
State_WI	21.9391	3.3166
State_WV	21.2375	3.3253
State_WY	20.0103	3.3239
t0	-10.1283	1.5545
t2	1.1524	0.1815
b1	0.8752	0.0243
j1	-1.3414	0.2704
j2	-1.3016	0.2224
R-square	0.9666	
Root MSE	9.3394	
Observations	4567	

Model 6		
Variable	Coefficient	S.E.
State_AK	68.0983	4.8708
State_AL	70.0627	4.8869
State_AR	69.3268	4.8655
State_AZ	70.5461	4.8661
State_CA	70.4218	4.9037
State_CO	69.8248	4.9467
State_CT	70.1264	4.9560
State_DC	69.8024	4.9293
State_DE	69.8608	4.9324
State_FL	70.6838	4.8716
State_GA	70.1762	4.9259
State_HI	67.5865	4.9040
State_IA	69.8836	4.9124
State_ID	68.7233	4.9201
State_IL	70.3744	4.9345
State_IN	70.0431	4.9376
State_KS	69.1838	4.9171
State_KY	69.5242	4.9171
State_LA	69.9121	4.9466
State_MA	70.3122	4.9548
State_MD	70.2614	4.9348
State_ME	68.7766	4.9195
State_MI	69.8819	4.9552
State_MN	70.0764	4.9105
State_MO	69.7402	4.9144
State_MS	70.1096	4.9029
State_MT	68.5028	4.8888
State_NC	70.0974	4.9015
State_ND	68.8297	4.9008
State_NE	69.3169	4.8851
State_NH	69.4363	4.8960
State_NJ	70.1194	4.9633
State_NM	69.8534	4.9026
State_NV	69.7741	4.9211
State_NY	69.9672	4.9672
State_OH	70.1541	4.9305
State_OK	69.5545	4.9309
State_OR	69.2116	4.9088
State_PA	70.3375	4.9543
State_RI	69.9466	4.9329
State_SC	70.1774	4.8797
State_SD	69.0783	4.8834
State_TN	69.6894	4.8964
State_TX	70.8674	4.8732
State_UT	69.4429	4.8863
State_VA	70.1628	4.9326
State_VT	67.8890	4.9203
State_WA	69.4145	4.9331
State_WI	69.4335	4.9177
State_WV	68.9390	4.9132
State_WY	67.3690	4.8829
t0	-30.2125	2.3673
t2	3.4253	0.2850
b1	0.8199	0.0478
d1	0.1843	0.1651
d2	0.4681	0.0726
R-square	0.9425	
Root MSE	12.2483	
Observations	4567	

Model 7

Variable	Coefficient	S.E.
State_AK	32.9102	2.6235
State_AL	34.0669	2.7254
State_AR	33.8438	2.6948
State_AZ	34.2783	2.7475
State_CA	34.1784	2.7625
State_CO	33.8629	2.7479
State_CT	33.9715	2.7669
State_DC	33.9732	2.7261
State_DE	33.9674	2.7253
State_FL	34.3485	2.7639
State_GA	34.1048	2.7569
State_HI	32.9555	2.6727
State_IA	34.0259	2.7253
State_ID	33.3378	2.6796
State_IL	34.1271	2.7718
State_IN	34.0090	2.7508
State_KS	33.7194	2.7060
State_KY	33.8415	2.7138
State_LA	33.9520	2.7729
State_MA	34.0373	2.7750
State_MD	34.0977	2.7607
State_ME	33.4256	2.6726
State_MI	33.8339	2.7745
State_MN	34.0356	2.7370
State_MO	33.9426	2.7337
State_MS	34.0962	2.7307
State_MT	33.4258	2.6469
State_NC	34.0766	2.7367
State_ND	33.6488	2.6754
State_NE	33.7940	2.6973
State_NH	33.8766	2.7025
State_NJ	33.9433	2.7998
State_NM	34.0438	2.7222
State_NV	33.9905	2.7232
State_NY	33.8453	2.8111
State_OH	34.0392	2.7488
State_OK	33.7927	2.7157
State_OR	33.7039	2.6892
State_PA	34.0933	2.7813
State_RI	33.9590	2.7275
State_SC	34.1241	2.7177
State_SD	33.6997	2.6770
State_TN	33.9789	2.7113
State_TX	34.4291	2.7524
State_UT	33.9061	2.6837
State_VA	34.0636	2.7508
State_VT	33.0951	2.6638
State_WA	33.7650	2.7259
State_WI	33.7719	2.7117
State_WV	33.6192	2.6880
State_WY	33.1590	2.6433
t0	-13.9640	1.2216
t2	1.5895	0.1453
b1	0.9280	0.0373
h1	0.2215	0.0454
h2	0.3292	0.0523
R-square	0.9903	
Root MSE	37.8452	
Observations	4552	

Model 8

Variable	Coefficient	S.E.
State_AK	35.3125	4.9526
State_AL	37.5093	5.0265
State_AR	39.9268	6.7158
State_AZ	37.7874	4.9858
State_CA	37.5128	4.9174
State_CO	37.2238	4.9732
State_CT	37.3223	4.9562
State_DC	37.1079	5.0228
State_DE	36.9748	4.9447
State_FL	38.0022	5.0012
State_GA	37.7061	5.0454
State_HI	34.8410	4.9367
State_IA	40.4395	6.6952
State_ID	36.2677	4.9392
State_IL	37.5012	4.9429
State_IN	37.3513	4.9613
State_KS	36.7697	4.9970
State_KY	36.8543	4.9687
State_LA	37.2923	4.9540
State_MA	37.4709	4.9703
State_MD	37.5591	5.0236
State_ME	36.1441	5.0254
State_MI	37.2267	4.9576
State_MN	37.3694	4.9796
State_MO	37.2742	5.0676
State_MS	37.4815	5.0134
State_MT	35.7994	4.9576
State_NC	37.4502	5.0010
State_ND	39.5098	6.7049
State_NE	40.0185	6.7035
State_NH	36.8973	4.9704
State_NJ	37.3833	4.9410
State_NM	37.0723	4.9427
State_NV	37.2553	5.0116
State_NY	37.3281	4.9554
State_OH	37.3592	4.9516
State_OK	39.8591	6.5531
State_OR	36.3622	4.9248
State_PA	37.7060	5.0665
State_RI	37.2153	4.9853
State_SC	37.6689	5.0245
State_SD	39.7644	6.7388
State_TN	37.2144	4.9961
State_TX	38.0718	4.9929
State_UT	39.8857	6.6641
State_VA	37.4447	5.0106
State_VT	35.2688	4.9384
State_WA	36.9972	4.9326
State_WI	36.9340	4.9507
State_WV	36.1957	4.9344
State_WY	38.3688	6.7926
t0	-15.8165	2.4326
t2	1.6788	0.2216
s1	0.8278	0.5701
s2	-0.0520	0.0253
b1	0.8318	0.0247
w1	0.2903	0.0482
w2	0.2849	0.0495
R-square	0.9677	
Root MSE	9.1764	
Observations	4567	

Model 9

Variable	Coefficient	S.E.
State_AK	1.8320	0.3592
State_AL	3.6712	0.4368
State_AR	2.3075	0.1980
State_AZ	4.0871	0.5048
State_CA	3.9824	0.5123
State_CO	3.6027	0.4206
State_CT	3.7619	0.4606
State_DC	3.5902	0.3880
State_DE	3.6529	0.4065
State_FL	4.1604	0.5430
State_GA	3.7659	0.4377
State_HI	1.5444	0.3470
State_IA	2.7473	0.1963
State_ID	2.9995	0.3601
State_IL	3.9148	0.5004
State_IN	3.7293	0.4461
State_KS	3.2786	0.3736
State_KY	3.3538	0.4161
State_LA	3.7792	0.4412
State_MA	3.7945	0.4852
State_MD	3.8027	0.4474
State_ME	2.6020	0.3471
State_MI	3.6610	0.4664
State_MN	3.7560	0.4508
State_MO	3.3604	0.4222
State_MS	3.7118	0.4328
State_MT	2.3461	0.3493
State_NC	3.7650	0.4384
State_ND	1.9761	0.1599
State_NE	2.3871	0.1875
State_NH	3.3717	0.4212
State_NJ	3.7457	0.5071
State_NM	3.6508	0.4188
State_NV	3.6601	0.3946
State_NY	3.5950	0.5251
State_OH	3.7773	0.4583
State_OK	2.5978	0.1798
State_OR	3.1231	0.3872
State_PA	3.7882	0.4607
State_RI	3.6310	0.4157
State_SC	3.8521	0.4282
State_SD	2.0854	0.1686
State_TN	3.5423	0.4293
State_TX	4.2796	0.5281
State_UT	2.5229	0.1775
State_VA	3.7735	0.4327
State_VT	2.1092	0.3380
State_WA	3.6787	0.4133
State_WI	3.5068	0.4234
State_WV	2.8448	0.3730
State_WY	0.8432	0.1415
s1	-0.2059	0.0909
s2	0.0703	0.0183
b1	0.9662	0.0441
w1	0.2678	0.0544
w2	0.2130	0.0771
R-square	0.9622	
Root MSE	9.9251	
Observations	4567	

Model 10			Model 11			Model 12		
Variable	Coefficient	S.E.	Variable	Coefficient	S.E.	Variable	Coefficient	S.E.
State_AK	26.8779	3.2441	State_AK	26.9609	3.3219	State_AK	22.6847	2.8044
State_AL	28.9128	3.2286	State_AL	28.9971	3.3067	State_AL	25.1718	2.7852
State_AR	28.2708	3.2280	State_AR	28.3566	3.3060	State_AR	24.5079	2.7841
State_AZ	29.2751	3.2005	State_AZ	29.3271	3.2730	State_AZ	25.5209	2.7633
State_CA	29.1775	3.2217	State_CA	29.2619	3.2998	State_CA	25.4578	2.7795
State_CO	28.8056	3.2357	State_CO	28.8851	3.3155	State_CO	25.0437	2.7958
State_CT	28.9574	3.2353	State_CT	29.0422	3.3141	State_CT	25.1276	2.7915
State_DC	28.6617	3.2477	State_DC	28.7447	3.3246	State_DC	24.8483	2.8024
State_DE	28.6829	3.2497	State_DE	28.7667	3.3274	State_DE	24.8631	2.8045
State_FL	29.4184	3.1982	State_FL	29.5033	3.2764	State_FL	25.7442	2.7570
State_GA	29.0838	3.2322	State_GA	29.1664	3.3097	State_GA	25.3558	2.7893
State_HI	26.5230	3.2463	State_HI	26.6045	3.3212	State_HI	22.8186	2.8026
State_IA	28.7965	3.2449	State_IA	28.8814	3.3228	State_IA	24.9829	2.7995
State_ID	27.9617	3.2382	State_ID	28.0400	3.3156	State_ID	24.0953	2.7966
State_IL	29.1492	3.2343	State_IL	29.2323	3.3130	State_IL	25.3361	2.7902
State_IN	28.9459	3.2399	State_IN	29.0312	3.3180	State_IN	25.1426	2.7954
State_KS	28.3278	3.2402	State_KS	28.4099	3.3172	State_KS	24.5349	2.7971
State_KY	28.4573	3.2392	State_KY	28.5421	3.3170	State_KY	24.6568	2.7946
State_LA	28.9455	3.2337	State_LA	29.0298	3.3095	State_LA	25.2766	2.7917
State_MA	29.0544	3.2337	State_MA	29.1391	3.3117	State_MA	25.2375	2.7903
State_MD	29.0454	3.2391	State_MD	29.1288	3.3173	State_MD	25.2423	2.7944
State_ME	27.6359	3.2462	State_ME	27.7199	3.3232	State_ME	23.7763	2.8028
State_MI	28.8579	3.2290	State_MI	28.9423	3.3073	State_MI	25.0237	2.7851
State_MN	28.9328	3.2397	State_MN	29.0141	3.3178	State_MN	25.0833	2.7955
State_MO	28.6276	3.2382	State_MO	28.7101	3.3157	State_MO	24.8275	2.7937
State_MS	28.9273	3.2314	State_MS	29.0130	3.3094	State_MS	25.1949	2.7877
State_MT	27.4010	3.2435	State_MT	27.4784	3.3209	State_MT	23.5121	2.8019
State_NC	28.9526	3.2336	State_NC	29.0372	3.3116	State_NC	25.1977	2.7895
State_ND	27.8562	3.2459	State_ND	27.9402	3.3240	State_ND	23.9670	2.8017
State_NE	28.3603	3.2390	State_NE	28.4443	3.3174	State_NE	24.5407	2.7947
State_NH	28.4725	3.2400	State_NH	28.5546	3.3183	State_NH	24.6115	2.7965
State_NJ	29.0300	3.2255	State_NJ	29.1130	3.3032	State_NJ	25.2245	2.7820
State_NM	28.7391	3.2366	State_NM	28.7988	3.3131	State_NM	25.0030	2.7989
State_NV	28.7745	3.2286	State_NV	28.8261	3.2988	State_NV	24.9894	2.7899
State_NY	28.9253	3.2207	State_NY	29.0122	3.2984	State_NY	25.1382	2.7770
State_OH	28.9768	3.2388	State_OH	29.0620	3.3170	State_OH	25.1625	2.7942
State_OK	28.5192	3.2425	State_OK	28.6046	3.3198	State_OK	24.7573	2.7998
State_OR	28.0822	3.2355	State_OR	28.1656	3.3131	State_OR	24.1740	2.7918
State_PA	29.1095	3.2347	State_PA	29.1915	3.3127	State_PA	25.3055	2.7906
State_RI	28.7823	3.2463	State_RI	28.8678	3.3245	State_RI	24.9457	2.8015
State_SC	29.0509	3.2219	State_SC	29.1369	3.3002	State_SC	25.3066	2.7785
State_SD	28.0382	3.2439	State_SD	28.1215	3.3218	State_SD	24.1938	2.7990
State_TN	28.6923	3.2264	State_TN	28.7762	3.3044	State_TN	24.9281	2.7831
State_TX	29.5088	3.1998	State_TX	29.5947	3.2791	State_TX	25.7912	2.7588
State_UT	28.3967	3.2283	State_UT	28.4689	3.3040	State_UT	24.6060	2.7886
State_VA	28.9706	3.2412	State_VA	29.0544	3.3193	State_VA	25.1686	2.7964
State_VT	26.9756	3.2467	State_VT	27.0602	3.3232	State_VT	23.1031	2.8024
State_WA	28.7046	3.2355	State_WA	28.7882	3.3135	State_WA	24.7820	2.7913
State_WI	28.5722	3.2397	State_WI	28.6559	3.3170	State_WI	24.7357	2.7956
State_WV	27.8668	3.2499	State_WV	27.9526	3.3273	State_WV	24.0587	2.8047
State_WY	26.6265	3.2436	State_WY	26.7093	3.3234	State_WY	22.8521	2.8023
t0	-10.4435	1.5277	t0	-10.4301	1.5426	t0	-8.7000	1.3241
t2	1.1916	0.1790	t2	1.1901	0.1809	t2	1.0093	0.1565
b1	0.8698	0.0246	b1	0.8696	0.0248	b1	0.8585	0.0256
w1	0.3033	0.0680	w1	0.2950	0.0568	w1	0.1364	0.1326
w2	0.3054	0.0696	w2	0.2997	0.0579	w2	0.2487	0.0578
a1	-0.0079	0.0469	a1	0.0571	0.1556	a1	0.5213	0.2100
a2	-0.0087	0.0460	a2	0.0489	0.1600	a2	0.2962	0.1318
R-square	0.9667		R-square	0.9667		R-square	0.9674	
Root MSE	9.3217		Root MSE	9.3210		Root MSE	9.2236	
Observations	4567		Observations	4567		Observations	4567	

Model 13		
Variable	Coefficient	S.E.
State_AK	16.9355	3.4692
State_AL	19.5465	3.4812
State_AR	18.8769	3.4785
State_AZ	19.8707	3.4590
State_CA	19.8457	3.4740
State_CO	19.4279	3.4851
State_CT	19.4921	3.4733
State_DC	19.2066	3.4915
State_DE	19.2292	3.4927
State_FL	20.1358	3.4541
State_GA	19.7321	3.4849
State_HI	17.2231	3.5022
State_IA	19.3406	3.4896
State_ID	18.4468	3.4805
State_IL	19.6902	3.4773
State_IN	19.5082	3.4830
State_KS	18.8991	3.4878
State_KY	19.0255	3.4828
State_LA	19.6669	3.4911
State_MA	19.6161	3.4719
State_MD	19.5957	3.4835
State_ME	18.1559	3.4833
State_MI	19.3781	3.4644
State_MN	19.4252	3.4825
State_MO	19.1811	3.4843
State_MS	19.5727	3.4851
State_MT	17.8518	3.4861
State_NC	19.5672	3.4842
State_ND	18.3113	3.4863
State_NE	18.8928	3.4850
State_NH	18.9573	3.4832
State_NJ	19.5782	3.4643
State_NM	19.3860	3.4980
State_NV	19.3380	3.4839
State_NY	19.5042	3.4576
State_OH	19.5297	3.4802
State_OK	19.1450	3.4922
State_OR	18.4967	3.4722
State_PA	19.6612	3.4766
State_RI	19.3154	3.4861
State_SC	19.6827	3.4732
State_SD	18.5425	3.4874
State_TN	19.2969	3.4759
State_TX	20.1672	3.4534
State_UT	18.9864	3.4800
State_VA	19.5268	3.4854
State_VT	17.4780	3.4810
State_WA	19.0949	3.4701
State_WI	19.1108	3.4794
State_WV	18.4304	3.4940
State_WY	17.2778	3.4922
t0	-7.4256	1.6301
t2	0.8685	0.1917
b1	0.8611	0.0265
a1	0.6316	0.1190
a2	0.4650	0.0756
R-square	0.9669	
Root MSE	9.2939	
Observations	4567	

Model 14		
Variable	Coefficient	S.E.
State_AK	9.5460	9.9804
State_AL	11.9991	9.6295
State_AR	11.1770	9.7777
State_AZ	12.4550	9.5622
State_CA	12.2251	9.6924
State_CO	12.1578	9.4485
State_CT	12.6507	9.2511
State_DC	12.1266	9.3667
State_DE	12.0615	9.4223
State_FL	12.6138	9.5721
State_GA	12.3328	9.5209
State_HI	9.0987	10.0746
State_IA	11.8945	9.6341
State_ID	10.9464	9.7317
State_IL	12.4810	9.4842
State_IN	12.2584	9.4792
State_KS	11.3863	9.6608
State_KY	11.5100	9.6686
State_LA	12.5174	9.3058
State_MA	12.6781	9.3068
State_MD	12.3900	9.4628
State_ME	10.6601	9.6958
State_MI	12.4468	9.3113
State_MN	12.0540	9.6219
State_MO	11.6962	9.6647
State_MS	12.1683	9.5118
State_MT	10.0709	9.9861
State_NC	11.9679	9.7043
State_ND	10.8321	9.7155
State_NE	11.3572	9.7069
State_NH	11.6095	9.5981
State_NJ	12.9440	9.1431
State_NM	11.8357	9.6202
State_NV	11.9364	9.5677
State_NY	12.7671	9.1862
State_OH	12.1716	9.5718
State_OK	11.5981	9.6537
State_OR	10.9724	9.8202
State_PA	12.5207	9.4291
State_RI	12.2865	9.3524
State_SC	12.1198	9.6376
State_SD	10.9780	9.7599
State_TN	11.6459	9.7493
State_TX	12.5037	9.7180
State_UT	11.2319	9.8406
State_VA	12.1622	9.5705
State_VT	10.0472	9.5578
State_WA	11.9331	9.5323
State_WI	11.6491	9.6614
State_WV	10.7205	9.8812
State_WY	9.3713	9.9413
t0	-4.5443	3.9531
t2	0.5808	0.4116
c1	-0.5533	0.3046
c2	0.0326	0.0347
b1	0.8778	0.0274
w1	0.2925	0.0515
w2	0.2981	0.0458
R-square	0.9679	
RMSE	9.1536	
Observations	4567	

Model 15		
Variable	Coefficient	S.E.
State_AK	3.5952	8.7544
State_AL	5.8859	8.4632
State_AR	5.0183	8.5812
State_AZ	6.3802	8.4188
State_CA	6.0203	8.5219
State_CO	6.1577	8.2975
State_CT	6.8051	8.1319
State_DC	6.2628	8.2291
State_DE	6.1560	8.2770
State_FL	6.4954	8.4329
State_GA	6.2742	8.3679
State_HI	3.1812	8.8352
State_IA	5.8288	8.4582
State_ID	4.8809	8.5461
State_IL	6.4442	8.3412
State_IN	6.2457	8.3280
State_KS	5.2980	8.4810
State_KY	5.3820	8.4899
State_LA	6.6533	8.1816
State_MA	6.7704	8.1837
State_MD	6.3927	8.3180
State_ME	4.5793	8.5064
State_MI	6.5401	8.1854
State_MN	5.9745	8.4505
State_MO	5.5684	8.4866
State_MS	6.1664	8.3629
State_MT	4.0863	8.7600
State_NC	5.8276	8.5199
State_ND	4.7759	8.5172
State_NE	5.2583	8.5151
State_NH	5.5735	8.4313
State_NJ	7.0991	8.0396
State_NM	5.8012	8.4432
State_NV	5.8771	8.4047
State_NY	6.8951	8.0825
State_OH	6.0791	8.4095
State_OK	5.4982	8.4742
State_OR	4.8834	8.6184
State_PA	6.5106	8.2895
State_RI	6.4248	8.2167
State_SC	6.0074	8.4715
State_SD	4.8780	8.5500
State_TN	5.4874	8.5649
State_TX	6.2748	8.5507
State_UT	5.1468	8.6223
State_VA	6.0770	8.4067
State_VT	3.9987	8.3895
State_WA	5.8807	8.3737
State_WI	5.5418	8.4838
State_WV	4.6024	8.6206
State_WY	3.3305	8.6977
t0	-2.4685	3.3432
t2	0.3694	0.3463
c1	-0.5198	0.2720
c2	-0.0379	0.1269
c3	-0.0058	0.0500
c4	-0.0017	0.0066
b1	0.9300	0.0357
w1	0.2836	0.0619
w2	0.2764	0.0364
R-square	0.9691	
RMSE	8.9888	
Observations	4567	

Model 16

Variable	Coefficient	S.E.
State_AK	17.0018	5.5430
State_AL	18.7653	5.6419
State_AR	17.7498	5.6256
State_AZ	19.7100	5.6038
State_CA	19.9473	5.5696
State_CO	19.3527	5.6854
State_CT	19.6768	5.6324
State_DC	18.4803	5.6653
State_DE	18.4964	5.6645
State_FL	20.3097	5.5588
State_GA	19.4903	5.6243
State_HI	17.0232	5.6632
State_IA	18.5713	5.6617
State_ID	17.5102	5.6606
State_IL	19.9904	5.6222
State_IN	19.6328	5.6665
State_KS	17.9009	5.6789
State_KY	18.5673	5.6702
State_LA	18.8248	5.5636
State_MA	19.8956	5.5876
State_MD	19.4440	5.6400
State_ME	17.9484	5.6745
State_MI	20.1710	5.6255
State_MN	19.2714	5.6407
State_MO	19.0063	5.6191
State_MS	18.8587	5.6263
State_MT	17.7958	5.6067
State_NC	19.1364	5.6124
State_ND	17.2322	5.6265
State_NE	17.7791	5.6733
State_NH	18.6044	5.6911
State_NJ	20.1089	5.6170
State_NM	18.6012	5.6099
State_NV	18.5251	5.6166
State_NY	20.0549	5.5557
State_OH	19.7788	5.6608
State_OK	17.6217	5.6679
State_OR	17.9494	5.6623
State_PA	20.2618	5.6445
State_RI	18.5871	5.6076
State_SC	18.9771	5.6178
State_SD	17.2530	5.6742
State_TN	18.2918	5.5841
State_TX	20.2237	5.5971
State_UT	17.5312	5.5984
State_VA	19.2248	5.6467
State_VT	17.3072	5.6419
State_WA	18.8195	5.6074
State_WI	18.6382	5.6283
State_WV	17.7771	5.6332
State_WY	15.9743	5.6779
t0	-6.8242	2.6931
t2	0.5852	0.3178
n1	-0.3077	0.0844
pc1	0.5128	0.2432
bp	0.7638	0.0559
d3	0.2386	0.0588
R-square	0.9406	
Root MSE	12.4556	
Observations	4567	

Model 17

Variable	Coefficient	S.E.
State_AK	44.3754	7.8220
State_AL	45.0719	7.7439
State_AR	45.0296	7.7613
State_AZ	45.4897	7.7481
State_CA	45.2471	7.8135
State_CO	44.7237	7.7036
State_CT	44.9710	7.8123
State_DC	44.8480	7.7886
State_DE	44.8834	7.7766
State_FL	45.7236	7.8081
State_GA	45.2116	7.7504
State_HI	43.5360	7.5748
State_IA	44.9209	7.7604
State_ID	44.4862	7.6268
State_IL	45.2043	7.8320
State_IN	44.7918	7.7378
State_KS	44.8676	7.6937
State_KY	44.4355	7.7336
State_LA	46.0003	7.9379
State_MA	45.0429	7.8516
State_MD	45.1639	7.8065
State_ME	43.7347	7.6623
State_MI	44.6835	7.7889
State_MN	44.8802	7.7795
State_MO	44.9420	7.7386
State_MS	45.1696	7.7728
State_MT	44.2548	7.6490
State_NC	45.0362	7.7549
State_ND	44.4879	7.7784
State_NE	44.8758	7.7590
State_NH	44.2717	7.7343
State_NJ	45.2459	7.8555
State_NM	45.0833	7.8327
State_NV	45.1552	7.7471
State_NY	45.2907	7.8897
State_OH	44.6637	7.7159
State_OK	44.7409	7.7293
State_OR	43.9776	7.6495
State_PA	44.8583	7.7437
State_RI	44.8480	7.8233
State_SC	45.0818	7.7231
State_SD	44.4404	7.7294
State_TN	45.3233	7.8145
State_TX	45.3508	7.7418
State_UT	44.9916	7.7777
State_VA	45.0741	7.7745
State_VT	43.1094	7.6813
State_WA	44.5409	7.7176
State_WI	44.7343	7.7540
State_WV	44.1364	7.7090
State_WY	43.7190	7.6626
nt0	0.3855	0.0833
mt	-0.0801	0.0329
t1	-19.5476	3.7152
t2	2.3839	0.4551
nt7	-0.1655	0.0584
b1	0.5856	0.0469
pc1	-0.0096	0.0031
w1	0.5435	0.1101
w2	0.6873	0.0460
R-square	0.9507	
Root MSE	303.5030	
Observations	4592	

Model 18

Variable	Coefficient	S.E.
L7.ReportedD	0.6986	0.0069
_cons	5.0046	0.4210
R-square	0.6893	
Root MSE	26.5453	
Observations	4639	

Model 19

Variable	Coefficient	S.E.
L7.ReportedD	0.6681	0.0067
_cons	6.6079	0.4096
R-square	0.6820	
Root MSE	25.8280	
Observations	4639	

Model 20

Variable	Coefficient	S.E.
L7.DailyDead	0.7360	0.0064
_cons	3.4596	0.3862
R-square	0.7428	
Root MSE	24.1541	
Observations	4639	

Model 21

Variable	Coefficient	S.E.
L7.DailyDead	0.7854	0.0035
_cons	3.1725	0.2138
R-square	0.9148	
Root MSE	13.3709	
Observations	4639	

Model 22

Variable	Coefficient
State_AK	0.2006
State_AL	1.5370
State_AR	0.8090
State_AZ	2.1963
State_CA	2.0034
State_CO	1.3761
State_CT	1.6062
State_DC	1.1951
State_DE	1.2206
State_FL	2.5509
State_GA	1.8229
State_HI	0.1409
State_IA	1.3681
State_ID	0.5918
State_IL	1.9454
State_IN	1.5884
State_KS	0.8553
State_KY	0.9743
State_LA	1.5894
State_MA	1.7707
State_MD	1.7541
State_ME	0.4285
State_MI	1.4539
State_MN	1.5664
State_MO	1.1551
State_MS	1.5599
State_MT	0.3378
State_NC	1.5993
State_ND	0.5339
State_NE	0.8839
State_NH	0.9883
State_NJ	1.7275
State_NM	1.2831
State_NV	1.3298
State_NY	1.5576
State_OH	1.6383
State_OK	1.0367
State_OR	0.6691
State_PA	1.8697
State_RI	1.3484
State_SC	1.7655
State_SD	0.6404
State_TN	1.2324
State_TX	2.7912
State_UT	0.9140
State_VA	1.6275
State_VT	0.2214
State_WA	1.2469
State_WI	1.0928
State_WV	0.5399
State_WY	0.1559

Model 23

Variable	Coefficient	S.E.
LogPopulatio	0.3652	0.0430
PercentBlack	0.0197	0.0030
PercentHispa	0.0172	0.0049
PercentAge8	0.1163	0.0627
_cons	-2.6082	0.2842
R-square	0.7938	
Root MSE	0.2845	
Observations	51	

Model 24

Variable	Coefficient
State_AK	30.8645
State_AL	236.4667
State_AR	124.4626
State_AZ	337.8853
State_CA	308.2178
State_CO	211.7076
State_CT	247.1032
State_DC	183.8660
State_DE	187.7866
State_FL	392.4490
State_GA	280.4530
State_HI	21.6738
State_IA	210.4832
State_ID	91.0498
State_IL	299.2881
State_IN	244.3758
State_KS	131.5789
State_KY	149.8908
State_LA	244.5222
State_MA	272.4093
State_MD	269.8608
State_ME	65.9177
State_MI	223.6777
State_MN	240.9784
State_MO	177.7016
State_MS	239.9922
State_MT	51.9721
State_NC	246.0431
State_ND	82.1384
State_NE	135.9798
State_NH	152.0495
State_NJ	265.7632
State_NM	197.3949
State_NV	204.5898
State_NY	239.6327
State_OH	252.0446
State_OK	159.4940
State_OR	102.9358
State_PA	287.6393
State_RI	207.4532
State_SC	271.6207
State_SD	98.5245
State_TN	189.5980
State_TX	429.4077
State_UT	140.6124
State_VA	250.3867
State_VT	34.0594
State_WA	191.8292
State_WI	168.1248
State_WV	83.0635
State_WY	23.9773

Model 25

Variable	Coefficient
State_AK	0.1660
State_AL	1.2716
State_AR	0.6693
State_AZ	1.8169
State_CA	1.6574
State_CO	1.1384
State_CT	1.3288
State_DC	0.9887
State_DE	1.0098
State_FL	2.1103
State_GA	1.5081
State_HI	0.1165
State_IA	1.1318
State_ID	0.4896
State_IL	1.6094
State_IN	1.3141
State_KS	0.7075
State_KY	0.8060
State_LA	1.3149
State_MA	1.4648
State_MD	1.4511
State_ME	0.3545
State_MI	1.2028
State_MN	1.2958
State_MO	0.9556
State_MS	1.2905
State_MT	0.2795
State_NC	1.3231
State_ND	0.4417
State_NE	0.7312
State_NH	0.8176
State_NJ	1.4291
State_NM	1.0615
State_NV	1.1001
State_NY	1.2886
State_OH	1.3553
State_OK	0.8577
State_OR	0.5535
State_PA	1.5467
State_RI	1.1155
State_SC	1.4606
State_SD	0.5298
State_TN	1.0195
State_TX	2.3091
State_UT	0.7561
State_VA	1.3464
State_VT	0.1831
State_WA	1.0315
State_WI	0.9041
State_WV	0.4467
State_WY	0.1289

Model 26

Variable	Coefficient	S.E.
State_AK	19.1718	5.5455
State_AL	22.8740	5.5486
State_AR	22.0474	5.5568
State_AZ	23.5269	5.5377
State_CA	23.6248	5.5105
State_CO	22.8014	5.5331
State_CT	23.1423	5.5123
State_DC	22.3685	5.5664
State_DE	22.3263	5.5622
State_FL	23.5596	5.5434
State_GA	23.2117	5.5427
State_HI	19.7769	5.6115
State_IA	22.5782	5.5493
State_ID	20.8793	5.5795
State_IL	23.5277	5.5071
State_IN	23.0678	5.5314
State_KS	21.8978	5.5721
State_KY	22.2463	5.5515
State_LA	23.1644	5.5624
State_MA	23.3854	5.4903
State_MD	23.1839	5.5355
State_ME	21.1217	5.5520
State_MI	23.1641	5.5079
State_MN	22.9096	5.5305
State_MO	22.6315	5.5517
State_MS	22.8856	5.5500
State_MT	20.5095	5.5764
State_NC	23.0000	5.5447
State_ND	21.0049	5.5624
State_NE	21.9977	5.5622
State_NH	22.1678	5.5486
State_NJ	23.7615	5.5012
State_NM	22.4717	5.5759
State_NV	22.4211	5.5878
State_NY	23.7318	5.4834
State_OH	23.1541	5.5151
State_OK	22.1275	5.5729
State_OR	21.6158	5.5637
State_PA	23.4885	5.5143
State_RI	22.5008	5.5332
State_SC	22.7793	5.5456
State_SD	21.4924	5.5670
State_TN	22.5039	5.5482
State_TX	23.4079	5.5367
State_UT	21.9073	5.5616
State_VA	22.9497	5.5357
State_VT	20.5659	5.5622
State_WA	22.4466	5.5406
State_WI	22.2606	5.5282
State_WV	21.2802	5.5830
State_WY	19.8045	5.5516
t0	-7.7761	2.6721
t2	0.8279	0.3255
b1	0.5529	0.0634
w1	0.3298	0.0830
w2	0.5021	0.0666
R-square	0.8082	
Root MSE	22.9496	
Observations	4565	

Additional References

24. E. C. Schneider, Failing the test—the tragic data gap undermining the US pandemic response. *N. Engl. J. Med.* **383**, 299-302. doi: 10.1056/NEJMp2014836 (2020).

25. M. Salerno, F. Sessa, A. Piscopo, A. Montana, M. Torrisi, F. Patanè, P. Murabito, G. L. Volti, C. Pomara, No autopsies on COVID-19 deaths: a missed opportunity and the lockdown of science. *J. Clin. Med.* **9**, 1472 (2020).

26. D.M. Weinberger, J. Chen, T. Cohen, F.W. Crawford, F. Mostashari, D. Olson, V.E. Pitzer, N.G. Reich, M. Russi, L. Simonsen, A. Watkins, Estimation of excess deaths associated with the COVID-19 pandemic in the United States, March to May 2020. *JAMA Intern. Med.,* doi:10.1001/jamainternmed. 2020 .3391 Publishe (2020).

26. U.S. Center for Disease Control. Planning scenarios. https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html, retrieved 30 June 2020.

28. C. Courtemanche, J. Garuccio, A. Le, J. Pinkston, A. Yelowitz, Strong social distancing measures in the United States reduced the COVID-19 growth rate: study evaluates the impact of social distancing measures on the growth rate of confirmed COVID-19 cases across the United States. *Health Affairs* **10**, 1377 (2020).

29. A. Goolsbee, N. B. Luo, R. Nesbitt, C. Syverson, Lockdown Policies at the State and Local Level. doi: http://dx.doi.org/10.2139/ssrn.3682144 (20 Aug 2020).

30. R. Chetty, J. N. Friedman, N. Hendren, M. Stepner, How did covid-19 and stabilization policies affect spending and employment? a new real-time economic tracker based on private sector data. doi: http://dx.doi.org/10.3386/w27431 (25 June 2020).