Longitudinal proteomic profiling of dialysis patients with COVID-19 reveals markers of severity and predictors of death

Jack Gisby*[1], Candice L. Clarke*[1,2], Nicholas Medjeral-Thomas*[1,2], Talat H. Malik[1], Artemis Papadaki[1], Paige M. Mortimer[1], Norzawani B. Buang[1], Shanice Lewis[1], Marie Pereira[1], Frederic Toulza[1], Ester Fagnano[1], Marie-Anne Mawhin[1], Emma E. Dutton[1], Lunnathaya Tapeng[1], Arianne C. Richard[3], Paul D. W. Kirk[4,5], Jacques Behmoaras[1], Eleanor Sandhu[1,2], Stephen P. McAdoo[1,2], Maria F. Prendecki[1,2], Matthew C. Pickering[1], Marina Botto[1], Michelle Willicombe*[1,2], David C. Thomas*[1,2], James E. Peters*[1,6]
* equal contributions

Author Affiliations
1) Centre for Inflammatory Disease, Dept of Immunology and Inflammation, Imperial College London.
2) Renal and Transplant Centre, Hammersmith Hospital, Imperial College Healthcare NHS Trust, London, United Kingdom.
3) CRUK Cambridge Institute and Cambridge Institute for Medical Research, University of Cambridge.
4) MRC Biostatistics Unit, Forvie Way, University of Cambridge.
5) Cambridge Institute of Therapeutic Immunology & Infectious Disease, University of Cambridge.
6) Health Data Research UK, London, UK.

Correspondence to: James E. Peters. Email: j.peters@imperial.ac.uk

Abstract

End-stage kidney disease (ESKD) patients are at high risk of severe COVID-19. We measured 436 circulating proteins in serial blood samples from hospitalised and non-hospitalised ESKD patients with COVID-19 (n=256 samples from 55 patients). Comparison to 51 non-infected patients revealed 221 differentially expressed proteins, with consistent results in a separate subcohort of 46 COVID-19 patients. 203 proteins were associated with clinical severity, including IL6, markers of monocyte recruitment (e.g. CCL2, CCL7), neutrophil activation (e.g. proteinase-3) and epithelial injury (e.g. KRT19). Machine learning identified predictors of severity including IL18BP, CTSD, GDF15, and KRT19. Survival analysis with joint models revealed 69 predictors of death. Longitudinal modelling with linear mixed models uncovered 32 proteins displaying different temporal profiles in severe versus non-severe disease, including integrins and adhesion molecules. These data implicate epithelial damage, innate immune activation, and leucocyte-endothelial interactions in the pathology of severe COVID-19 and provide a resource for identifying drug targets.

Introduction

Coronavirus disease (COVID-19), caused by the SARS-CoV-2 virus, displays wide clinical heterogeneity from asymptomatic to fatal disease. Patients with severe disease exhibit marked inflammatory responses and immunopathology. The mechanisms underlying this remain incompletely characterised and the key molecular mediators are yet to be determined. The first treatment shown to reduce mortality from COVID-19 in randomised trials was dexamethasone [1], a corticosteroid which has broad non-specific effects on the immune system. Even with corticosteroid treatment, mortality in severe COVID-19 remains significant. There is a wide armamentarium of existing drugs that target inflammation more selectively, providing potential repurposing opportunities for the treatment of COVID-19. Recently, the REMAP-CAP trial has demonstrated efficacy of anti-IL6 receptor blockade in patients admitted to intensive care units with severe disease [2]. In order to select the most promising agents for future trials, we urgently need to better understand the molecular drivers of severe disease. Proteins are the effector molecules of biology and the targets of most drugs. Therefore, proteomic profiling to identify the key mediators of severe disease provides a valuable tool for identifying and prioritising potential drug targets [3].

Risk factors for severe or fatal COVID-19 include age, male sex, non-European ancestry, obesity, diabetes mellitus, cardiovascular disease, and immunosuppression [4]. End-stage kidney disease (ESKD) is one of the strongest risk factors for severe COVID-19 (estimated hazard ratio for death 3.69) [4], and ESKD patients hospitalised with COVID-19 have a mortality of approximately 30% [5–8]. ESKD patients have a high prevalence of vascular and

cardiometabolic disease (e.g. hypertension, ischaemic heart disease, diabetes), either as a
result of the underlying cause of their renal disease or as a consequence of renal failure. In
addition, ESKD results in both relative immunosuppression and chronic low-grade
inflammation, which may impact viral defence and the host inflammatory response.

Here we performed proteomic profiling of serial blood samples of ESKD patients with
COVID-19, leveraging the unique opportunity for longitudinal sampling in both the outpatient
and inpatient settings afforded by a large multi-ethnic haemodialysis cohort (**Figure 1a**).
These data revealed 221 proteins that are dysregulated in COVID-19 versus matched non-
infected ESKD patients. Using linear mixed models, joint models and machine learning, we
identified proteins that are markers of COVID-19 severity and risk of death. Finally, we
characterised the temporal dynamics of the blood proteomic response during COVID-19
infection in ESKD patients, uncovering 32 proteins that display altered trajectories in patients
with severe versus non-severe disease.

Results
We recruited 55 ESKD patients with COVID-19 (subcohort A; **Table 1**). All patients were
receiving haemodialysis prior to acquiring COVID-19. Blood samples were taken as soon as
feasible following COVID-19 diagnosis. At time of initial sample, 30 patients were outpatients
attending haemodialysis sessions and 25 were hospitalised inpatients (**Methods, Figure 1**).
Following the initial blood sample, serial sampling was performed for 51/55 patients. We also
recruited 51 non-infected haemodialysis patients as ESKD controls, mirroring the age, sex
and ethnicity distribution of the COVID-19 cases (**Figure 1 figure supplement 1a-c**). We
used the Olink proteomics platform to measure 436 proteins (**Supplementary File 1a**) in
256 plasma samples from the COVID-19 patients and the 51 control samples. The proteins
measured consisted of 5 multiplex 'panels' focussed on proteins relevant to immuno-
inflammation, cardiovascular and cardiometabolic disease. The 436 proteins assayed
showed strong enrichment for immune-related proteins (**Supplementary File 1b).**

In addition, we performed the Olink proteomic assays in 52 serum samples from a separate
set of 46 COVID-19 positive ESKD patients (subcohort B), and 11 serum samples from
ESKD COVID-19 negative controls (a subset of the controls described above). For the large
majority of patients only a single timepoint was available. A higher proportion of these
patients (41/46, 89%) were hospitalised and had severe disease (**Table 2**) than in subcohort
A (**Figure 1, Table 1**).

Proteomic differences between COVID-19 positive and negative ESKD patients

Principal components analysis (PCA) of proteomic data from subcohort A demonstrated

differences between samples from COVID-19 positive cases and controls, although the two

groups did not separate into discrete clusters (**Figure 2a-b**). To examine the effects of

COVID-19 on the plasma proteome, we performed a differential expression analysis in

subcohort A between COVID-19 cases (n=256 samples passing quality control from 55

patients) and non-infected ESKD controls (n=51) using linear mixed models, which account

for serial samples from the same individual (**Methods**). This revealed 221 proteins

associated with COVID-19 (5% false discovery rate, FDR); the vast majority were

upregulated, with only 40 downregulated (**Figure 3a**, **Supplementary File 1c**). In order to

provide a succinct and standardised nomenclature, we report proteins by the symbols of the

genes encoding them (see **Supplementary File 1a** for a mapping of symbols to full protein

names**)**. The most strongly upregulated proteins (in terms of fold change) were DDX58,

CCL7, IL6, CXCL11, KRT19 and CXCL10, and the most strongly downregulated were

SERPINA5, CCL16, FABP2, PON3, ITGA11 and MMP12 (**Figure 3 figure supplement 1**).

Notably, many of the upregulated proteins were chemotaxins.

We observed that a high proportion of the measured proteins were associated with COVID-

19. Given the highly targeted nature of the Olink panels that we used (enriched for immune

and inflammation-related proteins), this was not surprising. Nevertheless, to ensure that the

Benjamini-Hochberg adjustment of p-values was controlling the false discovery rate at the

5% level, we performed two additional analyses (**Methods**). First, we estimated the FDR

using an alternative method (the plug-in method of Tibshirani [9]); this confirmed appropriate

FDR control. Second, we used permutation to estimate the distribution of the number of

proteins expected to be declared significant under the null hypothesis of no association

between any proteins and COVID-19. This showed that the probability of observing the

number of differentially abundant proteins we identified was highly unlikely under the null

(empirical $p < 1 \times 10^{-5}$; **Figure 3 figure supplement 2**).

Although our COVID-19 negative controls were well matched in terms of age, sex and

ethnicity (**Figure 1 figure supplement 1a-c**), perfect matching of comorbidities was not

feasible in the context of the healthcare emergency at the time of patient recruitment. There

was a higher prevalence of diabetes in the COVID-19 cases compared to the controls (61.8

% versus 47.1%, respectively; **Table 1**). To evaluate whether differing rates of diabetes had

impacted the proteins identified as differentially abundant between cases and controls, we

performed a sensitivity analysis adding diabetes as an additional covariate in the linear

mixed model. This did not materially affect our findings; estimated effect sizes and -log10 p-

values from models with and without the inclusion of diabetes were highly correlated

(Pearson r > 0.99, and r = 0.95, respectively; **Figure 3 figure supplement 3a-b**). Full results
from both models are shown in **Supplementary File 1c**. Similarly, there were also
differences in the underlying cause of ESKD in cases compared controls (**Table 1**). We
therefore performed a further sensitivity analysis adjusting for underlying cause of renal
failure. This did not make any meaningful difference to our results (**Figure 3 figure**
supplement 3c-d, Supplementary File 1c).

We also considered the possibility that timing of haemodialysis might affect the plasma
proteome. To minimise the impact of this, all samples were taken prior to haemodialysis. For
the large majority (86.6%) of samples, the most recent haemodialysis was between 48 and
72 hours prior to blood draw. This consistency in timing of blood sampling reduces the
potential for impact of this issue. Nevertheless, to evaluate whether timing of haemodialysis
might have impacted our results, we performed a sensitivity analysis including time from last
haemodialysis as a covariate. Our results were not materially affected by this, with -log10 p-
values and estimated effect sizes very highly correlated with those obtained without inclusion
of this covariate (Pearson r >0.99 for effect size estimates and for -log10 p-values; **Figure 3**
figure supplement 4a-b, Supplementary File 1c).

We used the smaller subcohort B (n=52 serum samples from 46 patients with COVID-19;
Methods) for validation. We first projected the data from subcohort B into the PCA space of
subcohort A to examine how well the separation of cases and controls in the PCA space
replicated (**Methods**). This revealed clearer separation of infected and non-infected patients
than in subcohort A (**Figure 2c-d**), perhaps reflecting the higher proportion of hospitalised
patients (41 of 46 patients) in subcohort B (**Table 2**). We next performed differential
abundance analysis in subcohort B and found 201 proteins that were dysregulated in cases
versus controls (5% FDR) (**Supplementary File 1c**). Of the 221 differentially abundant
proteins from subcohort A, 150 (69.7%) were also identified in subcohort B at 5% FDR
(**Figure 4a**). Effect sizes in each dataset showed a strong correlation (r=0.80, **Figure 4b**).
This demonstrates that our findings are highly reproducible despite differences in sample
sizes and blood materials (plasma versus serum in subcohort A and B, respectively).

<u>Proteins associated with COVID-19 severity</u>
Examination of the principal components plot labelling samples by clinical severity at the
time of sampling (defined by WHO severity scores, graded as mild, moderate, severe or
critical) demonstrated a gradient of COVID-19 severity, best captured by principal
components 1 and 3 (**Figure 2 figure supplement 1a)**. To determine the proteomic effects

174 of COVID-19 severity, we tested for associations between proteins and WHO severity score

175 at the time of blood sampling, using linear mixed models with severity encoded as an ordinal

176 predictor (**Methods**). This analysis revealed 203 proteins associated with severity (**Figure**

177 **3b, Supplementary File 1d**). The majority of these were upregulated in more severe

178 disease, with only 42 down-regulated. A sensitivity analysis adjusting for time since last

179 haemodialysis, made no significant impact on our results (**Figure 3 figure supplement 4c-**

180 **d, Supplementary File 1d**). Consistent with previous reports, we found that severe COVID-

181 19 was characterised by elevated IL6. In addition, we observed a signature of upregulated

182 monocyte chemokines (e.g. CCL2, CCL7, CXCL10), neutrophil activation and degranulation

183 (e.g. PRTN3, MPO) and epithelial injury (e.g. KRT19, AREG, PSIP1, GRN). (**Figure 3b,c,**

184 **Figure 5)**. SERPINA5 and leptin showed the greatest downregulation as COVID-19 severity

185 increased (**Figure 3b,c)**.

187 We next asked how does the COVID-19 severity protein signature relate to the proteins that

188 are differentially abundant between cases and controls? The majority (140/203; 69%) of

189 severity-associated proteins were also identified as differentially abundant in the COVID-19

190 positive versus negative analysis (**Figure 6a**). Log fold changes for proteins in COVID-19

191 versus non-infected patients were correlated with effect sizes in the severity analysis, such

192 that the proteins most upregulated in cases versus controls also tended to show the greatest

193 upregulation in severe disease (**Figure 6b**). However, there were some notable exceptions

194 (eg CCL20, IL17C, OSM) that were strongly associated with severity but not differentially

195 expressed in infected versus non-infected patients (**Figure 6c**).

197 <u>Supervised learning to predict COVID-19 severity</u>

198 Principal components analysis revealed that some samples from patients who had mild or

199 moderate disease at the time of sampling clustered with samples from patients with severe

200 disease (**Figure 2 figure supplement 1a**). Examination of the same PCA plot this time

201 labelling samples according to the patient's overall clinical course (measured by peak WHO

202 severity score over the duration of the illness) (**Figure 2 figure supplement 1b)**, revealed

203 that these samples came from individuals who subsequently developed severe or critical

204 disease. This suggested that molecular changes may predate clinical deterioration. To

205 evaluate this further, we used supervised learning approaches to test if the proteomic

206 signature of the first blood sample for each patient in our dataset could identify whether the

207 patient either had clinically severe COVID-19 at the time of sampling or would develop

208 severe disease in the future. Whereas differential expression analyses consider each protein

209 marker separately, machine learning techniques allow examination of all proteins

210 concurrently, thus capturing non-linear relationships in the dataset. Using Random Forests,

we trained a classifier on the first sample for each COVID-19 patient to predict the overall clinical course, defined by peak WHO severity. For the purposes of this analysis, we binarised clinical course into either WHO mild/moderate or severe/critical.

The Random Forests method achieved 71% accuracy in predicting peak severity. By contrast, using only clinically available predictors (demographics, comorbidities and clinical laboratory results), the Random Forests method achieved 66% accuracy in predicting peak severity. Combining clinical parameters plus proteins did not improve accuracy (71%) compared to using proteomic predictors alone, suggesting that the information contained in the clinical predictors is captured at the proteomic level. While we do not believe that proteomic profiling is likely to enter clinical practice for risk stratification during this pandemic, the features selected by the classifier can highlight proteins of biological importance. We therefore interrogated the model to identify key proteins by calculating feature importance metrics (**Methods, Supplementary File 1e**). The most important proteins for indicating the presence of current or future severe disease were IL18BP, CTSD (Cathepsin D), GDF15, KRT19, TNFSF11 and IL1RL1 (ST2) (**Figure 7a)**. It is notable that through this distinct analytical approach, KRT19 again emerged as a key biomarker of severe disease.

Proteins associated with risk of death
9/55 patients in subcohort A died. We therefore sought to identify proteins associated with risk of death. To leverage the dynamic nature of repeated protein measurements for prediction of death, we utilised joint models, which combine linear mixed models and Cox proportional hazards models [10,11] (**Methods**). This analysis identified 44 proteins for which increased concentration was associated with increased risk of death (**Figure 7b, Supplementary File 1f**), including CST3, IL22RA1, AZU1, CCL28 and SPON1, and 25 proteins for which increased concentration was associated with reduced risk of death, including CD84, TNFSF12, TANK, PRKCQ and ADM.

Associations with clinical laboratory tests
A number of routine clinical laboratory tests have well characterised associations with COVID-19 (e.g. elevated inflammatory markers, d-dimer and reduced lymphocyte count) [12]. We therefore compared our proteomic data from COVID-19 patients at each timepoint to contemporaneous clinical laboratory measurements using linear mixed models (**Methods**). We found associations between plasma proteins and all clinical laboratory measurements except troponin (**Figure 8**, **Supplementary File 1g**). Many of these proteins were also markers of severity (e.g. IL6, KRT19, IFN-gamma and CXCL10 were strongly

248 associated with raised CRP and ferritin and reduced lymphocyte counts). Of note CCL7, a
249 monocyte chemokine that was also identified as an important marker of severity by the
250 Random Forests classifier, was associated with lower monocyte count and raised
251 inflammatory markers. Elevated neutrophil count was associated with Oncostatin-M, which
252 regulates IL6, GCSF and GMCSF production, and with the proteases MMP9 and defensin.
253

<u>Longitudinal analysis reveals proteins with distinct temporal profiles according to severity</u>
The immune response to infection is dynamic, and therefore snapshot measurements
provide only partial insights. Leveraging the dense serial sampling in our dataset (**Figure 1**),
we modelled the temporal trajectory of each protein and asked whether or not any protein
trajectories differed in patients with a severe/critical versus mild/moderate overall clinical
course. This was achieved using linear mixed models that included a term for time from first
symptoms and a time x severity interaction term (**Methods**).
261

178 proteins displayed a significant association with time from first symptoms (5% FDR),
demonstrating the temporal variability in plasma proteins across the disease course
(**Supplementary File 1h**). Moreover, we identified 32 proteins for which there was
significant interaction between time and severity, i.e. proteins displaying differential temporal
trajectories between mild/moderate and severe/critical infections (**Supplementary File 1h,**
Figure 9). Among the proteins with the strongest temporal differences according to clinical
course were the integrins ITGA11 and ITGB6, the adhesion molecule ICAM1, TNFRSF10B
(a receptor for TRAIL) and PLAUR, the receptor for urokinase plasminogen activator. Most
of these proteins exhibited rising profiles in the more severe patients but flat profiles in milder
cases. ACE2, the receptor for SARS-CoV-2, also displayed this pattern (**Figure 9**). In
contrast, abundance of ITGA11, which was also identified as reduced in the analysis of
infected versus non-infected patients, fell over time in the severe group.
274

<u>Testing for proteins associated with ethnicity</u>
In the UK, individuals from ethnic minorities are at higher risk of severe disease and death
from COVID-19 [4]. We therefore examined whether any of the proteins we measured
exhibited differences across ethnicities, analysing COVID-19 positive cases and controls
separately (**Methods**). In COVID-19 negative ESKD patients, no proteins were significantly
associated with ethnicity in a multivariable model adjusting for age and sex. In COVID-19
positive ESKD patients, there is the potential for protein associations with ethnicity to be
confounded by disease severity. To account for this, we included severity as well as age and
sex as covariates. A single protein, LY75, was associated with ethnicity in this multivariable
model (nominal P 0.0001, Benjamini-Hochberg adjusted P 0.04, with higher levels in white

patients). Using the same within cases analysis strategy in subcohort B, we found no
proteins were significantly associated with ethnicity after multiple testing correction, although
the nominal P value for LY75 was 0.025. While these analyses failed to identify substantial
ethnicity-related variation in the proteins we measured, an important caveat is that there
were relatively modest numbers of individuals from each ethnic group, and so statistical
power was limited. Larger multi-ethnic studies are needed to adequately address this
question.

Comparisons to other proteomic studies in COVID-19

Other studies have used a variety of proteomic platforms to investigate COVID-19. We
compared our findings to those of 3 published studies [13–15] and a preprint by Filbin *et al*
[16]. Of the 221 proteins that were differentially abundant in our analysis of COVID-19
positive versus negative ESKD patients, 116 associations had been previously reported
(**Supplementary File 1i**). Of the 203 proteins associated with severity, 165 had previously
been reported (**Supplementary File 1j**).

We focussed in more detail on the study by Filbin *et al* [16] because of the large sample size
and the breadth of proteomic assay used. This study comprised 384 patients with acute
respiratory distress (306 COVID-19 positive and 79 COVID-19 negative), and measured
1,472 proteins using the Olink Explore platform. 417 of these were also measured in our
study. Of the 221 proteins differentially abundant in our case/control analysis, 210 were
measured in their study. Of these 100 (47.6%) were significant in their analysis of COVID-19
positive versus negative respiratory distress. In addition, we observed strong correlation
(r=0.69) between the estimated log fold changes in our and their studies (**Figure 4 figure**
supplement 1). Of the 203 proteins associated with severity in our study, 192 were
measured in their study. 157 of these were significantly associated with severity, giving a
concordance of 81.8%. Thus, despite the differences in study design and clinical
populations, we observed notable similarities in our results and those reported by Filbin *et al*
[16].

Discussion

In this study we performed plasma proteomic profiling of haemodialysis patients with COVID-
19. A strength of our study was that we were able to perform serial blood sampling in both
the outpatient and inpatient setting, including longitudinal samples from the same individual
before and after hospitalisation. This was possible because haemodialysis patients are
unable to fully isolate as they must continue to attend for regular dialysis sessions.

Moreover, haemodialysis patients represent an important group since ESKD is one of the

strongest risk factors for death from COVID-19 [4,6–8]. Data from the UK Renal Registry

shows that 7- and 14-day mortality for COVID-19 infected in-centre haemodialysis patients

was 11% and 19%, respectively [17]. Data from the Scottish Renal Registry estimates 30-

day mortality following a positive COVID-19 test as 22%, and as of 31 May 2020, 28.2% of

renal replacement therapy patients who had a positive COVID-19 test had died [18]. In our

local population of 1,352 in-centre haemodialysis patients, 315 patients had tested positive

for COVID-19 by the end of our study period (31 May 2020), of whom 53% required

hospitalisation and 85 (27%) died. The OpenSAFELY study [4] examined ~17 million UK

primary care records and linked these to the UK COVID-19 mortality register. Patients with

estimated glomerular filtration rate (eGFR) <30ml/min/1.73m^2 had a hazard ratio (HR) for

death of 3.56 after adjustment for age and sex.

In part, the high mortality from COVID-19 in ESKD patients likely reflects the fact that these

patients are enriched for cardiometabolic traits that predispose to severe COVID-19.

However, in multivariable analyses adjusting for these factors, impaired renal function

remains an independent risk factor for severe COVID-19 [4]. Moreover, there is an inverse

relationship between renal function and risk of death from COVID-19 across the spectrum of

chronic kidney disease. These observations support the notion that the state of ESKD *per se*

is an important determinant of outcome in COVID-19. ESKD is well recognised as an

immunosuppressed state [19–21], with defects in both innate and adaptive immunity [22–

25]. Accordingly, ESKD confers increased vulnerability to viral infections including influenza

and respiratory syncytial virus [26–29]. In addition, ESKD results in a chronic low-grade

inflammatory state [30]. This tendency to a pro-inflammatory state, combined with reduced

ability to respond to viruses, may contribute to the abnormal host response to SARS-CoV-2

infection, producing the immunopathology that leads to severe COVID-19.

Our comparison of COVID-19 positive and negative haemodialysis patient plasma samples

revealed 221 proteins that were differentially abundant in COVID-19. The majority of these

were upregulated, with strong representation of viral response proteins (e.g. DDX58, IFNG),

cytokines/chemokines (e.g. IL6, CCL7, CXCL10 and CXCL11) and epithelial proteins (e.g.

KRT19, PSIP1) (**Figure 3a**). The COVID-19 negative controls in this analysis were carefully

matched to cases in terms of age, sex and ethnicity. However, complete matching of clinical

characteristics was not feasible; there were differences in the prevalence of diabetes and the

underlying causes of ESKD between COVID-19 positive cases and controls (**Table 1**).

Sensitivity analyses adjusting for these covariates gave highly consistent results, indicating

that our findings are robust. In addition, we validated our findings when we analysed serum
samples from a separate subcohort of COVID-19 positive ESKD patients.

ESKD is itself likely to significantly impact the plasma proteome. Previous cross-sectional
studies have shown that the levels of many circulating proteins have an inverse relationship
with eGFR [31,32]. A longitudinal study using an Olink proteomics panel (although not one
used in our study) found that for 74% of the 84 proteins measured, protein levels rose as
eGFR fell [33]. For many proteins, it is unclear whether this inverse relationship with renal
function reflects cause or effect. Some proteins may be increased in chronic kidney disease
due to reduced renal clearance, some may be elevated secondary to tissue injury or chronic
inflammation, and others may be drivers of renal injury. Regardless, this observation of
widespread changes in the blood proteome of kidney disease patients emphasises the
importance of using COVID-19 ESKD patients rather than healthy individuals as our control
group.

Analysis within COVID-19 cases revealed 203 proteins associated with disease severity, the
strongest of which was IL6 (**Figure 3b**). Association of IL6 with severe disease is well-
established and has already received considerable attention [34,35]. Despite promising initial
case reports of IL6R receptor blockade in COVID-19, convincing efficacy was not
demonstrated in early randomised trials [36]. More recently, the REMAP-CAP trial has
shown the benefit of anti-IL6R therapy when given to critically ill patients on admission to
intensive care units [2], indicating that IL6 does contribute to critical illness from COVID-19.
Our finding that IL6 was most strongly upregulated in severe disease demonstrates the
value of plasma proteomic profiling in identifying putative drug targets.

Members of the CCL and CXCL chemokine families (e.g. CCL2, CCL7, CCL20 and
CXCL10) were strongly associated with severity. Likewise, higher levels of CCL2, CCL7,
CCL20 and CXCL10 were associated with lower blood lymphocyte count and higher
inflammatory markers (CRP and ferritin) (**Supplementary File 1g**), which are clinical
markers of severe disease and poorer outcome in COVID-19 [37]. Of note, CCL20 is a
chemoattractant for lymphocytes [38], and its negative association with lymphocyte count
may reflect a direct effect on migration of lymphocytes from the blood into the tissues rather
than simply marking severe disease. CCL2 (also known as MCP-1) and CCL7 (MCP-3), are
both chemokines for monocytes, and CXCL10 has pleiotropic immunological effects
including chemotaxis. These chemokines were also negatively correlated with blood
monocyte count, suggesting recruitment of these innate immune cells into damaged tissues.

The neutrophil proteases PRTN3 (proteinase-3) and MPO (myeloperoxidase) (**Figure 5)** and

the neutrophil-derived protein AZU1 were associated with severe disease (**Supplementary**

File 1d), indicating that neutrophil activation and degranulation are features of severe

COVID-19. Degranulation of neutrophils releasing PRTN3 and MPO could potentially

contribute to oxidative damage in the lungs and thus more severe disease.

A striking finding of our study was the association of disease severity with upregulation of

epithelial proteins (e.g. KRT19) and epithelial tissue repair pathways (e.g. PSIP1, AREG,

GRN (progranulin)), most likely reflecting lung and vascular damage. KRT19 was notably

prominent in our analyses, as well as the study by Filbin *et al.* [16](**Supplementary File 1j**).

KRT19 is an intermediate filament protein, important for the structural integrity of epithelial

cells [39]. These data suggest that severe COVID-19 is characterised by destruction of the

lung epithelium and vascular endothelium. Vascular injury might thus explain the high level

of vascular thrombosis seen in patients in severe disease. In summary, our data reveal that

severe COVID-19 is characterised proteomically by a signature of innate immune activation

and epithelial injury.

69% of proteins associated with severity were also differentially abundant in the case versus

control analysis (**Figure 6a**), and for the large majority of proteins the within-cases severity

analysis effect size was proportional to the fold change between cases and controls (**Figure**

6b). This suggests that, in general, the distinction in the plasma proteome between severe

and mild COVID-19 is a quantitative difference in the COVID-19 signature, rather than there

being an orthogonal signature involving a different set of proteins. Consistent with this

concept, examination of PCA plots coloured by severity revealed that while there was a

gradient of COVID-19 severity, the samples from severe or critical patients did not form a

discrete cluster distinct from those from patients with milder disease (**Figure 2 figure**

supplement 1). However, there were a few exceptions where proteins that were associated

with severity were not upregulated in the case-control analysis. These included OSM, IL17C,

and CCL20 (**Figure 6c**). These proteins therefore reflect biological processes specifically of

severe disease and may represent therapeutic targets.

Survival analysis identified 44 proteins associated with increased risk of death (**Figure 7b**).

As expected, many of these were also associated with disease severity, high CRP and lower

lymphocyte count (**Figure 7 figure supplement 1**). In contrast, 25 proteins were associated

with reduced risk of death (**Figure 7b**). One such protein is the multi-functional cytokine

TNFSF12 (TWEAK). Although TWEAK can exert pro-inflammatory effects, it also can inhibit

the innate immune response [40] and promote tissue repair and endothelial cell proliferation

and survival [41], which may be beneficial responses in COVID-19. This illustrates that
although proteins associated with inflammation are often thought to be destructive, the
inflammatory response also induces a programme for limiting injury and initiating tissue
repair. Insufficient activation of such homeostatic mechanisms may contribute to why some
individuals get severe COVID-19.

The host immune response to COVID-19 is a dynamic process, and clinical deterioration
typically occurs 7-10 days after first symptoms. Temporal information may therefore be
important in determining optimum timing of therapeutic intervention (e.g. blockade of a
particular cytokine). By taking serial samples and examining their patterns within individuals
over time, we were able to model protein trajectories and found that many proteins display
temporal variability during COVID-19. Longitudinal measurements also allow molecular
comparison of severe versus mild disease trajectories. By modelling the interaction term
between time from first symptoms and overall disease course, we found 32 proteins that
displayed distinct temporal profiles in severe vs mild disease. These results point to
enhanced leucocyte-endothelial cell interactions indicated by upregulation of cell adhesion
molecules (e.g. ITGB6, ICAM1) in severe disease. This endothelial activation may contribute
to COVID-19-associated thrombosis discussed above. Management of thrombosis in
COVID-19 currently consists of anticoagulation. Our results suggest that disrupting
leucocyte-endothelial interactions may be a complementary therapeutic strategy.

Several proteins associated with either risk of death or clinical severity lie in pathways
targeted by existing drugs. PARP1 was identified as an important marker of current or future
severe COVID-19, and also was associated with risk of death. PARP1 is associated with
inflammatory and vascular disease [42]. PARP1 inhibitors are in use for cancer [43], and our
data suggest that re-purposing of PARP1 inhibition in COVID-19 should be explored further.
IL33 was associated with both risk of death and clinical severity, and its receptor IL1RL1
(ST2) was associated with clinical severity and identified as an important predictor of severe
clinical course. Monoclonal antibodies against IL33 and its receptor are in late-stage
development for asthma [44], and could also be explored in COVID-19. As discussed above,
MPO was associated with clinical severity. MPO inhibitors [45] might have a role in reducing
neutrophil-mediated tissue injury in COVID-19. Finally, inhibitors of monocyte chemokines
(e.g. CCL2) and their receptors have been developed [46,47], although drugging these
pathways is made more challenging by molecular cross-talk. An important caveat is that we
cannot determine whether the associations we observed are drivers of pathology in COVID-
19 or simply reflect the downstream consequences of inflammation and tissue injury. Future

studies using Mendelian randomisation analysis will provide a useful tool for assessing
causality and prioritizing drug targets.

Other groups have studied the plasma or serum proteome in COVID-19 [13–16,48], using
either mass spectrometry or immunoassays including the Olink platform. Mass spectrometry
is less sensitive than immunoassays and so it likely to be unable to detect many of the
cytokines measured here. Conversely, it can provide complementary information by
measuring many proteins that our immunoassays did not target. A limitation of our study was
that we used Olink panels that measured specific proteins selected on their relevance to
inflammation, immunity, cardiovascular and metabolic disease. This bias precluded formal
pathway enrichment analysis. In general, our results had greater similarities to studies that
used immunoassays over mass spectrometry (**Supplementary File 1i-j**). 47.6% of proteins
differentially expressed in COVID-19 positive versus negative ESKD patients in our study
were differentially expressed in COVID-19 positive versus negative acute respiratory distress
syndrome patients in the study of Filbin et al [16], who used a different Olink proteomics
platform. Moreover, we observed consistent effect sizes (**Figure 4 figure supplement 1**).
These similarities are striking given the difference in clinical populations and control groups;
in Filbin *et al*'s report the controls included patients with non-COVID-19 respiratory infections
whereas our control group did not have active infection. The concordance in proteins
associated with COVID-19 severity within cases was even higher (81.8%). The similarities
suggests a similar plasma proteomic signature of COVID-19 across different clinical
populations, particularly the signature associated with severity.

In summary, this study reveals proteins associated with COVID-19 infection and severity,
and demonstrates altered dynamic profiles between patients with severe disease and those
with a more indolent course. Our results emphasise the importance of studying and targeting
mechanisms that reduce the lung epithelial and endothelial damage to both alleviate the
severity of the infection and to reduce the chance of long-lasting complications. These data
provide a valuable resource for therapeutic target prioritisation.

Materials and Methods

<u>Subjects and samples</u>
Ethical approval: All participants (patients and controls) were recruited from the Imperial
College Renal and Transplant Centre and its satellite dialysis units, London, and provided
written informed consent prior to participation. Study ethics were reviewed by the UK
National Health Service (NHS) Health Research Authority (HRA) and Health and Care

Research Wales (HCRW) Research Ethics Committee (reference 20/WA/0123: The impact

of COVID-19 on patients with renal disease and immunosuppressed patients). Ethical

approval was given.

Subcohort A: We recruited 55 COVID-19 positive haemodialysis patients, either as

outpatients or as inpatients (**Table 1**). All patients were receiving in-centre outpatient

haemodialysis prior to COVID-19 diagnosis. COVID-19 was confirmed in all cases with

positive nasal PCR for the SARS-CoV-2 virus. Patients were recruited during the first UK

national lockdown, with recruitment from 8[th] April – 30[th] May 2020. Blood was collected in

EDTA tubes and centrifuged to obtain plasma, and stored at –80^{o}C. Sample processing was

performed within 4 hours of venepuncture. The initial sample was taken as an outpatient for

30 patients and as an inpatient for 25. Where feasible, serial blood samples were taken. In

total, 259 samples were taken (3 subsequently failed QC – see below). The median number

of serial samples was 5 (range 1-10) (**Figure 1 figure supplement 1d**). 8 patients who were

recruited as outpatients were subsequently admitted to hospital with COVID-19 over the

course of the study. 27 of 55 (49.1%) patients had severe or critical disease (defined by

peak WHO severity). 9 (16.4%) patients died.

In addition, we recruited 51 COVID-19 negative haemodialysis controls. COVID-19 negative

haemodialysis controls were selected to mirror the cases in terms of demographic features

(age, sex, ethnicity) (**Figure 1 figure supplement 1a-c**). These control patients had no

clinical features of any other infection.

Subcohort B: We also recruited a separate set of 46 COVID-19 positive haemodialysis

patients (**Table 2**). These patients were recruited from the same centre, but slightly earlier

than subcohort A (recruitment commenced on 30[th] March 2020). For these patients, blood

was collected in serum tubes and centrifuged to obtain serum. At this time, we had very

limited access to laboratory facilities and so plasma was not collected from these patients. 5

were outpatients and 41 were inpatients, reflecting the fact that UK policy was weighted

towards inpatient testing at the time these patients were recruited. 33 of 46 patients (71.7%)

had severe or critical disease (by peak WHO severity), and 9 (19.6%) patients died. For 40

patients, only one sample from a single timepoint was collected, and for 6 patients, 2

samples were collected. To provide controls for subcohort B, we used serum samples from

11 non-infected haemodialysis patients (collected at the same time as plasma from a subset

of the ESKD control group described above).

<u>Clinical severity scores</u>

Severity scoring was performed based on WHO classifications (WHO clinical management
of COVID-19: Interim guidance 27 May 2020) adapted for clinical data available from
electronic medical records. 'Mild' was defined as COVID-19 symptoms but no evidence of
pneumonia and no hypoxia. 'Moderate' was defined as symptoms of pneumonia or hypoxia
with oxygen saturation (SaO2) greater than 92% on air, or an oxygen requirement no greater
than 4L/min. 'Severe' was defined as SaO2 less than 92% on air, or respiratory rate more
than 30 per minute, or oxygen requirement more than 4L/min. 'Critical' was defined as organ
dysfunction or shock or need for high dependency or intensive care support (i.e. the need for
non-invasive ventilation or intubation). Severity scores were charted throughout a patient's
illness. We defined the overall severity/clinical course for each patient as the peak severity
score that occurred during the patient's illness.
553
Proteomic assays
Plasma and serum proteomic measurements were performed using Olink proximity
extension immunoassays (https://www.olink.com/products/). Five 92-protein multiplex Olink
panels were run ('inflammation', 'immune response', 'cardiometabolic', 'cardiovascular 2' and
'cardiovascular 3'), resulting in 460 measurements per sample. Since a small number of
proteins were measured on more than one panel, we measured a total of 436 unique
proteins. The Olink assays were run using 88 samples/plate. All plates were run in a single
batch. Plate layouts was carefully designed to avoid confounding of potential plate effects
with biological or clinical variables of interest. To achieve this, we used an experimental
design that combined ensuring case/control balance across plates with random selection of
samples from each category and random ordering of allocation to wells. This is outlined in
more detail as follows. We ensured that each plate contained a mixture of control and case
samples. Specifically, a fixed proportion of each plate was designated for control samples.
The allocation of specific control samples to each plate was performed using randomisation.
For the case samples, we again used randomisation for plate assignment, with the constraint
that once one sample from a given patient was allocated to a plate, all other longitudinal
samples from that patient were assigned to same plate. Finally, once all the samples had
been allocated to plates, the layout of samples within each plate was determined through a
further randomisation step for well allocation.
573
574 Protein annotation
575 We used the Human Protein Atlas version 20.0 [49] for protein annotation (**Figure 1 figure**
576 **supplement 2**). We performed enrichment analysis of the 436 proteins that we measured
577 using string-db [50].
578

<u>Normalisation and quality assessment and control</u>

The data was normalised using standard Olink workflows to produce relative protein
abundance on a log2 scale ('NPX'). Quality assessment was performed by a) examination of
Olink internal controls, and b) inspection of boxplots, relative log expression (RLE) plots [51]
and PCA. Following these steps, 3 poor quality samples were removed. In addition, 5
samples failed quality control on a single proteomic panel only, with the remaining panels
passing QC. For these samples, proteins on the panel that failed QC were set to missing,
and the data for the remaining proteins was retained.

Principal components analysis revealed no substantial impact of plate effects (**Figure 2**
figure supplement 2). 13 proteins were assayed more than once due to their inclusion in
multiple Olink panels. For plasma, the median correlation between the assays was 0.986
with an IQR of 0.974-0.993 and a range of 0.925 to 0.998. For serum, the median correlation
between the assays was 0.991 with an IQR of 0.952-0.995 and a range of 0.737-0.999. We
removed duplicate assays at random prior to subsequent analyses.

For 11 ESKD controls, we had contemporaneous plasma and serum samples. To assess the
comparability of these two matrices, we calculated the Pearson's correlation coefficient
between the assays for each protein (**Supplementary File 1k**). 344/436 (78.9%) of proteins
had a Pearson's r > 0.5. We also report the variance of each protein in plasma and serum
since low correlation may reflect low variance. The proteins with the lowest estimated
Pearson correlation coefficient were AZU1, STK4 and TANK. We highlight that this
comparison had small sample size (only 11 samples) and that the samples were from control
patients without infection. Caution should be made in extrapolating these findings to the
context of active infection where protein dynamic ranges may be different.

<u>Missing values</u>

Following QC, 0.22% data points were missing for the plasma dataset and 0.35% for the
serum dataset. For analyses that required no missing values (PCA and supervised learning),
we imputed missing values as follows. The dataset was first scaled and centred, and missing
values imputed using caret's k-nearest neighbours (kNN) method [52]. The 5 closest
samples (by Euclidean distance) were used to estimate each missing value.

<u>Principal Components Analysis</u>

Singular value decomposition was used to perform PCA on the proteomic data from
subcohort A (plasma samples). We then used the loadings from subcohort A together with

the proteomic data from subcohort B to calculate principal components scores. This enabled projection of subcohort B data into the PCA space of subcohort A.

<u>Differential protein abundance analysis: COVID-19 positive versus negative</u>

Differential protein abundance analyses between COVID-19 positive and negative samples were performed using linear mixed models, to account for the use of serial samples from the same individuals (R lme4 package [53]). This analysis compared 256 samples from 55 COVID-19 patients with 51 non-infected patients (1 sample per non-infected patients). Age, sex and ethnicity were included as covariates. We used a random intercept term to estimate the variability between individuals in the study and account for repeated measures. The regression model in R notation was:

$$NPX \sim covid\_status + sex + age + ethnicity + (1 \mid individual)$$

where NPX represents the protein abundance and covid_status was a categorical variable (infected/non-infected). Sex and ethnicity were also categorical variables. Age was a quantitative variable. We calculated P values using a type 3 F test in conjunction with Satterthwaite's method for estimating the degrees of freedom for fixed effects [54]. The regression model was fitted for each of the 436 proteins individually. Multiple testing correction was performed using the Benjamini-Hochberg method and a 5% FDR used for the significance threshold.

The same approach was used for subcohort B. This analysis comprised 52 serum samples from 46 COVID-19 positive patients versus 11 samples from non-infected patient samples (1 sample per non-infected patient).

As sensitivity analyses, we repeated the differential abundance analyses between case and controls for the subcohort A adjusting for additional covariates and comparing this to the basic model (i.e. using age, sex, and ethnicity alone). This was performed for each of the following parameters: diabetes status, cause of ESKD, and time to last haemodialysis.

<u>Testing for associations between proteins and clinical severity</u>

For testing the association of plasma proteins with the 4-level WHO severity rating (mild, moderate, severe and critical) within COVID-19 positive cases from subcohort A (n=256 samples from 55 patients), we used a similar linear mixed modelling approach to the COVID-19 positive versus negative differential abundance analysis; for this analysis, the covid_status term was replaced by a severity variable encoded using orthogonal polynomial contrasts to account for ordinal nature of severity levels. As before, age, sex and ethnicity

were included as covariates. As a sensitivity analysis, we repeated the analysis with time to

last haemodialysis (days) as an additional covariate.

<u>Testing for associations between proteins and clinical laboratory tests</u>

The linear mixed modelling strategy was also employed for testing association of temporal

clinical laboratory variables and protein levels, with the value of the clinical variable (as a

quantitative trait) used in place of covid_status. Only COVID-19 positive patients were

included in this analysis. Contemporaneous lab measurements were not available for all

samples. This varied according to the clinical lab parameter. Some (eg troponin, d-dimer)

were measured less frequently than full blood count and CRP. Details of the proportion of

missing values for each lab parameter are included in (**Supplementary File 1g**). We also

calculated correlations between clinical laboratory variables and protein levels using the R

package rmcorr, which determines the overall within-individual relationship among paired

measures that have been taken on two or more occasion [55].

<u>Testing for associations between proteins and ethnicity</u>

We performed testing of protein levels and ethnicity separately in COVID-19 negative ESKD

patients and COVID-19 positive ESKD patients. These analyses were limited to individuals

who were White, South Asian (Indian, Pakistani or Bangladeshi ancestry) or Black as there

were too few individuals from other ethnic groups for meaningful interpretation. For COVID-

19 negative patients (1 sample per patient), we performed linear regression for each protein

with ethnicity as the predictor variable, and age and sex as covariates. For COVID-19

positive patients we used a linear mixed model to account for serial samples from the same

individual, again with age, and sex as covariates.

<u>Multiple testing correction</u>

We used the Benjamini-Hochberg method to control the FDR at 5% for all statistical

analyses.

<u>Alternative estimation of the FDR using the plug-in method</u>

To provide additional support that the Benjamini-Hochberg procedure was providing

adequate control of the FDR, we also used the plug-in method of Tibshirani [9] as an

alternative method to estimate the FDR, as described below.

1. We defined R as the number of associations declared significant in the real data.

2. We defined C as the test statistic used as the significance threshold used in the real data

(i.e. that corresponding to an adjusted p-value of 0.05).

3. The expected number of proteins that we would find significant under the null hypothesis

that no proteins are differentially abundant between COVID-19 positive versus negative

patients (i.e. false positives) was estimated using a permutation strategy. We randomly

permuted each individuals' COVID-19 status label 100,000 times and, in each case,

repeated the differential abundance analysis on the permuted data. The estimated the

number of false positives ($\hat{V}$) was then estimated by the number of associations with test

statistic > *C* in 100,000 permutations of the data, divided by the number of permutations.

4. The estimated FDR was then calculated as $\hat{V}/R$.

We implemented a similar approach for the testing the association of proteins with severity

scores within cases.

Using this method, the estimated FDR for the case versus control analysis was 0.062 and for

the severity analysis 0.057, indicating that we had appropriately controlled the FDR.

Empirical p-value calculation

As a complementary analysis, based on the approach of Filbin *et al.* [16], we estimated the

empirical P-value for the likelihood of observing as many significant proteins as we identified

in the real data if the null hypothesis of no differentially abundant proteins in cases versus

controls were true. We again used 100,000 permutations of the case control labels to

estimate the null distribution. We performed Benjamini-Hochberg adjustment on the nominal

p-values of each permutation, and counted the number of proteins that were significant

(adjusted p-value <0.05) in each permutation.

The distribution of the number of proteins declared significant is shown in **Figure 3 figure**

supplement 2a; on no occasion in 100,000 permutations did we observe more proteins

declared significant than in the real data. We can thus state that the empirical P-value (the

fraction of permutation runs where we observed $\geq$ the number of associations in the real

data) is less than $1/100{,}000 = 1 \times 10^{-5}$.

We also applied this method to the association testing of proteins with severity scores within

cases (**Figure 3 figure supplement 2b)**. Again, on no occasion in 100,000 permutations did

we observe more proteins declared significant than in the real data (empirical p-value $<1 \times 10^{-5}$

).

Supervised learning

Random forest models were fit using R's randomForest and caret packages [52,56]. Data
was centred, scaled and imputed as in *Data Preparation* with the caveat that, during cross-
validation, the pre-processing procedure was first applied on the resampled (training) data
before the same method was applied without re-calculation to the holdout (test) set. To
estimate model accuracy, we used 4-fold cross-validation. The cross-validation procedure
was repeated 100 times. The model's parameters were kept constant at 500 trees and an
mtry value (number of proteins randomly sampled as candidates at each node) calculated as
the square root of the number of features. After parameter estimation, we fit a final model
trained using the entirety of the dataset. This model was used for subsequent feature
extraction. Random forest feature extraction was carried out using the R
randomForestExplainer package. We made use of the following importance measures:
accuracy decrease (the average decrease in prediction accuracy upon swapping out a
feature), number of trees (the number of trees with a node corresponding to a feature) and
mean minimal depth (the average depth at which a node corresponding to a feature occurs).
Three models were generated with different input features: i) proteomic data alone; ii) clinical
parameters alone; iii) proteomic data and clinical parameters. Clinical parameters included
sex, age, ethnicity, cause of ESKD, comorbidities, smoking status, radiological evidence of
pulmonary infiltrates, and clinical laboratory tests.
740
Survival analysis using joint modelling
Following scaling and centering, we fit linear mixed models for each protein to capture the
temporal trajectories of each individual. A polynomial spline of degree 2 was used to model
protein concentration with respect to time (from symptom onset, measured in days); the
spline was fitted for samples that were taken between 1 and 28 days from first symptoms,
inclusive. Proteomic data after that point was censored. We estimated both random
intercepts and random slopes for each individual, as per the following R formula notation:

748   NPX ~ time + (time | individual)

749 These were joined to a Cox regression model using the jointModel package [11] in order to
750 estimate the association of each protein with risk for death. P values were calculated using a
751 Wald test for the association between the linear mixed model and Cox regression.
752 Benjamini-Hochberg adjustment was applied, with an adjusted p-value of 0.05 used as the
753 significance threshold.
754
Longitudinal analysis
We also used linear mixed models to estimate the temporal profile of each protein. For this
longitudinal analysis we explicitly modelled the time from first symptoms. We set up the

model to test for each protein a) whether the protein significantly change over time and b)
whether the protein changes over time differently in individuals with a mild versus severe
disease course. The latter was performed statistically by testing for an interaction effect
between time and clinical course. For the purposes of this analysis, we binarised patients
into severe or non-severe clinical course according to the peak WHO severity disease of
their illness. Patients with a peak WHO score of mild or moderate were considered non-
severe and those with a peak score of severe or critical were considered severe.

We then used R's bs function to fit a polynomial spline of degree 2 to model protein
concentration with respect time (from symptom onset, measured in days) [57]. The spline
was fit for samples that were taken between 1 and 21 days from symptom onset, inclusive.
We estimated random slopes with respect to time, in addition to random intercepts, to
account for each individual's unique disease course. For each protein, we fitted the following
model (R notation):
772   NPX ~ time * severity + sex + age + ethnicity + (time | individual)
To identify proteins that changed significantly over time, we examined the P-values for the
main effect of time. To identify proteins with distinct temporal profiles between severe and
non-severe cases, we examined the P values for the time x severity interaction term. For
each of these two research questions, P-values were adjusted for the multiple proteins
tested using the Benjamini-Hochberg method and 5% FDR used as the significance
threshold.

Data availability statement
All data generated during this study are included in the manuscript and supporting files.
Underlying source data for all analyses (individual-level proteomic and clinical phenotyping
data) are available without restriction as **Source Data Files 1-4**. In addition, these data have
been deposited in the Dryad Digital Repository (doi:10.5061/dryad.6t1g1jwxj).

Code availability: code is available in the following GitHub repository:
https://github.com/jackgisby/longitudinal_olink_proteomics

Author contributions:
Conceived and designed the study: MCP, MB, MW, DCT, JEP.
Funding acquisition: JB, MCP, MB, MW, DCT, JEP.
Patient recruitment and sample collection: CLC, NM-T, ES, SPM, MFP, MW, DCT.
Led and coordinated patient recruitment: MW.
Sample processing: THM, PMM, NBB, SL, MP, FT, EF, M-A M, ED, LT, MB.
Clinical phenotyping: CLC, NM-T, SL, ES, MFP, MCP, JEP.
Data analysis: JG, AP.
Supervised the analysis: JEP.
Statistical support: PK, ACR.
Wrote the paper: JG, JEP

Results interpretation and editing the paper: MB, MCP, DT, ACR, JB.
All authors critically reviewed and approved the manuscript before submission.

Acknowledgements
The authors thank the patients who volunteered for this study and the staff at Imperial College Healthcare NHS Trust (the Imperial College Healthcare NHS Trust renal COVID-19 group and dialysis staff):

Appelbe M, Ashby DR, Brown EA, Cairns T, Charif R, Condon M, Corbett RW, Duncan N, Edwards C, Frankel A, Griffith M, Harris S, Hill P, Kousios A, Levy JB, Loucaidou M, Lightstone L, Liu L, Lucisano G, Lynch K, Mclean A, Moabi D, Muthusamy A, Nevin M, Palmer A, Parsons D, Prout V, Salisbury E, Smith C, Tam F, Tanna A, Tansey K, Tomlinson J, Webster P.

We also acknowledge the efforts of renal specialist doctors in training for assistance with recruiting patients to this study.

We also thank:
- Hari and Rachna Murgai and Milan and Rishi Khosla for generous support with sample transport.
- Dr Kerry Rostron for exceptional support with laboratory facilities in challenging circumstances and the Department administrators for their help.
- Dr Brian Tom and Dr Jessica Barrett (MRC Biostatistics Unit, University of Cambridge) for statistical advice.
- Prof Sir John Savill (Melbourne Academic Centre for Health) for comments on the manuscript.

Funding statement
This research was partly funded by Community Jameel and the Imperial President's Excellence Fund and by a UKRI-DHSC COVID-19 Rapid Response Rolling Call (MR/V027638/1) (to JEP). We also acknowledge a contribution from UKRI/NIHR through the UK Coronavirus Immunology Consortium (UK-CIC) and the National Institute for Health Research (NIHR) Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. JEP is supported by UKRI Innovation Fellowship at Health Data Research UK (MR/S004068/2). DCT is supported by a Stage 2 Wellcome-Beit Prize Clinical Research Career Development Fellowship (20661206617/A/17/Z and 206617/A/17/A) and the Sidharth Burman endowment. MCP is a Wellcome Trust Senior Fellow in Clinical Science (212252/Z/18/Z). NM-T and ES are supported by Wellcome Trust and Imperial College London Research Fellowships, and CLC by an Auchi Clinical Research Fellowship. PDWK is supported by the UK Medical Research Council (MC_UU_00002/13).

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Competing interests:
None of the authors have any patents (planned, pending or issued) relevant to this work. Dr. McAdoo reports personal fees from Celltrion, Rigel, GSK and Cello, unrelated to the submitted work. Dr Peters has received travel and accommodation expenses and hospitality

850   from Olink to speak at Olink-sponsored academic meetings. None of the other authors have
851   any competing interests.
852
853

Figure Legends

Figure 1 – Study design.
a) Schematic representing a summary of the patient cohorts, sampling and the major analyses. Blue and red stick figures represent outpatients and hospitalised patients, respectively. **b)** Timing of serial blood sampling in relation to clinical course of COVID-19 (subcohort A). Black asterisks indicate when samples were obtained. Three patients were already in hospital prior to COVID-19 diagnosis (indicated by red bars).

Figure 1 figure supplement 1 – Baseline characteristics of subcohort A.
The number of COVID-19 positive and negative patients in subcohort A (plasma), stratified by: **a)** sex, **b)** age, and **c)** ethnicity. **d)** Serial samples obtained for COVID-19 patients.

Figure 2 – Principal components analysis.
PC = principal component. Each point represents a sample. Colouring indicates COVID-19 status. The directions and relative sizes of the 6 largest PC loadings are plotted as arrows (middle column).
a-b) Subcohort A. Due to serial sampling, there are multiple samples for most patients. The proportion of variance explained in subcohort A by each PC is shown in parentheses on the axis labels. **c-d**) Subcohort B. Samples are projected into the PCA coordinates from subcohort A.

Figure 2 figure supplement 1 – Principal components analysis in relation to clinical severity.
a) Colouring indicates WHO severity at time of sampling. **b)** Colouring indicates overall clinical course (indicated by peak WHO severity) for the patient from which that sample was taken.

Figure 2 figure supplement 2 – Principal components analysis in relation to assay plate.
Principal components analysis of the subcohort A coloured by plate.

Figure 3 – Identification of dysregulated proteins.
a) Proteins up-regulated (red) or down-regulated (blue) in COVID-19 positive patients versus negative ESKD patients. n= 256 plasma samples from 55 COVID-19 positive patients, versus n= 51 ESKD controls (1 sample per control patient).
b) Proteins associated with disease severity. Associations of protein levels against WHO severity score at the time of sampling. Linear gradient indicates the effect size. A positive effect size (red) indicates that an increase in protein level is associated with increasing disease severity and a negative gradient (blue) the opposite. n= 256 plasma samples from 55 COVID-19 positive patients. For **a)** and **b)**: P-values from linear mixed models after Benjamini-Hochberg adjustment; significance threshold= 5% FDR; dark-grey = non-significant.
c) Heatmap showing protein levels for selected proteins with strong associations with severity. Each column represents a sample (n=256 COVID-19 samples and 51 non-infected samples). Each row represents a protein. Proteins are annotated using the symbol of their encoding gene. For the purposes of legibility, not all significantly associated proteins are shown; the heatmap is limited to the 17% most up- or down-regulated proteins (by effect size) of those with a significant association. Proteins are ordered by hierarchical clustering. Samples are ordered by WHO severity at the time of blood sample ('Severity'). 'Overall course' indicates the peak WHO severity over the course of the illness.

Figure 3 figure supplement 1. Differential abundance analysis between ESKD patients with and without COVID-19.

Heatmap showing selected proteins with the largest fold changes in differential abundance analysis (subcohort A). As for Figure 3, the heatmap is limited to the 17% most up- or down-regulated proteins (by fold change) of those with a significant association.

Figure 3 figure supplement 2 – Permutation analysis to estimate the null distribution.
Histogram showing the distribution of the number of associations declared significant (FDR 5%) after random permutation of class labels (100,000 replications). **a)** the COVID-19 +ve versus -ve differential abundance analysis. **b)** the COVID-19 severity differential abundance analysis. The vertical red line denotes the number of proteins we found significant in the analysis with the true sample labels.

Figure 3 figure supplement 3 – Sensitivity analyses adjusting for diabetes status and cause of ESKD.
As sensitivity analyses, the COVID-19 positive versus negative differential abundance regressions were repeated adding diabetes status (a-b) and cause of ESKD (c-d) as additional covariates. The basic model included age, sex and ethnicity as covariates. Each point represents a protein. A comparison of -log10 p-values and effect sizes are shown for all 436 proteins. r indicates Pearson's correlation coefficient.

Figure 3 figure supplement 4 – Sensitivity analysis adjusting for time since last haemodialysis.
Comparison of results obtained with and without adding time since last haemodialysis as an additional covariate to the regression models. **a-b)** COVID-19 positive versus negative differential expression analysis. **c-d)** Severity analysis. Each point represents a protein. r indicates Pearson's correlation coefficient.

Figure 4 – Validation.
a) Overlap between the significant associations in the differential abundance analysis between ESKD patients with and without COVID-19 in subcohorts A and B. 5% FDR was used as the significance threshold in both analyses.
b) Comparison of estimated effect sizes for all 436 proteins in the differential abundance analyses (COVID-19 positive versus negative) in subcohort A and B. Each point represents a protein. Pearson's r is shown.
Differential abundance analyses were performed using linear mixed models. Subcohort A analysis (plasma samples): 256 samples from 55 COVID-19 patients versus 51 non-infected patient samples (single time-point). Subcohort B (serum samples): 52 samples from 55 COVID-19 patients and 11 non-infected patient samples (single time-point).

Figure 4 figure supplement 1– Comparison with the report of Filbin *et al.* [16]
Comparison of log2 fold change for COVID-19 positive versus negative ESKD patients in our study versus COVID-19 positive versus negative respiratory distress patients in the report by Filbin *et al.* [16]. Colours indicate whether a protein was significantly differentially abundant in each study. Pearson's r is shown.

Figure 5 - Selected proteins strongly associated with COVID-19 severity.
Violin plots showing distribution of plasma protein levels according to COVID-19 status at the time of blood draw. Boxplots indicate median and interquartile range. n=256 samples from 55 COVID-19 patients and 51 samples from non-infected patients. WHO severity indicates the clinical severity score of the patient at the time the sample was taken. Mild n=135 samples; moderate n=77 samples; severe n=29 samples; critical n= 15 samples. Upper panel: monocyte chemokines. Middle panel: markers of epithelial injury. Lower panel: 2 neutrophil proteases, and IL6.

Figure 6 – comparison of proteins differentially expressed in COVID-19 with those associated with clinical severity.

a) Overlap between the proteins significantly differentially expressed in COVID-19 (n=256 COVID-19 samples and 51 non-infected samples) versus those associated with severity (within cases analysis, n=256 samples) (subcohort A). 5% FDR was used as the significant cut-off in both analyses.

b) Comparison of effect sizes for each protein in the COVID-19 positive versus negative analysis (x-axis) and severity analysis (y-axis). Each point represents a protein. Pearson's r is shown.

c) Examples of proteins specifically associated with severity but not significantly differentially abundant in the comparison of all cases versus controls. Violin plots showing distribution of plasma protein levels according to COVID-19 status at the time of blood draw. Boxplots indicate median and interquartile range. n=256 samples from 55 COVID-19 patients and 51 samples from non-infected patients. WHO severity indicates the clinical severity score of the patient at the time the sample was taken. Mild n=135 samples; moderate n=77 samples; severe n=29 samples; critical n= 15 samples.

Figure 7 – Prediction of severe COVID-19 and death.
a) The 12 most important proteins for predicting overall clinical course (defined by peak COVID-19 WHO severity) using Random Forests supervised learning. If a variable is important for prediction, it is likely to appear in many decision trees (number of trees) and be close to the root node (i.e. have a low minimal depth). The mean minimal depth across all trees (white box) was used as the primary feature selection metric.
b) Proteins that are significant predictors of death (Benjamini-Hochberg adjusted p <0.05). n=256 samples from 55 COVID-19 positive patients, of whom 9 died. Risk coefficient estimates are from a joint model. Bars indicate 95% confidence intervals. For proteins with a positive risk coefficient, a higher concentration corresponds to a high risk of death, and vice versa for proteins with negative coefficients.

Figure 7 figure supplement 1 – proteins associated with risk of death: correlation to clinical severity and clinical laboratory measurements.
Proteins significantly associated with risk of death (5% FDR) are shown. The estimated effect size from the linear mixed model testing association with severity are also shown. Correlations between protein levels and contemporaneous clinical laboratory marker values were calculated using rmcorr [55] for each of the proteins significant (5% FDR) in the joint model. The rows and columns of the clinical marker correlation matrix are ordered by hierarchical clustering.

Figure 8 – Associations of clinical laboratory markers with plasma proteins.
Proteins that are positively (red) or negatively (blue) associated with clinical laboratory parameters (5% FDR). P-values from differential abundance analysis using linear mixed models after Benjamini-Hochberg adjustment. Dark-grey = non-significant. Two associations were found for d-dimer (not shown- see **Supplementary File 1g**).

Figure 9 – Modelling of temporal protein trajectories.
The top 18 proteins displaying the most significantly (5% FDR) different longitudinal trajectories between patients with a mild or moderate (n= 28) versus severe or critical (n= 27) overall clinical course (defined by peak WHO severity). Means and 95% confidence intervals for each group, predicted using linear mixed models (**Methods**), are plotted. The remainder of significant proteins are shown in **Figure 9 figure supplement 1.** Individual data points are shown in **Figure 9 figure supplement 2**.

Figure 9 figure supplement 1– display of modelled temporal trajectories for other proteins with a significant time x severity interaction. Proteins significant at 5% FDR but not shown in Figure 9 are displayed here.

1017 **Figure 9 figure supplement 2– raw data points for modelling of temporal protein**
1018 **trajectories.** The 8 most significant proteins from Figure 9 are displayed.
1019
1020
1021

1022

1023

	COVID-19 positive ESKD patients (n=55)			ESKD controls (n=51)
	Overall	Patients with peak severity mild or moderate (n=28)	Patient with peak severity severe or critical (n=27)	
Age				
Median	72.2	73.4	68.5	70.1
(Q1-Q3)	62.5-77.3	65.5-76.4	61.8-78.8	62.2-75.1
Sex				
M	39 (70.9%)	18 (64.3%)	21 (77.8%)	36 (70.6%)
F	16 (29.1%)	10 (35.7%)	6 (22.2%)	15 (29.4%)
Ethnicity				
White	16 (29.1%)	5 (17.9%)	11 (40.7%)	13 (25.5%)
Black	8 (14.5%)	5 (17.9%)	3 (11.1%)	8 (15.7%)
South Asian	18 (32.7%)	10 (35.7%)	8 (29.6%)	20 (39.2%)
Asian (other)	4 (7.3%)	1 (3.6%)	3 (11.1%)	3 (5.9%)
Other	9 (16.4%)	7 (25.0%)	2 (7.4%)	7 (13.7%)
Diabetes	34 (61.8%) *	16 (57.1%)	18 (66.7%)	24 (47.1%) *
Current Smoker	1 (1.8%)	1 (3.6%)	0	0
ESKD Cause				
DN	29 (52.7%)	14 (50.0%)	15 (55.6%)	20 (39.2%)
Genetic	1 (1.8%)	1 (3.6%)	0	1 (2.0%)
GN	3 (5.5%)	1 (3.6%)	2 (7.4%)	9 (17.6%)
HTN/Vascular	5 (9.1%)	3 (5.5%)	2 (7.4%)	7 (13.7%)
Other	8 (14.5%)	5 (17.9%)	3 (11.1%)	4 (7.8%)
Unknown	9 (16.4%)	4 (14.3%)	5 (18.5%)	10 (19.6%)
Hospitalisation due to COVID-19†	33 (60%)	6 (21.4%)	27 (100%)	N/A
Fatal COVID-19	9 (16.3%)	0 (0%)	9 (33.3%)	N/A

1024

1025 **Table 1 – Characteristics of subcohort A.**
1026 GN = Glomerulonephritis. HTN = Hypertension. IQR = inter-quartile range. 'South Asian' represents
1027 individuals with Indian, Pakistani or Bangladeshi ancestry. Subsets defined according to peak WHO
1028 severity over the course of the illness. N/A = not applicable.
1029 *One patient had type 1 diabetes, the remainder type 2.
1030

1031

	COVID-19 positive ESKD patients (n=46)	COVID-19 negative ESKD controls (n=11)*
Age		
Median	64.3	71.6
(Q1-Q3)	60.3-73.0	(61.7-73.9)
Sex		
M	32 (69.6%)	8 (72.3%)
F	14 (30.4%)	3 (27.3%)
Ethnicity		
White	11 (23.9%)	3 (27.3%)
Black	8 (17.4%)	3 (27.3%)
South Asian	12 (26.1%)	3 (27.3%)
Asian (other)	7 (15.2%)	0
Other	8 (17.4%)	2 (18.2%)
Diabetes	29 (63.0%)	6 (54.5%)
Current Smoker	2 (4.3%)	0 (%)
ESKD Cause		
DN	19 (41.3%)	5 (45.5%)
Genetic	1 (2.2%)	0
GN	7 (15.2%)	1 (9.1%)
HTN/Vascular	3 (6.5%)	1 (9.1%)
Other	3 (6.5%)	2 (18.2%)
Unknown	13 (28.3%)	2 (18.2%)
Hospitalisation due to COVID-19	41 (89.1%)	N/A
Severe or critical COVID-19	33 (71.7%)	N/A
Fatal COVID-19	9 (19.6%)	N/A

1032
1033 **Table 2 – Characteristics of subcohort B.**
1034 GN = Glomerulonephritis. HTN = Hypertension. IQR = inter-quartile range. 'South Asian' represents
1035 individuals with Indian, Pakistani or Bangladeshi ancestry. Subsets defined according to peak WHO
1036 severity over the course of the illness. N/A = not applicable. *These 11 controls are a subset of the
1037 control patients used in subcohort A.
1038
1039

Supplementary File 1 table legends.

Supplementary File 1a. Protein Annotation.
List of the 436 proteins measured. GeneID = gene symbol of the gene encoding the protein (used as the main identifier in the manuscript); UniProt = UniProt ID; Olink Assay Name = protein id used by Olink; Protein Name = full protein name; Panel name = the name of the 92 protein multiplex Olink panel on which the protein was measured.

Supplementary File 1b. Enrichment of Reactome terms for the entire set of proteins measured.
The results of enrichment testing for genes corresponding to all 436 measured proteins against the background of the genome. The analysis was performed against the Reactome pathways using string-db. The list of Reactome terms is ordered by the number of proteins associated with the term.

Supplementary File 1c. Differential abundance analysis for COVID-19 positive vs negative ESKD patients in subcohort A and B.
Summary statistics for all 436 proteins are shown. Pvalue = nominal p-value from linear mixed model. Adjusted Pvalue = p-values after Benjamini-Hochberg correction. Fold change = estimated fold change from regression coefficient. Proteins are ordered based on results in subcohort A: first by whether they are significant or not (at 5% FDR), then by fold change (from positive to negative). Note the associations are not ordered by p-value so strong associations do not necessarily appear at the top of the table. Significant adjusted p-values are coloured in green and non-significant in grey. Estimated fold changes are coloured in a gradient from red to blue for up or downregulated in COVID-19 +ve versus –ve, respectively.
Sample size for subcohort A: n= 256 plasma samples from 55 COVID-19 positive ESKD patients, versus n= 51 ESKD controls (1 sample per control patient).
Sample size for subcohort B: 52 samples from 55 COVID-19 patients and 11 non-infected patient samples (single time-point).

Supplementary File 1d Associations of proteins and COVID-19 severity (subcohort A).
Summary statistics for all 436 proteins are shown. Pvalue = nominal p-value from linear mixed model. Adjusted Pvalue = p-values after Benjamini-Hochberg correction. Fold change = estimated fold change from regression coefficient. Proteins are ordered first by whether they are significant or not (at 5% FDR), then by linear gradient (effect size) from positive to negative. Note the associations are not ordered by p-value so strong associations do not necessarily appear at the top of the table.

Supplementary File 1e. **Predictors of clinical course from Random Forests.**
Importance metrics for each protein for prediction according to a random forest model trained to predict current or future severe/critical disease using the first sample of each patient. Proteins are ordered by mean minimal depth across all trees – this was used as the primary importance metric.

Supplementary File 1f. Proteomic predictors of fatal COVID-19.
Summary statistics from joint models for fatal disease. Results for all 436 proteins are shown. "Is significant" indicates significance (green) or not (grey) at 5% FDR. The association coefficient for each protein indicates the direction and magnitude of the estimated log relative risk for death (red indicates higher protein levels increase risk of death, blue the opposite). 95% confidence intervals are plotted.

Supplementary File 1g. Associations of proteins and clinical laboratory measurements.
Clinical variable = clinical lab tests: white cell count, lymphocyte count, neutrophil count, monocyte count, C-reactive protein, ferritin, d-dimer, troponin.

Supplementary File 1h. Longitudinal proteomic profiling with linear mixed models.
Summary statistics from the linear mixed models used to identify proteins with differential temporal trajectories between mild/moderate (n=28) and severe/critical COVID-19 patients (n=27). Summary statistics for all 436 proteins are shown.

Pvalue = nominal p-value from linear mixed model for the interaction term between time from symptom onset (days) and overall WHO severity (as a binary variable: mild-moderate or severe-critical).

Adjusted Pvalue = p-values after Benjamini-Hochberg correction. "Is significant" indicates significance (green) or not (grey) at 5% FDR.

Supplementary File 1i. Comparison to other proteomic studies of COVID-19 positive vs negative patients.
Proteins that were differentially abundant in COVID-19 +ve vs -ve patients in our data are listed (5% FDR). TRUE indicates that the protein was reported as differentially abundant in the relevant previous proteomic study. The final column summarises whether the association was previously reported in any of the 4 studies. We have not harmonised significance thresholds between studies: we simply report whether the authors declared the protein significant by the threshold of their study.

Supplementary File 1j. Comparison to other proteomic studies of COVID-19 severity.
Proteins that were associated with severity in our data are listed (5% FDR). TRUE indicates that the protein was reported as associated with severity in the relevant previous proteomic study. The final column summarises whether the association was previously reported in any 1 or more of the 4 studies. We have not harmonised significance thresholds between studies: we simply report whether the authors declared the protein significant by the threshold of their study.

Results are shown for all 436 proteins against all 8 lab measurements.

Adjusted p-value = p-value from linear mixed model after Benjamini-Hochberg correction.

Gradient indicates effect size and direction. A positive gradient (red) indicates higher concentrations of proteins are associated with higher clinical laboratory measurements. "Is significant" indicates significance (green) or not (grey) at 5% FDR.

Contemporaneous clinical laboratory tests were not available for all plasma samples. The proportion of samples for which contemporaneous lab tests were available were: white cell count 66%, neutrophils 66%, monocytes 66%, lymphocytes 66%, CRP 64%, ferritin 36%, troponin 35%, d-dimer 30%.

Supplementary File 1k. Per protein correlations between plasma and serum levels derived from the same blood sample in 11 COVID-19 negative ESKD patients.
Plasma and serum were taken from 11 non-infected ESKD patients that were measured in both subcohort A (plasma) and B (serum). Pearson's r was calculated for the 11 paired measurements for each protein. Proteins are ordered by r value; this column is coloured from red to blue for positive and negative r values, respectively. 95% confidence intervals are reported. We also report the variance of the NPX levels for each protein in plasma and in serum.

References

1.	Horby, P. *et al.* Dexamethasone in Hospitalized Patients with Covid-19 — Preliminary Report. *N. Engl. J. Med.* **0**, NEJMoa2021436 (2020) doi:10.1056/NEJMoa2021436.

2.	Gordon, A. C. *et al.* Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19 - Preliminary report. *medRxiv* (2021) doi:10.1101/2021.01.07.21249390.

3.	Suhre, K., McCarthy, M. I. & Schwenk, J. M. Genetics meets proteomics: perspectives for large population-based studies. *Nat. Rev. Genet.* (2020) doi:10.1038/s41576-020-0268-2.

4.	Williamson, E. J. *et al.* Factors associated with COVID-19-related death using OpenSAFELY. *Nature* **584**, 430–436 (2020) doi:10.1038/s41586-020-2521-4.

5.	Docherty, A. B. *et al.* Features of 16,749 hospitalised UK patients with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol. *medRxiv* 2020.04.23.20076042 (2020) doi:10.1101/2020.04.23.20076042.

6.	Corbett, R. W. *et al.* Epidemiology of COVID-19 in an Urban Dialysis Center. *J. Am. Soc. Nephrol.* **31**, 1815–1823 (2020) doi:10.1681/ASN.2020040534.

7.	Ng, J. H. *et al.* Outcomes of patients with end-stage kidney disease hospitalized with COVID-19. *Kidney Int.* (2020) doi:10.1016/j.kint.2020.07.030.

8.	Valeri, A. M. *et al.* Presentation and outcomes of patients with ESKD and COVID-19. *J. Am. Soc. Nephrol.* **31**, 1409–1415 (2020) doi:10.1681/ASN.2020040470.

9.	Hastie T, Tibshirani R, F. J. High-Dimensional Problems: p ≫ N. in *The Elements of Statistical Learning* 649–698 (Springer Series in Statistics, 2001).

10.	Ibrahim, J. G., Chu, H. & Chen, L. M. Basic concepts and methods for joint models of longitudinal and survival data. *J. Clin. Oncol. Off. J. Am. Soc. Clin. Oncol.* **28**, 2796–2801 (2010) doi:10.1200/JCO.2009.25.0654.

11.	Rizopoulos, D. JM: An R package for the joint modelling of longitudinal and time-to-event data. *J. Stat. Softw.* **35**, 1–33 (2010) doi:10.18637/jss.v035.i09.

12.	Guan, W. *et al.* Clinical characteristics of coronavirus disease 2019 in China. *N. Engl. J. Med.* **382**, 1708–1720 (2020) doi:10.1056/NEJMoa2002032.

13.	Shen, B. *et al.* Proteomic and Metabolomic Characterization of COVID-19 Patient Sera. *Cell* **182**, 59-72.e15 (2020) doi:10.1016/j.cell.2020.05.032.

14.	Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* **584**, 463–469 (2020) doi:10.1038/s41586-020-2588-y.

15.	Arunachalam, P. S. *et al.* Systems biological assessment of immunity to mild versus severe COVID-19 infection in humans. *Science (80-.).* **6261**, 1–18 (2020) doi:10.1126/science.abc6261.

16.	Filbin, M. R. *et al.* Plasma proteomics reveals tissue-specific cell death and mediators of cell-cell interactions in severe COVID-19 patients. *bioRxiv* 2020.11.02.365536 (2020) doi:10.1101/2020.11.02.365536.

17.	COVID-19 Data. *The Renal Association* https://renal.org/health-professionals/covid-19/covid-19-data (2020).

18.	Scottish Renal Registry COVID-19 report. *Public Health Scotland* https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/scottish-renal-registry-covid-19-report/ (2020).

19. Eiselt, J. *et al.* Previous Vaccination and Age are More Important Predictors of
Immune Response to Influenza Vaccine than Inflammation and Iron Status in Dialysis
Patients. *Kidney Blood Press. Res.* **41**, 139–147 (2016) doi:10.1159/000443416.

20. Girndt, M., Sester, U., Sester, M., Kaul, H. & Köhler, H. Impaired cellular immune
function in patients with end-stage renal failure. *Nephrology, dialysis,*
transplantation : official publication of the European Dialysis and Transplant
Association - European Renal Association vol. 14 2807–2810 (1999)
doi:10.1093/ndt/14.12.2807.

21. Sarnak, M. J. & Jaber, B. L. Mortality caused by sepsis in patients with end-stage
renal disease compared with the general population. *Kidney Int.* **58**, 1758–1764
(2000) doi:10.1111/j.1523-1755.2000.00337.x.

22. Alexiewicz, J. M., Smogorzewski, M., Fadda, G. Z. & Massry, S. G. Impaired
phagocytosis in dialysis patients: studies on mechanisms. *Am. J. Nephrol.* **11**, 102–
111 (1991) doi:10.1159/000168284.

23. Massry, S. & Smogorzewski, M. Dysfunction of polymorphonuclear leukocytes in
uremia: role of parathyroid hormone. *Kidney Int. Suppl.* **78**, S195-6 (2001)
doi:10.1046/j.1523-1755.2001.59780195.x.

24. Girndt, M., Sester, M., Sester, U., Kaul, H. & Köhler, H. Molecular aspects of T- and
B-cell function in uremia. *Kidney Int. Suppl.* **78**, S206-11 (2001) doi:10.1046/j.1523-
1755.2001.59780206.x.

25. Meier, P., Dayer, E., Blanc, E. & Wauters, J.-P. Early T cell activation correlates with
expression of apoptosis markers in patients with end-stage renal disease. *J. Am.*
Soc. Nephrol. **13**, 204–212 (2002).

26. Betjes, M. G. H. Immune cell dysfunction and inflammation in end-stage renal
disease. *Nat. Rev. Nephrol.* **9**, 255–265 (2013) doi:10.1038/nrneph.2013.44.

27. Boattini, M. *et al.* Influenza and respiratory syncytial virus infections in the oldest-old
continent. *Eur. J. Clin. Microbiol. Infect. Dis. Off. Publ. Eur. Soc. Clin. Microbiol.* **39**,
2085–2090 (2020) doi:10.1007/s10096-020-03959-9.

28. Prasad, N. *et al.* Respiratory Syncytial Virus–Associated Hospitalizations Among
Adults With Chronic Medical Conditions. *Clin. Infect. Dis.* (2020)
doi:10.1093/cid/ciaa730.

29. Usvyat, L. A. *et al.* Seasonal variations in mortality, clinical, and laboratory parameters
in hemodialysis patients: a 5-year cohort study. *Clin. J. Am. Soc. Nephrol.* **7**, 108–
115 (2012) doi:10.2215/CJN.03880411.

30. Carrero, J. J. & Stenvinkel, P. Inflammation in end-stage renal disease--what have we
learned in 10 years? *Semin. Dial.* **23**, 498–509 (2010) doi:10.1111/j.1525-
139X.2010.00784.x.

31. Naseeb, U. *et al.* Complementary LC-MS/MS Proteomic Analysis of Uremic Plasma
Proteins. *J. Coll. Physicians Surg. Pak.* **25**, 606–609 (2015).

32. Christensson, A. *et al.* The Impact of the Glomerular Filtration Rate on the Human
Plasma Proteome. *Proteomics. Clin. Appl.* **12**, e1700067 (2018)
doi:10.1002/prca.201700067.

33. Lind, L. *et al.* Longitudinal effects of aging on plasma proteins levels in older adults -
associations with kidney function and hemoglobin levels. *PLoS One* **14**, e0212060
(2019) doi:10.1371/journal.pone.0212060.

34.   Wu, C. *et al.* Risk Factors Associated With Acute Respiratory Distress Syndrome and
1229   Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China.
1230   *JAMA Intern. Med.* **180**, 934–943 (2020) doi:10.1001/jamainternmed.2020.0994.

1231   35.   Li, X. *et al.* Risk factors for severity and mortality in adult COVID-19 inpatients in
1232   Wuhan. *J. Allergy Clin. Immunol.* **146**, 110–118 (2020) doi:10.1016/j.jaci.2020.04.006.

1233   36.   Furlow, B. COVACTA trial raises questions about tocilizumab's benefit in COVID-19.
1234   *Lancet Rheumatol.* **2**, e592 (2020) doi:10.1016/S2665-9913(20)30313-1.

1235   37.   Gupta, R. K. *et al.* Development and validation of the ISARIC 4C Deterioration model
1236   for adults hospitalised with COVID-19: a prospective cohort study. *Lancet. Respir.*
1237   *Med.* (2021) doi:10.1016/S2213-2600(20)30559-2.

1238   38.   Schutyser, E., Struyf, S. & Van Damme, J. The CC chemokine CCL20 and its
1239   receptor CCR6. *Cytokine Growth Factor Rev.* **14**, 409–426 (2003) doi:10.1016/s1359-
1240   6101(03)00049-2.

1241   39.   Saha, S. K. *et al.* KRT19 directly interacts with β-catenin/RAC1 complex to regulate
1242   NUMB-dependent NOTCH signaling pathway and breast cancer properties.
1243   *Oncogene* **36**, 332–349 (2017) doi:10.1038/onc.2016.221.

1244   40.   Maecker, H. *et al.* TWEAK attenuates the transition from innate to adaptive immunity.
1245   *Cell* **123**, 931–944 (2005) doi:10.1016/j.cell.2005.09.022.

1246   41.   Burkly, L. C., Michaelson, J. S., Hahm, K., Jakubowski, A. & Zheng, T. S. TWEAKing
1247   tissue remodeling by a multifunctional cytokine: role of TWEAK/Fn14 pathway in
1248   health and disease. *Cytokine* **40**, 1–16 (2007) doi:10.1016/j.cyto.2007.09.007.

1249   42.   Henning, R. J., Bourgeois, M. & Harbison, R. D. Poly(ADP-ribose) Polymerase
1250   (PARP) and PARP Inhibitors: Mechanisms of Action and Role in Cardiovascular
1251   Disorders. *Cardiovasc. Toxicol.* **18**, 493–506 (2018) doi:10.1007/s12012-018-9462-2.

1252   43.   Rouleau, M., Patel, A., Hendzel, M. J., Kaufmann, S. H. & Poirier, G. G. PARP
1253   inhibition: PARP1 and beyond. *Nat. Rev. Cancer* **10**, 293–301 (2010)
1254   doi:10.1038/nrc2812.

1255   44.   Corren, J. New Targeted Therapies for Uncontrolled Asthma. *J. allergy Clin. Immunol.*
1256   *Pract.* **7**, 1394–1403 (2019) doi:10.1016/j.jaip.2019.03.022.

1257   45.   Galijasevic, S. The development of myeloperoxidase inhibitors. *Bioorg. Med. Chem.*
1258   *Lett.* **29**, 1—7 (2019) doi:10.1016/j.bmcl.2018.11.031.

1259   46.   Vergunst, C. E. *et al.* Modulation of CCR2 in rheumatoid arthritis: a double-blind,
1260   randomized, placebo-controlled clinical trial. *Arthritis Rheum.* **58**, 1931–1939 (2008)
1261   doi:10.1002/art.23591.

1262   47.   Haringman, J. J. *et al.* A randomized controlled trial with an anti-CCL2 (anti-monocyte
1263   chemotactic protein 1) monoclonal antibody in patients with rheumatoid arthritis.
1264   *Arthritis Rheum.* **54**, 2387–2392 (2006) doi:10.1002/art.21975.

1265   48.   Rodriguez, L. *et al.* Systems-Level Immunomonitoring from Acute to Recovery Phase
1266   of Severe COVID-19. *Cell reports. Med.* **1**, 100078 (2020)
1267   doi:10.1016/j.xcrm.2020.100078.

1268   49.   Uhlen, M. *et al.* Tissue-based map of the human proteome. *Science (80-.).* **347**,
1269   1260419–1260419 (2015) doi:10.1126/science.1260419.

1270   50.   Szklarczyk, D. *et al.* STRING v11: Protein-protein association networks with increased
1271   coverage, supporting functional discovery in genome-wide experimental datasets.
1272   *Nucleic Acids Res.* **47**, D607–D613 (2019) doi:10.1093/nar/gky1131.

51.   Gandolfo, L. C. & Speed, T. P. RLE plots: Visualizing unwanted variation in high
dimensional data. *PLoS One* **13**, 1–9 (2018) doi:10.1371/journal.pone.0191629.

52.   Kuhn, M. Building predictive models in R using the caret package. *J. Stat. Softw.* **28**,
1–26 (2008) doi:10.18637/jss.v028.i05.

53.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models
Using lme4. *J. Stat. Softw.* **67**, (2015) doi:10.18637/jss.v067.i01.

54.   Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in
Linear Mixed Effects Models. *J. Stat. Softw.* **82**, (2017) doi:10.18637/jss.v082.i13.

55.   Bakdash, J. Z. & Marusich, L. R. Repeated measures correlation. *Front. Psychol.* **8**,
1–13 (2017) doi:10.3389/fpsyg.2017.00456.

56.   Leo,   B.   Random   forests.   *Mach.   Learn.*   **45**,   5–32   (2001)
doi:10.1023/A:1010933404324.

57.   Perperoglou, A., Sauerbrei, W., Abrahamowicz, M. & Schmid, M. A review of spline
function procedures in R. *BMC Med. Res. Methodol.* **19**, 1–16 (2019)
doi:10.1186/s12874-019-0666-3.

1288

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 1 figure supplement 1

Figure 2 figure supplement 1

Figure 2 figure supplement 2

Figure 3 figure supplement 1

Figure 3 figure supplement 2

Figure 3 figure supplement 3

Figure 3 figure supplement 4

COVID-19 +ve vs -ve differential abundance analysis

COVID-19 Severity differential abundance analysis

Figure 4 figure supplement 1

Figure 7 figure supplement 1

Figure 9 figure supplement 1

Figure 9 figure supplement 2