

25 **Abstract**

26

27 **Objectives** To analyse the changes in reported frequency of retinal detachment admissions and
28 vitreo-retinal surgery procedures performed between 2000-2018 in England. To obtain information
29 useful to contribute towards the planning of service delivery.

30 **Methods** Analysis of England's Hospital Episode Statistics from the Health and Social Care
31 Information Centre and population data from the United Kingdom's Office for National Statistics.

32 **Results** Episodes of "retinal detachments with breaks" increased year on year from 3,447
33 (7.0/100M) in 2000 to 10,971 (19.7/100M) in 2018 ($p < 0.001$), whereas records of "tractional retinal
34 detachment" increased from 290 (0.6/100M) to 910 (1.6/100M) in the same period ($p < 0.0001$). The
35 number of reported pars plana vitrectomies irrespective of indication increased over fourfold from
36 5,761 to 26,900 ($p < 0.0001$), while the number of scleral buckling records decreased by two thirds
37 from 2,897 to 780 ($p < 0.0001$). During the same period the population of England increased from
38 49.2 million-55.6 million, proportionally at a slower rate than that for recorded hospital episodes.

39 **Conclusions** The frequency of admissions to hospital for surgically treated retinal detachment seems
40 to have been increasing significantly since 2000. This effect is more marked in cases of
41 rhegmatogenous retinal detachment. This may be partially explained by repeat surgery in cases of
42 recurrent retinal detachment. Other possible explanations may be increased incidence of disease
43 (due to increased rates of cataract surgery, increasing longevity and rates of myopia), improvements
44 in patient access, and increased public awareness. It is possible that this observation is due in part to
45 local changes in methodology of hospital coding.

46

47

48 Introduction

49 Rhegmatogenous retinal detachment (RRD) is a serious ophthalmic condition where the
50 neurosensory retina becomes separated from the retinal pigment epithelium through ingress of fluid
51 from a retinal break. Without treatment, RRD leads to blindness in the affected eye. Advances in
52 surgical treatment allow for the prevention of vision loss in most cases. Recent refinements in
53 surgical technique including self-sealing small gauge vitrectomy and wide-angle viewing systems
54 have significantly improved the ability of surgeons to treat RRD quickly and safely.

55

56 In the United Kingdom, the annual incidence of RRD is reported to have increased from 6.1-9.8 cases
57 per 100,000 people during the 1970's, to 11.8-17.9 cases per 100 000 people in the 1990s¹. A more
58 recent report estimated an incidence increase from 13.4/100,000 in the 1990s to 15.4/100,000 in
59 the 2010s². A meta-analysis of 5 studies carried out in Europe³ estimated the average incidence of
60 RD in the continent at 13.3/100,000.

61

62 In England, all instances of hospital admission, day case intervention and emergency attendances are
63 recorded in a database containing details of diagnoses and procedures called the Hospital Episode
64 Statistics (HES)⁴. Each instance of care is recorded as one episode, regardless if a given patient is
65 treated more than once for the same condition. Diagnoses are recorded using the World Health
66 Organisation International Classifications of Diseases version 10 (ICD-10) classification system and
67 procedures are coded with the NHS's Office for Population Census and Surveys Classification of
68 Surgical Operations version 4 (OPCS-4). These data are identified and coded from medical records
69 following the discharge of each patient from each episode. English NHS healthcare providers use
70 these data to claim refund from the regional funding bodies for the care they deliver. The same data
71 are also processed and used for other purposes, such as health services delivery planning and
72 research.

73

74 We set out to analyse England's HES to seek changes in trends of admission to healthcare facilities
75 for treatment of RRD from 2000 to 2018. We also set out to study changes in frequency and type of
76 vitreo-retinal operations carried out during this period. The purpose was to gain insight into the
77 epidemiology of surgical retinal disease with the aim to inform service planning and resource
78 deployment.

79

80 **Materials and Methods**

81 The publicly available HES Hospital Admitted Patient Care Activity records were downloaded from
82 the NHS Digital⁴ internet portal both for procedures and diagnosis. The reported year follows the UK
83 fiscal year which runs from April until March and is reported based on the year of period end,
84 therefore 2004 comprises the period from April 2003 until March 2004. England's population data
85 were obtained from the mid-year estimate from the Office for National Statistics⁵ corresponding to
86 the fiscal year where the estimate lies. For example, the mid-year estimate for 2010 was used to
87 calculate the rate of admission for the fiscal year 2010-11.

88 *Diagnoses*

89 The ICD-10 codes for retinal detachment with breaks (H33.0) and tractional retinal detachment
90 (TRD) (H33.4) were analysed separately. The code H33.0 explicitly excludes cases where the retinal
91 detachment was associated with diabetes which should allow for a targeted analysis of RRD. There is
92 no similar distinction within code H33.4 between diabetes-related tractional detachment and other
93 tractional pathology such as retinopathy of prematurity or proliferative vitreo-retinopathy.
94 Associated codes within the same H33.x root include codes for retinoschisis and retinal cysts H33.1,
95 serous retinal detachment H33.2, retinal breaks without detachment H33.3. We analysed changes in
96 H33.2 and H33.3 as comparators as they should be susceptible to variation due to population
97 change. The code for other retinal detachments H33.5 was not analysed.

98 *Procedures*

99 The OPCS-4 codes for interventions on the retina and the lens were studied. Records were analysed
100 from the database of main procedure descriptor for pars plana vitrectomy (C79.2) and cataract
101 surgery using insertion of prosthetic replacement for lens (C75.1) as recommended by the NHS
102 Classification Service. For scleral buckle operations we aggregated all codes describing the
103 implantation of a scleral indenting device or otherwise creating scleral indent on the retina: overlay

104 scleroplasty (C54.1), imbrication of sclera (C54.2), buckling of sclera and implant (C54.3), buckling of
105 sclera and local or encircling explant (C54.4), buckling of sclera (C54.5), other specified buckling
106 operations for attachment of retina (54.8) and unspecified buckling operations for attachment of
107 retina (54.9).

108 Additionally, as there is no nationally mandated coding record of pars plana vitrectomy in primary
109 position for all vitrectomy operations, we built a construct adding surgical manoeuvres which
110 require pars plana vitrectomy access to the vitreous cavity, to the vitrectomy code (C79.2).
111 Therefore the following codes were grouped with C79.2 and analysed as a construct: injection of
112 vitreous substitute into vitreous body NEC (C79.3), internal tamponade of retina using liquid (C79.6),
113 removal of internal tamponade agent from vitreous body (C79.7), other specified operations on
114 vitreous body (C79.8), peel of epiretinal fibroglial membrane (C80.1), peel of internal limiting
115 membrane (C80.2), delamination of epiretinal fibrovascular membrane (C80.3), segmentation of
116 epiretinal fibrovascular membrane (C80.4), removal of subretinal vascular membrane (C80.5),
117 removal of subretinal membrane NEC (C80.6), other specified operations on retinal membrane
118 (C80.8) unspecified operations on retinal membrane (C80.9), pigment epithelium translocation of
119 retina (C83.1), macular translocation three hundred and sixty degrees (C83.2), limited macular
120 translocation (C83.3), other specified translocation of retina (C83.8), unspecified translocation of
121 retina (C83.9), epiretinal dissection (C84.1), excision of lesion of retina NEC (C84.2), biopsy of lesion
122 of retina (C84.3), retinal vascular sheathotomy (C84.4), drainage of subretinal fluid through retina
123 (C84.5), retinotomy NEC (C84.6), other specified other operations on retina (C84.8). As laser therapy
124 can be administered either transpupillarily or with endolaser probes, the codes for laser intervention
125 were not included.

126 *Statistical Analysis and variable associations*

127 Descriptive and linear regression analysis was carried out for diagnostic and intervention variables.

128 Linear regression analysis was carried out for variables potentially associated with RD including

129 cataract surgery volumes. The ratio between buckling operations and admissions for RRD was
130 calculated.

131 Consent was not required as this is a study of a database in the public domain and no patient
132 identifiable information was used.

133

134 **Results**

135 *Diagnoses*

136 During the business year ending 2001 there were 3,447 admissions to hospitals in England for RRD
137 with breaks (Figure 1). This figure more than trebled with an average annual growth of 7.2% and a
138 slope of 464.7 cases per year to 10,971 in 2018 ($R = 0.98$, $R^2 = 0.97$, $p < 0.001$). TRDs during 2000-01
139 were recorded in 290 instances. These too increased steadily at an annual rate of 7.9%, more than
140 trebling in the study period to 910 ($R=0.93$, $R^2=0.87$, $p<0.0001$). By comparison, serous retinal
141 detachments were recorded with a frequency of 3,941 in 2000 but reduced slightly throughout the
142 study period with a regression slope of -32.1 ($R=-0.56$, $R^2 = 0.32$, $p=0.01$). Retinal breaks without a
143 detachment were recorded at 3,141 in 2000, dropping to 1,842 in 2004 and then remaining
144 relatively stable from 2006 with an average of 2433.1 per annum ($R=-0.25$, $R^2=0.06$, $p=0.93$).

145 Calculating the rate of admission for the mid-year population level showed 7.0 episodes per 100,000
146 population for RD with breaks during 2000-01 with year-on-year increases to 19.7/100,000 in 2017-
147 18 with a slope of 0.79 and $R=0.98$. The slopes for admissions for TRD (0.05/year), serous retinal
148 detachment (-0.12/year) and retinal breaks without detachment (-0.06/year) show little change
149 compared to population growth.

150 To estimate the effect of re-admission for recurrent retinal detachment, we assumed an average
151 RRD recurrence rate of 13.0%⁶ and a single operation to repair the recurrence. Deducting these from
152 the total of admissions to estimate the value of primary retinal detachments, the regression slope is
153 reduced to 404.3 cases per annum of growth (Figure 2). Assuming 2.5 operations to repair a
154 recurrent retinal detachment further lowers the slope to 312.6 from 2,318.1 cases in 2000 to 7,378.0
155 cases in 2018.

156 *Procedures*

157 During 2000-01 there were 5,761 pars plana vitrectomy procedures carried out and recorded as
158 main intervention. Adding the codes of other surgical manoeuvres recorded as main intervention
159 and which obligatorily required pars plana vitrectomy, the total estimate of pars plana vitrectomy
160 rises to 8,777. Both figures have increased steadily with a yearly rate of 8.5% for vitrectomy (slope =
161 1,384.1, $R^2=0.97$, $p<0.0001$) and 6.9% for the composite of vitrectomy codes (slope=1,363.3,
162 $R^2=0.99$, $p<0.0001$). The final figure for 2017-18 is 26,900 for vitrectomy coded singly and 29,923 for
163 vitrectomy composite respectively (Figure 3). Similarly, for 2000-01 there were 2,897 scleral buckling
164 operations which decreased steadily at an average yearly rate of 7.0% to 780 cases with a slope of -
165 114.5 ($R^2=0.87$, $p<0.0001$)

166 *Associations*

167 The ratio of scleral buckle operations to RRD admissions changed from 84:100 in 2000-01 to 7.1:100
168 in 2017-18, at a rate of -0.04 buckling operation per admission for RRD per year. There was a strong
169 and statistically significant correlation between the number of cataract surgery procedures and
170 number of admissions for RRD with breaks ($r=0.932$, $r^2=0.868$, $p<0.0001$).

171 **Discussion**

172 We found that the frequency of admissions to hospital for the types of retinal detachment which are
173 treated by surgery (both TRD and RRD) increased steadily in England from 2000-2018. During the
174 same period, England's population increased from 49.2million to 54.3million, at a proportionally
175 slower rate than that for recorded hospital episodes. It is important to make the distinction between
176 incidence, prevalence and frequency of a disease. As the HES collect episodes, it is not possible to
177 determine neither incidence nor prevalence without obtaining the raw HES data to eliminate
178 duplicates as any episode of care for the same condition will be counted twice. However, given that
179 the methodology of data collection for HES can be assumed to be reasonably constant, trend
180 analysis for defined variables is useful in gaining understanding of the behaviour of healthcare
181 activity.

182 *Rhegmatogenous retinal detachment*

183 There have been several national and regional population studies published over recent years which
184 have estimated the annual incidence of RRD, reporting results ranging from 9.7 to
185 20.7/100,000/year in developed Western and Asian nations^{1-2,7-14}. Broadly, two types of
186 methodology were used: hospital records analysis and national reimbursement databases. Hospital
187 records analysis have been reported for defined populations where only a small number of hospitals
188 provide retinal detachment care. Manners⁷ analysed all admissions to Western Australian hospitals
189 between 2000-2013 reporting stable rates between 12.8 and 16.2/100,000 people. Also in Australia,
190 Howie⁸ reported rates for the island-state of Tasmania where a single surgeon provided care
191 between 2005 and 2010, finding a rate of 9.7/100,000 inhabitants. In Scotland⁹, a consortium
192 formed by 6 hospitals prospectively recorded all primary cases between 2007-2009 finding an
193 incidence of 12.05/100,000. A similar consortium formed in the Netherlands¹⁰ reported
194 18.2/100,000 cases during 2009. A further report from a single centre in the city of Odense in
195 Denmark¹¹ estimated an incidence of 20.7/100,000 however being a tertiary referral centre, it is

196 possible that it also treated patients from outside its reference area. This type of study is better
197 suited to define clinical variables and patient characteristics.

198 Analysis of national databases have also been reported studying recent years in England^{1,2},
199 Denmark¹², and Korea¹³. These are more likely to give more accurate estimates and allow to describe
200 time-trends over a period. In England, multiple database linkage analysis was carried out covering
201 the period from 1968 through to 2011. This study also sought to find associations with
202 socioeconomic and health status variables. Analysing the general three-digit code for retinal
203 detachment H33 without distinction, it estimated that the global incidence of all types of retinal
204 detachments increased from 13.4/100,000 in 1999 to 15.4/100,000 in 2011. A similar nationwide
205 study in Denmark¹² also found a steady increase of incidence from 2000 to 2011 which they linked to
206 an increase in rates of cataract surgery. The mean incidence over the period was reported at
207 13.7/100,000. In Korea¹³, a five-year study covering 2007 to 2011 found an incidence of
208 10.4/100,000 which remained stable. A further Taiwanese¹⁴ database covering 4% of the population
209 reported an estimated incidence of 16.4/100,000 cases from 2000-2012.

210 It is quite remarkable that in a period of 18 years, the rate for admissions for RRD in England went
211 from 7.0/100,000 to 19.7/100,000 people. Even when including an estimate for recurrent cases, this
212 represents a steep increase which almost doubles the figure of admissions to 13.7/100,000.

213 Several risk factors predisposing to RRD have been identified including male gender, increasing age,
214 previous cataract surgery, and myopia. Several of the database studies cited above have found that
215 age is an important risk factor for RRD^{2,12,14}. Shah found an almost 20-fold increased risk of RD in the
216 over 60s age group compared with the under 30s age group². As the average age of England's
217 population increases, so has the average age of admissions for RRD. While male gender is a
218 significant risk factor for RRD, there have been no recent significant changes in the gender
219 composition of England's population⁵. There are no accurate population-wide studies of the
220 prevalence of myopia in England. However, analysis of population cohorts has shown a sustained

221 worldwide increase in the prevalence of myopia particularly amongst young subjects. This effect is
222 more marked across Asia, Europe and the United States¹⁵. An eye with a refractive error of more
223 than -3 dioptres has a tenfold increased risk of RD compared to an emmetropic eye⁹. The increase in
224 the number of performed cataract operations can be partly attributed to population aging, and
225 hence a higher prevalence of cataracts. In addition, there has been an inclination to perform cataract
226 surgery at an earlier stage with the advent of phacoemulsification¹⁰. A review of the literature found
227 that the risk of RD following phacoemulsification is approximately ten times the risk for the general
228 population¹⁶.

229 It is interesting to note that the rate of retinal breaks remained relatively constant during the period
230 analysed. This observation is surprising because by definition RRDs occur after retinal tear
231 formation. Therefore if the natural rate of RD was increasing, it could be expected that the rate of
232 retinal tears would also have increase in parallel or faster. A potential explanation for this
233 observation is a proposed catastrophic behaviour of RRD development, where they would form
234 suddenly in “leaps and bounds”¹⁷, therefore only low-risk tears would be diagnosed prior to
235 developing into RD. Another potential explanation would be improvements in access to care for
236 patients both by improved referral pathways and by offering surgery to RDs which previously would
237 have been deemed inoperable. Further, we observed that when a correction was introduced to
238 account for estimated rates of recurrent RD, the slope of the growth in RD episodes became less
239 steep. We believe that it is likely that improved instrumentation has led to practitioners persevering
240 in achieving retinal attachment through multiple operations. Operating on recurrent RD would then
241 partially explain the marked increase in HES recorded episodes. It is worth noting that regardless of
242 the incidence of primary RD, a proportion of patients will unfortunately experience RD recurrences.
243 This fact needs to be considered when planning the delivery of service.

244 *Tractional retinal detachment*

245 In our study we also observed a big increase in TRD admissions. The prevalence of diabetes mellitus
246 increased year on year since 2005-2018 from 3.6%-5.8%, which can partly account for the trend we
247 observed¹⁸. However a group of researchers¹⁹ found reducing rates of vitrectomy for all indications
248 in diabetic patients in a well-defined geographic area in the North-East of England. They attributed
249 this to better screening and early treatment of proliferative diabetic retinopathy. In recent years,
250 public health intervention on diabetic eye disease in England has been robust, likely reducing the
251 proportion of patients developing severe disease. Therefore as the likelihood of developing TRD for
252 any given patient is decreasing, the number of new diabetics is increasing at a large rate.
253 Additionally, the code for TRD allows for coding cases caused by proliferative vitreo-retinopathy,
254 retinopathy of prematurity and rarer forms of TRD, making it more difficult to analyse diabetes cases
255 in isolation. Another factor which may contribute to explain this increase are improvements in
256 surgical techniques which have resulted in lower thresholds for intervention.

257 *Increase in vitrectomy*

258 We found that the number of reported pars plana vitrectomies irrespective of indication increased
259 over fourfold between 2000-2018. When analysing data since the introduction of vitrectomy
260 surgery, the rate of increase in England was near tenfold up to 2003¹. Advances in surgical
261 instrumentation, massification of the technology, the introduction of optical coherence tomography
262 have all contributed to the expanding application of vitrectomy surgery to a larger number of
263 conditions. Unfortunately, as ICD-10 codes for macular disease make no distinction between
264 conditions treated medically or surgically, analysis of the data on HES does not help to elucidate
265 changes in rates of macular surgery.

266 *Decrease in scleral buckling*

267 Once the mainstay treatment for RD, the number of scleral buckling performed decreased by two
268 thirds. This observation was already apparent in the data analysed up to 2004¹. This change

269 coincides with significant improvements in operating microscopes and surgical instrumentation, as
270 well as important advances in surgical techniques and training. A landmark randomised controlled
271 trial²⁰ published in 2007 showed no significant difference in the likelihood of achieving retinal
272 attachment overall except in pseudophakic eyes where it found an advantage for vitrectomy.
273 Despite scleral buckling allowing better visual outcomes in phakic eyes, vitrectomies have
274 overwhelmingly become the preferred surgical strategy for RRD. Presumably the effectiveness,
275 safety and speed of the operation have contributed to this. This poses challenges for training as a
276 reduced frequency of buckling surgery will potentially undermine the success rate of the operation
277 which remains the state of the art for certain types of RRD. Encouragingly, the ratio of scleral
278 buckling seems to have stabilised from 2015.

279 *Limitations of this study*

280 The use of national datasets has the advantage of providing large scale data over a long time span.
281 However, they are not without their limitations²¹. For one, registry studies offer no clinical data, such
282 as lens status or axial length, which are significant risk factors for RRD. Secondly, their validity
283 depends on the accuracy with which clinical information is reported and coded. The risk of missing
284 cases due to the lack of coding in this study is presumably low because coding is necessary for
285 reimbursing expenses in the NHS since the introduction of the “payment by result” policy in 1990.
286 However, not all NHS hospitals may be equally adept at obtaining appropriate reimbursement and
287 practices in coding may vary slightly from hospital to hospital. For example, not all hospitals may
288 code “pars plana vitrectomy” as the defining procedure, but another step done during the
289 operation. For this reason, adding OPCS manoeuvres which can only be done as part of a vitrectomy
290 operation gives a higher number, likely to be closer to the true number of vitrectomies carried out.
291 Given that these differences and potential errors in coding are likely to be systematic, HES studies
292 are useful in understanding trends. Another issue with coding is the way TRD is defined, not only
293 used for diabetic related RD but for other aetiologies of TRD. Shah² found that the increase in RD

294 admissions in England was likely to be associated with diabetes through linkage analysis with the
295 Oxford database. There is however no biological plausibility to diabetes leading to an increase in
296 RRD. As the study used the generic three-digit code for RD (H33), rather than breaking it down into
297 subcategories (H33.X). We found that both TRD and RRD increased significantly. It is possible that
298 some RD “with breaks” which should be coded H33.0 may have been coded as H33.4 “tractional RD”
299 and vice-versa, even if the code for RRD specifically excludes diabetes as an aetiology. Another
300 limitation of analysing aggregated HES reports is that as episodes are recorded per admission, data
301 about individual patients cannot be identified. Therefore multiple entries are made for the same
302 patient who has had more than one admission for vitreo-retinal surgery. The reported frequency of
303 RD may be underestimated by failing to record cases who opt not to have surgery. In addition, a
304 small number of patients may have undergone surgery abroad or in private hospitals not supplying
305 data to NHS Digital. Given the make-up of England’s healthcare provision, these numbers are likely
306 to be very small. For all the above reasons, caution must be taken in making clinical inferences from
307 analysing administrative data such as England’s HES.

308 In conclusion, the hospital admission rates for all types of RD seem to have increased significantly in
309 recent years, an effect more marked in cases of RRD. This observation is probably attributable to
310 increasing age, rise in the number of performed cataract operations and increase in the prevalence
311 of diabetes for the tractional type of retinal detachment. Other possible explanations may be
312 increased prevalence of myopia, changes in methodology of hospital coding, improvements in
313 patient access and public awareness, increased repeated procedures, and lower thresholds for
314 operating on a chronic RRD with poor prognosis. Despite the disruption that Covid-19 has brought to
315 medical care, a good understanding of these trends should allow a more accurate prediction of
316 future demand for vitreo-retinal surgery across England.

317

318 **Conflict of Interest**

319 The authors declare no potential of conflicts of interest with respect to the research, authorship,
320 and/or publication of this article.

321 **Funding**

322 The authors received no financial support for the research, authorship, and/or publication of this
323 article.

324

325 **References**

- 326 1. El-Amir AN, Keenan TD, Abu-Bakra M, Tanner V, Yeates D, Goldacre MJ. Trends in rates of
327 retinal surgery in England from 1968 to 2004: studies of hospital statistics. Br J Ophthalmol.
328 2009;93(12):1585-90.
- 329 2. Shah V, Hall N, Goldacre MJ. Retinal detachment in England: database studies of trends over
330 time and geographical variation. Br J Ophthalmol. 2015;99(5):639-43
- 331 3. Li JQ, Welchowski T, Schmid M, Holz FG, Finger RP. Incidence of Rhegmatogenous Retinal
332 Detachment in Europe - A Systematic Review and Meta-Analysis. Ophthalmologica.
333 2019;242(2):81-86.
- 334 4. NHS Digital. [https://digital.nhs.uk/data-and-information/data-tools-and-services/data-](https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics)
335 [services/hospital-episode-statistics](https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics) [Accessed 2nd May 2020].
- 336 5. Office for National Statistics. <http://www.ons.gov.uk> [Accessed on 3rd May 2020].
- 337 6. Jackson TL, Donachie PH, Sallam A, Sparrow JM, Johnston RL. United Kingdom National
338 Ophthalmology Database study of vitreoretinal surgery: report 3, retinal detachment.
339 Ophthalmology. 2014 Mar;121(3):643-8.
- 340 7. Manners S, Ng JQ, Kemp-Casey A, Chow K, Kang CY, Preen DB. Retinal detachment surgery in
341 Western Australia (2000-2013): a whole-population study. Br J Ophthalmol.
342 2017;101(12):1679-1682.
- 343 8. Howie AR, Darian-Smith E, Allen PL, Vote BJ. Whole population incidences of patients
344 presenting with rhegmatogenous retinal detachments within Tasmania, Australia. Clin Exp
345 Ophthalmol. 2016;44(2):144-6.
- 346 9. Mitry D, Charteris DG, Yorston D, Siddiqui MA, Campbell H, Murphy AL et al; Scottish RD
347 Study Group. The epidemiology and socioeconomic associations of retinal detachment in
348 Scotland: a two-year prospective population-based study. Invest Ophthalmol Vis Sci.
349 2010;51(10):4963-8.

- 350 10. Van de Put MAJ, Hooymans JMM, Los LI; Dutch Rhegmatogenous Retinal Detachment Study
351 Group. The incidence of rhegmatogenous retinal detachment in The Netherlands.
352 Ophthalmology. 2013;120(3):616-622.
- 353 11. Poulsen CD, Peto T, Grauslund J, Green A. Epidemiologic characteristics of retinal
354 detachment surgery at a specialized unit in Denmark. Acta Ophthalmol. 2016;94(6):548-55.
- 355 12. Hajari JN, Bjerrum SS, Christensen U, Kiilgaard JF, Bek T, la Cour M. A nationwide study on
356 the incidence of rhegmatogenous retinal detachment in Denmark, with emphasis on the risk
357 of the fellow eye. Retina. 2014;34(8):1658-65.
- 358 13. Park SJ, Choi NK, Park KH, Woo SJ. Five year nationwide incidence of rhegmatogenous retinal
359 detachment requiring surgery in Korea. PLoS One. 2013;13(11):e80174.
- 360 14. Chen SN, Lian leB, Wei YJ. Epidemiology and clinical characteristics of rhegmatogenous
361 retinal detachment in Taiwan. Br J Ophthalmol. 2016;100(9):1216-20.
- 362 15. Williams KM, Bertelsen G, Cumberland P, Wolfram C, Verhoeven VJ, Anastasopoulos E et al;
363 European Eye Epidemiology (E(3)) Consortium. Increasing Prevalence of Myopia in Europe
364 and the Impact of Education. Ophthalmology. 2015;122(7):1489-97.
- 365 16. Qureshi MH, Steel DHW. Retinal detachment following cataract phacoemulsification-a
366 review of the literature. Eye (Lond). 2020 Apr;34(4):616-631.
- 367 17. Lakawicz JM, Bottega WJ, Prenner JL, Fine HF. An analysis of the mechanical behaviour of a
368 detaching retina. Math Med Biol. 2015 Jun;32(2):137-61.
- 369 18. Diabetes UK Consortium. Available from: <http://www.diabetes.org.uk> [Accessed 3rd May
370 [2020](#)].
- 371 19. Vaideanu D, Sandhu SS, Ling J, Richardson J, Steel DH. Rate of diabetic vitrectomy in a
372 defined geographical part of North East England. Ophthalmic Epidemiol. 2014;21(3):178-83.
- 373 20. Heimann H, Bartz-Schmidt KU, Bornfeld N, Weiss C, Hilgers RD, Foerster MH; Scleral Buckling
374 versus Primary Vitrectomy in Rhegmatogenous Retinal Detachment Study Group. Scleral

375 buckling versus primary vitrectomy in rhegmatogenous retinal detachment: a prospective
376 randomized multicenter clinical study. *Ophthalmology*. 2007;114(12):2142-54.

377 21. Sinha S, Peach G, Poloniecki JD, Thompson MM, Holt PJ. Studies using English administrative
378 data (Hospital Episode Statistics) to assess health-care outcomes--systematic review and
379 recommendations for reporting. *Eur J Public Health*. 2013 Feb;23(1):86-92.

380 **Figure Legends**

381 Figure 1: Completed consultant episodes at hospitals in England for retinal detachment codes

382 H33.x, 2000-2018.

383 Figure 2: Hospital admissions for retinal detachment and deduction of RD recurrence estimate

384 by mid-year population in England 2000-2018.

385 Figure 3: Number of recorded pars plana vitrectomy procedures and composite of vitrectomy

386 and associated manoeuvres.

387 Figure 4: Ratio of admissions for retinal detachment episodes and composite scleral buckle

388 operations.







