Interventions targeting nonsymptomatic cases can be important to prevent

local outbreaks: COVID-19 as a case-study

AUTHORS

F.A. Lovell-Read[1], S. Funk[2], U. Obolski[3,4], C.A. Donnelly[5,6], R.N. Thompson[1,2,7]

AFFILIATIONS

[1]Mathematical Institute, University of Oxford (Andrew Wiles Building, Radcliffe Observatory

Quarter, Woodstock Road, Oxford, OX2 6GG, UK)

[2]Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene &

Tropical Medicine (Keppel Street, London, WC1E 7HT, UK)

[3]Porter School of the Environment and Earth Sciences, Tel Aviv University (Tel Aviv, Israel,

6997801)

[4]School of Public Health, Tel Aviv University (Tel Aviv, Israel, 6997801)

[5]Department of Statistics, University of Oxford (24-29 St. Giles', Oxford, OX1 3LB, UK)

[6]MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease

Epidemiology, Imperial College London (Norfolk Place, London, W2 1PG, UK)

[7]Christ Church, University of Oxford, (St. Aldate's, Oxford, OX1 1DP, UK)

CORRESPONDING AUTHOR

F.A. Lovell-Read. Address: Merton College, Merton Street, Oxford, OX1 4JD.

E-mail: francesca.lovell-read@merton.ox.ac.uk. Tel: +44 (0)7511 676 592.

ABSTRACT

Background

During infectious disease epidemics, a key question is whether cases travelling to new locations will trigger local outbreaks. The risk of this occurring depends on a range of factors, such as the transmissibility of the pathogen, the susceptibility of the host population and, crucially, the effectiveness of local surveillance in detecting cases and preventing onward spread. For many pathogens, presymptomatic and/or asymptomatic (together referred to here as nonsymptomatic) transmission can occur, making effective surveillance challenging. In this study, using COVID-19 as a case-study, we show how the risk of local outbreaks can be assessed when nonsymptomatic transmission can occur.

Methods

We construct a branching process model that includes nonsymptomatic transmission, and explore the effects of interventions targeting nonsymptomatic or symptomatic hosts when surveillance resources are limited. Specifically, we consider whether the greatest reductions in local outbreak risks are achieved by increasing surveillance and control targeting nonsymptomatic or symptomatic cases, or a combination of both.

Findings

Seeking to increase surveillance of symptomatic hosts alone is typically not the optimal strategy for reducing outbreak risks. Adopting a strategy that combines an enhancement of surveillance of symptomatic cases with efforts to find and isolate nonsymptomatic hosts leads to the largest reduction in the probability that imported cases will initiate a local outbreak.

Interpretation

During epidemics of COVID-19 and other infectious diseases, effective surveillance for

nonsymptomatic hosts can be crucial to prevent local outbreaks.

Funding

UKRI-BBSRC, Wellcome Trust, UKRI-MRC, FCDO, EDCTP2, Christ Church (Oxford).

1. INTRODUCTION

Emerging epidemics represent a substantial challenge to human health worldwide.[1,2] When cases

are clustered in specific locations, two key questions are: i) Will exported cases lead to local

outbreaks in new locations? and ii) Which surveillance and control strategies in those new

locations will reduce the risk of local outbreaks?

Branching process models are used for a range of diseases to assess whether cases that are newly

arrived in a host population will generate a local outbreak driven by sustained local

transmission.[3-6] These models can also be used to predict the effectiveness of potential control

interventions. For example, early in the COVID-19 pandemic, Hellewell *et al.*[7] used simulations

of a branching process model to predict whether or not new outbreaks would fade out under

different contact tracing strategies. Thompson[8] estimated the probability of local outbreaks

analytically using a branching process model and found that effective isolation of infectious

hosts leads to a substantial reduction in the outbreak risk.

A factor that can hinder control interventions during any epidemic is the potential for individuals

to transmit a pathogen while not showing symptoms. For COVID-19, the incubation period has

been estimated to last approximately five or six days on average,[9,10] and presymptomatic

transmission can occur during that period.[11-14] Additionally, asymptomatic infected individuals

(those who never develop symptoms) are also thought to contribute to transmission.[11,15,16]

Motivated by the need to assess the risk of outbreaks outside China early in the COVID-19

pandemic, we show how the risk that imported cases will lead to local outbreaks can be

estimated using a branching process model, including nonsymptomatic individuals in the model

explicitly. We explore the effects of interventions that aim to reduce this risk. Under the

assumption that detected infected hosts are isolated effectively, we consider whether it is most

effective to dedicate resources to enhancing surveillance targeting symptomatic individuals, to

instead focus on increasing surveillance for nonsymptomatic individuals, or to use a combination

of these approaches.

We show that the maximum reduction in outbreak risk almost always corresponds to a mixed

strategy involving enhanced surveillance of both symptomatic and nonsymptomatic hosts. This

remains the case even if the surveillance effort required to find nonsymptomatic infected

individuals is significantly larger than the effort required to find symptomatic individuals. This

highlights the benefits of not only seeking to find and isolate symptomatic hosts, but also

dedicating resources to finding nonsymptomatic cases during infectious disease epidemics.

2. METHODS

<u>2.1 Model</u>

We consider a branching process model in which infectious individuals are classified as

asymptomatic (A), presymptomatic (I_1), or symptomatic (I_2). Hosts in any of these classes may

generate new infections. The parameter ξ represents the proportion of new infections that are

asymptomatic, so that a new infection either involves increasing A by one (with probability ξ), or

increasing I_1 by one (with probability $1 - \xi$).

Presymptomatic hosts may go on to develop symptoms (transition from I_1 to I_2) or be detected

and isolated (so that I_1 decreases by one). Symptomatic individuals (I_2) can be isolated (so that

I_2 decreases by one) or be removed due to recovery or death (so that I_2 decreases by one).

Similarly, asymptomatic hosts may be detected and isolated, or recover (so that A decreases by

one in either case).

A schematic showing the different possible events in the model is shown in Figure 1A. The

analogous compartmental differential equation model to the branching process model that we

consider is given by

$$109 \qquad \frac{\mathrm{d}A}{\mathrm{d}t} = \xi(\eta\beta A + \alpha\beta I_1 + \beta I_2) - \frac{\varepsilon\gamma}{1 - \mathrm{f}(\rho_1)}A - \nu A,$$

$$110 \qquad \frac{\mathrm{d}I_1}{\mathrm{d}t} = (1 - \xi)(\eta\beta A + \alpha\beta I_1 + \beta I_2) - \frac{\varepsilon\gamma}{1 - \mathrm{f}(\rho_1)}I_1 - \lambda I_1,$$

$$111 \qquad \frac{\mathrm{d}I_2}{\mathrm{d}t} = \lambda I_1 - \frac{\gamma}{1 - \mathrm{f}(\rho_2)}I_2 - \mu I_2.$$

In our model, the parameter β and its scaled counterparts $\alpha\beta$ and $\eta\beta$ represent the rates at which

symptomatic, presymptomatic and asymptomatic hosts generate new infections, respectively.

Since we are modelling the beginning of a potential local outbreak, we assume that the size of

the susceptible population remains approximately constant, and do not track depletion of

susceptible individuals explicitly. The parameter λ governs the rate at which presymptomatic

individuals develop symptoms, so that the expected duration of the presymptomatic period is

$1/\lambda$ days in the absence of interventions. Similarly, without interventions, the expected durations

of the symptomatic and asymptomatic infectious periods are $1/\mu$ days and $1/\nu$ days,

respectively.

The baseline rate at which symptomatic individuals are detected and isolated is determined by

the parameter γ. Its scaled counterpart $\epsilon\gamma$ is the analogous quantity for nonsymptomatic

individuals, where the scaling factor $\epsilon < 1$ reflects the fact that nonsymptomatic individuals are

more challenging to detect than symptomatic individuals. We assume that the sensitivity of

surveillance is identical for presymptomatic and asymptomatic individuals, and therefore use the

same isolation rate for both of these groups.

The parameters ρ_1 and ρ_2 represent the surveillance intensification effort targeted at

nonsymptomatic and symptomatic hosts, respectively. The function $f(\rho) = \frac{\delta\rho}{1+\rho}$ governs the

reduction in the expected time to isolation for a given surveillance effort, ρ. The functional form

of $f(\rho)$ is chosen as it generates a reduced expected time to isolation when the surveillance effort

increases, yet the isolation rate saturates and cannot increase indefinitely. The function $f(\rho)$ is

shown in Figure 1B for different values of the parameter δ.

Figure 1: The branching process model used in our analyses. A. Schematic showing the different event types in

the branching process model. The parameters of the model are described in the text and in Table 1. B. The

relationship between the surveillance intensification effort (ρ) and the reduction in the expected time to

isolation ($f(\rho)$), shown for different values of the parameter δ (solid lines). The parameter δ represents the

upper bound of $f(\rho)$ (dotted lines). This general functional relationship between surveillance effort and

isolation effectiveness is assumed to hold for surveillance of both nonsymptomatic and symptomatic

individuals, although nonsymptomatic hosts are more challenging to detect than symptomatic hosts ($\varepsilon < 1$).

<u>2.2 Reproduction number</u>

The basic reproduction number, R_0, represents the expected number of secondary infections

generated by a single infected individual introduced into a fully susceptible population in the

absence of intensified surveillance:

151
$$R_0 = \frac{\xi\eta\beta}{\nu + \epsilon\gamma} + (1 - \xi)\left[\frac{\alpha\beta}{\lambda + \epsilon\gamma} + \frac{\lambda}{\lambda + \epsilon\gamma}\frac{\beta}{\gamma + \mu}\right].$$

This expression is the sum of the expected number of transmissions from a host who begins in

the asymptomatic class and from a host who begins in the presymptomatic class, weighted by the

respective probabilities ξ and $1 - \xi$ that determine the chance that the host experiences a fully

asymptomatic course of infection. The expected number of transmissions from a host who begins

in the presymptomatic class comprises transmissions occurring during the incubation period and

transmissions occurring during the symptomatic period, accounting for the possibility that the

host is isolated prior to developing symptoms.

The proportion of infections arising from presymptomatic hosts in the absence of intensified

surveillance is then given by

162
$$K_p = \frac{\frac{(1 - \xi)\alpha}{\lambda + \epsilon\gamma}}{\frac{\xi\eta}{\nu + \epsilon\gamma} + (1 - \xi)\left[\frac{\alpha}{\lambda + \epsilon\gamma} + \frac{\lambda}{\lambda + \epsilon\gamma}\frac{1}{\gamma + \mu}\right]}, \quad (1)$$

163 and the equivalent quantity for asymptomatic hosts is given by

164
$$K_a = \frac{\frac{\xi\eta}{\nu + \epsilon\gamma}}{\frac{\xi\eta}{\nu + \epsilon\gamma} + (1 - \xi)\left[\frac{\alpha}{\lambda + \epsilon\gamma} + \frac{\lambda}{\lambda + \epsilon\gamma}\frac{1}{\gamma + \mu}\right]}. \quad (2)$$

2.3 Baseline values of model parameters

Since this research was motivated by the need to estimate outbreak risks outside China in the

initial stages of the COVID-19 pandemic, we used a baseline set of parameter values in our

analyses that was informed by studies conducted during this pandemic (Table 1). Where

possible, these parameter values were obtained from existing literature. However, we also

performed sensitivity analyses to determine how our results varied when the parameter values

were changed (see Supplementary Material).

The value of the parameter governing the baseline rate at which symptomatic individuals are

isolated, γ, was chosen to match empirical observations which indicate that individuals who seek

medical care prior to recovery or death do so around four to six days after symptom onset.[17]

Specifically, we assumed that time to first medical visit could be used a proxy for time to

isolation, and chose γ so that the expected time period to isolation conditional on isolation

occurring during the symptomatic period was given by $\frac{1}{\gamma+\mu} = 4 \cdot 6$ days.[17] This is different to the

time period that we refer to as the expected time to isolation for symptomatic hosts, which is $\frac{1}{\gamma}$

days (see Methods).

Table 1. Parameters of the model and the values used in the baseline version of our analysis.

Parameter	Meaning	Baseline value	Justification
R_0	Expected number of secondary infections caused by a single infected individual (when $\rho_1 = \rho_2 = 0$)	$R_0 = 3$	Within estimated range for SARS-CoV-2[18-20]
ξ	Proportion of infections which are asymptomatic	$\xi = 0 \cdot 2$	[21]
β	Rate at which symptomatic individuals generate new infections	$\beta = 0 \cdot 336 \ \text{days}^{-1}$ (3 s.f.)	Chosen so that $R_0 = 3$
α	Relative rate at which presymptomatic individuals generate new infections (compared to symptomatic individuals)	$\alpha = 2 \cdot 78 \ \text{days}^{-1}$ (3 s.f.)	Chosen so that 48·9% of transmissions arise from presymptomatic hosts (i.e. $K_p = 0 \cdot 489$)[11]
η	Relative rate at which asymptomatic individuals generate new infections (compared to symptomatic individuals)	$\eta = 0 \cdot 519 \ \text{days}^{-1}$ (3 s.f.)	Chosen so that 10·6% of transmissions arise from asymptomatic hosts (i.e. $K_a = 0 \cdot 106$)[11]
γ	Isolation rate of symptomatic individuals without intensified surveillance	$\gamma = 0 \cdot 0924$ days^{-1} (3 s.f)	Chosen so that $\frac{1}{\gamma+\mu} = 4 \cdot 6 \ \text{days}$[17]
ε	Relative isolation rate of nonsymptomatic individuals without intensified surveillance (compared to symptomatic individuals)	$\epsilon = 0 \cdot 1$	Assumed (for different values, see Figure S7)
λ	Rate at which presymptomatic individuals develop symptoms	$\lambda = 0 \cdot 5 \ \text{days}^{-1}$	[14]
μ	Recovery rate of symptomatic individuals	$\mu = 1/8 \ \text{days}^{-1}$	[22-24]

ν	Recovery rate of asymptomatic individuals	$\nu = 0 \cdot 1 \text{ days}^{-1}$	Chosen so that, in the absence of interventions, the expected duration of infection is identical for all infected hosts $(\frac{1}{\nu} = \frac{1}{\lambda} + \frac{1}{\mu})$
δ	Upper bound on the fractional reduction in the time to isolation (if no other event occurs)	$\delta = 0 \cdot 8$	Assumed (for different values, see Figure S11)
ρ_1	Surveillance intensification effort targeted at nonsymptomatic hosts	ρ_1 allowed to vary in the range [0,20]	N/A – range of values explored
ρ_2	Surveillance intensification effort targeted at symptomatic hosts	ρ_2 allowed to vary in the range [0,20]	N/A – range of values explored

2.4 Probability of a local outbreak

The probability that a single imported infectious host initiates a local outbreak was calculated

analytically using the branching process model.

The probability of a local outbreak not occurring, starting from i presymptomatic hosts, j

symptomatic hosts, and k asymptomatic hosts was denoted by $q_{i,j,k}$. Starting from one

presymptomatic host (so that $i = 1$ and $j = k = 0$), there are four possibilities for the next event.

That host could: i) generate a new asymptomatic infection (with probability $\frac{\xi\alpha\beta}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}$); ii)

generate a new presymptomatic infection (with probability $\frac{(1-\xi)\alpha\beta}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}$); iii) develop symptoms

(with probability $\frac{\lambda}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}$), or; iv) be isolated (with probability $\frac{\frac{\varepsilon\gamma}{1-f(\rho_1)}}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}$). Consequently,

$$q_{1,0,0} = \frac{\xi\alpha\beta}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}q_{1,0,1} + \frac{(1-\xi)\alpha\beta}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}q_{2,0,0} + \frac{\lambda}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}q_{0,1,0} + \frac{\frac{\varepsilon\gamma}{1-f(\rho_1)}}{\alpha\beta+\lambda+\frac{\varepsilon\gamma}{1-f(\rho_1)}}q_{0,0,0}.$$

If there are no infectious hosts present in the population (i.e. $i = j = k = 0$), then a local

outbreak will not occur and so $q_{0,0,0} = 1$. Assuming that transmission chains arising from two

infectious individuals are independent gives $q_{1,0,1} = q_{1,0,0}\,q_{0,0,1}$ and $q_{2,0,0} = q_{1,0,0}^2$. Hence,

$q_{1,0,0} = a\xi q_{1,0,0}\, q_{0,0,1} + a(1 - \xi)q_{1,0,0}^2 + bq_{0,1,0} + (1 - a - b),$ (3)

where $a = \dfrac{\alpha\beta}{\alpha\beta + \lambda + \frac{\varepsilon\gamma}{1 - f(\rho_1)}}, \; b = \dfrac{\lambda}{\alpha\beta + \lambda + \frac{\varepsilon\gamma}{1 - f(\rho_1)}}.$

Similarly, considering the probability of a local outbreak failing to occur starting from a single

symptomatic host gives

$q_{0,1,0} = \dfrac{\xi\beta}{\beta + \frac{\gamma}{1 - f(\rho_2)} + \mu}\, q_{0,1,1} + \dfrac{(1 - \xi)\beta}{\beta + \frac{\gamma}{1 - f(\rho_2)} + \mu}\, q_{1,1,0} + \dfrac{\frac{\gamma}{1 - f(\rho_2)} + \mu}{\beta + \frac{\gamma}{1 - f(\rho_2)} + \mu}\, q_{0,0,0}.$

As before, noting that $q_{0,0,0} = 1$ and assuming that different infection lineages are independent

leads to

$q_{0,1,0} = c\xi q_{01,0}q_{0,0,1} + c(1 - \xi)q_{1,0,0}q_{0,1,0} + (1 - c),$ (4)

where $c = \dfrac{\beta}{\beta + \frac{\gamma}{1 - f(\rho_2)} + \mu}.$

Finally, considering the probability of a local outbreak failing to occur starting from a single

asymptomatic host gives

$q_{0,0,1} = d\xi q_{0,0,1}^2 + d(1 - \xi)q_{1,0,0}q_{0,0,1} + (1 - d),$ (5)

where $d = \dfrac{\eta\beta}{\eta\beta + \nu + \frac{\varepsilon\gamma}{1 - f(\rho_1)}}.$

Equations (3), (4) and (5) may be combined to give a single quartic equation for $q_{0,0,1}$, yielding

four sets of solutions for $q_{1,0,0}, q_{0,1,0}$ and $q_{0,0,1}$. It is straightforward to verify that $q_{1,0,0} =$

$q_{0,1,0} = q_{0,0,1} = 1$ is always a solution, and further solutions can be found numerically. The

appropriate solution to take is the minimal non-negative real solution $q_{1,0,0} = q_{1,0,0}^*, \; q_{0,1,0} =$

$q_{0,1,0}^*, \; q_{0,0,1} = q_{0,0,1}^*$ (see Supplementary Material). Then, the probability of a local outbreak

occurring beginning from a single presymptomatic host is given by

223
$$p_{1,0,0} = 1 - q^*_{1,0,0},$$

with equivalent expressions holding for $p_{0,1,0}$ and $p_{0,0,1}$ (the probability of a local outbreak

occurring beginning from a single symptomatic host or a single asymptomatic host,

respectively).

Throughout, we consider the probability p of a local outbreak starting from a single

nonsymptomatic host entering the population, accounting for the possibility that the

nonsymptomatic host is either presymptomatic or asymptomatic:

231
$$p = (1 - \xi)p_{1,0,0} + \xi p_{0,0,1}.$$

2.5 Role of the funding source

The funders had no role in study design, preparation of the manuscript or the decision to publish.

3. RESULTS

3.1 Probability of a local outbreak

We considered the effect of R_0 and the duration of the presymptomatic and asymptomatic

periods on the probability of a local outbreak when a nonsymptomatic host enters a new host

population (Figure 2). We considered presymptomatic periods of length $1/\lambda = 1$ day, $1/\lambda = 2$

days and $1/\lambda = 4$ days; in each case, the duration of the asymptomatic period ($1/\nu$ days) was

adjusted so that the relative proportion of infections arising from asymptomatic hosts compared

to presymptomatic hosts remained fixed ($K_a/K_p = 0 \cdot 218$, as in the baseline case). If instead

nonsymptomatic infections are not accounted for, the infectious period follows an exponential

distribution and the probability of a local outbreak is given by $p = 1 - 1/R_0$ (red dash-dotted

line in Figure 2A). Including nonsymptomatic infection in the model therefore led to an

increased risk of a local outbreak in the absence of surveillance intensification (Figure 2A).

We then considered the dependence of the probability of a local outbreak on the intensity of

surveillance targeting nonsymptomatic and symptomatic hosts (Figure 2B-D). The maximum

value of the surveillance intensification effort that we considered (given by ρ_1 or ρ_2 values of

20) corresponded to a 76% reduction in the expected time to isolation (blue line in Figure 1B),

i.e. a 76% reduction in $\frac{1}{\varepsilon\gamma}$ or $\frac{1}{\gamma}$.

The length of the presymptomatic and asymptomatic periods significantly affected the

dependence of the probability of a local outbreak on the level of surveillance targeted at

nonsymptomatic and symptomatic hosts. In Figure 2B, in which the duration of the

presymptomatic period was 1 day, increasing surveillance targeted at nonsymptomatic hosts (ρ_1)

had a limited effect on the probability of a local outbreak, while increasing surveillance targeted

at symptomatic hosts (ρ_2) had a more significant effect. For example, increasing the surveillance

effort targeted at nonsymptomatic hosts to $\rho_1 = 5$ (a 67% reduction in the time to isolation) only

reduced the probability of a local outbreak from 0·730 to 0·716 (3 s.f.), whereas the equivalent

effort targeted at symptomatic hosts ($\rho_2 = 5$) reduced the probability to 0·630 (3 s.f.). As shown

in Figures 3C and D, however, when the presymptomatic and asymptomatic periods were longer,

the benefit of directing surveillance resources towards detecting nonsymptomatic individuals

increased. This was because longer presymptomatic and asymptomatic periods increased the

proportion of infections generated by nonsymptomatic individuals ($K_p + K_a$, see equations (1)

and (2)); a presymptomatic period of 1 day, 2 days and 4 days corresponded to values of $K_p +$

K_a equal to 0·424, 0·595 and 0·746, respectively.

Figure 2. The effect of the duration of the presymptomatic and asymptomatic periods on the probability of
a local outbreak (p), starting from a single nonsymptomatic host. A. The probability of a local outbreak as
a function of the basic reproduction number R_0, for presymptomatic periods of lengths $1/\lambda = 1$ day
(purple), $1/\lambda = 2$ days (blue) and $1/\lambda = 4$ days (green) in the absence of enhanced surveillance ($\rho_1 =$
$\rho_2 = 0$). In each case, the duration of the asymptomatic period ($1/\nu$) is adjusted so that the relative
proportion of infections arising from asymptomatic hosts compared to presymptomatic hosts remains
constant ($K_a/K_p = 0.218$, as in the baseline case). The red dash-dotted line indicates the probability of a
local outbreak in the absence of nonsymptomatic transmission. The vertical grey dotted line indicates
$R_0 = 3$, the baseline value used throughout. B. The probability of a local outbreak as a function of the
surveillance intensification efforts ρ_1 and ρ_2, for $1/\lambda = 1$ day. C. The analogous figure to B but with
$1/\lambda = 2$ days. D. The analogous figure to B but with $1/\lambda = 4$ days. Red dotted lines indicate contours of
constant local outbreak probability (i.e. lines on which the probability of a local outbreak takes the values
shown). The value of β is varied in each panel to give the appropriate value of R_0. All other parameter
values are held fixed at the values in Table 1 (except where stated).

3.2 Optimising surveillance enhancement

We next considered in more detail the impact of surveillance targeted at nonsymptomatic hosts

(ρ_1) relative to the impact of surveillance targeted at symptomatic hosts (ρ_2). For our baseline

parameter values, we considered the probability of a local outbreak starting from a single

imported nonsymptomatic individual for a range of values of ρ_1 and ρ_2. We calculated the

steepest descent contours (white lines in Figure 3A) numerically using a gradient maximisation

approach, in which at each point the contour direction was determined by minimising the local

outbreak probability over a fixed search radius (see Supplementary Material). These contours

indicate how ρ_1 and ρ_2 should be altered to maximise the reduction in the probability of a local

outbreak. In this case, enhancing surveillance targeting both symptomatic and nonsymptomatic

hosts is always optimal (the steepest descent contours are neither horizontal nor vertical).

We then considered a scenario in which, at any time, it is only possible to direct resources

towards enhancing surveillance of either nonsymptomatic individuals or symptomatic

individuals (e.g. tracing and testing of nonsymptomatic contacts of known infectious individuals,

or screening for symptomatic individuals at public events). In Figure 3B, the blue region

represents values of ρ_1 and ρ_2 for which enhancing surveillance targeting symptomatic hosts (i.e.

increasing ρ_2) leads to a larger reduction in the local outbreak probability than enhancing

surveillance targeting nonsymptomatic hosts (i.e. increasing ρ_1). In contrast, in the green region,

enhancing surveillance of nonsymptomatic individuals is more effective than enhancing

surveillance of symptomatic individuals. The white line represents the steepest descent contour

starting from $\rho_1 = \rho_2 = 0$, under the constraint that surveillance can only be enhanced for

symptomatic or nonsymptomatic hosts at any time.

Practical deployment of surveillance is often subject to logistical constraints, and policy-makers

may wish to design surveillance strategies to achieve a specific objective – for example, to

maximise the effectiveness of limited resources or to minimise the cost of achieving a desired

outcome. We therefore also considered the following two examples of such objectives.

Objective 1: Minimise the probability of a local outbreak for a fixed total surveillance effort.

First, we considered the question: given a fixed maximum surveillance effort ($\rho_1 + \rho_2 = C$),

how should surveillance be targeted at nonsymptomatic and symptomatic hosts? This involves

setting the values of ρ_1 and ρ_2 to minimise the local outbreak probability. The optimal strategies

in this case are shown in Figure 3C. The red dotted lines represent contours along which the total

surveillance effort $\rho_1 + \rho_2$ is held constant (i.e. different values of C). On each contour, the red

circle indicates the point at which the local outbreak probability is minimised.

If surveillance resources are increased (i.e. C increases), a further question is how surveillance

should then be increased. In Figure 3C, the white line represents the contour of steepest descent,

under the constraint that the total change in surveillance effort ($\rho_1 + \rho_2$) is held constant at each

step (rather than a constant search radius, as in Figure 3A – for more details, see the

Supplementary Material). This contour coincides exactly with that shown in Figure 3B.

These results indicate that, if surveillance resources are such that C is greater than $2 \cdot 8$

(corresponding to a 59% reduction in time to isolation of symptomatic hosts), the optimal

surveillance strategy involves both enhanced surveillance of symptomatic individuals and

nonsymptomatic individuals (the red dots correspond to positive values of ρ_1, unless C is less

than 2·8).

Objective 2: Minimise the total surveillance effort to achieve a pre-specified reduction in the

probability of a local outbreak

Second, we considered the question: given a pre-specified acceptable risk level (i.e. probability

of a local outbreak), how should the surveillance level targeted at nonsymptomatic and

symptomatic hosts be chosen? This involves choosing ρ_1 and ρ_2 to minimise $\rho_1 + \rho_2$ along a

given contour corresponding to a fixed local outbreak probability (red dotted lines in Figure 3D).

On each contour, the red circle indicates the point along that contour at which the total

surveillance effort $\rho_1 + \rho_2$ is minimised. These optimal points also lie exactly along the line on

which enhancing surveillance targeted at symptomatic hosts is equally effective compared to

enhancing surveillance targeted at nonsymptomatic hosts.

As long as the target local outbreak probability was less than 0.69, optimal surveillance involved

enhanced surveillance of nonsymptomatic individuals as well as symptomatic individuals. For

example, in order to reduce the local outbreak probability to 0.6, the optimal approach was to

deploy resources such that $\rho_1 = 12 \cdot 4$ (a 74% reduction in time to isolation of nonsymptomatic

individuals) and $\rho_2 = 18 \cdot 0$ (a 76% reduction in time to isolation of symptomatic individuals).

Plots analogous to Figure 3D in which the parameters were varied from their baseline values are

shown in the Supplementary Material. In each case we considered, our main message was

unchanged. There always existed a threshold local outbreak probability such that, below this

threshold, the optimal strategy for further reduction in the local outbreak probability involved

enhancing surveillance targeting both nonsymptomatic and symptomatic individuals.

Figure 3. Optimal surveillance strategies to reduce the probability of a local outbreak (p) starting from a
single nonsymptomatic host. A. The local outbreak probability for different values of ρ_1 and ρ_2, with the
steepest descent contours overlaid (white lines). For the maximum reduction in the probability of a local
outbreak at each point, surveillance must be enhanced for both nonsymptomatic and symptomatic
individuals. B. Values of ρ_1 and ρ_2 for which increasing surveillance for nonsymptomatic hosts (i.e.
increasing ρ_1) is more effective at reducing the local outbreak probability than increasing surveillance for
symptomatic hosts (i.e. increasing ρ_2) (green region) and vice versa (blue region). The white line
represents the steepest descent contour starting from $\rho_1 = \rho_2 = 0$, under the constraint that surveillance
can only be enhanced for symptomatic or nonsymptomatic hosts at any time. The diagonal section of the
steepest descent contour is made up of small horizontal and vertical sections. C. Strategies for minimising
the local outbreak probability for a given fixed total surveillance effort ($\rho_1 + \rho_2 = C$). Red dotted lines
indicate contours on which $\rho_1 + \rho_2$ is constant (i.e. lines on which $\rho_1 + \rho_2$ takes the values shown); red
circles indicate the points along these contours at which the local outbreak probability is minimised. The
white line indicates the optimal surveillance enhancement strategy if the maximum possible surveillance
level (i.e. the maximum value of $\rho_1 + \rho_2 = C$) is increased. D. Strategies for minimising the surveillance

effort required to achieve a pre-specific risk level (an "acceptable" local outbreak probability). Red dotted
lines indicate contours of constant local outbreak probability (i.e. lines on which the probability of a local
outbreak takes the values shown); red circles indicate the points along these contours at which the total
surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal strategy to follow if the pre-
specified risk level is increased or reduced.

381 **4. DISCUSSION**

A key component of infectious disease epidemic management is inferring the risk of outbreaks in

different locations.[3-6,25] Different surveillance and control strategies can be introduced to reduce

the risk that imported cases will lead to local outbreaks.[7,8,26] However, for a range of pathogens,

public health measures are hindered by nonsymptomatic hosts who can transmit the pathogen yet

are challenging to detect.[11,25,26]

Here, we showed how the probability of a local outbreak can be estimated using a branching

process model that accounts for nonsymptomatic transmission (Figure 1). The model can be used

to assess the local outbreak probability for different surveillance strategies that target

nonsymptomatic or symptomatic individuals (Figure 2). Previous studies have shown that

detection of nonsymptomatic infections can be a key component of epidemic forecasting[25] and

containment,[26] and have demonstrated the benefits of identifying and isolating infectious

nonsymptomatic hosts to reduce transmission.[11,12] We focused instead on investigating how

surveillance should be targeted at nonsymptomatic or symptomatic hosts in order to reduce the

probability that cases imported to new locations will trigger a local outbreak (Figure 3A,B). We

also showed how the optimal surveillance level targeting these two groups can be assessed when

surveillance resources are limited and policy-makers have specific objectives (Figure 3C,D). In

each case, our main conclusion was that enhanced surveillance of nonsymptomatic hosts ($\rho_1 >$

0) can be an important component of reducing the local outbreak risk during epidemics.

Our goal here was to use the simplest possible model to explore the effects of surveillance of

nonsymptomatic and symptomatic individuals on the risk of local outbreaks. However, for our

modelling approach to be used to make precise quantitative predictions during epidemics, it

would be necessary to update the model to include the range of different specific surveillance

and control interventions that are in place. Detection of nonsymptomatic individuals is facilitated

by contact tracing and testing, which are carried out routinely during epidemics and can be

included in models explicitly.[7,26,27] Reductions in contacts due to social distancing strategies and

school or workplace closures could also be accounted for, although such interventions are often

introduced after a local outbreak has begun rather than in the initial phase of a potential local

outbreak as considered here. We modelled the level of surveillance targeted at nonsymptomatic

and symptomatic hosts in a simple way, using a function describing the relationship between

surveillance effort and effectiveness (Figure 1B). If different public health measures are included

in the model explicitly, then it would be possible to increase the accuracy of assessments of the

relative public health benefits of specific interventions that only target symptomatic individuals

(e.g. screening for passengers with heightened temperatures at airports) compared to

interventions that also target nonsymptomatic hosts (e.g. travel bans or quarantine of all inbound

passengers). Of course, this would require data from which the relative effectiveness of different

measures could be inferred.

In this article, we chose a baseline set of parameter values that were consistent with findings of

studies conducted during the COVID-19 pandemic, although constructing a detailed transmission

model for this pandemic was not our main focus. For example, we set the relative rates at which

presymptomatic and asymptomatic individuals generate new infections compared to

symptomatic individuals so that 48·9% of transmissions arise from presymptomatic infectors,

and 10·6% arise from asymptomatic infectors.[11] While this is in line with other reported

estimates,[28,29] there is substantial variation between studies. We therefore also conducted

sensitivity analyses in which we explored a range of different values of model parameters

(Supplementary Material). In each case we considered, our main conclusion was unchanged:

surveillance of nonsymptomatic individuals can contribute to reducing the risk of local

outbreaks. This result is therefore expected to hold for epidemics of any pathogen for which

nonsymptomatic individuals contribute significantly to transmission.

Despite the necessary simplifications, we have shown how the risk of local outbreaks can be

estimated during epidemics using a branching process model that includes nonsymptomatic

infectious hosts explicitly. Determining the extent to which nonsymptomatic individuals

contribute to transmission is essential early in emerging epidemics of a novel pathogen. As we

have shown, if transmissions occur from nonsymptomatic infectors, dedicating surveillance

resources towards finding nonsymptomatic cases can be an important component of public health

measures that aim to prevent local outbreaks.

DECLARATION OF INTERESTS

We declare no competing interests.

AUTHORS' CONTRIBUTIONS

Conceptualisation: All authors. Methodology: FALR, UO, RNT. Investigation: FALR. Writing –

original draft: FALR, RNT. Writing – review and editing: All authors. Supervision: RNT.

FUNDING

FALR acknowledges funding from the Biotechnology and Biological Sciences Research Council

(UKRI-BBSRC), grant number BB/M011224/1. SF and RNT acknowledge funding from the

Wellcome Trust, grant number 210758/Z/18/Z. RNT also acknowledges funding from Christ

Church (University of Oxford) via a Junior Research Fellowship. CAD acknowledges funding

from the MRC Centre for Global Infectious Disease Analysis (reference MR/R015600/1), jointly

funded by the UK Medical Research Council (MRC) and the UK Foreign, Commonwealth &

Development Office (FCDO) under the MRC/FCDO Concordat agreement, and is also part of

the EDCTP2 programme supported by the European Union.

ACKNOWLEDGEMENTS

Thank you to Andrew Wood for proofreading.

REFERENCES

463
1.   Bloom DE, Cadarette D. Infectious Disease Threats in the Twenty-First Century:

Strengthening the Global Response. *Front. Immunol.* 2019; **10**(549).

2.   Morens DM, Fauci AS. Emerging Infectious Diseases: Threats to Human Health and

Global Stability. *PLos Pathog.* 2013; **9**(7): e1003467.

3.   Boldog P, Tekeli T, Vizi Z, Denes A, Bartha FA, Rost G. Risk Assessment of Novel

Coronavirus COVID-19 Outbreaks Outside China. *J. Clin. Med.* 2020; **9**(2): 571.

4.   Daon Y, Thompson RN, Obolski U. Estimating COVID-19 outbreak risk through air

travel. *J. Travel Med.* 2020; **27**(5).

5.   Lashari AA, Trapman P. Branching process approach for epidemics in dynamic

partnership network. *J. Math. Biol.* 2018; **76**(1): 265-94.

6.   Thompson RN, Thompson CP, Pelerman O, Gupta S, Obolski U. Increased frequency of

travel in the presence of cross-immunity may act to decrease the chance of a global pandemic.

Philos. Trans. R. Soc. Lond., B, Biol. Sci. 2019; **374**(1775): 20180274.

7.   Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks by

isolation of cases and contacts. *Lancet Glob. Health* 2020; **8**(4): e488-e96.

8.   Thompson RN. Novel Coronavirus Outbreak in Wuhan, China, 2020: Intense

Surveillance Is Vital for Preventing Sustained Transmission in New Locations. *J. Clin. Med.*

2020; **9**(2): 498.

9.   Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease 2019

(COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann.*

Intern. Med. 2020; **172**(9): 577-82.

10.   McAloon C, Collins A, Hunt K, et al. Incubation period of COVID-19: a rapid systematic

review and meta-analysis of observational research. *BMJ Open* 2020; **10**(8): e039652.

11.   Ferretti L, Wymant C, Kendall M, et al. Quantifying SARS-CoV-2 transmission suggests

epidemic control with digital contact tracing. *Science* 2020; **368**(6491): eabb6936.

12.   Moghadas SM, Fitzpatrick MC, Sah P, et al. The implications of silent transmission for

the control of COVID-19 outbreaks. *Proc. Natl. Acad. Sci. U. S. A.* 2020; **117**(30): 17513-5.

13.   Tindale LC, Stockdale JE, Coombe M, et al. Evidence for transmission of COVID-19

prior to symptom onset. *eLife* 2020; **9**: e57149.

14.   Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic Transmission of

SARS-CoV-2 — Singapore, January 23–March 16, 2020. *Morb. Mortal. Weekly Rep.* 2020;

69(14): 411-5.

15.   Bai Y, Yao L, Wei T, et al. Presumed Asymptomatic Carrier Transmission of COVID-19.

J. Am. Med. Assoc. 2020; **323**(14): 1406-7.

16.   Furukawa NW, Brooks JT, Sobel J. Evidence Supporting Transmission of Severe Acute

Respiratory Syndrome Coronavirus 2 While Presymptomatic or Asymptomatic. *Emerging*

Infectious Diseases 2020; **26**(7): e201595.

17.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus–Infected Pneumonia. *N. Engl. J. Med.* 2020; **382**(13): 1199-207.

18.   Lai C-C, Shih T-P, Ko W-C, Tang H-J, Hsueh P-R. Severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the

challenges. *Int. J. Antimicrob. Agents* 2020; **55**(3): 105924.

19.   Liu Y, Gayle AA, Wilder-Smith A, Rocklov J. The reproductive number of COVID-19 is

higher compared to SARS coronavirus. *J. Travel Med.* 2020; **27**(2): taaa021.

20.   Zhao S, Lin Q, Ran J, et al. Preliminary estimation of the basic reproduction number of

novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early

phase of the outbreak. *Int. J. Infect. Dis.* 2020; **92**: 214-7.

21.   Buitrago-Garcia D, Egli-Gany D, Counotte MJ, et al. Occurrence and transmission

potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic

review and meta-analysis. *PLoS Med.* 2020; **17**(9): e1003346.

22.   Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 Infections and

Transmission in a Skilled Nursing Facility. *N. Engl. J. Med.* 2020; **382**(22): 2081-90.

23.   Bullard J, Dust K, Funk D, et al. Predicting Infectious Severe Acute Respiratory

Syndrome Coronavirus 2 From Diagnostic Samples. *Clin. Infect. Dis.* 2020: ciaa638.

24.   Wolfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized

patients with COVID-2019. *Nature* 2020; **581**(7809): 465-9.

25.   Thompson RN, Gilligan CA, Cunniffe NJ. Detecting Presymptomatic Infection Is

Necessary to Forecast Major Epidemics in the Earliest Stages of Infectious Disease Outbreaks.

PLoS Comput. Biol. 2016; **12**(4): e1004836.

26.   Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease

outbreak controllable. *Proc. Natl. Acad. Sci. U. S. A.* 2004; **101**(16): 6146-51.

27.   Webb G, Browne C, Huo X, Seydi O, Seydi M, Magal P. A Model of the 2014 Ebola

Epidemic in West Africa with Contact Tracing. *PLoS Curr.* 2015; **7**.

28.   He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility

of COVID-19. *Nat. Med.* 2020; **26**(5): 672-5.

29.   Ren X, Li Y, Yang X, et al. Evidence for pre-symptomatic transmission of coronavirus

disease 2019 (COVID-19) in China. *Influenza Other Respi. Viruses* 2020: 1-9.

30.   Norris JR. Markov Chains (Cambridge Series in Statistical and Probabilistic

Mathematics). Cambridge: Cambridge University Press; 1997.

SUPPLEMENTARY MATERIAL

1. Probability of a local outbreak

In the Methods section, we outlined an approach for deriving the probability of a local outbreak starting from a

single infectious host in either the presymptomatic, symptomatic or asymptomatic classes. Here, we provide

more details about that derivation. The probability of a local outbreak not occurring, starting from i

presymptomatic hosts, j symptomatic hosts and k asymptomatic hosts, is denoted by $q_{i,j,k}$. If we consider the

temporal evolution of (i, j, k) to be a Markov process on the state space $M = \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$, then $q_{i,j,k}$ is the

hitting probability of the state $(0,0,0)$ starting from the state (i, j, k). The vector of hitting probabilities $q^* =$

$\{q^*_{i,j,k} \mid (i, j, k) \in M\}$ is therefore the minimal non-negative (real) solution to the following system of

equations:

$q_{0,0,0} = 1,$

$q_{i,j,k} = \sum_{(l,m,n) \in M} p_{(i,j,k),(l,m,n)} \, q_{l,m,n}$ for $(i, j, k) \neq (0,0,0),$

where $p_{(i,j,k),(l,m,n)}$ is the transition probability from state (i, j, k) to state (l, m, n).[30] Here, minimality means

that if $\hat{q} = \{\hat{q}_{i,j,k} \mid (i, j, k) \in M\}$ is another non-negative real solution, then $q^*_{i,j,k} \leq \hat{q}_{i,j,k}$ for all $(i, j, k) \in M$.

From this, equations (3), (4) and (5) in the main text are obtained. These equations may be reduced to a single

quartic equation for $q_{0,0,1}$, yielding four solutions for $q_{0,0,1}$ and four corresponding solutions for each of $q_{1,0,0}$

and $q_{0,1,0}$. One solution is always given by $q_{1,0,0} = q_{0,1,0} = q_{0,0,1} = 1$; the other solutions may be found

numerically. As described above, we take the minimal non-negative real solution $q_{1,0,0} = q^*_{1,0,0}, q_{0,1,0} =$

$q^*_{0,1,0}, q_{0,0,1} = q^*_{0,0,1}$, and observe that the probability of a local outbreak occurring starting from i

presymptomatic hosts, j symptomatic hosts and k asymptomatic hosts is simply $1 - q_{i,j,k}$, giving the result

stated in the main text. If required, this result can be confirmed for specific model parameter values via repeated

simulation of the branching process model, where the probability of a local outbreak corresponds to the

proportion of simulations that progress beyond the initial stochastic phase.

2. Computation of steepest descent contours

The steepest descent contours shown in Figure 3A were computed using a gradient maximisation approach, in

which at each point the contour direction was determined by minimising the local outbreak probability over a

fixed search radius (Figure S1A,B). Starting from ρ_1, ρ_2, at each step we considered increasing ρ_1 by an amount

$\Delta\rho_1$ and increasing ρ_2 by an amount $\Delta\rho_2$ subject to the constraint $(\Delta\rho_1)^2 + (\Delta\rho_2)^2 = r^2$, where r is a small

pre-specified constant. To achieve this, we scanned over the circular arc $\Delta\rho_1 = r \cos\theta$, $\Delta\rho_2 = r \sin\theta$, for $\theta \in$

$[0, \pi/2]$ (Figure S1A). In practice, this range was discretised into 33 search directions evenly spaced between 0

and $\pi/2$. We then selected the pair of $\Delta\rho_1, \Delta\rho_2$ values for which the local outbreak probability evaluated at $\rho_1 +$

$\Delta\rho_1, \rho_2 + \Delta\rho_2$ was minimised. The process was then repeated beginning from $\rho_{1_{new}} = \rho_1 + \Delta\rho_1, \rho_{2_{new}} = \rho_2 +$

$\Delta\rho_2$ (Figure S1B). The white line in Figure 3B, which divides the region in which increasing ρ_1 has a greater

effect on the local outbreak probability from the region in which increasing ρ_2 has a greater effect on the local

outbreak probability, was computed in an analogous way, with the additional restriction that we only considered

the search directions $\theta = 0$ and $\theta = \pi/2$ (i.e. intensifying only surveillance of nonsymptomatic or symptomatic

hosts).

In Figure 3C, the white line represents the contour of steepest descent under the constraint that the total change

in surveillance effort ($\Delta\rho_1 + \Delta\rho_2 = S$) is held constant at each step, rather than fixing the search radius

$(\Delta\rho_1)^2 + (\Delta\rho_2)^2 = r^2$ as in Figure 3A. Therefore, instead of scanning over a circular arc, at each step we scan

along the line $\Delta\rho_1 = s, \Delta\rho_2 = S - s$, where c varies in the range $[0, S]$ (Figure S1D). Otherwise, the process is

completely analogous to that described above.

Figure S1. Computation of the steepest descent contours shown in the main text. A. To compute the steepest

descent contours shown in Figure 3A of the main text, we increment ρ_1 and ρ_2 by scanning over a constant

search radius $(\Delta\rho_1)^2 + (\Delta\rho_2)^2 = r^2$ (blue arc), and moving to the point $(\rho_{1new}, \rho_{2new})$ along that arc at which

the local outbreak probability is minimised. B. The process shown in A is repeated to generate the complete

contour (red dashed line). C. The analogous figure to B, in which the search direction is limited to directly to

the right ($\theta = 0$) or directly upwards ($\theta = \pi/2$). This procedure is used to generate the contour in Figure 3B in

the main text. D. The analogous figure to B, in which the total change in surveillance effort ($\Delta\rho_1 + \Delta\rho_2 = S$) is

held constant at each step, rather than the search radius. This procedure is used to generate the contour in

Figures 3C and D in the main text.

3. Robustness of results to parameter values used

In this section, we demonstrate how our results are affected by varying the parameters from their baseline

values given in Table 1. We performed sensitivity analyses on the values of $R_0, \xi, K_p, K_a, \gamma + \mu, \epsilon, \lambda, \nu$ and δ.

For each of these, we present plots analogous to Figure 3D for six different values of the relevant parameter. In

each case we considered, our qualitative message was unchanged – for any reduction in the local outbreak

probability below a particular threshold value, the optimal strategy involved surveillance targeting both

nonsymptomatic and symptomatic individuals.

Figure S2. Varying the basic reproduction number R_0 from its baseline value ($R_0 = 3$). Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal strategy to follow if the pre-specified risk level is reduced. Apart from R_0 and β (which is changed in each panel to set the value of R_0), all parameters are held fixed at their baseline values given in Table 1. A. $R_0 = 1 \cdot 5$. B. $R_0 = 2$. C. $R_0 = 2 \cdot 5$. D. $R_0 = 3$ (baseline). E. $R_0 = 3 \cdot 5$. F. $R_0 = 4$.

Figure S3. Varying the proportion of infections from asymptomatic infectors, ξ, from its baseline value ($\xi = 0.2$). Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal strategy to follow if the pre-specified risk level is reduced. Apart from ξ and β (which is changed in each panel to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A. $\xi = 0$. B. $\xi = 0 \cdot 1$. C. $\xi = 0 \cdot 2$ (baseline). D. $\xi = 0 \cdot 3$. E. $\xi = 0 \cdot 4$. F. $\xi = 0 \cdot 5$.

618
619

Figure S4. Varying the proportion of infections arising from presymptomatic hosts in the absence of intensified surveillance (K_p, given by expression (1) in the main text) from its baseline value ($K_p = 0.489$). In each case, the proportions of infections arising from asymptomatic and symptomatic hosts are adjusted so that they remain in the same ratio as in the baseline case. Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal strategy to follow if the pre-specified risk level is reduced. Apart from K_p and K_a, as well as α and η (which are changed in each panel to set the values of K_p and K_a), and β (which is

630 changed in each panel to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A.
631 $K_p = 0 \cdot 2$. B. $K_p = 0 \cdot 3$. C. $K_p = 0 \cdot 4$. D. $K_p = 0 \cdot 5$. E. $K_p = 0 \cdot 6$. F. $K_p = 0 \cdot 7$.
632

Figure S5. Varying the proportion of infections arising from asymptomatic hosts in the absence of intensified
surveillance (K_a, given by expression (2) in the main text) from its baseline value ($K_a = 0.106$). In each case,
the proportions of infections arising from presymptomatic and symptomatic hosts are adjusted so that they
remain in the same ratio as in the baseline case. Plots are analogous to Figure 3D, showing strategies for
minimising the surveillance effort required to achieve a pre-specified risk level (an "acceptable" local outbreak
probability). Red dotted lines represent contours along which the probability of a local outbreak is constant, as
labelled; red circles indicate the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is

minimised. The white line indicates the optimal strategy to follow if the pre-specified risk level is reduced.
Apart from K_p and K_a, as well as α and η (which are changed in each panel to set the values of K_p and K_a), and
β (which is changed in each panel to set $R_0 = 3$), all parameters are held fixed at their baseline values given in
Table 1. A. $K_a = 0 \cdot 01$. B. $K_a = 0 \cdot 05$. C. $K_a = 0 \cdot 1$. D. $K_a = 0 \cdot 15$. E. $K_a = 0 \cdot 2$. F. $K_a = 0 \cdot 25$.
645

Figure S6. Varying the expected time period to isolation conditional on isolation occurring during the
symptomatic period, $1/(\gamma + \mu)$, from its baseline value ($1/(\gamma + \mu) = 4.6$ days). This is achieved by varying
the parameter γ, whilst holding the recovery rate of symptomatic individuals μ equal to its baseline value ($\mu = 1/8$ days^{-1}). Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort
$1/8$ days^{-1}). Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort
required to achieve a pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines

represent contours along which the probability of a local outbreak is constant, as labelled; red circles indicate
the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line
indicates the optimal strategy to follow if the pre-specified risk level is reduced. Apart from γ and β (which is
changed in each panel to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A.
$1/(\gamma + \mu) = 2$ days. B. $1/(\gamma + \mu) = 3$ days. C. $1/(\gamma + \mu) = 4$ days. D. $1/(\gamma + \mu) = 5$ days. E. $1/(\gamma + \mu) =$
6 days. F. $1/(\gamma + \mu) = 7$ days.
658

Figure S7. Varying ϵ, the relative isolation rate of nonsymptomatic individuals without intensified surveillance
(compared to symptomatic individuals), from its baseline value ($\epsilon = 0.1$). Plots are analogous to Figure 3D,
showing strategies for minimising the surveillance effort required to achieve a pre-specified risk level (an

"acceptable" local outbreak probability). Red dotted lines represent contours along which the probability of a
local outbreak is constant, as labelled; red circles indicate the points along these contours at which the total
surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal strategy to follow if the pre-
specified risk level is reduced. Apart from ϵ and β (which is changed in each panel to set $R_0 = 3$), all
parameters are held fixed at their baseline values given in Table 1. A. $\epsilon = 0 \cdot 01$. B. $\epsilon = 0 \cdot 02$. C. $\epsilon = 0 \cdot 05$.
D. $\epsilon = 0 \cdot 1$ (baseline). E. $\epsilon = 0 \cdot 2$. F. $\epsilon = 0 \cdot 3$.
669

Figure S8. Varying the duration of the presymptomatic period, $1/\lambda$, from its baseline value ($1/\lambda = 0.5$ days).
Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a
pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along

which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these
contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal
strategy to follow if the pre-specified risk level is reduced. Apart from λ and β (which is changed in each panel
to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A. $1/\lambda = 0.5$ days. B.
$1/\lambda = 1$ day. C. $1/\lambda = 2$ days (baseline). D. $1/\lambda = 4$ days. E. $1/\lambda = 6$ days. F. $1/\lambda = 8$ days.
679

Figure S9. Varying the duration of the symptomatic period, $1/\mu$, from its baseline value ($1/\mu = 8$ days). The
parameter γ is varied simultaneously such that $1/(\gamma + \mu)$, the expected time period to isolation conditional on
isolation occurring during the symptomatic period, remains at its baseline value ($1/(\gamma + \mu) = 4.6$ days). Plots
are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a pre-

specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along
which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these
contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal
strategy to follow if the pre-specified risk level is reduced. Apart from μ and γ, and β (which is changed in each
panel to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A. $1/\mu = 5$ days. B.
$1/\mu = 6$ days. C. $1/\mu = 7$ days. D. $1/\mu = 8$ days (baseline). E. $1/\mu = 9$ days. F. $1/\mu = 10$ days.
691

Figure S10. Varying $1/\nu$, the duration of the asymptomatic period, from its baseline value ($1/\nu = 10$ days).
Plots are analogous to Figure 3D, showing strategies for minimising the surveillance effort required to achieve a
pre-specified risk level (an "acceptable" local outbreak probability). Red dotted lines represent contours along

which the probability of a local outbreak is constant, as labelled; red circles indicate the points along these
contours at which the total surveillance effort $\rho_1 + \rho_2$ is minimised. The white line indicates the optimal
strategy to follow if the pre-specified risk level is reduced. Apart from ν and β (which is changed in each panel
to set $R_0 = 3$), all parameters are held fixed at their baseline values given in Table 1. A. $1/\nu = 7$ days. B.
$1/\nu = 8$ days. C. $1/\nu = 9$ days. D. $1/\nu = 10$ days (baseline). E. $1/\nu = 11$ days. F. $1/\nu = 12$ days.
701

Figure S11. Varying the upper bound on the fractional reduction in the time to isolation (if no other event
occurs), δ, from its baseline value ($\delta = 0.8$). Plots are analogous to Figure 3D, showing strategies for
minimising the surveillance effort required to achieve a pre-specified risk level (an "acceptable" local outbreak
probability). Red dotted lines represent contours along which the probability of a local outbreak is constant, as

labelled; red circles indicate the points along these contours at which the total surveillance effort $\rho_1 + \rho_2$ is
minimised. The white line indicates the optimal strategy to follow if the pre-specified risk level is reduced.
Apart from δ, all parameters are held fixed at their baseline values given in Table 1. A. $\delta = 0 \cdot 5$. B. $\delta = 0 \cdot 6$.
C. $\delta = 0 \cdot 7$. D. $\delta = 0 \cdot 8$ (baseline). E. $\delta = 0 \cdot 9$. F. $\delta = 0 \cdot 95$.