Identifying Neuroanatomical and Behavioral Features for Autism Spectrum Disorder Diagnosis in Children using Machine Learning

Yu Han[1]*, Donna M. Rizzo[2], John P. Hanley[3], Emily L. Coderre[1], Patricia A. Prelock[1]

1 Department of Communication Sciences and Disorders, University of Vermont, Burlington, VT, United States.
2 Department of Civil and Environmental Engineering, University of Vermont, Burlington, VT, United States.
3 Department of Microbiology and Molecular Genetics, University of Vermont, Burlington, VT, United States.

* yu.han@uvm.edu (YH)

Abstract

Autism spectrum disorder (ASD) is a neurodevelopmental disorder that can cause significant social, communication, and behavioral challenges. Diagnosis of ASD is complicated and there is an urgent need to identify ASD-associated biomarkers and features to help automate diagnostics and develop predictive ASD models. The present study adopts a novel evolutionary algorithm, the conjunctive clause evolutionary algorithm (CCEA), to select features most significant for distinguishing individuals with and without ASD, and is able to accommodate datasets having a small number of samples with a large number of feature measurements. The dataset is unique and comprises both behavioral and neuroimaging measurements from a total of 28 children from 7 to 14 years old. Potential biomarker candidates identified include brain volume, area, cortical thickness, and mean curvature in specific regions around the cingulate cortex, frontal cortex, and temporal-parietal junction, as well as behavioral features associated with theory of mind. A separate machine learning classifier (i.e., k-nearest neighbors algorithm) was used to validate the CCEA feature selection and for ASD prediction. Study findings demonstrate how machine learning tools might help move the needle on improving diagnostic and predictive models of ASD.

Introduction

Autism Spectrum Disorder (ASD) is a neurodevelopmental disorder that can cause significant social, communication, and behavioral challenges. According to the most recent report from the Center for Disease Control and Prevention (CDC), the number of children in the U.S. diagnosed with ASD are about 1 in every 54 [80]. While genetic and environmental factors have been linked to the development of ASD, at present there is no identified cause or cure of ASD.

ASD is characterized by impairments in social interaction and the presence of restricted and repetitive behaviors, interests, or activities [1–4]. At the core of social impairment is often a deficit in theory of mind (ToM), the ability to recognize and understand the thoughts, feelings and perspectives of others. Some symptoms of ASD are not evident until age two or later. In fact, a child may appear to be developing according to typical milestones until the age of two and then stop learning new skills and may even lose skills [5, 6]. Currently, the diagnosis of autism is based on behavioral symptoms alone, often assessed through the Autism Diagnostic Observation Schedule-Second edition (ADOS-2) and the Autism Diagnostic Interview-revised (ADI-R) [7, 8]. A typical diagnostic appointment consists of evaluations lasting several hours at a designated clinical office. The rigorous and time-consuming nature of ASD diagnostic examinations often leads to a demand that exceeds the capacity to see patients. As a result, many diagnostic centers have expanding wait lists for appointments. This bottleneck can translate to delays in diagnosis of 13 months and longer [5, 9–15]. It is also believed that a substantial number of individuals on the spectrum remain undetected [82]. With growing awareness of ASD, there is a high demand for a faster

and more automated ASD diagnostic approach that might allow for more efficient

diagnosis and early identification of high-risk populations [16].

Building an automated diagnostic and predictive model of ASD is timely as many

studies have adopted machine learning approaches to identify significant biomarkers

that include both behavioral and biological features. For instance, Duda and colleagues

(2016) applied machine learning to distinguish ASD from attention deficit hyperactivity

disorder (ADHD) using the Social Responsiveness Scale for children between 5 to 13

years old [17]. Bone et al. (2015) trained their models to diagnose children with ASD

versus neurotypical (NT) children using the same Social Responsiveness Scale and the

ADI-R score for children between 5 to 17 years old [18]. Other studies aggregated items

from the ADOS and scores from the Autism Quotient (AQ) to accurately classify an ASD

group [19]. Although behavioral outcome measures are commonly used for ASD

diagnosis and assessment with solid reliability and validity, there are some limitations

when using them   to classify participants. Specifically, behavioral outcome measures

are typically prone to performance demands, as well as environmental and personal

factors that could influence responses on any given day. Consequently, the ability to

identify more consistent ASD markers using neuroimaging measures to supplement

behavioral measures becomes important.

As a result of the wide range and subjective nature of behavioral measures used

in diagnosing ASD, many studies are exploring brain-based biological markers (e.g.,

measurable via magnetic resonance imaging (MRI)) to identify a common etiology

across individuals with ASD. Currently, these less subjective markers are attractive not

only for diagnostic purposes, but as possible targets for interventions [20]. Independent

structural MRI studies have found differences in whole brain volume and the developmental trajectories between individuals with and without ASD [21–29]. Other structural brain abnormalities associated with ASD include cortical folding signatures that appear in the following regions of the brain: temporal-parietal junction, anterior insula, posterior cingulate, lateral and medial prefrontal cortices, corpus callosum, intraparietal sulcus, and occipital cortex [21–29]. Evidence also shows that an accelerated expansion of cortical surface area, but not cortical thickness, causes an early overgrowth of the brain in children with ASD [30], while other studies suggest that individuals with ASD tend to have thinner cortices and reduced surface area as an effect of aging [31]. With these clear brain differences among those with and without ASD, it is informative and critical to look for brain-based ASD biomarkers.

Machine learning (ML) has been introduced to the neuroimaging field to identify the atypical brain regions in individuals with ASD. The support vector machine (SVM) is an algorithm that avoids overfitting and is known for high classification accuracy without requiring large sample sizes. It has been used to classify ASD from NT participants using extracted features from functional connectivity metrics and grey matter volume [32–35]. Other ASD applications of ML classifiers include deep neural networks [37] and the random forest (RF) algorithm; the latter uses random ensembles of independently grown decision trees [38]. Although these methods have demonstrated high accuracy for classifying ASD, to our knowledge most of them have not been used to identify input variables most closely associated with ASD (i.e., feature selection). In addition, the majority of studies use data from the Autism Brain Imaging Data Exchange (ABIDE) dataset, which includes 1112 existing resting-state functional MRI (rs-fMRI) datasets

with corresponding structural MRI and phenotypic information from 539 individuals with ASD and 573 age-matched NT controls between the ages of 7 and 64 collected from 24 international brain imaging laboratories [39]. For example, Guo et al. (2017) selected features associated with ASD using a deep neural network from brain resting-state functional connectivity patterns using the ABIDE dataset [83]. However, findings using datasets from various sites have several limitations.

Specifically, classification across a heterogeneous population as is true in autism is challenging [40, 41], particularly when neuroimaging data are pooled from multiple acquisition sites such as the ABIDE dataset, which has considerable variation in demographic and phenotypic profiles. Data variance introduced via scanner hardware, imaging protocols, operator characteristics, regional demographics, and other site-specific acquisition factors can also affect the classification performance. It is often difficult to collect neuroimaging data from individuals with ASD given the scanner noises and high expectations for individuals to remain still. In fact, most individual site datasets have small sample sizes that can lead to overfitting and classification inaccuracies using traditional ML algorithms. Moreover, while the ultimate goal for ML-based diagnostic classification in neuroimaging is to identify discriminative features that provide insight into atypical structure and connectivity patterns in the affected population [42], many of the ML algorithms applied to ASD were designed to classify large amounts of data (e.g., ABIDE) rather than optimize the selection of input features.

ASD drivers or markers are likely the result of a complex interaction of factors with no single factor (i.e., main effect or univariate model) driving the system. As such, traditional statistical tools (e.g., logistic regression) that search for univariate drivers of

ASD are unlikely to find consistent patterns. Thus, ML techniques that explore large search spaces for multivariate interactions are both needed and becoming popular in helping to elucidate the complex interactions in systems such as ASD. Our study employs one such ML tool: an evolutionary algorithm [44] called the conjunctive clause evolutionary algorithm (CCEA) [36]. The CCEA was specifically designed to efficiently explore large search spaces for complex interactions between features and some associated nominal outcome (e.g., ASD or NT). In addition, the CCEA has built-in tools to prevent overfitting to produce easily interpretable parsimonious models.

This study examines the utility of using the CCEA for feature selection in ASD, particularly to address traditional statistical challenges associated with datasets having small sample sizes and a large number of feature measurements. Additionally, the selected features are validated and used for diagnostic classification by applying a separate and more traditional ML classifier (i.e., the k-nearest neighbors (KNN) algorithm). The dataset in the present study has a relatively large number of features, consisting of both behavioral and neuroimaging measurements. Although some behavioral features may seem less objective compared to an assessment of neuroanatomical features, they are often easier to obtain and more pragmatic for children with ASD especially if they have strong psychometric properties. Clinicians seldom place a child with ASD in a scanner to obtain neuroanatomical information before conducting behavioral assessment for the purpose of diagnosis and treatment. The combination of behavioral measures with neuroanatomical information supports the value of making brain-behavior connections that will advance our understanding of ASD. In the present study, the behavioral measurements include scores of language

ability, intellectual ability, and ToM. The neuroimaging measurements include brain volume, brain surface area, cortical thickness, and cortical curvature extracted from MRI whole-brain T1-weighted scans. These features were collected from a total of 28 children ages 7 to 14, of which 9 children had been diagnosed with ASD. Only a subset of these 28 children were used for feature selection in the CCEA (7 children with ASD and 14 NT children), as another subset (2 children with ASD and 5 NT children) were enrolled at a later time (i.e., after the CCEA was trained). While this later cohort was not included in the CCEA feature selection analysis, it was included in the subsequent validation and predictive k-nearest neighbors (KNN) modeling.

Using the CCEA, we aim to identify discriminative biomarkers and behavioral features to help develop an automated diagnostic and predictive system for ASD. We believe this is a pioneering study for:

- Selecting discriminative biomarkers among children from 7 to 14 years old and classifying ASD.
- Including both behavioral and neuroimaging measurements in the feature selection model for better prediction and understanding of ASD.
- Identifying models (sets of features) that most strongly correlate to children with ASD given a dataset with a relatively small sample size (i.e., N=28) and a large number of features (i.e., 247 neuroimaging features and 13 behavioral features).
- Developing a predictive ML model using input features selected by the CCEA.

Materials and Methods

Participants

A total of 9 children with ASD (1 female) and 19 NT children (7 female), ages 7-

14, were enrolled in the study. In addition to the behavioral assessments described below, the ASD group also completed the ADOS-2 and the Social Communication Questionnaire-Lifetime version (SCQ) [45] to confirm their ASD diagnosis. Although diagnosis of ASD is typically done at an early age, the characteristics of ASD are long-term, and classification with additional neurobiological information at any age recognizes the potential for brain-behavior comparisons with NT populations. Potential changes behaviorally and neurobiologically at any age may also inform types, duration, and intensity of intervention that may influence these changes. Therefore, although we tested older children, this study informs an increased understanding of likely biomarkers of ASD.

Ethical Statement

The present study was approved by the University of Vermont (UVM) Research Protection Office and IRB committee. The specific staff member who approved the study protocol, 19-0005, was Karen Crain. Consent forms were obtained from each participant's legal guardian and assent forms were obtained from each participant.

Behavioral Measurements

All children participated in 2-3 hours of baseline behavioral assessments that included the Comprehensive Assessment of Spoken Language (CASL) [81], the Universal Nonverbal Intelligence Test-2 (UNIT-2) [46, 47], the Theory of Mind Task Battery (ToMTB) [48] and the Theory of Mind Inventory-2 (ToMI-2) [49, 50]. Measures of language and cognition are typical in the assessment of ASD, as the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) requires an assessment of language and intellectual functioning beyond the diagnosis of ASD.

The CASL is an orally administered research-based assessment consisting of 15 subtests measuring language for individuals ranging from 3 to 21 years of age. For the present study, only those basic subsets that establish the CASL language core are used: Antonyms, Sentence Completion, Syntax Construction, Paragraph Comprehension, and Pragmatic Judgment. Reliability for the core subtests ranges from 0.80 to 0.90 with composite scores ranging from 0.92 to 0.96. Test-retest reliability for individual subtests range from 0.65 to 0.95 with core composite coefficients from 0.92 to 0.93. The UNIT-2 is a multidimensional assessment of intelligence for individuals with speech, language, or hearing impairments. It consists of nonverbal tasks that test symbolic memory, non-symbolic quantity, analogic reasoning, spatial memory, numerical series, and cube design. Reliability studies include internal consistency, test-retest, and scorer reliability. Subtest coefficients range from 0.88 to 0.96 and 0.93 to 0.98 for the composites. Content, criterion and construct validity were largely correlated with other cognitive measures and demonstrated the value of the UNIT-2 for the assessment of diverse groups of children.

ToM is a core deficit in ASD that is often used to explain the social impairments of the disorder. ToM is the ability to reason about the thoughts and feelings of self and others, including the ability to predict what others will do or how they will feel in a given situation on the basis of their inferred beliefs [51, 52]. Scores from both the ToMTB and ToMI-2 were included to provide representative measures of a child's social cognition level. The ToMTB and ToMI-2 are two norm-referenced tools and behavioral tasks used as outcome measures to assess ToM [51, 52]. Scores from both the ToMTB and ToMI-2 provide valid representations of a child's social cognition level. The ToMI-2 is a

parent-informant measure of a child's functional level of ToM. Each of the 60 items assesses a particular ToM dimension using items that range from simple content to those that evaluate more complex skills. Each item is rated on a 20-unit continuous scale anchored by "Definitely Not" and "Definitely." Respondents indicate their response with a vertical hash mark at the point on the scale that best reflects their attitude. Item, subscale, and composite scores range from 0-20 with higher values reflecting greater parental confidence that the child possesses a particular ToM skill. The ToMI-2 is designed to be a socially and ecologically valid index of ToM as it occurs in everyday social interactions. It has demonstrated excellent test-retest reliability, internal consistency, and criterion-related validity for both NT and ASD children as well as contrasting-group validity and statistical evidence of construct validity (i.e., factor analysis). The ToMTB directly assesses a child's understanding of a series of scenarios tapping ToM. It consists of 15 test questions within nine tasks, arranged in ascending difficulty. Tasks are presented as short vignettes that appear in a story-book format. Each page has color illustrations and accompanying text. For all tasks, children are presented with one correct response option and three plausible distractors. Memory control questions are included that must be passed for credit on the test questions. The ToMTB demonstrates excellent internal consistency and inter task agreement (α = 0.91 at T1 and α =0.94 at T2) and strong construct validity (r = 0.66, p < .01) [48, 50, 53].

In selecting potential features for the CCEA, we included 13 behavioral features. These included the total score of the CASL, full scale score of the UNIT-2, abbreviated score of the UNIT-2, total score of the ToMTB, total composite mean of the ToMI-2 (i.e., assessing overall ToM ability), early subscale mean of the ToMI-2 (i.e., assessing early-

developing ToM abilities such as regulating desire-based emotion and recognition of happiness and sadness), basic subscale mean of the ToMI-2 (i.e., assessing basic ToM ability such as recognition of surprise), advanced subscale mean of the ToMI-2 (i.e., assessing advanced ToM ability such as recognition of embarrassment). Drawn from a larger study about emotion recognition and ToM, we also included scores from single ToMI-2 items assessing recognition of simple emotions such as happiness and sadness, as well as more complex emotions such as surprise and embarrassment, which ASD children often find difficult to recognize and process [54–56]. Table 1 provides an overview of scores on the 13 behavioral measures. Results from independent sample t-tests found that NT participants scored significantly higher (p<0.05) than ASD participants on the CASL, UNIT-2 full scale, ToMTB, ToMI-2 total, ToMI-2 early subscale, ToMI-2 basic subscale, ToMI-2 advanced subscale, as well as ToMI-2 single items of surprise, embarrassment and desire-based emotion. In its psychometric development, the ToMI and ToMTB were tested against children with and without autism with scores on both measures, discriminating those children with a diagnosis of ASD from those without such a diagnosis. These measures of ToM also showed a delay in development of early developing and basic ToM skills and a failure to achieve more advanced ToM skills [48,49,50].

Table 1. Participant Behavioral Assessments Scores: NT vs. ASD

			NT Group (n=19)	ASD Group (n=9)	Group Difference
Age			10.2 (7-14)	11 (8-13)	$p = 0.36$
CASL			113 (89-132)	78.8 (40-121)	$p < 0.001$**
UNIT-2 Full Scale			111.9 (100-135)	96.2 (56-127)	$p < 0.05$*
Unit-2 Abbreviation			110.9 (91-124)	102 (67-121)	$p=0.16$
ToMTB			13.3 (11-15)	11.1 (5-14)	$p < 0.05$*
ToMI-2		Total	17.2 (11.5-19.3)	12.4 (9.1-17.6)	$p < 0.001$**
		Early Subscale	17.7 (13.8-20)	14.6 (11.6-17.9)	$p < 0.05$*
		Basic Subscale	18 (13.2-19.7)	14 (8.5-19.2)	$p < 0.001$**
		Advanced Subscale	16 (8.3-18.9)	9.4 (6-16.4)	$p < 0.001$**
		Happy	18.5 (14.1-20)	17.3 (12.5-20)	$p = 0.24$
		Sad	17.5 (7.1-20)	16.9 (13.1-20)	$p = 0.62$
		Surprise	17.8 (12.2-20)	15.3 (10.3-20)	$p < 0.05$*
		Embarrassment	15.3 (5.8-20)	10.3 (4.8-20)	$p < 0.05$*
		Desire-Based	18.9 (15-20)	16.7 (14.6-20)	$p < 0.05$*

MRI Acquisition and Preprocessing

All data were acquired using the MRI Center for Biomedical Imaging 3T Philips Achieva dStream scanner and 32-channel head coil at the University of Vermont (UVM). Parameters for T1 acquisition were TR 6.4s, TE 2.9s, flip angle 8 degree, 1mm isotropic imaging resolution with a 256 x 240 *mm2* field of view and 225 slices. Participants watched three videos at home before coming to the MRI center. The first was a cartoon video explaining what an MRI is, and what one might experience while lying in an MRI

scanner [58]. The second video, recorded at the UVM MRI mock scanner room, helped

visualize the real setting and procedures a child would experience. The third video

explained the procedures of wearing earplugs. All participants practiced laying still and

became familiar with the scanner noise in the mock scanner room. The T1 structural

scan was preprocessed using the Human Connectome Project (HCP) minimal

preprocessing pipelines, including spatial artifact/distortion removal, surface generation,

cross-modal registration, and alignment to standard space. These pipelines are

specially designed to capitalize on the high-quality data offered by the HCP. The final

standard space makes use of a recently introduced CIFTI file format and the associated

grayordinates spatial coordinate system. This allows for combined cortical surface and

subcortical volume analyses while reducing the storage and processing requirements

for high spatial and temporal resolution data [60]. Brain anatomical features were

extracted using FreeSurfer aparcstats2tabl script [59], which reads brain parcellation

statistics directly from T1 scan and exports them to excel sheets. These extracted

anatomical features include volume, cortical thickness, mean curvature, and area of all

ROIs for each subject. These ROIs were defined using the automatic segmentation

procedures that assign one of 37 labels to each brain voxel, including left and right

caudate, putamen, pallidum, thalamus, lateral ventricles, hippocampus, and amygdala

[61]. There were 276 brain features included in total.

Conjunctive Clause Evolutionary Algorithm

We used an evolutionary algorithm to identify the features associated with ASD.

The CCEA is a machine learning tool that searches for both the combinations of

features associated with a given category (e.g., ASD) as well as their corresponding

range of feature values [36]. The CCEA can find feature interactions even in the absence of main-effects, and can, therefore, identify feature combinations that would be difficult to discover using traditional statistics. The CCEA selects for the best conjunctive clauses (CCs) of the form:

$$CC_k = F_i \in a_i \wedge F_j \in a_j ...,\qquad(1)$$

where F_i represents a risk factor i whose value lies in the range a_i; and the symbol $\wedge$ represents a conjunction (i.e., logical AND). One benefit of the CCEA is that it produces parsimonious models that are correlated with a select category (e.g., ASD). The models generated by the evolutionary algorithm can be described by their order or total number of features in the conjunctive clause. One example of a parsimonious second order conjunctive clause is: a person with a right hemisphere isthmus cingulate volume of 3,300 – 4,100 mm^3 AND a right hemisphere posterior cingulate volume of 4,100 – 6,200 mm^3 is more likely to have ASD than someone who does not meet these criteria. The CCEA was able to generate models that ranged from 1st to 5th-order in the present study. Since we are only interested in the most parsimonious (i.e., lowest order models) to draw meaningful conclusions and to avoid the overfitting that often occurs with higher-order models, we focused on second-order models (i.e., those having only two features) with the highest fitness (PMF) in which all of the features identified were brain anatomical features. Because of our desire to explore the predictive capability of the behavioral features, we also expanded our analysis to include third-order models (i.e., model combinations with three features) in which behavioral features were included.

The fitness of each conjunctive clause (CC) is evaluated using the hypergeometric probability mass function (PMF), and only the most-fit conjunctive

clauses are saved. The hypergeometric PMF is not a *p*-value and thus, is not

constrained by issues associated with what threshold is "significant" [62–64]. To prevent

overfitting, the CCEA performs feature sensitivity on each conjunctive clause to ensure

each feature contributes to the overall fitness. The sensitivity of each feature is

calculated by taking the difference between the conjunctive clause fitness and the

fitness when that feature is removed. Thus, a feature's sensitivity may be viewed as the

amount of fitness that it contributes to the conjunctive clause. Positive predictive value

(PPV) is the number of true positives divided by the sum of true and false positives; and

class coverage is the number of true positives divided by the sum of true positives and

false negatives (i.e., the percent of ASD individuals that match the conjunctive clause).

In this work, the CCEA was run five times using the training set to ensure a more

thorough search of the fitness landscape.

K-nearest Neighbors Algorithm and Leave-One-Out Cross Validation

In order to further validate the CCEA's selection of features capable of

discriminating between children with and without ASD, we built a separate KNN

classification model and used leave-one-out cross validation on all 28 subjects. The

KNN is a classification algorithm that assumes that things that exist in close proximity

(i.e., nearer to each other) are more similar. In this study, each subject was classified

into one of two output classes (i.e., ASD or NT) based on a plurality vote of its

neighbors, with the subject being assigned to the class most similar to its k-nearest

neighbors. When k = 1, then the subject is assigned to the class of a single nearest

neighbor [65]. Leave-one-out cross validation is a special case of cross validation where

the number of folds equals the number of subjects in the data set. Thus, the KNN

algorithm is applied once for each subject, using all other instances as the training set

and using the selected subject as a single-item test set [66]. After model validation, we

trained three separate KNN classifiers using a balanced dataset (6 NT and 6 ASD

subjects) and feature sets identified by the CCEA to classify the remaining 16 subjects.

See Table 2.

Table 2: Subject Inclusion and Distribution

Models		Subjects	Include Subjects Used In CCEA? N=21 (14 NT & 7 ASD)	Include Subjects from the Later Cohort That Was Not Used In CCEA? N=7 (5 NT & 2 ASD)
CCEA		NT = 14 ASD = 7	All	None
KNN Validation Model		NT = 19 ASD = 9	All	All
KNN Prediction Model	Training	NT = 6 ASD = 6	NT = 6 ASD = 4	NT= None ASD = 2
	Testing	NT = 13 ASD = 3	NT = 8 ASD = 3	NT = 5 ASD = None

Results

CCEA Feature Selection: 14 NT and 7 ASD

When using the CCEA for feature selection, 2438 CCs (i.e., models or sets of

features) were generated ranging from first-order to fifth-order. The PPV of the 2438

models ranged from 46.47% to 100% and their class coverage ranged from 42.86% to

100%. Among these models, we looked for the most parsimonious (i.e., lowest order

models) to draw meaningful conclusions and to avoid the overfitting that often occurs with higher-order models. As a result, we selected 8 second-order models (i.e., those having only two features) with the highest fitness (PMF) among the total 520 second-order models. These 8 "best performing" models each have 100% PPV and 100% class coverage, see Table 3. All of the features identified were brain anatomical features.

Table 3. Second-order CC model features and range of values

CC #	Feature One	Value	Feature Two	Value
113	lh posteriorcingulate volume	[3500,4600]	lh rostralmiddlefrontal volume	[20000,25000]
679	lh posteriorcingulate volume	[3500,4600]	rh rostralmiddlefrontal volume	[21000,26000]
1449	lh posteriorcingulate volume	[3500,4600]	lh medialorbitofrontal thickness	[2.6,2.8]
2199	lh posteriorcingulate volume	[3500,4600]	rh rostralmiddlefrontal area	[6400,8500]
887	rh isthmuscingulate volume	[3300,4100]	rh posteriorcingulate volume	[4100,6200]
1488	rh isthmuscingulate volume	[3300,4100]	lh medialorbitofrontal thickness	[2.6,2.8]
2200	rh isthmuscingulate volume	[3300,4100]	rh rostralmiddlefrontal area	[6400,8500]
2269	rh isthmuscingulate volume	[3300,4100]	rh posteriorcingulate area	[1400,2900]

Using CC 113 (Table 3) as an example, this second-order model can be interpreted as: any subject whose posterior cingulate gyrus volume was within the range of 3500 to 4600 mm^3 AND left rostral middle frontal gyrus volume was within the range of 20,000 to 25,000 mm^3 would be classified as having ASD. The volume of the left

hemisphere posterior cingulate gyrus and the volume of the right hemisphere isthmus of

the cingulate gyrus were the two features to appear most frequently (i.e., four times)

across all second-order models, suggesting that the volume of cingulate gyrus is a

potentially important biomarker for ASD. Figure 1 provides a 2D visualization for the

range of feature values (numerical boundaries) associated with these models and the

placement of each subject within this range.

(a) CC113

(b) CC679

(c) CC1449

(d) CC2199

(e) CC887

(f) CC1488

(g) CC2200

(h) CC2269

Fig 1. 2D visualization of second-order CC models. Green dots represent ASD subjects and group together within the rectangle defining the range of values in Table 3.

Because of our desire to explore the predictive capability of the behavioral features, we expanded our analysis to include third-order models (i.e., model combinations with three features). There were 651 third-order models in total; while some consisted only of 310 anatomical brain features, others included two behavioral features plus one brain anatomical feature. We selected the 6 best-performing third-order models with the highest fitness (PMF); each had 100% PPV and 100% class coverage, see Table 4. Each of these third-order models contained two behavioral features and one brain anatomical feature.

Table 4. Third-order CC model features and range of values

CC #	Feature One	Value	Feature Two	Value	Feature Three	Value
46	ToMTB total	[5,13]	ToMI early subscale mean	[12,18]	lh parsorbitalis meancurv	[0.17,0.2]
191	ToMTB total	[5,13]	ToMI early subscale mean	[12,18]	rh superiorparietal thickness	[2.4,2.7]
264	ToMTB total	[5,13]	ToMI early subscale mean	[12,18]	rh parsorbitalis meancurv	[0.18,0.2]
317	ToMTB total	[5,13]	ToMI early subscale mean	[12,18]	rh inferiortemporal meancurv	[0.15,0.18]
1163	ToMTB total	[5,13]	ToMI total composite mean	[9,18]	lh postcentral thickness	[2.2,2.5]
1446	ToMTB total	[5,13]	ToMI early subscale mean	[12,18]	lh medialorbitofrontal thickness	[2.6,2.8]

Using CC 46 (Table 4) as an example, any subject who had a total score on ToMTB within the range of 5 to 13 AND an early subscale mean score on ToMI-2 within the range of 12 to 18 AND a mean curvature value of the left hemisphere pars orbitalis

within the range of 0.17 to 2 would be classified as having ASD. The ToMTB total score

feature occurred in all of our best-fit, third-order models; and the ToMI-2 early subscale

mean score occurred in all but one (CC 1163) of the models, where the ToMI-2 total

composite mean played a role. Such a finding further suggests that the ToMTB and

ToMI-2 might be effective for ASD testing and diagnosis. See Figure 2.

Fig 2. 3D visualization of third-order CC models. Green dots represent ASD subjects and group together within the pink cube defining the range of values in Table 4.

KNN Leave-One-Out Cross Validation

As mentioned earlier, a cohort of new subjects comprising 2 ASD and 5 NT children were enrolled at a later time. This later cohort was combined with the 21 subjects used in CCEA to cross-validate the KNN classifiers. Using the 8 unique features of the second-order model, the KNN (k=3) achieved 89.29% classification accuracy, where 17 of the 19 NT subjects and 8 of the 9 ASD were classified accurately; see Table 5, diagonals of the second-order confusion matrix.

Using the 9 unique features of the third-order model, the KNN (k=7) validation accuracy fell to 78.57% compared to when using the 8 unique features from the second-order model, where 15 of the 19 NT and 7 of the 9 ASD subjects were classified accurately. See Table 5, diagonals of third-order confusion matrix.

Given the better ASD classification performance of the second-order neuroanatomical features, and that the behavioral measurements/features are relatively easier to collect among children with ASD, it was important to explore whether the ASD prediction results might be improved when the behavioral features were combined with the second-order features. As a result, we added the three behavioral features (i.e.,ToMTB total score, ToMI-2 total composite mean and ToMI-2 early subscale mean) from the third-order models to the 8 second-order brain anatomical features and cross-validated a new KNN (k=2) classifier. With the total of 11 unique features, a validation accuracy of 85.71% was achieved with 16 of the 19 NT subjects and 8 of the 9 ASD being classified accurately. See the confusion matrix of Table 5.

Table 5. Cross Validation Confusion Matrices

N=28 (9 ASD & 19 NT)	Second-order Model Features *ACC Score: 89.29%*		Third-order Model Features *ACC Score: 78.57%*		Second-order Model Features Plus Three Behavioral Features *ACC Score: 85.71%*	
	Predicted ASD	**Predicted NT**	**Predicted ASD**	**Predicted NT**	**Predicted ASD**	**Predicted NT**
True ASD	8	1	7	2	8	1
True NT	2	17	4	15	3	16

Classification of ASD and NT subjects using the KNN model

To further examine whether the KNN classifier could discriminate subjects with ASD and NT, we developed three classification models – one using the 8 unique features from the second-order models, one using the 9 features from the third-order models, and one using the 11-feature model (i.e., eight second-order neuroanatomical features and three behavioral features). The best classification accuracy was achieved using a balanced training set that consisted of 6 NT and 6 ASD subjects, among which only 2 of the 6 ASD subjects were not part of the original CCEA feature selection. The remaining 16 subjects were used for testing (13 NT and 3 ASD).

The KNN (k=1) results for the second-order model features are shown in Table 6, columns 2 and 3; a classification accuracy of 87.5% was achieved, with all 3 of the ASD subjects and 11 of the 13 NT being classified accurately. Both of the misclassified NT subjects were part of the original CCEA feature selection.

The KNN (k=3) classification accuracy for the third-order model features was 81.25%, with all 3 of the ASD subjects and 10 of the 13 NT subjects correctly classified. Of the 3 misclassified NT subjects, 2 were included in the original CCEA feature

selection. See Table 6, columns 4 and 5.

Lastly, the KNN (k=3) predictions for the combined 11-feature model (8 neuroanatomical features and three behavioral features) are shown in Table 6, columns 6 and 7; classification accuracy is 93.75%, with all 3 of the ASD subjects and 12 out of 13 NT subjects classified accurately. The one misclassified NT subject was not part of the original CCEA feature selection analysis.

Table 6. Classification Confusion Matrices

N=16 (3 ASD & 13 NT)	Second-order Model Features ACC Score:87.5%		Third-order Model Features ACC Score: 81.25%		Second-order Model Features Plus Three Behavioral Features ACC Score: 93.75%	
	Predicted ASD	Predicted NT	Predicted ASD	Predicted NT	Predicted ASD	Predicted NT
True ASD	3	0	3	0	3	0
True NT	2	11	3	10	1	12

Discussion

This study used a new ML feature selection tool, the CCEA, to identify biomarkers and behavioral features capable of successfully discriminating between children (7 to 14 year of age) with and without ASD given a small dataset collected from a single research site. ML tools have long been applied to ASD research, but it remains a far-reaching goal to build a diagnostic system for ASD that incorporates both feature selection and prediction. Previous studies face the challenge of using datasets across different research sites for classification purposes, rather than identifying input variables most closely associated with ASD (i.e., feature selection) [17, 18, 32–35, 37–39, 67, 68]. Additionally, traditional ML algorithms do not work well with ASD datasets given the

large amount of variance and the heterogeneous nature of the disorder [40, 41]. Meanwhile, it requires tremendous effort to include ASD individuals in a research study given the social, cognitive, and language challenges of this population. Thus, nearly all ASD datasets have a large number of features with relatively small sample sizes which, despite being unsuitable for many ML algorithms, often leads to overfitting and poor classification accuracy. However, the CCEA in this work was able to address such issues by efficiently exploring large search spaces for feature interactions associated with some nominal outcome (e.g., ASD or NT). It also adopted built-in tools to prevent overfitting to produce parsimonious models.

The present study demonstrated exceptionally good performance (i.e., 100% ASD PPV and 100% class coverage) of the features identified by the CCEA. The selected CCEA features from the parsimonious second and third order models included volume, area, cortical thickness, and mean curvature in specific regions around the cingulate cortex, frontal cortex, and temporal-parietal junction areas as biomarkers for ASD (e.g., the pericalcarine cortex, posterior cingulate cortex, isthmus of the cingulate gyrus, pars orbitalis, etc.). Such findings are consistent with previous literature suggesting that individuals with ASD have abnormalities in these brain regions [17, 21–28, 69–72], which facilitates our identification of potential ASD biomarkers. Additionally, third-order models from the training set include measurements from the ToMI-2 and the ToMTB as important features [50, 53]. This provides evidence that outcome measures from ToMI-2 and ToMTB are able to distinguish ASD from NT which further validates the use of these tools in ASD assessments.

It is impressive that the KNN classifiers are able to achieve such high

classification accuracy given our sample size, and validation of the discriminant features

selected by the CCEA models. In particular, the KNN classifiers perform better using the

second-order neuroanatomical features than the third-order feature models, which

emphasizes the importance of focusing on parsimonious models selected by the CCEA.

In addition, the KNN achieved the highest classification accuracy when adding the

behavioral features from the third-order models to the neuroanatomical features from

the second-order models. In most cases, neuroimaging measurements are conducted

along with behavioral assessments. As the third-order models are included to explore

the potential role that behavioral features might play along the side of neuroanatomical

features, it is convincing to see the highest classification accuracy achieved when

combining the behavioral features and the neuroanatomical features. These findings

highlight the heterogeneous and multi-facet characteristics of ASD itself. Thus, although

it is more difficult to implement MRI among children with ASD, such findings support the

idea that neuroanatomical measurements increase confidence in diagnosis. It also

suggests that a good ASD prediction model should consider including both behavioral

and neuroanatomical features to establish brain-behavior connections to advance our

understanding of ASD.

     This study further demonstrates the robustness of the CCEA as a feature

selection methodology. The accuracy of these features when used as input variables in

the KNN classifier suggests their potential to help clinicians and researchers target

specific domains in ToM in treating the social challenges most often seen in children

with ASD. The implications of our findings for clinical researchers reinforce earlier

findings regarding the brain-behavior connections for children with and without ASD

related to ToM understanding [49, 51, 53, 73–75]. Knowing these connections may guide future researchers in the assessment of change following intervention at both a behavioral and neurobiological level. This may also lead to knowledge about which interventions may be most effective for children with specific neurobiological markers.

The present study has established important biomarker candidates of ASD. These biomarker candidates support previous research adopting traditional neuroimaging measurements identifying similar brain regions to explain the abnormalities in ASD [17, 21–28, 71]. Importantly, ML methodologies can perform as well as the traditional approaches in the field of neuroscience and specifically in our assessment of ASD in selecting neuroanatomical biomarkers. Although ML techniques have been adopted to help with diagnosis and treatment development in medicine [76–79], the heterogeneity in ASD creates challenges. Typically, large, diverse, and comprehensive datasets are required to extract solid biomarkers, which can be time-consuming and may be less accurate with traditional approaches. Under such circumstances, ML techniques as described in this study can help advance the development of an automatic diagnostic and predictive system for ASD. The present study provides a new direction for adopting ML techniques in ASD research and other areas of medicine with similar heterogeneity in disease conditions.

Acknowledgments

We thank Jay V. Gonyea, Administrative Director, and Scott Hipko, Senior Research Technologist, in the MRI Research Unit at the University of Vermont, for their support in acquiring the MRI scans. We thank Dr. Richard Watts, Ph.D., Director of the FAS Brain Imaging Center at Yale University, and Dr. Joseph Orr, Ph.D., Assistant Professor at Texas A&M University, for sharing their knowledge in MRI data pre-processing.

References

1. Xia-An Bi, Yang Wang, Qing Shu, Qi Sun, Qian Xu. Classification of Autism Spectrum Disorder Using Random Support Vector Machine Cluster. Front Genet. 2018 Feb 6;9:18. doi: 10.3389/fgene.2018.00018.

2. American Psychiatric Association. Diagnostic and Statistical Manual. 5th ed DSM-5. Washington, DC. 2013.

3. Michael Fitzgerald. The Clinical Gestalts of Autism: Over 40 years of Clinical Experience with Autism. Open access peer-reviewed chapter. 2017. doi: 10.577265906cf0.

4. Sebastien Levy, Marlena Duda, Nick Haber, Dennis P. Wall. Sparsifying machine learning models identify stable subsets of predictive features for behavioral detection of autism. Mol Autism. 2017; 8: 65.

5. Patrick F Bolton, Jean Golding, Alan Emond, Colin D Steer. Autism spectrum disorder and autistic traits in the Avon Longitudinal Study of Parents and Children: precursors and early signs. J Am Acad Child Adolesc Psychiatry. 2012 Mar;51(3):249-260.e25.

6. Jamie M. Kleinman, Pamela E. Ventola, Juhi Pandey et al. Diagnostic Stability in Very Young Children with Autism Spectrum Disorders. J Autism Dev Disord. 2008 Apr; 38(4): 606–615.

7. Catherine Lord, Michael Rutter, Ann Le Couteur. Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. Journal of Autism and Developmental Disorders volume 24. 1994. Pages 659–685.

8. C Lord, S Risi, L Lambrecht, E H Cook Jr, B L Leventhal, P C DiLavore, A Pickles, M Rutter. The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. J Autism Dev Disord. 2000 Jun;30(3):205-23.

9. Centers for Disease Control and Prevention. Prevalence of autism spectrum disorders–Autism and Developmental Disabilities Monitoring Network, 14 sites, United States, 2008. MMWR Surveill Summ. 2012 Mar 30;61(3):1-19.

10. Raphael Bernier, Alice Mao, Jennifer Yen. Psychopathology, families, and culture: autism. Child Adolesc Psychiatr Clin N Am. 2010 Oct;19(4):855-67.

11. David S Mandell, Maytali M Novak, Cynthia D Zubritsky. Factors associated with age of diagnosis among children with autism spectrum disorders. Pediatrics.

2005 Dec; 116(6):1480-6. doi: 10.1542/peds.2005-0185.
September 20, 2021.

12. David S Mandell, John Listerud, Susan E Levy, Jennifer A Pinto-Martin. Race differences in the age at diagnosis among medicaid-eligible children with autism. J Am Acad Child Adolesc Psychiatry. 2002 Dec;41(12):1447-53.

13. Morrier, Michael J, Hess, Kristen L, Heflin, L. Juane. Ethnic Disproportionality in Students with Autism Spectrum Disorders. Multicultural Education, p31-38 Fall 2008.

14. Rachel A Rhoades, Angela Scarpa, Brenda Salley. The importance of physician knowledge of autism spectrum disorder: results of a parent survey. BMC Pediatr. 2007 Nov 20;7:37. doi: 10.1186/1471-2431-7-37.

15. Lisa D Wiggins, Jon Baio, Catherine Rice. Examination of the time between first evaluation and first autism spectrum diagnosis in a population-based sample. J Dev Behav Pediatr. 2006 Apr;27(2 Suppl):S79-87. doi: 10.1097/00004703-200604002-00005.

16. Milan N. Parikh, Hailong Li, Lili He. Enhancing Diagnosis of Autism With Optimized Machine Learning Models and Personal Characteristic Data. Front Comput Neurosci. 2019; 13: 9.

17. M Duda, R Ma, N Haber, D P Wall. Use of machine learning for behavioral distinction of autism and ADHD. Transl Transl Psychiatry. 2016 Feb 9;6(2):e732. doi: 10.1038/tp.2015.221.

18. Daniel Bone, Matthew S Goodwin, Matthew P Black, Chi-Chun Lee, Kartik Audhkhasi, Shrikanth Narayanan. Applying machine learning to facilitate autism diagnostics: pitfalls and promises. J Autism Dev Disord. 2015 May;45(5):1121-36. doi: 10.1007/s10803-014-2268-6.

19. Ashwood, K. L., Gillan, N. et al. Predicting the diagnosis of autism in adults using the Autism-Spectrum Quotient (AQ) questionnaire. Psychological medicine. 2016. 46(12), 2595–2604. https://doi.org/10.1017/S0033291716001082

20. E Feczko, N M Balba et al. Subtyping cognitive profiles in Autism Spectrum Disorder using a Functional Random Forest algorithm. Neuroimage. 2018 May 15;172:674-688. doi: 10.1016/j.neuroimage.2017.12.044.

21. David G.Amaral, Cynthia Mills Schumann, Christine WuNordahl. Neuroanatomy of autism. Trends in Neurosciences. 2008. Volume 31, Issue 3, Pages 137-145.

22. Paolo Brambilla, Antonio Hardan, Stefania Ucelli di Nemi, Jorge Perez, Jair C Soares, Francesco Barale. Brain anatomy and development in autism: review of

structural MRI studies. Brain Res Bull. 2003 Oct 15;61(6):557-69. doi: 10.1016/j.brainresbull.2003.06.001.

23. Donna L Dierker, Eric Feczko et al. Analysis of cortical shape in children with simplex autism. Cereb Cortex. 2015 Apr;25(4):1042-51. doi: 10.1093/cercor/bht294.

24. R Kucharsky Hiess, R Alter et al. Corpus callosum area and brain volume in autism spectrum disorder: quantitative analysis of structural MRI from the ABIDE database. J Autism Dev Disord. 2015 Oct;45(10):3107-14. doi: 10.1007/s10803-015-2468-8.

25. Nicholas Lange, Brittany G Travers, Erin D Bigler et al. Longitudinal volumetric brain changes in autism spectrum disorder ages 6-35 years. Autism Res. 2015 Feb;8(1):82-93. doi: 10.1002/aur.1427. Epub 2014 Nov 7.

26. Christine Wu Nordahl, Donna Dierker, Iman Mostafavi et al. Cortical folding abnormalities in autism revealed by surface-based morphometry. J Neurosci. 2007 Oct 24;27(43):11725-35.

27. Mahsa Shokouhi, Justin H G Williams, Gordon D Waiter, Barrie Condon. Changes in the sulcal size associated with autism spectrum disorder revealed by sulcal morphometry. Autism Res. 2012 Aug;5(4):245-52. doi: 10.1002/aur.1232.

28. Sofie L Valk, Adriana Di Martino, Michael P Milham, Boris C Bernhardt. Multicenter mapping of structural network alterations in autism. Hum Brain Mapp. 2015 Jun;36(6):2364-73. doi: 10.1002/hbm.22776.

29. Gregory L Wallace, Ian W Eisenberg, Briana Robustelli, Nathan Dankner, Lauren Kenworthy, Jay N Giedd, Alex Martin. Longitudinal cortical development during adolescence and young adulthood in autism spectrum disorder: increased cortical thinning but comparable surface area changes. J Am Acad Child Adolesc Psychiatry. 2015 Jun;54(6):464-9. doi: 10.1016/j.jaac.2015.03.007.

30. Sungji Ha, In-Jung Sohn, Namwook Kim, Hyeon Jeong Sim, Keun-Ah Cheon. Characteristics of Brains in Autism Spectrum Disorder: Structure, Function and Connectivity across the Lifespan. Exp Neurobiol. 2015 Dec; 24(4): 273–284.

31. C Ecker, A Shahidiani, Y Feng, E Daly et al. The effect of age, diagnosis, and their interaction on vertex-based measures of cortical thickness and surface area in autism spectrum disorder. J Neural Transm (Vienna). 2014 Sep;121(9):1157-70. doi: 10.1007/s00702-014-1207-1.

32. Guillaume Chanel, Swann Pichon, Laurence Conty, Sylvie Berthoze, Coralie Chevallier, Julie Gr`ezes. Classification of autistic individuals and controls using cross-task characterization of fMRI activity. NeuroImage: Clinical, Volume 10,

2016, Pages 78-88.

33. Heng Chen, Xujun Duan, Feng Liu, Fengmei Lu, Xujing Ma, Youxue Zhang, Lucina Q Uddin, Huafu Chen. Multivariate classification of autism spectrum disorder using frequency-specific resting-state functional connectivity–A multi-center study. Prog Neuropsychopharmacol Biol Psychiatry. 2016 Jan 4;64:1-9.

34. Ilaria Gori, Alessia Giuliano, Filippo Muratori et al. Gray matter alterations in young children with autism spectrum disorders: comparing morphometry at the voxel and regional level. J Neuroimaging. Nov-Dec 2015;25(6):866-74. doi: 10.1111/jon.12280.

35. Yan Jin, Chong-Yaw Wee, Feng Shi, Kim-Han Thung, Dong Ni, Pew-Thian Yap, Dinggang Shen. Identification of infants at high-risk for autism spectrum disorder using multiparameter multiscale white matter connectivity networks. Hum Brain Mapp. 2015 Dec;36(12):4880-96. doi: 10.1002/hbm.22957.

36. John P Hanley, Donna M Rizzo, Jeffrey S Buzas, Margaret J Eppstein. A Tandem Evolutionary Algorithm for Identifying Causal Rules from Complex Data. Evol Comput. Spring 2020;28(1):87-114.

37. Paola Odriozola, Lucina Q Uddin, Charles J Lynch, John Kochalka, Tianwen Chen, Vinod Menon. Insula response and connectivity during social and non-social attention in children with autism. Soc Cogn Affect Neurosci. 2016 Mar;11(3):433-44. doi: 10.1093/scan/nsv126.

38. Colleen P Chen, Christopher L Keown, Afrooz Jahedi, Aarti Nair, Mark E Pflieger, Barbara A Bailey, Ralph-Axel M¨uller. Diagnostic classification of intrinsic functional connectivity highlights somatosensory, default mode, and visual regions in autism Neuroimage Clin. 2015 Apr 9;8:238-45. doi:10.1016/j.nicl.2015.04.002.

39. Cameron Craddock, Yassine Benhajali, Carlton Chu et al. The Neuro Bureau Preprocessing Initiative: open sharing of preprocessed neuroimaging data and derivatives. In: Front. Neuroinform. 2013.

40. Wolfgang Huf, Klaudius Kalcher, Roland N Boubela et al. On the generalizability of resting-state fMRI machine learning classifiers. Front Hum Neurosci. 2014 Jul 29;8:502. doi: 10.3389/fnhum.2014.00502.

41. Clare Kelly, Bharat B Biswal, R Cameron Craddock, F Xavier Castellanos, Michael P Milham. Characterizing variation in the functional connectome: promise and pitfalls. Trends Cogn Sci. 2012 Mar;16(3):181-8. doi:10.1016/j.tics.2012.02.001.

42. Pradyumna Lanka, D Rangaprakash, Michael N Dretsch, Jeffrey S Katz, Thomas S Denney Jr, Gopikrishna Deshpande. Supervised machine learning for diagnostic classification from large-scale neuroimaging datasets. Brain Imaging Behav. 2020 Dec;14(6):2378-2416. doi: 10.1007/s11682-019-00191-8.

43. Baihua Li, Arjun Sharma, James Meng, Senthil Purushwalkam, Emma Gowen. Applying machine learning to identify autistic adults using imitation: An exploratory study. PLOS ONE. 2017.

44. Zbigniew Michalewicz, Marc Schoenauer. Evolutionary Algorithms. Encyclopedia of Information Systems. 2003.

45. M. Rutter, A. Bailey, and C. Lord. The Social Communication Questionnaire Manual. Western Psychological Services, 2003.

46. Bruce A. Bracken, R. Steve McCallum. Universal nonverbal intelligence test (2nd ed.). Western Psychological Services, 2016.

47. R.S. McCallum. Handbook for Nonverbal Assessment. Springer, 2017.

48. T.L. Hutchins, P.A. Prelock, W. Chase. Test-retest reliability of a theory of mind task battery for children with autism spectrum disorders. Focus Autism Other Dev Disabil. 2008.

49. T.L. Hutchins and P.A. Prelock. Technical Manual for the Theory of Mind Inventory-2. Copyrighted manuscript. Theoryofmindinventory.com. 2016.

50. T.L. Hutchins, P.A. Prelock, L. Bonazinga. Psychometric evaluation of the theory of mind inventory (tomi): a study of typically developing children and children with autism spectrum disorder. J Autism Dev Disord. 2012.

51. S. Baron-Cohen. An essay on autism and theory of mind. MIT Press. 1995.

52. S. Baron-Cohen, A. M. Leslie, U. Frith. Does the autistic-child have a theory of mind. Cognition. 1985.

53. Matthew D Lerner, Tiffany L Hutchins, Patricia A Prelock. Brief report: preliminary evaluation of the theory of mind inventory and its relationship to measures of social skills. J Autism Dev Disord. 2011 Apr;41(4):512-7. doi: 10.1007/s10803-010-1066-z.

54. Julie Hadwin, Josef Perner. Pleased and surprised: Children's cognitive theory of emotion. British Journal of Developmental Psychology. 1991.

55. A. Hillier, L. Allinson. Understanding embarrassment among those with autism: Breaking down the complex emotion of embarrassment among those with autism.

Journal of Autism and Developmental Disorders. 2002.

56. T. Ruffman, T.R. Keenan. The belief-based emotion of surprise: The case for a lag in under- standing relative to false belief. Developmental Psychology. 1996.

57. B. Seider. A developmental analysis of elementary school-aged children's concepts of pride and embarrassment. Child Development. 1988.

58. Getting an MRI: A Cartoon for Kids. https://www.youtube.com/watch?v=Q Pa6KFL1Nw%5C&t=139s%5Ccf2.

59. FreeSurfer. https://surfer.nmr.mgh.harvard.edu/fswiki/aparcstats2table.

60. Matthew F Glasser, Stamatios N Sotiropoulos, J Anthony Wilson et al. The minimal preprocessing pipelines for the Human Connectome Project. Neuroimage. 2013 Oct 15;80:105-24. doi: 10.1016/j.neuroimage.2013.04.127.

61. Bruce Fischl, David H Salat, Evelina Busa et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron. 2002 Jan 31;33(3):341-55.
62. Regina Nuzzo. Scientific method: Statistical errors. Nature. 2014.

63. R.L. Wasserstein, A.L. Schirm, N.A. Lazar. Moving to a World Beyond p <0.05. The American Statistician. 2019.

64. R. L. Wasserstein, A. L. Schirm, and N. A. Lazar. The ASA 92s statement on p-values: context, process, and purpose. The American Statistician. 2016.

65. Mucherino A., Papajorgji P.J., Pardalos P.M. k-Nearest Neighbor Classification. In: Data Mining in Agriculture. Springer Optimization and Its Applications. Springer, New York, NY. 2009. https://doi.org/10.1007/978-0-387-88615-2 4.

66. Mucherino A., Papajorgji P.J., Pardalos P.M. Cross-Validation. Encyclopedia of Machine Learning. Springer, Boston, MA. 2011. https://doi.org/10.1007/978-0-387-30164-8 190.

67. D. Li, W. Yang, S. Wang. Classification of foreign fibers in cotton lint using machine vision and multi-class support vector machine. Electron. Agric. 2010

68. Y. Zhang, L. Wu. Classification of fruits using computer vision and a multiclass support vector machine. Sensors. 2012.

69. K. Nickel et al. Volume Loss Distinguishes Between Autism and (Comorbid) Attention- Deficit/Hyperactivity Disorder FreeSurfer Analysis in Children. Frontiers in Psychiatry.2018.

70. M. C. Postema, D. van Rooij, E. Anagnostou. Altered structural brain asymmetry in autism spectrum disorder in a study of 54 datasets. Nat Commun. 2019.

71. G. L. Wallace. Longitudinal cortical development during adolescence and young adulthood in autism spectrum disorder: Increased cortical thinning but comparable surface area changes. Acad. Child Adolesc. Psychiatry.2015.

72. B.A. Zielinski et al. Longitudinal changes in cortical thickness in autism and typical development. Brain. 2014.

73. R.K. Kana, J.O. Maximo, D.L. Williams et al. Aberrant functioning of the theory-of-mind network in children and adolescents with autism. Molecular Autism. 2015.

74. Mayumi Takeuchi, Masafumi Harada, Hiromu Nishitani. Deficiency of "theory of mind" in autism estimated by fMRI. International Congress Series. April 2002, Pages 737-740.

75. E. O'Nions et al. Neural bases of theory of mind in children with autism spectrum disorders and children with conduct problems and callous-unemotional traits. Developmental Science. 2014.

76. Ahmed, Z., Mohamed, K., Zeeshan, S., Dong, X. Artificial intelligence with multi-functional machine learning platform development for better healthcare and precision medicine. The journal of biological databases and curation. 2020

77. Editorial. Ascent of machine learning in medicine. Nat. Mater. 2019.

78. Shah, P., Kendall, F., Khozin, S. et al. Artificial intelligence and machine learning in clinical development: a translational perspective. Digit. Med. 2019.

79. Davenport, T., Kalakota, R. The potential for artificial intelligence in healthcare. Future healthcare journal. 2019.

80. Centers for Disease Control and Prevention. 2020 Community Report on Autism.

81. Elizabeth Carrow-Woolfolk. Comprehensive Assessment of Spoken Language-Second Edition. Pearson.2017.

82. F. Thabtah and D. Peebles. A new machine learning model based on induction of rules for autism detection. Health Informatics Journal. 2019.

83. Guo X, Dominick KC, Minai AA, Li H, Erickson CA, Lu LJ. Diagnosing Autism Spectrum Disorder from Brain Resting-State Functional Connectivity Patterns Using a Deep Neural Network with a Novel Feature Selection Method. Front Neurosci. 2017 Aug 21;11:460. doi: 10.3389/fnins.2017.00460. PMID: 28871217; PMCID:

PMC5566619.