A genetic perspective on the relationship between non-cancerous gynecological diseases and endometrial cancer

Authors

Pik Fang Kho[1, 2], Sally Mortlock[3], Endometrial Cancer Association Consortium, International Endometriosis Genetics Consortium, Peter A.W. Rogers[4], Dale R. Nyholt[2,5], Grant W. Montgomery[3], Amanda B. Spurdle[1], Dylan M. Glubb[1*], Tracy A. O'Mara[1*].

Affiliations

[1]Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia.

[2]School of Biomedical Science, Faculty of Health, Queensland University of Technology, Brisbane, Queensland, Australia.

[3]The Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland, Australia.

[4]Department of Obstetrics and Gynaecology, Gynaecology Research Centre, Royal Women's Hospital, University of Melbourne, Parkville, Victoria, Australia.

[5]Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia.

[*]These authors contributed equally to the work.

Corresponding Author

Dr Tracy O'Mara, PhD, Molecular Cancer Epidemiology Group, QIMR Berghofer Medical Research Institute, 300 Herston Road, Brisbane QLD Australia 4006. Phone: +61 7 3362 0389. Email: Tracy.OMara@qimrberghofer.edu.au

Declarations

Funding

PFK is supported by an Australian Government Research Training Program PhD Scholarship and QIMR Berghofer Postgraduate Top-Up Scholarship, TAO'M, GWM and ABS are supported by NHMRC Investigator Fellowships (APP1173170, GNT1177194 and APP177524).

This work was supported by National Health and Medical Research Council (NHMRC) Project Grants (APP1109286, GNT1026033, GNT1105321 and GNT1147846). Funding sources had no role in study design, data curation and analysis, data interpretation, report writing and submission for publication.

Conflicts of interest/Competing interests

The authors declare no potential conflicts of interest.

Ethics approval

This work used summary-level GWAS meta-analysis results, and thus ethical approval was not required.

Consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material (data transparency)

Summary-level GWAS meta-analysis results for PCOS, uterine fibroids and endometrial cancer that support the findings of this study are available at the NHGRI-EBI GWAS Catalog (https://www.ebi.ac.uk/gwas/downloads/summary-statistics). Other data generated and/or analyzed during this study are included in this article and its supplementary information files or are available on reasonable request.

Code availability

Not applicable

Abstract

The non-cancerous gynecological diseases, endometriosis, polycystic ovary syndrome (PCOS) and uterine fibroids, have been proposed as endometrial cancer risk factors; however, disentangling their relationships is complicated due to their shared risk factors and comorbidity. Using genome-wide association study (GWAS) summary data, we have explored the relationship between these non-cancerous gynecological diseases and endometrial cancer risk, assessing genetic correlations, causal relationships and shared genetic risk regions. Firstly, we found significant genetic correlation between endometrial cancer and PCOS ($r_G = 0.36$, se = 0.12, P = 1.6×10^{-3}), and uterine fibroids ($r_G = 0.24$, se = 0.09, P = 5.4×10^{-3}). Adjustment for genetically predicted body mass index (BMI; a risk factor for PCOS, uterine fibroids and endometrial cancer) substantially attenuated the genetic correlation between endometrial cancer and PCOS, but not uterine fibroids. Despite the observed genetic correlation, genetic causal inference tests (latent causal variable and Mendelian randomization analyses) did not support a causal relationship between any of the non-cancerous gynecological diseases and endometrial cancer. Gene-based association analysis revealed four shared endometriosis and endometrial cancer risk loci (9p21.3, 15q15.1, 17q21.32 and 3q21.3) and two shared uterine fibroid and endometrial cancer risk loci (5p15.33 and 11p13). In summary, we have shown that PCOS and uterine fibroids are genetically correlated with endometrial cancer, although the genetic architecture shared between endometrial cancer and PCOS likely relates to BMI. Furthermore, shared genetic risk regions, and thus potentially shared causal genes, were identified between the risk for endometrial cancer and endometriosis, and uterine fibroids.

Keywords

Endometrial cancer, endometriosis, polycystic ovary syndrome, uterine fibroids, genetic correlation, Mendelian randomization, causal relationship, gene-based analysis

Introduction

Endometriosis, polycystic ovary syndrome (PCOS) and uterine fibroids are three prevalent non-cancerous gynecological diseases affecting 10-15% (Parasar et al. 2017), 6-9% (Azziz et al. 2011) and 5-69% (Stewart et al. 2017) of women of reproductive age, respectively.

However, the prevalence of these three diseases is likely to be underestimated because of underdiagnosis (Agarwal et al. 2019; De La Cruz and Buchanan 2017). Although the three non-cancerous gynecological diseases primarily affect premenopausal women and endometrial cancer is largely a postmenopausal malignancy, many risk factors are shared (e.g. chronic estrogen exposure, inflammation, insulin resistance and obesity (Harris and Terry 2016; Li et al. 2019; Wise et al. 2016)), supporting a relationship between them.

A number of studies have used observational data to assess associations between the three non-cancerous gynecological diseases and endometrial cancer risk, the findings of which have been heterogeneous (Harris and Terry 2016; Johnatty et al. 2020; Li et al. 2019; Wise et al. 2016). Indeed, the use of observational studies to evaluate these associations can be confounded by: (i) the failure to adequately account for potential confounders e.g. oral contraceptive use (a common treatment for endometriosis and PCOS, which is associated with reduced endometrial cancer risk); (ii) the reliance of disease status classification on self-reported data which is subject to misclassification bias from asymptomatic undiagnosed cases; (iii) misdiagnosis of early stage endometrial cancer as uterine fibroids due to shared clinical presentation; (iv) detection bias in cohort studies as a result of an increased surveillance for endometrial cancer among patients with non-cancerous gynecological diseases; and (v) the comorbidity of non-cancerous gynecological diseases e.g. patients with uterine fibroids can concurrently suffer from endometriosis (Choi et al. 2017; Johnatty et al. 2020; Matalliotaki et al. 2018; Nagai et al. 2015; Uimari et al. 2011) or PCOS (Wise et al. 2007). Thus, it remains difficult to determine from observational studies if there is a true causal association between these non-cancerous gynecological diseases and endometrial cancer.

Genome-wide association study (GWAS) data have demonstrated genetic overlap between endometrial cancer and endometriosis (Masuda et al. 2020; Painter et al. 2018), and uterine fibroids (Masuda et al. 2020), which may partly explain the comorbidities of these diseases. However, whether these comorbidities are due to causal relationships or shared genetic etiology remains to be explained. In this study, we have further used GWAS data to disentangle associations between non-cancerous gynecological disease and endometrial cancer. As inherited genetic variants are non-modifiable, genetic approaches to explore

relationships between phenotypes are less influenced by confounding inherent in observational studies. Firstly, we have performed genetic correlation analysis, using the largest currently available datasets to clarify the shared genetic risk between the non-cancerous gynecological diseases and endometrial cancer. To investigate causal relationships, we have assessed genetic causal inference through analyses that use genetic variants associated with the gynecological disease of interest as instruments. Lastly, it is possible that these diseases may not be causally related to endometrial cancer but rather develop through effects of disease-specific risk variants at shared risk loci. To assess this possibility, we have performed gene-based analysis of each GWAS to identify shared genetic risk loci.

Materials and Methods

GWAS data

GWAS summary data were publicly available for PCOS (Day et al. 2018) (https://doi.org/10.17863/CAM.27720) and uterine fibroids (Gallagher et al. 2019) (ftp://ftp.ebi.ac.uk/pub/databases/gwas/summary_statistics/GallagherCS_31649266_GCST009158). For endometriosis, we accessed data (Sapkota et al. 2017) via collaboration. For PCOS and uterine fibroids, publicly available summary data were based on analyses excluding the 23andMe, Inc., cohort in the original publication, due to data sharing agreements with the company (Day et al. 2018; Gallagher et al. 2019). For Mendelian randomization analyses, risk estimates and respective standard errors of genome-wide significant variants were accessed from the largest published GWAS for each disease. Details of studies and sample sizes used in each analysis are shown in **Table 1.** Detailed descriptions of the quality control procedures and GWAS analysis can be found in the corresponding publications.

GWAS summary data for endometrial cancer were derived from O'Mara et al. (2018). As the GWAS for endometrial cancer (O'Mara et al. 2018), endometriosis (Rahmioglu et al. 2018), and uterine fibroids (Gallagher et al. 2019) included participants from UK Biobank, we re-analyzed the endometrial cancer dataset, excluding these participants to avoid sample overlap bias in the two sample Mendelian randomization analysis. This revised endometrial cancer GWAS meta-analysis consisted of 12,270 cases and 46,126 controls of European descent.

Genetic variants with minor allele frequency (MAF) < 1% and imputation information score < 0.4 were filtered, leaving ~9 million genetic variants. This revised endometrial cancer GWAS was used only in Mendelian randomization analysis, while the published endometrial cancer GWAS (O'Mara et al. 2018) was used in all other analyses. Prior to genetic correlation and latent causal variable (LCV) analyses, genetic variants in the extended human major histocompatibility complex region (26–34 Mb on chromosome 6) were removed due to the complex linkage disequilibrium (LD) structure in this region.

Genetic correlation between non-cancerous gynecological diseases and endometrial cancer
We used LD Score regression (Bulik-Sullivan et al. 2015) to estimate the genetic correlation between each non-cancerous gynecological disease and endometrial cancer. Genetic correlation analyses were restricted to common HapMap3 variants (MAF > 0.01). To reduce bias from potential residual confounding in genetic correlation analyses, including unknown sample overlap, we used the estimated genetic covariance intercept, obtained without constraint. Genetic correlation values range from -1 to 1; positive values indicated that shared genetic variants have concordant effects across the genome, whereas negative values indicated divergent effects.

Evidence of genetic correlation may reflect a causal relationship or sharing of genetic pathways. Obesity is a major risk factor for endometrial cancer, and is prevalent amongst women with PCOS and uterine fibroids (Ilaria and Marci 2018; Sam 2007). For genetic correlation of PCOS or uterine fibroids with endometrial cancer, we thus additionally corrected for the effect of obesity, as measured by genetically predicted BMI. PCOS and uterine fibroids GWAS were conditioned using summary data from a large GWAS of BMI (Yengo et al. 2018) in GCTA-mtCOJO analysis (Zhu et al. 2018) before performing LD score regression analysis.

Genetic causal inference tests

We conducted LCV analysis using GWAS summary data to estimate the proportion of genetic components for non-cancerous gynecological diseases that also affect endometrial cancer risk, while correcting for heritability and genetic correlation between traits (O'Connor and Price 2018). This method is robust to confounding due to pleiotropy. The genetic causality proportion (GCP) derived from LCV analysis ranges from -1 to 1. GCP values close to 1 indicate that genetic components for non-cancerous gynecological disease may affect endometrial cancer risk, while GCP values close to -1 indicate that genetic components for endometrial cancer may affect non-cancerous gynecological disease risk.

To further explore potential causal relationships, we performed two-sample Mendelian randomization analysis using independent (LD $r^2 < 0.01$) genetic variants associated with the non-cancerous gynecological diseases at genome wide significance ($P < 5 \times 10^{-8}$) as instruments. The list of genetic instruments and the respective risk association estimates were extracted from the largest GWAS of endometriosis (Rahmioglu et al. 2018), PCOS (Day et al. 2018) and uterine fibroids (Gallagher et al. 2019). We filtered out independent genetic variants with ambiguous alleles and intermediate frequencies (i.e., variants with A/T or C/G alleles and minor allele frequency of more than 0.42), leaving 26 variants as genetic instruments for endometriosis, 14 for PCOS and 25 for uterine fibroids.

As non-cancerous gynecological diseases mostly affect premenopausal women and endometrial cancer primarily affects postmenopausal women, we performed a unidirectional analysis, assessing the effect of genetic predisposition to non-cancerous gynecological disease on endometrial cancer risk. We used inverse variance weighted (IVW) analysis as the primary analysis by regressing the genetic variant-endometrial cancer association on the genetic variant-non-cancerous gynecological disease association, weighted by inverse of their variance. This method has the most power to detect associations; however it has a strong assumption of no heterogeneity (potentially resulting from pleiotropy) amongst genetic variants (Hemani et al. 2018). Thus, this method assumes all genetic variants for the exposure of interest have a proportional effect on outcome risk.

We also performed several sensitivity analyses which are more robust to heterogeneity amongst genetic variants: MR-Egger, weighted median, and weighted mode analysis. MR-Egger analysis regresses genetic variant-outcome association on genetic variant-exposure association, without a constraint on the regression intercept (Bowden et al. 2015). If the MR-Egger regression intercept is non-zero, it provides evidence of horizontal pleiotropy amongst genetic variants (where a genetic variant belongs to more than one independent pathway that influences outcome risk). The MR-Egger regression slope represents valid effect estimates after controlling for pleiotropic effects, provided the Instrument Strength Independent of Direct Effect (InSIDE) assumption (where a genetic variant's association with the exposure of interest is independent from its direct effect on outcome) is met (Bowden et al. 2015). We also performed weighted median (Bowden et al. 2016) and weighted mode (Hartwig et al. 2017) analyses, which are more robust to violation of the InSIDE assumption. Weighted median analysis relies on the assumption that more than 50% of the weights come from valid genetic instruments (Bowden et al. 2016), while weighted mode analysis relies on the assumption that most of the weights come from valid genetic instruments (Hartwig et al. 2017).

LCV and two-sample Mendelian randomization analysis were performed using the CTG-VL platform (Cuéllar-Partida et al. 2019) and the "TwoSampleMR" (Hemani et al. 2018) package in R, respectively. Unless stated otherwise, results with a Bonferroni-corrected p-value for testing the three non-cancerous gynecological diseases ($P < 0.05/3 = 0.017$) were considered statistically significant.

Gene-based analysis

While genetic correlation analysis assesses the average genetic concordance across the genome for two traits, it does not reveal common local genomic regions that harbor trait-associated variation. Further, a lack of evidence for genetic correlation may reflect opposing pleiotropic effects across the genome. To identify genetic risk regions shared between the non-cancerous gynecological diseases and endometrial cancer, we performed gene-based

analysis using the fast and flexible set-based association test (fastBAT) (Bakshi et al. 2016). fastBAT performs enrichment analysis on sets of GWAS variants located within 50kb of gene regions. A random sample of 10,000 unrelated participants from the UK Biobank was used as the reference panel in these analyses. We applied a false discovery rate (FDR) < 0.05 for the gene-based analysis and adjacent identified genes were combined into a single locus if within 1 Mb of each other.

Results

We found endometrial cancer was significantly genetically correlated with PCOS (r_G = 0.36, se = 0.12, P = 1.6×10^{-3}) and uterine fibroids (r_G = 0.24, se = 0.09, P = 5.4×10^{-3}) but not with endometriosis (r_G = -0.02, se = 0.09, P = 0.83) (**Table 2**). After adjusting for genetically predicted BMI, the genetic correlation between PCOS and endometrial cancer decreased and was no longer significant (r_G = 0.19, se = 0.14, P = 0.17), indicating that the initial genetic correlation was, at least partly, mediated by genetically predicted BMI (**Table 2**). In contrast, there was no material difference in the genetic correlation between uterine fibroids and endometrial cancer after adjusting for genetically predicted BMI (**Table 2**).

The LCV analysis provided no evidence to suggest genetic components of the non-cancerous gynecological diseases affect endometrial cancer risk (**Table 3**). Mendelian randomization analyses also provided no evidence of causal relationships between genetic predisposition to the non-cancerous gynecological diseases and endometrial cancer risk (**Table 3, Figure 1**). No MR-Egger intercepts differed from zero (**Table 3**), indicating a lack of evidence for confounding by horizontal pleiotropy amongst genetic instruments.

Gene-based analyses by fastBAT revealed 24 genetic regions associated with endometrial cancer risk, 28 regions with endometriosis risk and 41 regions with uterine fibroids, while no associations with PCOS passed FDR < 0.05 (**Supplementary Table 1**). We found four shared genetic risk regions (3q21.3, 9p21.3, 15q15.1 and 17q21.32) for endometriosis and endometrial cancer, containing seven shared candidate susceptibility genes (**Table 4**). Three of these regions (9p21.3, 15q15.1 and 17q21.32) have independently been associated with endometrial cancer and endometriosis risk through GWAS (O'Mara et al. 2018; Rahmioglu et al. 2018). The LD of lead risk variants at each gene was compared and one region (17q21.32) demonstrated evidence of a shared genetic risk signal (r^2 > 0.9; **Table 4**). Additionally, we

found two shared genetic risk regions (5p15.33 and 11p13) for uterine fibroids and endometrial cancer, containing five shared candidate susceptibility genes (**Table 5**). Through GWAS, one of these regions (5p15.33) has been associated with uterine fibroids risk (Gallagher et al. 2019) and the other (11p13) has independently associated with uterine fibroids and endometrial cancer risk (Gallagher et al. 2019; O'Mara et al. 2018). The LD of lead risk variants at each gene was compared but there was no strong genetic correlation at either of these regions ($r^2 \leq 0.4$; **Table 5**), suggesting that these genetic risk signals may be independent.

Discussion

Using large-scale genome-wide datasets, we observed evidence of positive genetic correlation between endometrial cancer and PCOS, and uterine fibroids, but not endometriosis. The observed genetic correlation between endometrial cancer and PCOS was at least partly mediated by genetically predicted BMI, consistent with the role of BMI as a common risk factor for PCOS and endometrial cancer. Further genetic analyses provided no evidence for a causal relationship between the non-cancerous gynecological diseases and endometrial cancer but revealed several genetic risk regions shared between endometrial cancer and endometriosis, and uterine fibroids.

Previous studies have demonstrated biological links between endometriosis and endometrial cancer (reviewed in Bulun (2009)), but we found no evidence for genetic correlation. However, two previous studies have reported a positive genetic correlation (Masuda et al. 2020; Painter et al. 2018). These discrepancies with our study may be related to: i) the smaller sample sets used by these studies; ii) the ethnicity studied (Masuda et al. (2020) analyzed a Japanese population); or iii) the different genetic correlation analysis approaches used. For example, unlike Painter et al. (2018), we used an unconstrained LD score regression intercept to account for potential residual confounding, resulting in a conservative estimate of genetic correlation. Indeed, we found the estimated genetic covariance intercept to be significantly different from zero, suggesting the presence of bias from population stratification and/or sample overlap. The null results from the genetic causal inference analyses of endometriosis and endometrial cancer are concordant with observational studies

that observed no associations after controlling for ascertainment bias by excluding recent endometriosis diagnosis (Melin et al. 2007; Olson et al. 2002; Rowlands et al. 2011).

Despite the limited evidence for genetic correlation and causal relationships between endometriosis and endometrial cancer, we identified four shared genetic risk regions, three of which (9p21.3, 15q15.1 and 17q21.32) have independently been associated with risk of both diseases through GWAS (O'Mara et al. 2018; Rahmioglu et al. 2018). The shared candidate susceptibility genes at 9p21.3 (C*DKN2B-AS1*), 15q15.1 (*BMF*) and 17q21.32 (*CBX1*, *MIR1203*, *SKAP1* and *SNX11)* have been previously identified as candidate target genes at endometrial cancer GWAS risk loci through chromatin looping studies in endometrial cell lines (O'Mara et al. 2019). The remaining shared endometriosis and endometrial cancer risk region at 3q21.3 has not been previously identified by GWAS and may represent a future GWAS risk locus for both diseases. This region was recently reported as an endometrial cancer risk region in a cross-cancer meta-analysis of endometrial, breast, ovarian and prostate cancer (Kar et al. 2020). However, larger GWAS for endometriosis and endometrial cancer, with more statistical power, will be required to detect genome-wide significant associations at this region.

The null associations identified between PCOS and endometrial cancer risk from genetic causal inference analyses is concordant with observational studies that have found no association after accounting for the effect of obesity (Fearnley et al. 2010; Zucchetto et al. 2009). Indeed, these findings are consistent with our observation of substantial attenuation in genetic correlation between PCOS and endometrial cancer after adjusting for genetic components of BMI.

Although we detected a genetic correlation between uterine fibroids and endometrial cancer risk, consistent with observational studies (Fortuny et al. 2009; Rowlands et al. 2011; Wise et al. 2016), no evidence for a causal association was found. fastBAT analysis revealed two genetic risk regions (5p15.33 and 11p13) that may be shared by both diseases. The 5p15.33 region has been associated with uterine fibroids in GWAS (Gallagher et al. 2019) and with endometrial cancer in a candidate locus study (Carvajal-Carmona et al. 2015). Three

candidate risk genes were shared between uterine fibroids and endometrial cancer at 5p15.33. The most biologically relevant of these is *TERT,* which encodes telomerase reverse transcriptase and maintains chromosomal stability by elongating the telomere (Rubtsova et al. 2012). Relevantly, chromosomes in uterine fibroids (Bonatz et al. 1998; Rogalla et al. 1995) and in endometrial tumors (reviewed by Alnafakh et al. (2019)) have been shown to have shorter telomere length. Telomere shortening could induce genomic instability and thus abnormal cellular growth. Indeed, a Mendelian randomization study found genetically predisposed telomere length strongly predicted endometrial cancer risk (Telomeres Mendelian Randomization et al. 2017). The 11p13 shared risk region has been associated with both uterine fibroids (Gallagher et al. 2019) and endometrial cancer (O'Mara et al. 2018) risk in GWAS. *WT1* and *WT1-AS* were revealed as shared candidate susceptibility genes for uterine fibroids and endometrial cancer at this region. We have previously identified *WT1* and *WT1-AS* as candidate targets of endometrial cancer risk GWAS variants through chromatin looping studies (O'Mara et al. 2019). *WT1* has also been identified through chromatin looping as a candidate target of uterine fibroids risk variants (Rafnar et al. 2018). *WT1* encodes a transcription factor that is essential for urogenital development (reviewed by Roberts (2005)) and in the GTEx database is most highly expressed in the uterus (https://gtexportal.org/home/). These observations suggest that alteration of uterine *WT1* expression by risk variation associated with endometrial cancer and uterine fibroids may affect susceptibility to these diseases.

Despite the genetic causal inference test analyses providing no evidence for causal relationships, we cannot discard the possibility that non-cancerous gynecological diseases have modest causal effects on endometrial cancer. Nevertheless, our findings indicate that the shared comorbidities are more likely to be related through shared genetic risk elements and thus women who have had non-cancerous gynecological disease may be at a greater risk of developing endometrial cancer. Future research will be needed to clarify such relationships.

Comorbidities may complicate our interpretation of the relationships between the non-cancerous gynecological diseases and endometrial cancer revealed in observational studies. To reduce this confounding, prospective studies with long follow-up, large sample sizes and case identification using surgical confirmation would ideally be required. A strength of our

study is that genetic causal inference analysis is a cost-effective alternative approach that allows assessment of potential causal relationships while reducing bias from unmeasured confounding. Another strength is that we have demonstrated the use of gene-based analysis to detect genetic risk regions below a genome-wide significance threshold. A limitation of the gene-based analysis is that many nearby genes are likely to share GWAS variants, thus the role of the identified candidate susceptibility genes requires functional follow-up. Further, we have limited ability to identify candidate genetic risk regions and causal relationships for PCOS given the smaller dataset available. These analyses should be revisited when more genome-wide significant variants are revealed in future PCOS GWAS.

In conclusion, our study has revealed genetic risk variation shared between non-cancerous gynecological diseases and endometrial cancer, both across the genome and at local genomic regions, providing insights into the shared etiology of these diseases.

Acknowledgement

This work was conducted using the UK Biobank Resource (application number 25331). We thank the research participants and employees of 23andMe for making this work possible. We thank the many individuals who participated in the Endometrial Cancer Association Consortium and the International Endometriosis Genetics Consortium studies, and the numerous institutions and their staff who supported recruitment. A full list of consortium members and acknowledgements can be found in the Supplementary Note.

References

Agarwal SK et al. (2019) Clinical diagnosis of endometriosis: a call to action Am J Obstet Gynecol 220:354 e351-354 e312 doi:10.1016/j.ajog.2018.12.039

Alnafakh RAA, Adishesh M, Button L, Saretzki G, Hapangama DK (2019) Telomerase and Telomeres in Endometrial Cancer Front Oncol 9:344 doi:10.3389/fonc.2019.00344

Azziz R, Dumesic DA, Goodarzi MO (2011) Polycystic ovary syndrome: an ancient disorder? Fertility and sterility 95:1544-1548 doi:10.1016/j.fertnstert.2010.09.032

Bakshi A, Zhu Z, Vinkhuyzen AA, Hill WD, McRae AF, Visscher PM, Yang J (2016) Fast set-based association analysis using summary data from GWAS identifies novel gene loci for human complex traits Sci Rep 6:32894 doi:10.1038/srep32894

Bonatz G, Frahm SO, Andreas S, Heidorn K, Jonat W, Parwaresch R (1998) Telomere shortening in uterine leiomyomas American journal of obstetrics and gynecology 179:591-596 doi:10.1016/s0002-9378(98)70050-x

Bowden J, Davey Smith G, Burgess S (2015) Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression Int J Epidemiol 44:512-525 doi:10.1093/ije/dyv080

Bowden J, Davey Smith G, Haycock PC, Burgess S (2016) Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator Genet Epidemiol 40:304-314 doi:10.1002/gepi.21965

Bulik-Sullivan B et al. (2015) An atlas of genetic correlations across human diseases and traits Nat Genet 47:1236-1241 doi:10.1038/ng.3406

Bulun SE (2009) Endometriosis The New England journal of medicine 360:268-279 doi:10.1056/NEJMra0804690

Carvajal-Carmona LG et al. (2015) Candidate locus analysis of the TERT-CLPTM1L cancer risk region on chromosome 5p15 identifies multiple independent variants associated with endometrial cancer risk Hum Genet 134:231-245 doi:10.1007/s00439-014-1515-4

Choi EJ, Cho SB, Lee SR, Lim YM, Jeong K, Moon HS, Chung H (2017) Comorbidity of gynecological and non-gynecological diseases with adenomyosis and endometriosis Obstetrics & gynecology science 60:579-586 doi:10.5468/ogs.2017.60.6.579

Cuéllar-Partida G, Lundberg M, Kho PF, D'Urso S, Gutiérrez-Mondragón LF, Ngo TT, Hwang L-D (2019) Complex-Traits Genetics Virtual Lab: A community-driven web platform for post-GWAS analyses bioRxiv

Day F et al. (2018) Large-scale genome-wide meta-analysis of polycystic ovary syndrome suggests shared genetic architecture for different diagnosis criteria PLoS Genet 14:e1007813 doi:10.1371/journal.pgen.1007813

De La Cruz MS, Buchanan EM (2017) Uterine Fibroids: Diagnosis and Treatment Am Fam Physician 95:100-107

Fearnley EJ, Marquart L, Spurdle AB, Weinstein P, Webb PM, Australian Ovarian Cancer Study G, Australian National Endometrial Cancer Study G (2010) Polycystic ovary syndrome increases the risk of endometrial cancer in women aged less than 50 years: an Australian case-control study Cancer causes & control : CCC 21:2303-2308 doi:10.1007/s10552-010-9658-7

Fortuny J et al. (2009) Risk of endometrial cancer in relation to medical conditions and medication use Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology 18:1448-1456 doi:10.1158/1055-9965.EPI-08-0936

Gallagher CS et al. (2019) Genome-wide association and epidemiological analyses reveal common genetic origins between uterine leiomyomata and endometriosis Nat Commun 10:4857 doi:10.1038/s41467-019-12536-4

Harris HR, Terry KL (2016) Polycystic ovary syndrome and risk of endometrial, ovarian, and breast cancer: a systematic review Fertil Res Pract 2:14 doi:10.1186/s40738-016-0029-2

Hartwig FP, Davey Smith G, Bowden J (2017) Robust inference in summary data Mendelian randomization via the zero modal pleiotropy assumption Int J Epidemiol 46:1985-1998 doi:10.1093/ije/dyx102

Hemani G et al. (2018) The MR-Base platform supports systematic causal inference across the human phenome Elife 7 doi:10.7554/eLife.34408

Ilaria S, Marci R (2018) From obesity to uterine fibroids: an intricate network Curr Med Res Opin 34:1877-1879 doi:10.1080/03007995.2018.1505606

Johnatty SE et al. (2020) Co-existence of leiomyomas, adenomyosis and endometriosis in women with endometrial cancer Scientific reports 10:3621 doi:10.1038/s41598-020-59916-1

Kar SP et al. (2020) Combining genome-wide studies of breast, prostate, ovarian and endometrial cancers maps cross-cancer susceptibility loci and identifies new genetic associations bioRxiv

Li J et al. (2019) Impact of endometriosis on risk of ovarian, endometrial and cervical cancers: a meta-analysis Arch Gynecol Obstet 299:35-46 doi:10.1007/s00404-018-4968-1

Masuda T et al. (2020) GWAS of five gynecologic diseases and cross-trait analysis in Japanese Eur J Hum Genet 28:95-107 doi:10.1038/s41431-019-0495-1

Matalliotaki C et al. (2018) Co-existence of benign gynecological tumors with endometriosis in a group of 1,000 women Oncology letters 15:1529-1532 doi:10.3892/ol.2017.7449

Melin A, Sparen P, Bergqvist A (2007) The risk of cancer and the role of parity among women with endometriosis Hum Reprod 22:3021-3026 doi:10.1093/humrep/dem209

Nagai K et al. (2015) Disease history and risk of comorbidity in women's life course: a comprehensive analysis of the Japan Nurses' Health Study baseline survey BMJ open 5:e006360 doi:10.1136/bmjopen-2014-006360

O'Connor LJ, Price AL (2018) Distinguishing genetic correlation from causation across 52 diseases and complex traits Nat Genet 50:1728-1734 doi:10.1038/s41588-018-0255-0

O'Mara TA et al. (2018) Identification of nine new susceptibility loci for endometrial cancer Nat Commun 9:3166 doi:10.1038/s41467-018-05427-7

O'Mara TA, Spurdle AB, Glubb DM, Endometrial Cancer Association C (2019) Analysis of Promoter-Associated Chromatin Interactions Reveals Biologically Relevant Candidate Target Genes at Endometrial Cancer Risk Loci Cancers (Basel) 11 doi:10.3390/cancers11101440

Olson JE, Cerhan JR, Janney CA, Anderson KE, Vachon CM, Sellers TA (2002) Postmenopausal cancer risk after self-reported endometriosis diagnosis in the Iowa Women's Health Study Cancer 94:1612-1618

Painter JN et al. (2018) Genetic overlap between endometriosis and endometrial cancer: evidence from cross-disease genetic correlation and GWAS meta-analyses Cancer medicine doi:10.1002/cam4.1445

Parasar P, Ozcan P, Terry KL (2017) Endometriosis: Epidemiology, Diagnosis and Clinical Management Curr Obstet Gynecol Rep 6:34-41 doi:10.1007/s13669-017-0187-1

Rafnar T et al. (2018) Variants associating with uterine leiomyoma highlight genetic background shared by various cancers and hormone-related traits Nat Commun 9:3636 doi:10.1038/s41467-018-05428-6

Rahmioglu N et al. (2018) Large-scale genome-wide association meta-analysis of endometriosis reveals 13 novel loci and genetically-associated comorbidity with other pain conditions bioRxiv:406967 doi:10.1101/406967

Roberts SG (2005) Transcriptional regulation by WT1 in development Curr Opin Genet Dev 15:542-547 doi:10.1016/j.gde.2005.08.004

Rogalla P, Rohen C, Hennig Y, Deichert U, Bonk U, Bullerdiek J (1995) Telomere repeat fragment sizes do not limit the growth potential of uterine leiomyomas Biochem Biophys Res Commun 211:175-182 doi:10.1006/bbrc.1995.1793

Rowlands IJ, Nagle CM, Spurdle AB, Webb PM, Australian National Endometrial Cancer Study G, Australian Ovarian Cancer Study G (2011) Gynecological conditions and the risk of endometrial cancer Gynecologic oncology 123:537-541 doi:10.1016/j.ygyno.2011.08.022

Rubtsova MP, Vasilkova DP, Malyavko AN, Naraikina YV, Zvereva MI, Dontsova OA (2012) Telomere lengthening and other functions of telomerase Acta Naturae 4:44-61

Sam S (2007) Obesity and Polycystic Ovary Syndrome Obes Manag 3:69-73 doi:10.1089/obe.2007.0019

Sapkota Y et al. (2017) Meta-analysis identifies five novel loci associated with endometriosis highlighting key genes involved in hormone metabolism Nat Commun 8:15539 doi:10.1038/ncomms15539

Stewart EA, Cookson CL, Gandolfo RA, Schulze-Rath R (2017) Epidemiology of uterine fibroids: a systematic review BJOG 124:1501-1512 doi:10.1111/1471-0528.14640

Telomeres Mendelian Randomization C et al. (2017) Association Between Telomere Length and Risk of Cancer and Non-Neoplastic Diseases: A Mendelian Randomization Study JAMA Oncol 3:636-651 doi:10.1001/jamaoncol.2016.5945

Uimari O, Jarvela I, Ryynanen M (2011) Do symptomatic endometriosis and uterine fibroids appear together? Journal of human reproductive sciences 4:34-38 doi:10.4103/0974-1208.82358

Wise LA, Palmer JR, Stewart EA, Rosenberg L (2007) Polycystic ovary syndrome and risk of uterine leiomyomata Fertility and sterility 87:1108-1115 doi:10.1016/j.fertnstert.2006.11.012

Wise LA, Sponholtz TR, Rosenberg L, Adams-Campbell LL, Kuohung W, LaValley MP, Palmer JR (2016) History of uterine leiomyoma and risk of endometrial cancer in black women Cancer causes & control : CCC 27:545-552 doi:10.1007/s10552-016-0728-3

Yengo L et al. (2018) Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry Hum Mol Genet 27:3641-3649 doi:10.1093/hmg/ddy271

Zhu Z et al. (2018) Causal associations between risk factors and common diseases inferred from GWAS summary data Nat Commun 9:224 doi:10.1038/s41467-017-02317-2

Zucchetto A et al. (2009) Hormone-related factors and gynecological conditions in relation to endometrial cancer risk Eur J Cancer Prev 18:316-321

TABLES

Table 1. GWAS data analyzed in this study

Phenotype	N_{Cases}	$N_{Controls}$	Data available*	Reference
Endometrial cancer[a]	12,906	108,979	Published summary data	O'Mara et al. (2018)
Endometriosis	14,926	189,715	Published summary data	Sapkota et al. (2017)
Endometriosis	58,115	733,480	Only top variant statistics	Rahmioglu et al. (2018)
PCOS[b]	4,890	20,405	Published summary data	Day et al. (2018)
PCOS[c]	10,074	103,164	Only top variant statistics	Day et al. (2018)
Uterine fibroids[b]	20,406	223,918	Published summary data	Gallagher et al. (2019)
Uterine fibroids[c]	35,474	267,505	Only top variant statistics	Gallagher et al. (2019)

*Datasets with full summary statistics available were used in genetic correlation, LCV and gene-based analysis. Datasets with only top variant statistics available were used in Mendelian randomization analyses. [a]UK Biobank samples removed from dataset to minimize bias in Mendelian randomization analysis, resulting in 12,270 endometrial cancer cases and 46,126 controls. [b]23andMe samples removed from publicly available dataset due to data sharing issue. [c]23andMe samples included.

Table 2. Genetic correlation results for non-cancerous gynecological diseases and endometrial cancer obtained from LD score regression

Non-cancerous gynecological diseases	Covariate	r_G	SE	P
Endometriosis	-	-0.02	0.09	0.83
PCOS	-	**0.36**	**0.12**	**1.6×10⁻³**
PCOS	BMI	0.19	0.14	0.17
Uterine fibroids	-	**0.24**	**0.09**	**5.4×10⁻³**
Uterine fibroids	BMI	0.23	0.1	0.02

r_G: genetic correlation, SE: standard error, P: P value. Results passing **Bonferroni correction (P < 0.017) are in bold**

Table 3. Genetic causal inference results for non-cancerous gynecological diseases and endometrial cancer obtained from LCV and two-sample Mendelian randomization analyses

Trait 1	Trait 2	Mendelian randomization method	Beta	SE	P	GCP	SE	P
Endometriosis	Endometrial Cancer	IVW	0.09	0.09	0.34	0.06	0.54	1.00
		MR-Egger	0.44	0.30	0.15			
		MR-Egger (intercept)	-0.03	0.03	0.23			
		Weighted median	0.11	0.08	0.14			
		Weighted mode	0.13	0.11	0.26			
PCOS		IVW	-0.05	0.04	0.26	0.18	0.5	0.78
		MR-Egger	0.07	0.21	0.75			
		MR-Egger (intercept)	-0.02	0.03	0.57			
		Weighted median	-0.08	0.06	0.22			
		Weighted mode	-0.19	0.12	0.13			
Uterine fibroids		IVW	0.08	0.07	0.22	0.53	0.31	0.13
		MR-Egger	0.31	0.15	0.06			
		MR-Egger (intercept)	-0.02	0.01	0.12			
		Weighted median	0.06	0.07	0.37			
		Weighted mode	0.05	0.11	0.66			

Beta: effect size in natural log OR; SE: standard error; P: P-value; GCP: genetic causality proportion

Table 4. Shared candidate risk genes at shared endometrial cancer and endometriosis risk regions

Gene	Region	TopSNP$_{EC}$ (p-value)	TopSNP$_{Endom}$ (p-value)	LD between TopSNPs (r^2)*	fastBAT$_{EC}$ p-value	fastBAT$_{EC}$ FDR	fastBAT$_{Endom}$ p-value	fastBAT$_{Endom}$ FDR
RUVBL1	3q21.3	rs872267 (9.95E-07)	rs4857864 (5.35E-06)	0.47	8.49E-06	8.01E-03	8.26E-05	4.25E-02
CDKN2B-AS1	9p21.3	rs568447 (2.68E-06)	rs6475610 (1.73E-09)	1.0E-04	7.26E-05	3.56E-02	9.02E-09	3.55E-05
BMF	15q15.1	rs28371998 (6.81E-09)	rs7183386 (7.23E-06)	0.36	1.64E-07	5.74E-04	4.93E-06	9.35E-03
CBX1	17q21.32	rs7225865 (1.34E-08)	rs10445377 (3.20E-07)	0.99	3.35E-06	4.32E-03	2.63E-05	2.48E-02
MIR1203	17q21.32	rs4794505 (9.50E-09)	rs10445377 (3.20E-07)	0.99	5.88E-07	1.44E-03	4.93E-06	9.35E-03
SKAP1	17q21.32	rs882380 (4.66E-09)	rs10445377 (3.20E-07)	0.96	5.80E-06	5.93E-03	7.93E-06	1.25E-02
SNX11	17q21.32	rs17681336 (1.17E-08)	rs10445377 (3.20E-07)	0.99	1.55E-06	3.17E-03	4.51E-06	9.35E-03

TopSNP: lead variant for gene from fastBAT analysis; EC: endometrial cancer; Endom: endometriosis; LD: linkage disequilibrium; FDR: false discovery rate.

*LD was estimated using the EUR 1000Genomes reference panel.

Table 5. Shared candidate risk genes at endometrial cancer and uterine fibroids risk regions

Gene	Region	TopSNP$_{EC}$ (p-value)	TopSNP$_{UF}$ (p-value)	LD between TopSNPs (r^2)*	fastBAT$_{EC}$ p-value	fastBAT$_{EC}$ FDR	fastBAT$_{UF}$ p-value	fastBAT$_{UF}$ FDR
CLPTM1L	5p15.33	rs2736100 (5.17E-06)	rs72709458 (6.07E-15)	0.28	3.93E-07	1.20E-03	1.00E-09	1.53E-06
MIR4457	5p15.33	rs2736100 (5.17E-06)	rs72709458 (6.07E-15)	0.28	1.50E-06	3.17E-03	8.01E-11	1.63E-07
TERT	5p15.33	rs2736100 (5.17E-06)	rs72709458 (6.07E-15)	0.28	1.72E-06	3.25E-03	1.50E-11	4.06E-08
WT1	11p13	rs10835920 (1.33E-08)	rs11031731 (2.04E-21)	0.20	1.86E-05	1.52E-02	2.64E-16	3.22E-12
WT1-AS	11p13	rs10835920 (1.33E-08)	rs11031762 (9.95E-14)	0.40	1.91E-06	3.34E-03	1.04E-12	4.24E-09

TopSNP: lead variant for gene from fastBAT analysis; EC: endometrial cancer; UF: uterine fibroids; LD: linkage disequilibrium; FDR: false discovery rate.

*LD was estimated using the EUR 1000Genomes reference panel.

Figure 1. Association between genetic predisposition to non-cancerous gynecological diseases and endometrial cancer risk, obtained from two-sample Mendelian randomization analysis. The boxes represent odds ratio for endometrial cancer risk per standard deviation increment in genetically predisposition to non-cancerous gynecological disease. Error bars represent 95% confidence intervals.