

Original article

Non-occupational and occupational factors associated with specific SARS-CoV-2 antibodies among Hospital Workers – a multicentre cross-sectional study

Christian R. Kahlert^{1,2} MD[†], Raphael Persi^{1*} MMed[†], Sabine Güsewell³ PhD, Thomas Egger¹ MSc, Onicio B. Leal-Neto^{4,5} PhD, Johannes Sumer¹ MD, Domenica Flury¹ MD, Angela Brucher⁶ MD, Eva Lemmenmeier⁷ MD, J. Carsten Möller⁸ MD, Philip Rieder⁹ PhD, Reto Stocker⁹ MD, Danielle Vuichard-Gysin^{10,11} MD, Benedikt Wiggli¹² MD, Werner C. Albrich¹ MD, Baharak Babouee Flury¹ MD, Ulrike Besold¹³ MD, Jan Fehr¹⁴ MD, Stefan P. Kuster MD^{15,16}, Allison McGeer MD¹⁷, Lorenz Risch^{18,19,20} PhD, Matthias Schlegel¹ MD, Andrée Friedl¹² MD, Pietro Vernazza¹ MD[†], Philipp Kohler¹ MD[†]

[†]Contributed equally

Author affiliations:

¹ Cantonal Hospital St Gallen, Division of Infectious Diseases and Hospital Epidemiology, St Gallen, Switzerland

² Children's Hospital of Eastern Switzerland, Department of Infectious Diseases and Hospital Epidemiology, St. Gallen, Switzerland

³ Clinical Trials Unit, Cantonal Hospital of St. Gallen, St. Gallen, Switzerland

⁴ EpiTrack, Recife, Brazil

⁵ Department of Economics, University of Zurich, Zurich, Switzerland

⁶ Psychiatry Services of the Canton of St. Gallen (South), Switzerland

⁷ Clenia Littenheid AG, Private Clinic for Psychiatry and Psychotherapy, Littenheid,
Switzerland

⁸ Center for Neurological Rehabilitation, Zihlschlacht, Switzerland

⁹ Hirslanden Clinic, Zurich, Switzerland

¹⁰ Thurgau Hospital Group, Division of Infectious Diseases and Hospital Epidemiology,
Muensterlingen, Switzerland

¹¹ Swiss National Center for Infection Prevention (Swissnoso), Berne, Switzerland

¹² Kantonsspital Baden, Division of Infectious Diseases and Hospital Epidemiology, Baden,
Switzerland

¹³ Geriatric Clinic St. Gallen, St. Gallen, Switzerland

¹⁴ Department of Public and Global Health, University of Zurich, Zurich, Switzerland

¹⁵ Federal Office of Public Health, Bern, Switzerland

¹⁶ University Hospital and University of Zurich, Division of Infectious Diseases and Hospital
Epidemiology, Zurich, Switzerland

¹⁷ Sinai Health System, Toronto, Canada

¹⁸ Labormedizinisches Zentrum Dr Risch Ostschweiz AG, Buchs, Switzerland

¹⁹ Private Universität im Fürstentum Liechtenstein, Triesen, Liechtenstein

²⁰ Center of Laboratory Medicine, University Institute of Clinical Chemistry, University of Bern,
Inselspital, Bern, Switzerland

Keywords: COVID-19; Seroprevalence; Healthcare workers; Switzerland; Risk factors

Corresponding author:

Philipp Kohler, MD MSc

Division of Infectious Diseases and Hospital Epidemiology

Cantonal Hospital St. Gallen

It is made available under a [CC-BY-NC-ND 4.0 International license](#) .

Rorschacherstrasse 95, 9011 St. Gallen, Switzerland

Phone +41 71 494 11 48, fax +41 71 494 63 09

Email philipp.kohler@kssg.ch

Abstract : 233 words

Text: 2498 words

ABSTRACT

Objectives

Protecting healthcare workers (HCW) from Coronavirus Disease-19 (COVID-19) is critical to preserve the functioning of healthcare systems. We therefore assessed seroprevalence and identified risk factors for Severe Acute Respiratory Syndrome-Coronavirus-2 (SARS-CoV-2) seropositivity in this population.

Methods

Between June 22nd and August 15th 2020, HCW from institutions in Northern/Eastern Switzerland were screened for SARS-CoV-2 antibodies. We recorded baseline characteristics, non-occupational and occupational risk factors. We used pairwise tests of associations and multivariable logistic regression to identify factors associated with seropositivity.

Results

Among 4'664 HCW from 23 healthcare facilities, 139 (3%) were seropositive. Non-occupational exposures independently associated with seropositivity were contact with a COVID-19-positive household (adjusted OR=54, 95%-CI: 31-97) and stay in a COVID-19 hotspot (aOR=2.2, 95%-CI: 1.1-3.9). Blood group 0 vs. non-0 (aOR=0.4, 95%-CI: 0.3-0.7), active smoking (aOR=0.5, 95%-CI: 0.3-0.9) and living with children <12 years (aOR=0.3, 95%-CI: 0.2-0.6) were associated with decreased risk. Occupational risk factors were close contact to COVID-19 patients (aOR=2.8, 95%-CI: 1.5-5.5), exposure to COVID-19-positive co-workers (aOR=2.0, 95%-CI: 1.2-3.1), poor knowledge of standard hygiene precautions (aOR=2.0, 95%-CI: 1.3-3.2), and frequent visits to the hospital canteen (aOR=1.9, 95%-CI: 1.2-3.1).

Conclusions

Living with COVID-19-positive households showed by far the strongest association with SARS-CoV-2 seropositivity. We identified several potentially modifiable risk factors, which might allow mitigation of the COVID-19 risk among HCW. The lower risk among those living with children, even after correction for multiple confounders, is remarkable and merits further study.

TEXT

INTRODUCTION

Coronavirus disease 2019 (COVID-19) is currently afflicting healthcare systems around the globe. As of December 2nd 2020, over 1.5 million COVID-19 deaths have been reported worldwide [1]. In Switzerland, over 350'000 COVID-19 cases have been reported, more than 14'000 patients have been hospitalized and over 5'000 have died [2]. Seroprevalence studies among Swiss healthcare workers (HCW) performed in March and April 2020 have shown a low prevalence of 1% in the Eastern part of the country, and a higher prevalence of around 10% in the Western part [3,4]. The recent massive re-emergence of cases in many European countries including Switzerland is putting further strain on healthcare systems and hospital workers.

Studies from different countries suggest that HCW are at increased risk to acquire Severe Acute Respiratory Syndrome-Coronavirus-2 (SARS-CoV-2) when compared to the general population [5]. In the UK, HCW and their household contacts accounted for a sixth of all COVID-19 cases admitted to the hospital for those aged 18-65 years. This risk was increased for HCW involved in patient care [6]. In light of these data it is imperative to better understand risk factors for SARS-CoV-2 acquisition among HCW in order to better protect them from infection.

In this multicentre study from Switzerland, we aimed to assess the prevalence of specific antibodies against SARS-CoV-2 among HCW with and without patient contact. In addition, we identified non-occupational and occupational factors associated with seropositivity to inform prevention recommendations for this population.

METHODS

Study design and participants

We initiated a multicentre cross-sectional study between June 22nd and August 15th 2020 in healthcare institutions located in Northern and Eastern Switzerland. Acute care hospitals, rehabilitation clinics, geriatric and psychiatric clinics in the region were asked to participate.

Within every participating institution, employees aged 16 years or older were invited to enrol into the study via institutional webpages. Employees registered online and provided electronic consent. The study was approved by the ethics committee of Eastern Switzerland (#2020-00502).

Questionnaire and definitions

We implemented a multi-modular digital web-based questionnaire for institutions and participants. Questions about facility structure were asked in the institutional questionnaire. Participants received an invitation to the questionnaire by email and were asked about anthropometric data, occupational and non-occupational risk exposures, and previous SARS-CoV-2 nasopharyngeal swabs. Household contacts were defined as people living in the same household or intimate partners; close contact to COVID-19 patients was assumed for those with contact >15 minutes within 2 meters with or without personal protective measures (PPE); aerosol-generating procedures (AGP) were defined according to guidelines of the Swiss Center for Infection Prevention (Swissnoso). Poor knowledge of standard precautions was assumed for those who correctly identified less than 3 measures in a multiple choice question (among a choice of hand hygiene, surgical mask in case of respiratory symptoms, donning gowns in case of potential contamination with body fluids, cough etiquette, and vaccination). Low protection while caring for COVID-19 patients was assumed for those using less than 3 measures out of face masks, gloves, gowns, and goggles.

Sample processing

Upon registration, participants provided a venous blood sample, which was collected at local sites. Samples were analysed with an electro-chemiluminescence immunoassay (ECLIA, Roche Diagnostics, Rotkreuz, Switzerland, detection of total antibodies directed against the nucleocapsid-(N)-protein of SARS-CoV-2) run on a COBAS 6000 instrument, as described elsewhere [7]. A subgroup of samples with a positive signal in the ECLIA (at a cut-off index,

COI, ≥ 1) were also tested with an Enzyme-linked Immunosorbent Assay (ELISA, Euroimmune, Germany, detection each of IgG and IgA antibodies against S1 domain of the spike-(S)-protein including the immunologically relevant receptor binding domain). Cut-offs for seropositivity were applied as recommended by the manufacturers. Seropositivity was defined as positive result in the ECLIA followed by confirmation in the ELISA (either positive IgA or IgG).

Statistical analysis

The relative frequency of participants with ELISA confirmed positive and negative serology was compared between levels of baseline characteristics, non-occupational risk factors and occupational risk factors. Fisher's exact test was used for dichotomous factors or factors with a reference level, comparing each level to the reference. Individuals with missing data were removed from the analysis of the respective variable. Logistic regression was used for numeric and ordinal variables. Age, sex, body mass index (BMI), smoking status, comorbidities as well as non-occupational and occupational risk factors expected to influence seropositivity were entered into a multivariable logistic regression model. For sensitivity analysis, we fitted two additional models including place of residence (7 predefined regions) and institution either as fixed effects or as random effects to assess whether spatial proximity or clustering of observations confounded the effects of the risk factors. Analyses were performed with R statistical software, version 4.0.2.

RESULTS

Baseline characteristics

We included 17 institutions on 23 sites across Northern and Eastern Switzerland, thereof 19 inpatient sites (14 acute care; 1 geriatric clinic; 1 rehabilitation clinic; 3 psychiatric clinics) and 4 outpatient clinics (3 psychiatric facilities; 1 blood donation centre). The total of represented patient beds was 3'523 (thereof 106 ICU beds) (**Table 1**).

Of the 17'060 potentially eligible HCW, 4'664 (27%) participated in the study. Median age was 38 years (range 16-73); 3'654 (78%) were female. The majority were nurses (n=2'126; 46%) followed by physicians (n=776; 17%); 3'676 (79%) reported having patient contact (**Table 2**).

Seropositivity and self-reported PCR results

Overall, seropositivity was 3.0% (139/4'664). Among these 139, 88 (63%) were also tested with the confirmatory ELISA and all 88 samples had either positive IgA or IgG. On the institutional level, seropositivity ranged between 0.5% and 4.2% for inpatient, and 0% and 2.3% for outpatient facilities (**Table 1**). Seropositivity by district ranged from 0% to 13%. Seropositivity was lower in regions located in Eastern compared to Northern Switzerland (**Figure 1**).

A previous PCR result was reported by 864 of 4'664 (18.5%) participants. Of the 72 participants with positive PCR, 66 (92%) were also seropositive. On the other hand, 17/792 (2.2%) participants with negative PCR had a positive serology. Overall, 23/864 (2.7%) self-reported PCR results were discordant to serology results. Seroprevalence among those without previous PCR was 1.5% (56/3'800).

Non-occupational factors associated with seropositivity

Exposure to COVID-19 confirmed (55.7% vs. 2.1%, $p<0.001$) or symptomatic, not confirmed household contacts (5.5% vs. 1.9%, $p<0.001$) was strongly associated with seropositivity. Also, having visited a known COVID-19 hotspot in Austria (but not in Italy or France) was clearly associated with seropositivity (6.8% vs. 2.8%, $p=0.002$). Seroprevalence was lower among those with blood group 0 vs. non-0 (1.8% vs. 3.5%, $p=0.002$) and for those living with children aged 12 or younger (1.7% vs. 3.4%, $p=0.002$) (**Table 2**).

Occupational factors associated with seropositivity

Nurses had a higher (3.9%), physicians a lower (1.0%) seropositivity rate; no differences according to medical speciality were noted. Seroprevalence was higher among those with patient contact (3.1% vs. 1.7%, $p=0.037$), particularly for those with contact to confirmed COVID-19 patients (4.1% vs. 1.7%, $p<0.001$). Workers indicating low protection while caring for COVID-19 patients (5.8% vs. 3.5%, $p=0.019$) and those with poor knowledge of hygiene standards had higher seropositivity (4.1% vs. 2.6%, $p=0.018$) (**Figure 2, panels A/B**). The number of unprotected contacts to COVID-19 confirmed or symptomatic co-workers was associated with seropositivity (**Figure 2, panel C**). Also, workers who never/occasionally visited the hospital canteen had a lower seroprevalence compared to those with weekly/daily visits (1.9% vs. 3.5%, $p=0.004$) (**Figure 2, panel D**). This effect was consistent across institutions and professions (**Table S1**).

Multivariable analyses

In multivariable analysis, exposure to a COVID-19 positive household member remained the strongest risk factor for seropositivity with an adjusted odds ratio (aOR) of 54 (95% CI 31-97) (**Figure 3, Table S2**). Stay in a COVID-19 hotspot was associated with increased risk (aOR 2.2, 95% CI 1.1-3.9), whereas blood group 0 (aOR 0.4, 95% CI 0.3-0.7), active smoking (aOR 0.5, 95% CI 0.3-0.9) and living with children <12 years (aOR 0.3, 95% CI 0.2-0.6) were all associated with decreased risk after correcting for multiple confounder variables.

Significant occupational factors included close contact with a COVID-19 patient (aOR 2.8, 95% CI 1.5-5.5), exposure to a COVID-19 positive co-worker (aOR 2.0, 95% CI 1.2-3.1), poor knowledge of standard precautions (aOR 2.0, 95% CI 1.3-3.2), as well as having weekly/daily (vs. rarely/never) meals in the hospital canteen (aOR 1.9, 95% CI 1.2-3.1). Both models in the sensitivity analysis did not show any relevant impact of geographic region or institution on the significance level of the variables in the original model (**Table S2**).

DISCUSSION

In this cross-sectional study of a sample of 4'664 Swiss HCW, 3% of participants had specific SARS-CoV-2 antibodies. Our main findings are that exposure to a COVID-19 positive household member is the strongest risk factor for seropositivity, and that living with children under the age of 12, even after correction for multiple confounders, is clearly associated with decreased risk. Furthermore, we identified several exposures associated with seropositivity which might serve as leverage to further decrease the risk of SARS-CoV-2 acquisition among HCW.

We confirm findings from other studies showing that COVID-19 positive household contacts are the main source of SARS-CoV-2 infection for HCW [8,9]. Our findings are also in line with a Dutch study which concluded that nosocomial transmissions seemed rather uncommon and that multiple hospital introductions from the community are probably responsible for the large part of COVID-19 cases among patients and HCW, at least in a low prevalence setting [10]. Of course, this association might be overestimated given that the directionality of virus transmission cannot be definitely assessed with our study design.

An important finding of our study is that participants living with children <12 years were less likely to be seropositive. A study among over 300'000 HCW households from Scotland has recently found a similar association [11]. In contrast to the Scottish study, we corrected our result for important confounders, including HCW age, full-time working, and leisure activities.

Similarly, a population-based cohort from England using large public health databases with over 9 million adults showed that living with children <12 years was not associated with increased risk for SARS-CoV-2 infection, as opposed to living with children aged 12-18 [12]. An intriguing hypothesis for this finding is that certain childhood infections, particularly those with endemic coronaviruses such as HCoV-OC43 or HCoV-HKU1, might confer partial immunity (i.e. cross-immunity) to SARS-CoV-2. In line with this hypothesis, adults aged 15 to 44 years (having presumably an increased probability of living with young children) have been shown to

have higher antibody titers against the HCoV-OC43 N protein than older adults [13]. Also, supporting the notion of a rather immunological than a purely epidemiological phenomenon, a German study among over 4'000 COVID-19 patients suggested a less complicated disease course for those with frequent contact to children [14]. This hypothesis was partially confirmed by a recent study that demonstrated pre-existing humoral immunity (including neutralizing antibodies) to be particularly prevalent in children and adolescents [15]. This should be confirmed in prospective studies which evaluate the protective role of humoral and cellular immunity against endemic coronaviruses regarding SARS-CoV-2 acquisition.

Interestingly, stay in an Austrian ski resort where at least one COVID-19 superspreading event had occurred in February/March 2020 was an independent risk factor for seropositivity [16].

Several studies have by now identified an association between the AB0 blood group system and acquisition of COVID-19. Consistently, blood group 0 is considered to have a protective effect as shown in our study, whereas people with a non-0 blood group (mostly A) seem to carry an increased risk [17]. Whether the blood group also determines the course of the disease is less clear [18]. We also observed a lower seroprevalence among active smokers, confirming findings of a meta-analysis [19].

An important question is whether HCW caring for COVID-19 patients are in fact at increased risk for acquiring the disease themselves. A recent meta-analysis concluded that HCW do indeed have an increased risk compared to the general population [20]. Also, frontline HCW in Denmark showed higher seroprevalences than other HCW [21]. Our study confirms these findings, at least for those with close contact to COVID-19 patients. As opposed to other studies [20], a lower level of protection was not significantly associated with seropositivity in multivariable analysis, probably because of our restrictive definition of low protection. Due to the cross-sectional study design we cannot draw valid conclusions regarding the individual benefit of single protective measures such as gloves, gowns or goggles. However, participants performing AGPs as well as those working in intensive care or emergency rooms did not have an

increased risk for COVID-19, suggesting that current safety measures are sufficient for these high-risk HCW. Of note, poor knowledge of standard hygiene precautions was associated with detection of SARS-CoV-2 antibodies, supporting efforts to continuously educate HCW regarding basic infection prevention concepts.

Exposure to ill co-workers is a known risk factor for respiratory illness in HCW, not only for COVID-19 but also for other respiratory viral diseases [22]. Across all participating institutions, we identified visits to the hospital canteen as potential risk factor for seropositivity. We found one other study which reported staying in the same HCW break room and eating in proximity to other HCW as risk factor for SARS-CoV-2 transmission [23]. Visiting restaurants other than hospital canteens has previously been shown to be potentially associated with higher risk of SARS-CoV-2 acquisition [24–26]; however, this was not the case in our data. This discrepancy could be explained by the fact that i) the visitor turnover of hospital canteens is much higher than in other eating places and ii) that the probability of a HCW being infectious is higher than for an average visitor to other restaurants. We therefore suggest that hospitals should revisit and potentially reinforce the safety concepts of their canteens and food courts.

Our study has several limitations. First, causality cannot be inferred between exposures and seropositivity. Second, sampling bias may have arisen given that study participation was non-mandatory. Third, we relied on mostly self-reported data in our questionnaire which is subject to recall and other bias. Fourth, because of the low disease prevalence we have to assume that a certain proportion of our serology results are false positive. However, the reported specificity of >99% for the ECLIA [27], the positive confirmatory results in our tested subsample and the overall low proportion of discordant results between PCR and serology supports the validity of our testing approach. Strengths of the study are its large sample size, the inclusion of different types of healthcare institutions across a large geographic area, and consideration of not only occupational but a broad range of non-occupational risk factors. In particular the latter differentiates our study from most other seroprevalence studies performed among HCW.

To conclude, having a COVID-19 positive household member was by far the strongest predictor for SARS-CoV-2 seropositivity among our HCW. Furthermore, we identified several modifiable variables associated with seropositivity, including contact to COVID-19 co-workers, poor knowledge of standard hygiene precautions, and possibly frequent visits to the hospital canteen. Living with children below 12 years of age in the same household was independently associated with decreased risk, an extraordinary finding suggesting an increased role of cross-immunity.

Conflict of interest

None of the co-authors reports any conflict of interest.

Funding

This work was supported by the Swiss National Sciences Foundation (grant number 31CA30_196544; grant number PZ00P3_179919 to PK), the Federal Office of Public Health (grant number 20.008218/421-28/1), the Health Department of the Canton of St. Gallen, and the research fund of the Cantonal Hospital of St. Gallen.

Acknowledgments

We would like to warmly thank the large number of employees of the participating health care institutions who either took part in this study themselves or supported it. Furthermore, we thank the laboratory staff for shipment, handling and analysis of the blood samples. In particular, we acknowledge the organizational core team Simone Kessler and Susanne Nigg, who kept all strings between the participating centers and the laboratory and without whom this study would not have been possible.

Authors contributions

All authors contributed to the conceptualization of the study. CRK and PK supervised the study. CRK, OL, TE, PV and PK were responsible for data curation. TE was responsible for project administration. RP, JS, DF, AB, EL, CM, PR, RS, DV, BW, UB, LR and AF contributed to the investigations. LR provided laboratory resources. SG was responsible for the formal analysis and data visualizations. CRK, PV, and PK were responsible for funding acquisition. CRK, RP and PK wrote the original draft, which was critically reviewed and edited by all authors.

REFERENCES

- [1] COVID-19 Map. In: Johns Hopkins Coronavirus Resource Center. Available at: <https://coronavirus.jhu.edu/map.html>. Last accessed December 8th 2020. n.d.
- [2] Swiss Federal Office of Public Health - Current Situation in Switzerland -Daily Report. Available at: <https://www.covid19.admin.ch/en/overview>. Last accessed December 8th 2020 n.d.
- [3] Kohler PP, Kahlert CR, Sumer J, Flury D, Gusewell S, Leal-Neto OB, et al. Prevalence of SARS-CoV-2 antibodies among Swiss hospital workers: Results of a prospective cohort study. *Infect Control Hosp Epidemiol* 2020;1–5. <https://doi.org/10.1017/ice.2020.1244>.
- [4] Stringhini S, Wisniak A, Piumatti G, Azman AS, Lauer SA, Baysson H, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-POP): a population-based study. *The Lancet* 2020;396:313–9. [https://doi.org/10.1016/S0140-6736\(20\)31304-0](https://doi.org/10.1016/S0140-6736(20)31304-0).
- [5] Rudberg A-S, Havervall S, Manberg A, Jernbom Falk A, Aguilera K, Ng H, et al. SARS-CoV-2 exposure, symptoms and seroprevalence in healthcare workers in Sweden. *Nat Commun* 2020;11:5064. <https://doi.org/10.1038/s41467-020-18848-0>.
- [6] Shah ASV, Wood R, Gribben C, Caldwell D, Bishop J, Weir A, et al. Risk of hospital admission with coronavirus disease 2019 in healthcare workers and their households: nationwide linkage cohort study. *BMJ* 2020:m3582. <https://doi.org/10.1136/bmj.m3582>.
- [7] Baron RC, Risch L, Weber M, Thiel S, Grossmann K, Wohlwend N, et al. Frequency of serological non-responders and false-negative RT-PCR results in SARS-CoV-2 testing: a population-based study. *Clin Chem Lab Med* 2020. <https://doi.org/10.1515/cclm-2020-0978>.
- [8] Steensels D, Oris E, Coninx L, Nuyens D, Delforge M-L, Vermeersch P, et al. Hospital-Wide SARS-CoV-2 Antibody Screening in 3056 Staff in a Tertiary Center in Belgium. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.11160>.

- [9] Koh WC, Naing L, Chaw L, Rosledzana MA, Alikhan MF, Jamaludin SA, et al. What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. *PLOS ONE* 2020;15:e0240205. <https://doi.org/10.1371/journal.pone.0240205>.
- [10] Sikkema RS, Pas SD, Nieuwenhuijse DF, O’Toole Á, Verweij J, Linden A van der, et al. COVID-19 in health-care workers in three hospitals in the south of the Netherlands: a cross-sectional study. *Lancet Infect Dis* 2020;20:1273–80. [https://doi.org/10.1016/S1473-3099\(20\)30527-2](https://doi.org/10.1016/S1473-3099(20)30527-2).
- [11] Wood R, Thomson EC, Galbraith R, Gribben C, Caldwell D, Bishop J, et al. Sharing a household with children and risk of COVID-19: a study of over 300,000 adults living in healthcare worker households in Scotland. *MedRxiv* 2020:2020.09.21.20196428. <https://doi.org/10.1101/2020.09.21.20196428>.
- [12] Forbes H, Morton CE, Bacon S, McDonald HI, Minassian C, Brown JP, et al. Association between living with children and outcomes from COVID-19: an OpenSAFELY cohort study of 12 million adults in England. *MedRxiv* 2020:2020.11.01.20222315. <https://doi.org/10.1101/2020.11.01.20222315>.
- [13] Gao X, Zhou H, Wu C, Xiao Y, Ren L, Paranhos-Baccalà G, et al. Antibody against nucleocapsid protein predicts susceptibility to human coronavirus infection. *J Infect* 2015;71:599–602. <https://doi.org/10.1016/j.jinf.2015.07.002>.
- [14] Dugas M, Schrempf I-M, Ochs K, Frömmel C, Greulich L, Neuhaus P, et al. Association of contact to small children with a mild course of COVID-19. *Int J Infect Dis* 2020;100:314–5. <https://doi.org/10.1016/j.ijid.2020.09.003>.
- [15] Ng KW, Faulkner N, Cornish GH, Rosa A, Harvey R, Hussain S, et al. Preexisting and de novo humoral immunity to SARS-CoV-2 in humans. *Science* 2020. <https://doi.org/10.1126/science.abe1107>.

- [16] Kreidl P, Schmid D, Maritschnik S, Richter L, Borena W, Genger J-W, et al. Emergence of coronavirus disease 2019 (COVID-19) in Austria. *Wien Klin Wochenschr* 2020. <https://doi.org/10.1007/s00508-020-01723-9>.
- [17] Genomewide Association Study of Severe Covid-19 with Respiratory Failure. *N Engl J Med* 2020;383:1522–34. <https://doi.org/10.1056/NEJMoa2020283>.
- [18] Sardu C, Marfella R, Maggi P, Messina V, Cirillo P, Codella V, et al. Implications of ABO blood group in hypertensive patients with covid-19. *BMC Cardiovasc Disord* 2020;20:373. <https://doi.org/10.1186/s12872-020-01658-z>.
- [19] Simons D, Shahab L, Brown J, Perski O. The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: A living rapid evidence review with Bayesian meta-analyses (version 7). *Addict Abingdon Engl* 2020. <https://doi.org/10.1111/add.15276>.
- [20] Galanis P, Vraka I, Fragkou D, Bilali A, Kaitelidou D. Seroprevalence of SARS-CoV-2 antibodies and associated factors in health care workers: a systematic review and meta-analysis. *Epidemiology*; 2020. <https://doi.org/10.1101/2020.10.23.20218289>.
- [21] Iversen K, Bundgaard H, Hasselbalch RB, Kristensen JH, Nielsen PB, Pries-Heje M, et al. Risk of COVID-19 in health-care workers in Denmark: an observational cohort study. *Lancet Infect Dis* 2020;0. [https://doi.org/10.1016/S1473-3099\(20\)30589-2](https://doi.org/10.1016/S1473-3099(20)30589-2).
- [22] Buckrell S, Coleman BL, McNeil SA, Katz K, Muller MP, Simor A, et al. Sources of viral respiratory infections in Canadian acute care hospital healthcare personnel. *J Hosp Infect* 2020;104:513–21. <https://doi.org/10.1016/j.jhin.2020.01.009>.
- [23] Çelebi G, Pişkin N, Bekleviç AÇ, Altunay Y, Keleş AS, Tüz MA, et al. Specific risk factors for SARS-CoV-2 transmission among health care workers in a university hospital. *Am J Infect Control* 2020;48:1225–30. <https://doi.org/10.1016/j.ajic.2020.07.039>.

- [24] Lu J, Gu J, Li K, Xu C, Su W, Lai Z, et al. COVID-19 Outbreak Associated with Air Conditioning in Restaurant, Guangzhou, China, 2020. *Emerg Infect Dis* 2020;26:1628–31. <https://doi.org/10.3201/eid2607.200764>.
- [25] Fisher KA, Tenforde MW, Feldstein LR, Lindsell CJ, Shapiro NI, Files DC, et al. Community and Close Contact Exposures Associated with COVID-19 Among Symptomatic Adults ≥ 18 Years in 11 Outpatient Health Care Facilities - United States, July 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1258–64. <https://doi.org/10.15585/mmwr.mm6936a5>.
- [26] Lentz RJ, Colt H, Chen H, Cordovilla R, Popevic S, Tahura S, et al. Assessing coronavirus disease 2019 (COVID-19) transmission to healthcare personnel: The global ACT-HCP case-control study. *Infect Control Hosp Epidemiol* undefined/ed:1–7. <https://doi.org/10.1017/ice.2020.455>.
- [27] Mahase E. Covid-19: Two antibody tests are “highly specific” but vary in sensitivity, evaluations find. *BMJ* 2020;369. <https://doi.org/10.1136/bmj.m2066>.

TABLES

Table 1. Characteristics of institutions (n=17) including size, number of study participants and seropositivity.

	Sites (n)	Inpatients (yes vs no)	Beds (n)	ICU beds (n)	HCW (n)	HCW in study (n)	HCW in study (%)	Seropositive HCW (n)	Seropositive HCW (%)
TOTAL	23	NA	3'523	106	17'060	4664	27%	139	3.0%
Acute care	3	yes	765	36	5930	1074	18%	37	3.4%
Acute care	1	yes	370	10	2245	1023	46%	39	3.8%
Acute care	3	yes	304	7	1367	534	39%	9	1.7%
Acute care	1	yes	74	0	362	109	30%	3	2.8%
Acute care	1	yes	46	0	178	66	37%	1	1.5%
Acute care	1	yes	246	9	749	169	23%	7	4.1%
Acute care	1	yes	310	12	740	171	23%	3	1.8%
Acute care	1	yes	330	18	1788	448	25%	18	4.0%
Acute care	1	yes	129	6	525	159	30%	4	2.5%
Acute care	1	yes	100	8	632	109	17%	3	2.8%
Geriatric acute care	1	yes	98	0	265	123	46%	3	2.4%
Rehabilitation clinic	1	yes	135	0	510	168	33%	7	4.2%
Psychiatric clinic	1	yes	242	0	360	190	53%	1	0.5%
Psychiatric clinic	1	yes	150	0	391	108	28%	1	0.9%
Psychiatric clinic	1	yes	224	0	780	98	13%	1	1.0%
Psychiatry	3	no	NA	NA	178	88	49%	2	2.3%
Blood donation	1	no	NA	NA	60	27	45%	0	0.0%

Abbreviations: ICU, Intensive Care Unit; HCW, Healthcare Workers

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Table 2. Baseline, non-occupational and occupational factors by serostatus.

	Total n	Seropositive (n and %)	Seronegative (n and %)	OR with 95% CI	p-value
Gender					
Female	3654	105 (2.9%)	3549 (97.1%)	ref	-
Male	983	34 (3.5%)	949 (96.5%)	1.21 (0.79 - 1.81)	0.343
Age, median (IQR), OR per 10 years	38.3 (29.7-49.5)	35.5 (26.8-46.8)	38.4 (29.7-49.6)	0.83 (0.71 - 0.96)	0.012
BMI, median (IQR), OR per unit	23.4 (21.3-26.2)	24.2 (22.2-27.1)	23.4 (21.3-26.1)	1.03 (1.00 - 1.07)	0.078
Smoking status					
Never	2891	96 (3.3%)	2795 (96.7%)	ref	-
Active	951	27 (2.8%)	924 (97.2%)	0.58 (0.32 - 0.99)	0.049
Former	822	16 (1.9%)	806 (98.1%)	0.85 (0.53 - 1.33)	0.525
Comorbidity					
No	3021	80 (2.6%)	2941 (97.4%)	ref	-
Yes	1643	59 (3.6%)	1584 (96.4%)	1.37 (0.96 - 1.95)	0.072
Blood group (OR: one group vs all others)					
A	1396	51 (3.7%)	1345 (96.3%)	1.37 (0.95 - 1.97)	0.090
AB	161	6 (3.7%)	155 (96.3%)	1.27 (0.45 - 2.91)	0.482
B	354	14 (4.0%)	340 (96.0%)	1.38 (0.72 - 2.43)	0.254
O	1383	25 (1.8%)	1358 (98.2%)	0.51 (0.32 - 0.80)	0.002
I don't know	1305	41 (3.1%)	1264 (96.9%)	1.08 (0.73 - 1.58)	0.701
Influenza vaccine 2019/2020					
No	3159	102 (3.2%)	3057 (96.8%)	ref	-
Yes	1416	35 (2.5%)	1381 (97.5%)	0.76 (0.50 - 1.13)	0.189
BCG vaccine					
No	1586	55 (3.5%)	1531 (96.5%)	ref	-
Yes	1908	49 (2.6%)	1859 (97.4%)	0.73 (0.49 - 1.11)	0.134
I don't know	1104	34 (3.1%)	1070 (96.9%)	0.88 (0.56 - 1.39)	0.661
No of respiratory tract infections/year					
0 or 1	3862	105 (2.7%)	3757 (97.3%)	ref	-
2 to 4	776	31 (4.0%)	745 (96.0%)	1.49 (0.96 - 2.26)	0.062
5+	26	3 (11.5%)	23 (88.5%)	4.66 (0.88 - 15.8)	0.034
No of persons in household					
1 (OR per person)	814	17 (2.1%)	797 (97.9%)	0.94 (0.82 - 1.08)	0.383
2	1660	64 (3.9%)	1596 (96.1%)		
3	778	22 (2.8%)	756 (97.2%)		
4	957	29 (3.0%)	928 (97.0%)		
5+	455	7 (1.5%)	448 (98.5%)		

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

No of children ≤12 years					
0 (OR per person)	3526	120 (3.4%)	3406 (96.6%)	0.70 (0.52 - 0.90)	0.010
1	492	6 (1.2%)	486 (98.8%)		
2	509	12 (2.4%)	497 (97.6%)		
3+	137	1 (0.7%)	136 (99.3%)		
Confirmed COVID-19 case in household					
No	4585	95 (2.1%)	4490 (97.9%)	ref	-
Yes	79	44 (55.7%)	35 (44.3%)	59.1 (35.4 - 99.9)	< 0.001
Symptomatic household contact					
No	3269	62 (1.9%)	3207 (98.1%)	ref	-
Yes	1395	77 (5.5%)	1318 (94.5%)	3.02 (2.12 - 4.32)	< 0.001
Visit to a COVID-19 hotspot ¹					
No	4413	122 (2.8%)	4291 (97.2%)	ref	-
Yes	251	17 (6.8%)	234 (93.2%)	2.55 (1.42 - 4.35)	0.002
Leisure activities (currently; OR for with vs without activity)					
Visit to restaurant/bar	2783	84 (3.0%)	2699 (97.0%)	1.03 (0.72 - 1.49)	0.930
Sport club	833	28 (3.4%)	805 (96.6%)	1.17 (0.74 - 1.79)	0.499
Fitness/yoga classes	1462	49 (3.4%)	1413 (96.6%)	1.20 (0.82 - 1.73)	0.309
Theater/concerts	112	4 (3.6%)	108 (96.4%)	1.21 (0.32 - 3.27)	0.577
Cinema	290	14 (4.8%)	276 (95.2%)	1.72 (0.90 - 3.05)	0.071
Religious gatherings	228	6 (2.6%)	222 (97.4%)	0.87 (0.31 - 1.99)	1.000
Singing in choir	59	2 (3.4%)	57 (96.6%)	1.14 (0.13 - 4.41)	0.695
Active group musician	110	4 (3.6%)	106 (96.4%)	1.24 (0.33 - 3.33)	0.570
No of leisure activities above					
0 (OR per activity)	1045	25 (2.4%)	1020 (97.6%)	1.13 (0.95 - 1.34)	0.169
1	1875	55 (2.9%)	1820 (97.1%)		
2	1320	46 (3.5%)	1274 (96.5%)		
3	342	9 (2.6%)	333 (97.4%)		
4+	82	4 (4.9%)	78 (95.1%)		
No of shopping trips per week (currently)					
0 (OR per trip)	34	2 (5.9%)	32 (94.1%)	1.03 (0.87 - 1.21)	0.753
1	1212	34 (2.8%)	1178 (97.2%)		
2	1631	46 (2.8%)	1585 (97.2%)		
3	963	33 (3.4%)	930 (96.6%)		
4+	650	19 (2.9%)	631 (97.1%)		
Profession (OR: one profession vs all					

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

others)					
Nurse	2257	88 (3.9%)	2169 (96.1%)	1.87 (1.31 - 2.71)	< 0.001
Physician	776	8 (1.0%)	768 (99.0%)	0.30 (0.13 - 0.61)	< 0.001
Administration/Secretary	472	8 (1.7%)	464 (98.3%)	0.53 (0.22 - 1.09)	0.087
Physiotherapist	181	7 (3.9%)	174 (96.1%)	1.33 (0.52 - 2.87)	0.498
Other	769	16 (2.1%)	753 (97.9%)	0.65 (0.36 - 1.11)	0.130
Speciality (OR: one speciality vs all others)					
Internal Medicine	995	31 (3.1%)	964 (96.9%)	1.06 (0.68 - 1.61)	0.753
Surgery/Orthopedics	475	14 (2.9%)	461 (97.1%)	0.99 (0.52 - 1.74)	1.000
Intensive care	289	5 (1.7%)	284 (98.3%)	0.56 (0.18 - 1.35)	0.280
Emergency department	272	9 (3.3%)	263 (96.7%)	1.12 (0.50 - 2.23)	0.712
Other	585	18 (3.1%)	567 (96.9%)	1.04 (0.59 - 1.73)	0.896
Employment rate					
> 80%	2690	90 (3.3%)	2600 (96.7%)	ref	-
≤ 80%	1974	49 (2.5%)	1925 (97.5%)	0.74 (0.51 - 1.06)	0.098
Patient contact					
No	719	12 (1.7%)	707 (98.3%)	ref	-
Yes	3676	115 (3.1%)	3561 (96.9%)	1.23 (0.85 - 1.77)	0.263
Involved in AGP					
No	3228	90 (2.8%)	3138 (97.2%)	ref	-
Yes	1436	49 (3.4%)	1387 (96.6%)	1.90 (1.04 - 3.81)	0.037
No of correct standard precaution measures					
0 to 2	1073	44 (4.1%)	1029 (95.9%)	ref	-
3 or 4	2229	55 (2.5%)	2174 (97.5%)	0.59 (0.39 - 0.91)	0.012
5	1362	40 (2.9%)	1322 (97.1%)	0.71 (0.45 - 1.12)	0.146
Adherence to standard precautions					
almost always	2829	76 (2.7%)	2753 (97.3%)	ref	-
if I remember	1227	37 (3.0%)	1190 (97.0%)	1.13 (0.73 - 1.70)	0.604
often not possible	320	10 (3.1%)	310 (96.9%)	1.17 (0.53 - 2.30)	0.589
poorly	43	2 (4.7%)	41 (95.3%)	1.77 (0.20 - 7.02)	0.327
no answer	245	14 (5.7%)	231 (94.3%)	2.19 (1.13 - 3.99)	0.015
Caring for COVID-19 patients					
No	2348	40 (1.7%)	2308 (98.3%)	ref	-
Yes	2062	85 (4.1%)	1977 (95.9%)	2.48 (1.68 - 3.73)	< 0.001
Physical contact with COVID-19 patient					
No (only distant contact)	732	16 (2.2%)	716 (97.8%)	ref	-

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .

Yes	1329	69 (5.2%)	1260 (94.8%)	2.45 (1.39 - 4.56)	0.001
Exposure to coughing or sneezing by COVID-19 patient					
No	1544	52 (3.4%)	1492 (96.6%)	ref	-
Yes	517	33 (6.4%)	484 (93.6%)	1.96 (1.21 - 3.12)	0.005
Protection during close contact (n = 1329); OR for with vs without each protection					
Any face mask	1275	59 (4.6%)	1216 (95.4%)	0.21 (0.10 - 0.50)	< 0.001
Gloves	1125	49 (4.4%)	1076 (95.6%)	0.42 (0.24 - 0.76)	0.003
Gown	979	41 (4.2%)	938 (95.8%)	0.50 (0.30 - 0.86)	0.008
Goggles	931	39 (4.2%)	892 (95.8%)	0.54 (0.32 - 0.91)	0.015
None	47	8 (17.0%)	39 (83.0%)	4.10 (1.58 - 9.40)	0.002
No of protection measures above					
0 (OR per measure)	44	8 (18.2%)	36 (81.8%)	0.73 (0.61 - 0.87)	< 0.001
1	147	12 (8.2%)	135 (91.8%)		
2	116	6 (5.2%)	110 (94.8%)		
3	157	8 (5.1%)	149 (94.9%)		
4	865	35 (4.0%)	830 (96.0%)		
Contacts with COVID-19 positive co-worker					
No answer / don't know	1212	31 (2.6%)	1181 (97.4%)	1.15 (0.71 - 1.82)	0.564
None	2548	57 (2.2%)	2491 (97.8%)	ref	-
1-2 times	474	25 (5.3%)	449 (94.7%)	2.43 (1.44 - 4.01)	0.001
3 or more times	176	12 (6.8%)	164 (93.2%)	3.20 (1.53 - 6.17)	0.001
Frequency of meals in staff canteen					
never	765	10 (1.3%)	755 (98.7%)	ref	-
occasionally	659	17 (2.6%)	642 (97.4%)	2.00 (0.86 - 4.92)	0.083
weekly	1184	45 (3.8%)	1139 (96.2%)	2.98 (1.47 - 6.68)	0.001
daily	2027	66 (3.3%)	1961 (96.7%)	2.54 (1.29 - 5.57)	0.004

Abbreviations: OR, Odds Ratio; CI, Confidence Interval; IQR, Interquartile Range; BMI, Body Mass Index; BCG, Bacillus Calmette-Guerin; AGP, Aerosol-Generating Procedure; COVID-19, Coronavirus Disease-2019


¹ COVID-19 hotspots before April 2020 (i.e. Northern Italy, Austrian ski resorts, or Alsace)

FIGURE CAPTIONS


Figure 1. SARS-CoV-2 seropositivity by district (place of residence of healthcare workers) in Northern and Eastern Switzerland (in grey: no seroprevalence indicated for districts with less than 10 participants).

Figure 2. Figure shows (A) number of protective measures used (among face mask, gown, gloves, goggles) while caring for COVID-19 patients; (B) number of correctly identified elements of standard precautions (among hand hygiene, cough etiquette, mask in case of respiratory symptoms, vaccinations, donning of gowns if potential contact with body fluids); (C) number of contacts with COVID-19 positive co-workers; (D) frequency of meals in the hospital canteen.


Figure 3. Forest plot showing independent association of baseline, occupational and non-occupational risk factors with seropositivity based on multivariable logistic regression analysis.




(A) No of protective measures during COVID-19 patient care



(B) No of correctly identified standard precautions



(C) No of contacts with COVID-19 positive coworker



(D) Frequency of visits to the hospital canteen

