

1

2

3 **Non-COVID-19 patients in times of pandemic: decreased emergency department**
4 **visits and increased out-of-hospital mortality in Northern Italy**

5

6 **Authors:**

7 Luca Santi MD ¹, Davide Golinelli MD ², Andrea Tampieri MD ³, Gabriele Farina MD ¹, Manfredi Greco ²,
8 Simona Rosa MSc ², Michelle Beleffi MD ⁴, Bianca Biavati MD ⁴, Francesca Campinoti MD ⁴, Stefania
9 Guerrini MD ¹, Rodolfo Ferrari MD ³, Paola Rucci, PhD ², Maria Pia Fantini MD ², Fabrizio Giostra MD ¹.

10 ¹ Department of Emergency, Medicina d'Urgenza e Pronto Soccorso, Policlinico S. Orsola-Malpighi,
11 Bologna, Italy

12 ² Department of Biomedical and Neuromotor Sciences (DIBINEM), Alma Mater Studiorum - University of
13 Bologna. Via San Giacomo 12, 40126, Bologna, Italy.

14 ³ Department of Emergency, Medicina d'Urgenza e Pronto Soccorso. Ospedale S. Maria della Scaletta,
15 Imola, Italy

16 ⁴ Alma Mater Studiorum - Università di Bologna

17

18

19 **Authors' affiliations:**

20 Luca Santi, MD: Policlinico S. Orsola-Malpighi. Dipartimento dell'Emergenza-Urgenza. Unità operativa di
21 Medicina d'urgenza e Pronto Soccorso. Via Giuseppe Massarenti, 9, 40138, Bologna, Italy.
22 lucasanti@hotmail.it

23

24 Davide Golinelli, MD: Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum –
25 Università di Bologna. Via San Giacomo 12, 40126, Bologna, Italy. davide.golinelli@unibo.it
26 ORCID: 0000-0001-7331-9520

27 Andrea Tampieri MD: Ospedale S. Maria della Scaletta. Dipartimento di Emergenza-urgenza e Accettazione.
28 Unità Operativa di Pronto Soccorso e Medicina d'Urgenza. Viale Amendola, 2 – 40026 Imola, Italy.
29 a.tampieri@ausl.imola.bo.it

30

31 Gabriele Farina MD: Policlinico S. Orsola-Malpighi. Dipartimento dell'Emergenza-Urgenza. Unità operativa
32 di Medicina d'urgenza e Pronto Soccorso. Via Giuseppe Massarenti, 9, 40138, Bologna, Italy.
33 gabriele.farina@aosp.bo.it

34

35 Manfredi Greco: Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum – Università
36 di Bologna. Via San Giacomo 12, 40126, Bologna, Italy. manfredi.greco@studio.unibo.it

37

38 Simona Rosa, MSc: Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum –
39 Università di Bologna. Via San Giacomo 12, 40126, Bologna, Italy. simona.rosa@unibo.it

40

41 Michelle Beleffi MD: Alma Mater Studiorum - Università di Bologna. Via Zamboni, 33 - 40126 Bologna,
42 Italy. michellebeleffi@gmail.com

43

44 Bianca Biavati MD:Alma Mater Studiorum - Università di Bologna. Via Zamboni, 33 - 40126 Bologna, Italy.
45 biavati.bianca@gmail.com

46

47 Francesca Campinoti MD:Alma Mater Studiorum - Università di Bologna. Via Zamboni, 33 - 40126
48 Bologna, Italy. campinotifrancesca@gmail.com

49

50 Stefania Guerrini MD: Policlinico S. Orsola-Malpighi. Dipartimento dell'Emergenza-Urgenza. Unità
51 operativa di Medicina d'urgenza e Pronto Soccorso. Via Giuseppe Massarenti, 9, 40138, Bologna, Italy.

52 stefania.guerrini@aosp.bo.it

53

54 Rodolfo Ferrari MD: Ospedale S. Maria della Scaletta. Dipartimento di Emergenza-urgenza e Accettazione.
55 Unità Operativa di Pronto Soccorso e Medicina d'Urgenza. Viale Amendola, 2 – 40026 Imola, Italy.
56 r.ferrari@ausl.imola.bo.it

57

58 Paola Rucci, PhD: Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum –
59 Università di Bologna. Via San Giacomo 12, 40126, Bologna, Italy. paola.rucci2@unibo.it

60 ORCID: 0000-0002-7704-0831

61

62 Maria Pia Fantini, MD: Dipartimento di Scienze Biomediche e Neuromotorie, Alma Mater Studiorum –
63 Università di Bologna. Via San Giacomo 12, 40126, Bologna, Italy. mariapia.fantini@unibo.it

64 ORCID: 0000-0002-3257-6552

65

66 Fabrizio Giostra MD: Policlinico S. Orsola-Malpighi. Dipartimento dell'Emergenza-Urgenza. Unità
67 operativa di Medicina d'urgenza e Pronto Soccorso. Via Giuseppe Massarenti, 9, 40138, Bologna, Italy.

68 fabrizio.giostra@aosp.bo.it

69

70 **Corresponding author**

71 Luca Santi, MD: Policlinico S. Orsola-Malpighi. Dipartimento dell'Emergenza-Urgenza. Unità operativa di
72 Medicina d'urgenza e Pronto Soccorso. Via Giuseppe Massarenti, 9, 40138, Bologna, Italy. email:

73 lucasanti@hotmail.it Tel: + 39 0512143324

74

75

76

77

78 **Non-COVID-19 patients in times of pandemic: decreased Emergency Department visits and**
79 **increased out-of-hospital mortality in Northern Italy**

80

81

82 **Abstract**

83 **Objective.** The aim of this was to assess the short-term impact of the pandemic on non-COVID-19 patients living in a
84 one-million inhabitants area in Northern Italy (Bologna Metropolitan Area-BMA), analyzing time trends of Emergency
85 Department (ED) visits, hospitalizations and mortality.

86 **Methods.** We conducted a retrospective observational study using data extracted from BMA healthcare informative
87 systems. Weekly trends of ED visits, hospitalizations, in- and out-of-hospital, all-cause and cause-specific mortality
88 between December 1st, 2019 to May 31st, 2020, were compared with those of the same period of the previous year,
89 using Joinpoint regression models and incidence rate ratios.

90 **Results.** Non-COVID-19 ED visits and hospitalizations showed a stable trend until the first Italian case of COVID-19
91 has been recorded, on February 19th, 2020, when they dropped simultaneously. The reduction of ED visits was observed
92 in all age groups and across all severity and diagnosis groups. In the lockdown period a significant increase was found
93 in overall out-of-hospital mortality (43.2%) and cause-specific out-of-hospital mortality related to neoplasms (76.7%),
94 endocrine, nutritional and metabolic (79.5%) as well as cardiovascular (32.7%) diseases.

95 **Conclusions.** The pandemic caused a sudden drop of ED visits and hospitalizations of non-COVID-19 patients during
96 the lockdown period, and a concurrent increase in out-of-hospital mortality mainly driven by deaths for neoplasms,
97 cardiovascular and endocrine diseases. The findings of this study might be useful to understand both the population
98 reaction and the healthcare system response at the early phases of the pandemic in terms of reduced demand of care and
99 systems capability in intercepting it.

100

101

102

103 **Introduction**

104 **Background**

105 Globally, the Coronavirus Disease (COVID-19) pandemic represents a dramatic burden for healthcare services. Italy
106 was the first Western country affected by the pandemic. The first case of local transmission of Severe Acute Respiratory
107 Syndrome Coronavirus 2 (SARS-CoV-2) in Italy was confirmed in a thirty eight-year-old man in the municipality of
108 Codogno (Lombardy region) on February 19th, 2020 [1], while on February 21st, a resident of Vo', a small town near
109 Padua (Veneto Region), died of pneumonia due to SARS-CoV-2 infection [2,3]. In the following weeks, an exponential
110 growth in the number of cases and deaths in the neighboring regions of northern Italy was observed. As a consequence,
111 the national government enforced as containment measures a complete country lockdown on March 10th, 2020: at that
112 stage the confirmed cases in Italy were 10,149 and deaths 631 [4]. Hospitals and Emergency Departments (EDs) were
113 forced to rapidly adjust to this completely new situation in order to manage an extraordinarily high number of
114 contagious patients with respiratory symptoms [5]

115 However, something else changed due to the pandemic as rapidly as the spread of SARS-CoV-2. As a matter of fact, the
116 pandemic determined a sudden and significant reduction of non-COVID-19 patients seeking for treatment due to urgent
117 medical conditions [6]. This reduction in ED visits might impact the health status of non-COVID patients with acute
118 and chronic diseases [7].

119 In order to better understand this phenomenon and its possible consequences, we retrospectively analyzed amount and
120 type of ED visits, hospitalizations and mortality during lockdown and the preceding and following periods, in the
121 metropolitan area of Bologna (BMA) (~1,000,000 inhabitants), which is the principal town of one of the first and most
122 affected region of Italy and Europe, Emilia Romagna.

123 We hypothesized that trends of ED visits might have been influenced by age, as school closure and social distancing
124 measures might have differently impacted on young and elderly health needs. Moreover, the reduction of ED accesses
125 due to fear of contagion and favored by the stay-at-home message launched on general and social media might also
126 have differently affected the severity pattern of ED accesses. Indeed, we predicted a significant reduction of less severe
127 conditions referring to EDs, whereas no decrease of emergency conditions was expected, including the need of urgent
128 surgical intervention or ICU treatment. We also expected at least a progressive return of ED accesses and

129 hospitalizations to the pre-COVID-19 era levels during the post lockdown phase. No remarkable changes in all-cause
130 and cause-specific mortality, and in the place of death were hypothesized.

131 Our first aim was to compare time trends of ED visits, hospitalizations, in- and out-of-hospital all-cause and cause-
132 specific mortality of non-COVID-19 patients during the first months of the pandemic with the same period of 2019, as
133 possible indicators of unmet healthcare needs.

134 The secondary aim was to subdivide ED patients by age, diagnosis and illness severity, compare their trends, and
135 analyze hospitalizations' trends, focusing on the receiving hospital department (surgical department, medical
136 department, intensive care unit (ICU)).

137

138 **Methods**

139 **Study design, population and setting**

140 We conducted a retrospective observational study using anonymous data extracted from the healthcare administrative
141 databases and informative systems of BMA, Northern Italy.

142 The study population consisted of all age residents in BMA, which encompasses the city of Bologna and the
143 neighboring municipalities, for a total of 1,019,875 citizens as of January 1st, 2020 [8]. The demographic characteristics
144 of the population are reported in Table 1 [Suppl mat].

145 In the BMA, the first patient reported with SARS-CoV-2 infection was recorded on February 28th. On May 31st, the
146 number of infected patients was 5,021 with 684 deaths [4].

147 The BMA healthcare system consists of twelve hospitals equipped with EDs. Among these, four are located in the urban
148 area and two of these are large university hospitals, four are suburban whereas the last four are rural hospitals. The
149 volume of patients annually assessed in the different EDs ranges from ~3,000 up to ~150,000 in 2019.

150 This study follows the STROBE reporting guidelines for observational studies.

151

152 **Study data**

153 We examined the weekly trends of ED visits, hospitalizations of patients admitted from ED and in- and out-of-hospital
154 all-cause and cause-specific mortality between December 1st, 2019 to May 31st, 2020, and compared them with the
155 trends of the same period of the previous year.

156 The study period was subdivided into four periods, starting from the beginning of the flu peak up to four weeks after the
157 end of the national lockdown. Namely, the following are the four subgroups of the study period:

- 158 • flu period (from December 1st, 2019 to February 23rd, 2020the outbreak of the COVID-19 pandemic to the first
159 containment measures, i.e. from the beginning of the flu period to week 49th 2019 to week 8th 2020);
- 160 • pre-lockdown period (from the initial alarm period in the absence of containment institutional measures until the
161 lockdown institution, i.e. February 24th, 2020 to March 9th, 2020, weeks 9th to 10th 2020);
- 162 • lockdown period (March 10th, 2020 to May 3rd, 2020, weeks 11th to 18th 2020);
- 163 • post-lockdown period (i.e. the reopening period, from the May 4th, 2020 to May 31st, 2020, week 19th to 22nd 2020).

164 The data sources include the Emergency Department Database, the mortality registry updated to June 2020 and the Civil
165 Protection Database [4]. For the purpose of the present study, the data extracted from the diverse databases were the
166 following: weekly number of ED visits, type of ED visits (main diagnosis, severity code), patients characteristics (age,
167 gender), weekly number of hospitalizations, department of hospitalization (medical, surgical, ICU), weekly number of
168 deaths, cause of death (main diagnosis) and place of death (in-hospital, out-of-hospital).

169 The main diagnosis of ED visits and deaths was coded through the International Classification of Diseases Clinical
170 Modification Ninth (ICD-9-CM) and Tenth (ICD-10-CM) versions, respectively.

171 Non-COVID-19 cases were identified by excluding the COVID-19 cases diagnosed by the emergency physician as
172 suspected or confirmed (ICD-9-CM codes 480,480.3,079.82,V01.82,V01.79), in primary or secondary diagnosis of
173 discharged and hospitalized patients. For all mortality data, the specific codes related to COVID-19 and SARS-CoV-2
174 infection (U00-U85) were ruled out.

175

176 **Statistical analysis**

177 The demographic and clinical characteristics of the population were summarized using mean and standard deviation or
178 absolute and percentage frequencies.

179 We analyzed and displayed trends of ED visits, hospital admissions, and mortality, divided into the four time periods
180 and compared them with those of the same periods of the previous year.

181 The trends of ED visits, hospitalizations and all-cause mortality were analyzed using the Joinpoint Trend Analysis
182 Software 4.6.0.0 (Statistical Research and Applications Branch, National Cancer Institute, USA) [9]. Joinpoint models
183 allowed to identify changes in the slopes of ED visits and hospitalizations during the periods of interest.

184 We carried out a preliminary analysis comparing the time trend of ED visits, hospitalizations and mortality rates in 2019
185 with those of the years 2014-2018. This analysis showed no significant changes in ED visits, hospitalizations and
186 mortality overtime. Therefore, we considered 2019 as the reference year.

187 For what concerns age of non-COVID-19 ED patients during pandemic, it was classified into three groups: pediatric
188 patients (0-14 years), adults (15-64 year) and elderly (≥ 65 years). We also classified ED visits according to the triage
189 code (red, yellow, green and white) and by ICD-9-CM disease category and specific cause.

190 Then, we compared the trends of admissions to hospital departments between 2019 and 2020.

191 We also analyzed the trends of all-cause, cause-specific and in-hospital (public or private) and out-of-hospital (home,
192 social welfare structure, other) mortality between 2019 and 2020. The latter was also split into cause-specific out-of-
193 hospital mortality.

194 Incidence rate ratios (IRRs) of hospitalizations and mortality in 2020 vs. 2019 were estimated using Poisson regression
195 models.

196

197


198 **Results**

199 The empirical trends of ED visits, hospitalizations and mortality in 2019 and 2020 are reported in Figure 1.

200

201

202 **Figure 1. Trend of ED visits (a), hospitalizations (b) and mortality (c) during the study periods for years 2019 and**
 203 **2020. For a better comprehension of the burden of the pandemic on health care services, we displayed the trend**
 204 **of the three indicators including and excluding SARS-CoV-2 related data^a.**



205

206 ^aNote: SARS-CoV-2 cases weekly incidence is reported on the right y-axis.

207 ^x Using Joinpoint analysis, a significant change of trend ($p < 0.05$) was identified between week 8-9 and then week 12-13 2020 for ED
 208 visits and between week 8-9 and then 13-14 for hospitalizations. The analysis of mortality data showed a slight significant change
 209 during week 15 2020.

210 **ED visits**

211 Figure 1a shows that the number of ED visits was almost stable in the year 2019 until the end of 2020 flu period, when
212 it dropped simultaneously with the exponential growth of the local SARS-CoV-2 case curve. Then, during the lockdown
213 period the number of ED visits started to rise, but with a slower pace when compared to the previous decrease.

214 Joinpoint analysis showed a significant ($p<0.05$) reduction in ED visits starting from week nine (pre-lockdown period),
215 leading to the lowest numbers at the thirteenth week during lockdown, then followed again by a significant increase
216 ($p<0.05$) at the fourteenth week. During the lockdown period the overall reduction in ED visits compared to same
217 period of the previous year was -66.2% (Table 2 [Suppl mat]). ED accesses' decrease occurred during the steep climb of
218 COVID-19 spread until the peak. The subsequent slow increase in ED visits never returned to the previous year levels,
219 not even at the end of the study period, four weeks after the end of the national lockdown.


220 The impressive reduction in the number of ED visits was found in all age groups, throughout all study periods (except
221 for the flu period) (Table 3 [Suppl mat]). The highest decrease was observed in pediatric patients during lockdown (-
222 83.2%), although even adults and elderly patients ED accesses declined by 62.9% and 64.0%, respectively (Table 3
223 [Suppl mat]).

224 For what concerns triage severity codes of non-COVID-19 patients, the 4 codes were affected by the pandemic
225 similarly; as a matters of fact, a reduction regarded similarly the four triage codes (Figure 2) during the study periods
226 (except for the flu period) if compared to the previous year, with the most important decrease (between 60% and 70%)
227 observed during lockdown.

228

229 **Figure 2. Percentage change in ED visits by severity code compared to the previous year.**

230



231
232


233 Note: ED visits are categorized by severity codes according to the triage protocols in four levels of urgency: white code (not-urgent
234 condition), green code (postponable conditions); yellow code (medical condition requiring urgent care with no vital signs
235 impairment but at-risk for deterioration); red code (medical condition with acute impairment of vital signs requiring emergency care).

236

237

238 As for non-COVID ED visits for specific conditions, our analysis demonstrated a reduction which broadly affected all
239 disease categories, ranging from 41% to 83% in the lockdown period (Figure 3). The decrease of ED visits was more
240 pronounced for infectious and parasitic diseases (-83%), skin and subcutaneous tissue pathogenic conditions (-80%),
241 and congenital anomalies (-78%) (Figure 3). A remarkable decrease in ED visits was also observed for conditions
242 usually significantly impacting on BMA's EDs (data not shown), such as diseases of the cardiovascular system (-58%)
243 as well as endocrine, nutritional and metabolic diseases (-63%). Specifically, the remarkable reduction in ED visits
244 interested life-threatening illnesses, such as ischemic heart disease (-49.7%), intracranial hemorrhages (-33.3%) and
245 ischemic cerebrovascular disorders (-45.8 %) (Table 4 [Suppl mat]). Lower reductions or a modest increase were
246 recorded in complications of pregnancy, childbirth and the puerperium and some morbid conditions of perinatal origin
247 (+8% during pre-lockdown, -36% during lockdown, and -30% during post-lockdown period) (Figure 3).

248 **Figure 3. Percentage change in ED visits by disease category compared to the previous year.**



249

250 ^a Note: no remarkable changes of ED visits by disease category have been found during the flu period (data not shown).

251

252 **Hospitalizations**

253 Figure 1b shows hospitalizations trends in the study periods, which appears to be quite stable during year 2019 until the
254 end of 2020 flu period. Similarly to ED visits trends, hospital admissions started to drop simultaneously with the
255 exponential growth of the local SARS-CoV-2 case curve. The number of hospitalizations started to rise again during the
256 lockdown period, but with a slower pace.

257 Joinpoint analysis showed that non-COVID hospitalizations significantly ($p < 0.05$) declined between weeks eight and
258 fourteen before the peak of incidence of COVID-19 cases, and moderately increased afterwards (Figure 1).

259 The overall reduction of hospitalizations was -46% during the lockdown period compared to previous year (Table 2
260 [Suppl mat]). Admissions to all hospital departments declined during and after lockdown (Table 1). During lockdown,
261 admissions reduction was significant and interested likewise surgical and medical departments (-36.0%) as well as ICUs
262 (-36.4%).

263 **Table 1. Number and percentage change of hospitalizations in Medicine, Surgery and ICU departments, year 2020 compared to 2019.**

	2020 (n)				Change from previous year (%)				Incidence Rate Ratio [95% CI]			
	Flu-period	Pre-lockdown	Lockdown	Post-lockdown	Flu-period	Pre-lockdown	Lockdown	Post-lockdown	Flu-period	Pre-lockdown	Lockdown	Post-lockdown
Medicine	9,903	1,493	4,072	2,441	-1.5%	-11.9%	-36.0%	-22.3%	0.986 [0.961-1.013]	0.883 ^a [0.824-0.946]	0.654 ^a [0.629-0.679]	0.782 ^a [0.742-0.824]
Surgery	4,112	647	1,775	1,122	-5.9%	-7.3%	-36.0%	-19.4%	0.943 ^a [0.905-0.984]	0.927 [0.834-1.032]	0.647 ^a [0.610-0.686]	0.808 ^a [0.747-0.874]
Intensive Care Unit	709	100	286	186	2.3%	-21.3%	-36.4%	-18.4%	1.023 [0.922-1.135]	0.788 [0.606-1.023]	0.637 ^a [0.549-0.738]	0.816 ^a [0.673-0.990]

^a p<0.05

264

265

266 **Mortality**

267 The trend of non-COVID-19 mortality was similar in 2019 and 2020, with a slight decline after the winter season
268 (Figure 1c).

269 A small increase in all-cause overall mortality (7.5%) was observed only during lockdown (IRR 1.074, $p < 0.05$, 95%CI
270 [1.007-1.145]) (table 2), whereas in both pre- and post-lockdown periods no significant change in mortality was
271 detectable. Lockdown was associated with a significantly higher cause-specific mortality related to endocrine,
272 nutritional and metabolic diseases (47.9%, IRR 1.478, $p < 0.05$, 95%CI [1.094-1.998]) and diseases of the circulatory
273 system (15.5%, IRR 1.154, $p < 0.05$, 95%CI [1.031-1.291]). Conversely, mortality reduced only for infectious and
274 parasitic diseases (-36.2%, IRR 0.638, $p < 0.05$, 95%CI [0.437-0.931]). As for mental, behavioral and neuro-
275 developmental disorders, a significant increase in mortality during the pre-lockdown period was observed, while deaths
276 for neoplasms decreased during the post-lockdown weeks (Table 2).

277 We found a significant increase in out-of-hospital mortality, particularly during the lockdown weeks (43.2%, IRR 1.428,
278 $p < 0.05$, 95%CI [1.295-1.574]), whereas in the pre-lockdown period and in the post-lockdown period no remarkable
279 changes were identified. A concurrent reduction of in-hospital mortality was observed (-15.0%, IRR 0.759, $p < 0.05$,
280 95%CI [0.694-0.831]).

281 Analysis of cause-specific out-of-hospital mortality (Table 2) confirmed an increase in deaths for neoplasms, endocrine,
282 nutritional and metabolic diseases, and diseases of the circulatory systems during lockdown. Conversely, we found a
283 significant reduction only in genitourinary disorders.

284

285

286 **Table 2. Number and percentage change of all-cause mortality (overall and cause-specific), in-hospital and out-of-hospital mortality (overall and cause-specific),**
 287 **year 2020 compared to 2019.**

	2020 (n)				Change from previous year (%)				Incidence Rate Ratio [95% CI]			
	Flu Period	Pre- lockdown	Lock- down	Post- lockdown	Flu Period	Pre- lockdown	Lock- down	Post- lockdown	Flu Period	Pre-lockdown	Lock-down	Post-lockdown
Certain infectious and parasitic diseases	90	15	44	32	-12.6%	-25.0%	-36.2%	-3.0%	0.874 [0.659-1.159]	0.750 [0.384-1.465]	0.638 ^a [0.437-0.931]	0.970 [0.596-1.577]
Neoplasms	722	137	489	213	-2.2%	14.2%	2.1%	-22.0%	0.979 [0.883-1.084]	1.142 [0.894-1.458]	1.021 [0.900-1.158]	0.781 ^a [0.653-0.934]
Endocrine, nutritional and metabolic diseases	138	18	105	29	0.0%	20.0%	47.9%	-9.4%	1.000 [0.790-1.266]	1.200 [0.605-2.381]	1.478 ^a [1.094-1.998]	0.906 [0.548-1.498]
Mental, Behavioral and Neurodevelopmental disorders	131	35	83	45	-12.7%	84.2%	-3.5%	9.8%	0.874 [0.691-1.104]	1.842 ^a [1.054-3.219]	0.965 [0.714-1.305]	1.098 [0.719-1.676]
Diseases of the nervous system	118	23	72	24	18.0%	-8.0%	5.9%	-4.0%	1.180 [0.904-1.540]	0.920 [0.522-1.621]	1.059 [0.760-1.475]	0.960 [0.548-1.681]
Diseases of the circulatory system	915	161	650	229	-14.2%	-4.7%	15.5%	-16.7%	0.860 ^a [0.787-0.939]	0.953 [0.768-1.182]	1.154 ^a [1.031-1.291]	0.833 ^a [0.699-0.993]
Diseases of the respiratory system	325	49	218	60	-11.2%	-29.0%	19.8%	-26.8%	0.888 [0.765-1.031]	0.710 [0.493-1.024]	1.197 [0.984-1.458]	0.732 [0.525-1.021]
Diseases of the digestive system	100	14	64	28	-11.5%	0.0%	12.3%	0.0%	0.885 [0.676-1.158]	1.000 [0.477-2.098]	1.123 [0.786-1.604]	1.000 [0.592-1.688]
Diseases of the genitourinary system	62	19	36	24	-8.8%	58.3%	-25.0%	0.0%	0.912 [0.646-1.286]	1.583 [0.769-3.261]	0.750 [0.487-1.155]	1.000 [0.568-1.761]
Injury, poisoning and certain other consequences of external causes	120	24	77	35	-8.4%	9.1%	-2.5%	-16.7%	0.916 [0.715-1.173]	1.091 [0.612-1.945]	0.975 [0.712-1.334]	0.833 [0.532-1.305]
Other^a	103	16	55	21	21.2%	-5.9%	-6.8%	-34.4%	1.212 [0.909-1.614]	0.941 [0.476-1.863]	0.932 [0.646-1.346]	0.656 [0.379-1.138]
All-cause overall mortality	2,824	511	1,893	740	-7.7%	1.8%	7.5%	-16.6%	0.926 ^a [0.880-0.973]	1.018 [0.900-1.151]	1.074 ^a [1.007-1.145]	0.836 ^a [0.758-0.921]
In-hospital overall mortality	1,648	289	918	426	-6.5%	-4.9%	-15.0%	-19.3%	0.936 [0.876 - 1.000]	0.951 [0.810 - 1.117]	0.759 ^a [0.694 - 0.831]	0.808 ^a [0.711 - 0.917]
Out-of-hospital overall mortality	1,176	222	975	314	-9.2%	12.1%	43.2%	-12.5%	0.909 ^a [0.841 - 0.983]	1.121 [0.926 - 1.357]	1.428 ^a [1.295 a 1.574]	0.875 [0.752 - 1.018]
Certain infectious and parasitic diseases	15	1	7	3	66.7%	0.0%	133.3%	-25.0%	1.667 [0.729 - 3.808]	1.000 [0.063 - 15.988]	2.333 [0.603 - 9.023]	0.750 [0.168 - 3.351]

Neoplasms	232	42	228	85	-2.1%	50.0%	76.7%	-8.6%	0.979 [0.817 - 1.173]	1.500 [0.930 - 2.419]	1.766 ^a [1.4230 - 2.191]	0.914 [0.681 - 1.226]
Endocrine, nutritional and metabolic diseases	77	9	61	15	-2.5%	12.5%	79.4%	-6.3%	0.975 [0.712 - 1.334]	1.125 [0.434 - 2.916]	1.794 ^a [1.179 - 2.728]	0.937 [0.464 - 1.896]
Mental, Behavioral and Neurodevelopmental disorders	68	18	52	29	-17.1%	80.0%	40.5%	123.1%	0.829 [0.601 - 1.144]	1.780 [0.831 - 3.899]	1.405 [0.922 - 2.142]	2.230 ^a [1.160 - 4.290]
Diseases of the nervous system	73	14	52	12	40.4%	16.7%	36.8%	-14.3%	1.404 [0.984 - 2.003]	1.167 [0.540 - 2.522]	1.368 [0.901 - 2.079]	0.857 [0.397 - 1.853]
Diseases of the circulatory system	486	95	418	115	-15.9%	1.1%	32.7%	-21.2%	0.842 ^a [0.746 - 0.949]	1.011 [0.760 - 1.344]	1.326 ^a [1.146 - 1.534]	0.788 [0.617 - 1.006]
Diseases of the respiratory system	85	9	59	17	-13.3%	-43.8%	47.5%	-15.0%	0.868 [0.649 - 1.160]	0.563 [0.249 - 1.273]	1.477 [0.987 - 2.203]	0.850 [0.445 - 1.623]
Diseases of the digestive system	14	3	14	5	7.7%	-25.0%	40.0%	25.0%	1.077 [0.506 - 2.291]	0.750 [0.168 - 3.351]	1.400 [0.622 - 3.152]	1.250 [0.336 - 4.655]
Diseases of the genitourinary system	16	4	4	5	-20.0%	33.3%	-71.4%	-37.5%	0.800 [0.415 - 1.544]	1.333 [0.298 - 5.957]	0.286 ^a [0.094 - 0.868]	0.625 [0.205 - 1.911]
Injury, poisoning and certain other consequences of external causes	52	16	39	17	-27.8%	60.0%	11.4%	-15.0%	0.722 [0.506 - 1.032]	1.600 [0.726 - 3.525]	1.114 [0.706 - 1.759]	0.850 [0.445 - 1.623]
Other^b	58	11	41	11	5.5%	-8.3%	57.7%	-47.6%	1.055 [0.729 - 1.525]	0.917 [0.405 - 2.077]	1.577 [0.965 - 2.577]	0.524 [0.253 - 1.087]

^a p<0.05

^b Causes of deaths that accounted for ≤0.5% of the total (ICD-10-CM categories D50-89, H00-95, L00-99, M00-99, O00-9A, P00-96, Q00-99, R00-99, V00-Y99, Z00-99) were included in Other.

288
289
290
291

292 **Discussion**

293 Globally, Italy has been the first Western country to experience the disruptive effect of the pandemic in terms of cases,
294 deaths and the related burden on healthcare services [1-3, 10]. Since the beginning of the pandemic, a reduction of ED
295 visits and hospitalizations overall [6] and for specific diseases [11-15] have been reported worldwide.

296 In this paper we studied the pandemic consequences on non-COVID-19 patients in an area accounting for over one
297 million inhabitants. We highlighted how ED visits and hospitalizations reduced overall, and particularly during the
298 lockdown period. To the best of our knowledge, this is the first study to analyze also the trends of in- and out-of-
299 hospital, all-cause and cause-specific mortality, providing a comprehensive picture of the short-term consequences of
300 the pandemic on non-COVID-19 patients in Italy, the first hit European country.

301 Our findings indicate that the reduction of ED visits and hospitalizations started two weeks before the beginning of the
302 national lockdown, after school closures and the first Italian case hospitalized for COVID-19 in Codogno [1], when no
303 cases of local transmission had been still recorded in BMA. As already reported [2, 6, 18-20], a possible explanation is
304 that the population response is likely to be more affected by the national level authority risk message than the real local
305 situation [21]. Several possible reasons have been put forward to explain, at least partially, a reduction in ED visits and
306 hospitalizations [6, 7]. In addition to pathologies likely reduced by the lockdown and lifestyle changes, this may have
307 been caused by fear of the contagion, as well as by media campaigns encouraging to stay-at-home, together with the
308 increasingly stringent lockdown measures and postponement of elective procedures, or the sense of civic responsibility
309 of the population [7].

310 It should be highlighted that not all ED visits represent real emergencies, and inappropriate accesses to ED are a well-
311 known issue related to several reasons [7, 20, 22, 23]. That being said, the observed reduction of the less severe triage
312 codes was to somewhat expected, although not to this large extent. What we were not prepared for was the concurrent
313 decrease of the most serious cases. Furthermore, we showed how ED visits' reduction encompassed all pathological
314 conditions, thus including also acute and time-dependent diseases, similarly to what already reported in literature [24-
315 26]. The overall reduction of ED visits, regardless of age, severity, and causes, suggests the system's inability to guide
316 the patient in discriminating the real need for urgent care. Therefore, public awareness should be taken into due
317 consideration to be prepared for a recrudescence of the pandemic.

318 A clear reduction of non-COVID-19 patients hospitalization has also been recorded, both for medical and surgical
319 diseases and for conditions requiring intensive care support. Data on hospital admissions reported in previous studies
320 [6,12] appear to be not comparable, as obtained from selected populations [12] or different healthcare systems, such as
321 U.S. care system or other insurance-based systems [6]. In fact, unlike the U.S., the Italian National Healthcare Service
322 is a single-payer system funded by general taxation, similarly to the UK national health service [27]. This could have
323 an impact on a system's hospitalization rate [28].

324 Looking at the post-lockdown period, when the spread of COVID-19 was contained, we expected to see a progressive
325 increase in ED visits and a consequent increase in hospitalizations, as a consequence of the delay in obtaining medical
326 attention and care access due to lockdown. However, even if these indicators started to rise during the lockdown weeks
327 and throughout the post-lockdown period, they did not reach the pre-pandemic levels.

328 The all-cause mortality among non-COVID-19 patients showed only a minimal increase (+7.5%) during the lockdown
329 period. The significant increase in mortality for cardiovascular and endocrine systems' diseases that we have registered
330 during lockdown is in line with other studies [24-26, 29].

331 The key finding of our study is that, with the sudden drop of ED visits and hospitalizations, we found a statistically
332 significant increase of out-of-hospital all-cause mortality, mainly driven by an increase in deaths for neoplasms,
333 cardiovascular diseases and endocrine, nutritional and metabolic diseases. In-hospital mortality instead showed an
334 opposite and decreasing trend, due to the reduction of non-COVID-19 patients in healthcare facilities. We assume that
335 these data are attributable to a delay in seeking hospital care and to system's inability in providing the same standards
336 of care during a public health emergency, both in terms of quality and quantity. Our findings confirm those reported by
337 Walker, L.E et al [30]. related to a multi-hospital system, in which an increase of non-COVID-related overall mortality
338 was found to be mainly driven by out-of-hospital deaths, due to patients deferring care.

339 During the pre-lockdown weeks, we observed a significant increase in mortality for mental diseases. This might be due
340 to the sense of fear and anxiety which reigned during the pre-lockdown period, with a consequent exacerbation of
341 mental disorders [31, 32]. The reduction in overall mortality for neoplasms during the post-lockdown weeks, when
342 considered together with the increased out-of-hospital mortality for the same cause recorded in the lockdown period,
343 likely needs to be analyzed when medium- and long-term data will be available.

344 These data suggest that all-cause and cause-specific out-of-hospital mortality might be considered a useful indicator for
345 targeting the consequences of the pandemic on non-COVID-19 patients.

346

347 **Limitations**

348 This study has some limitations. First, mortality data might include under-diagnosed COVID-19 cases. In fact the type
349 of access and causes of death could have been coded incorrectly. However this is unavoidable due to the case definition
350 and the context of that time-period, when the pandemic disruptively changed routine practice. Furthermore, considering
351 the delayed onset of the pandemic in BMA and the preparedness efforts of local healthcare authorities, compared to
352 other Italian areas earlier affected, we assume that no significant underestimation of COVID-19 deaths occurred, and
353 that just a minority of COVID-19 deaths occurred out of the hospital, given that the vast majority of severely
354 symptomatic COVID-19 patients had been hospitalized.

355 Furthermore, we had access to cause-specific mortality data, but only aggregated into broad categories (e.g. death for
356 cardiovascular diseases, death for injury or poisoning, etc.). Data on more specific causes of death are still not available
357 and therefore no conclusions can be drawn on more specific causes of death.

358 Finally, the short time window of the post-lockdown period considered in this study does not allow us to generalize our
359 findings regarding possible medium and long term effects of the pandemic. Additional data are necessary to understand
360 the effects of diagnostic and therapeutic delays that the pandemic has inevitably entailed, both for chronic and for acute
361 patients.

362

363 **Conclusions**

364 In summary, our analysis showed a clear reduction of ED visits and hospitalizations for non-COVID-19 patients during
365 the first months of the pandemic in a wide and populated area, and a significant increase in non-COVID-19 out-of-
366 hospital deaths. Out-of-hospital mortality due to neoplasms, cardiovascular and endocrine systems' diseases showed a
367 significant increase during the lockdown weeks, while for other conditions the consequences of the pandemic on non-
368 COVID-19 patients are still unknown.

369 During the first COVID-19 wave, the reduction in ED visits and hospitalization probably eased the healthcare system in

370 facing an unprecedented emergency. In view of the recurrence of the crisis, the scenario described by our study
371 represents a teachable moment about population and healthcare systems' response at the early stages of the pandemic in
372 terms of reduced demand of care and systems' capability in intercepting it. According to our findings, we also advocate
373 the need for further investigations into the medium and long term effects of the pandemic on non-COVID-19 patients,
374 in order to strengthen population awareness and systems' preparedness.

375

376

377 **References**

- 378 1. Odone A, Delmonte D, Scognamiglio T, et al. (2020) COVID-19 deaths in Lombardy, Italy: data in context *Lancet*
379 *Public Health*. 5(6):e310. doi:10.1016/S2468-2667(20)30099-2
- 380 2. Gibertoni D, Kadjo YCA, Golinelli D, et al. (2020) Patterns of COVID-19 related excess mortality in the
381 municipalities of Northern Italy 2020.05.11.20097964; doi: <https://doi.org/10.1101/2020.05.11.20097964>
- 382 3. Lavezzo E, Franchin E, Ciavarella C, et al. (2020) Suppression of a SARS-CoV-2 outbreak in the Italian municipality
383 of Vo'. *Nature*. 584(7821):425-429. doi:10.1038/s41586-020-2488-1
- 384 4. Italian Civil Protection Department (2020) COVID-19: Situazione Italia.
385 <https://opendatadpc.maps.arcgis.com/apps/opsdashboard/index.html#/b0c68bce2cce478eaac82fe38d4138b1>. Accessed
386 14 October 2020.
- 387 5. Wang D, Hu B, Hu C, et al. (2020) Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel
388 Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 323(11):1061-1069. doi:10.1001/jama.2020.1585
- 389 6. Jeffery MM, D'Onofrio G, Paek H, et al. (2020) Trends in Emergency Department Visits and Hospital Admissions in
390 Health Care Systems in 5 States in the First Months of the COVID-19 Pandemic in the US. *JAMA Intern Med*.
391 180(10):1328-1333. doi:10.1001/jamainternmed.2020.3288
- 392 7. Rosenbaum L. (2020) The Untold Toll - The Pandemic's Effects on Patients without Covid-19. *N Engl J Med*.
393 382(24):2368-2371. doi:10.1056/NEJMms2009984
- 394 8. Emilia-Romagna Region Statistics Office [online] (2020) [https://statistica.regione.emilia-romagna.it/servizi-](https://statistica.regione.emilia-romagna.it/servizi-online/statistica-self-service/popolazione/popolazione-residente-dal-1861)
395 [online/statistica-self-service/popolazione/popolazione-residente-dal-1861](https://statistica-regione-emilia-romagna.it/servizi-online/statistica-self-service/popolazione/popolazione-residente-dal-1861). Accessed 14 October 2020

- 396 9. Kim HJ, Fay MP, Feuer EJ, et al. (2000) Permutation tests for joinpoint regression with applications to cancer rates.
397 Stat Med. 19(3):335-351. doi:10.1002/(sici)1097-0258(20000215)19:3<335::aid-sim336>3.0.co;2-z
- 398 10. Van Damme W, Dahake R, Delamou A, et al. (2020) The COVID-19 pandemic: diverse contexts; different
399 epidemics-how and why?. BMJ Glob Health. 5(7):e003098. doi:10.1136/bmjgh-2020-003098
- 400 11. Tartari F, Guglielmo A, Fuligni F, et al. (2020) Changes in emergency service access after spread of COVID-19
401 across Italy. J Eur Acad Dermatol Venereol. 34(8):e350-e351. doi:10.1111/jdv.16553
- 402 12. Baum A, Schwartz MD (2020) Admissions to Veterans Affairs Hospitals for Emergency Conditions During the
403 COVID-19 Pandemic. JAMA. 324(1):96-99. doi:10.1001/jama.2020.9972
- 404 13. Lazzerini M, Barbi E, Apicella A, et al. (2020) Delayed access or provision of care in Italy resulting from fear of
405 COVID-19. Lancet Child Adolesc Health. 4(5):e10-e11. doi:10.1016/S2352-4642(20)30108-5
- 406 14. Ferrero F, Ossorio MF, Torres FA, et al. (2020) Impact of the COVID-19 pandemic in the paediatric emergency
407 department attendances in Argentina. Arch Dis Child. archdischild-2020-319833. doi:10.1136/archdischild-2020-
408 319833
- 409 15. Schwarz V, Mahfoud F, Lauder L, et al. (2020) Decline of emergency admissions for cardiovascular and
410 cerebrovascular events after the outbreak of COVID-19. Clin Res Cardiol. 1-7. doi:10.1007/s00392-020-01688-9
- 411 16. Weber EJ, Hirst E, Marsh M. (2017) The patient's dilemma: attending the emergency department with a minor
412 illness. BMJ. 357:j1941. doi:10.1136/bmj.j1941
- 413 17. Buonanno P, Galletta S, Puca M. (2020) Estimating the severity of COVID-19: Evidence from the Italian epicenter.
414 PLoS One. 15(10):e0239569. Published 2020 Oct 1. doi:10.1371/journal.pone.0239569
- 415 18. Blangiardo M, Cameletti M, Pirani M, et al. (2020) Estimating weekly excess mortality at sub-national level in Italy
416 during the COVID-19 pandemic. PLoS One. 15(10):e0240286. doi:10.1371/journal.pone.0240286
- 417 19. Mafham MM, Spata E, Goldacre R, et al. (2020) COVID-19 pandemic and admission rates for and management of
418 acute coronary syndromes in England. Lancet. 396(10248):381-389. doi:10.1016/S0140-6736(20)31356-8
- 419 20. Kraaijevanger N, van Leeuwen H, Rijpsma D, et al. (2016) Motives for self-referral to the emergency department: a
420 systematic review of the literature. BMC Health Serv Res. 16(1):685. Published 2016 Dec 9. doi:10.1186/s12913-016-
421 1935-z
- 422 21. Westgard BC, Morgan MW, Vazquez-Benitez G, et al. (2020) An Analysis of Changes in Emergency Department
423 Visits After a State Declaration During the Time of COVID-19. Ann Emerg Med. 76(5):595-601.
424 doi:10.1016/j.annemergmed.2020.06.019

- 425 22 . Sancton K, Sloss L, Berkowitz J, et al. (2018) Low-acuity presentations to the emergency department: Reasons for
426 and access to other health care providers before presentation. *Can Fam Physician*. 64(8):e354-e360.
- 427 23. Chaiyachati K, Kangovi S. (2020) Inappropriate ED visits: patient responsibility or an attribution bias?. *BMJ Qual*
428 *Saf*. 29(6):441-442. doi:10.1136/bmjqs-2019-009729
- 429 24. De Filippo O, D'Ascenzo F, Angelini F, et al. (2020) Reduced Rate of Hospital Admissions for ACS during Covid-
430 19 Outbreak in Northern Italy. *N Engl J Med*. 383(1):88-89. doi:10.1056/NEJMc2009166
- 431 25. Solomon MD, McNulty EJ, Rana JS, et al. (2020) The Covid-19 Pandemic and the Incidence of Acute Myocardial
432 Infarction. *N Engl J Med*. 383(7):691-693. doi:10.1056/NEJMc2015630
- 433 26. Wu J, Mamas MA, Mohamed MO, et al. (2020) Place and causes of acute cardiovascular mortality during the
434 COVID-19 pandemic. *Heart*. heartjnl-2020-317912. doi:10.1136/heartjnl-2020-317912
- 435 27. Golinelli D, Toscano F, Bucci A, et al. (2017) Health Expenditure and All-Cause Mortality in the 'Galaxy' of Italian
436 Regional Healthcare Systems: A 15-Year Panel Data Analysis. *Appl Health Econ Health Policy*. 15(6):773-783.
437 doi:10.1007/s40258-017-0342-x
- 438 28. Brown LD. (2003) Comparing health systems in four countries: lessons for the United States. *Am J Public Health*.
439 93(1):52-56. doi:10.2105/ajph.93.1.52
- 440 29. Baldi E, Sechi GM, Mare C, et al. (2020) Out-of-Hospital Cardiac Arrest during the Covid-19 Outbreak in Italy. *N*
441 *Engl J Med*. 383(5):496-498. doi:10.1056/NEJMc2010418
- 442 30. Walker, L.E. et al. (2020) Impact of the SARS-CoV-2 Pandemic on Emergency Department Presentations in an
443 Integrated Health System, *Annals of Emergency Medicine*. 76 (4), S2 - S3. doi:10.1016/j.annemergmed.2020.09.014
- 444 31. Hoyer C, Ebert A, Szabo K, et al. (2020) Decreased utilization of mental health emergency service during the
445 COVID-19 pandemic. *Eur Arch Psychiatry Clin Neurosci*. 1-3. doi:10.1007/s00406-020-01151-w
- 446 32. Moreno C, Wykes T, Galderisi S, et al. (2020) How mental health care should change as a consequence of the
447 COVID-19 pandemic. *Lancet Psychiatry*. 7(9):813-824. doi:10.1016/S2215-0366(20)30307-2