A higher ratio of green spaces means a lower racial disparity in severe acute respiratory syndrome coronavirus 2 infection rates: A nationwide study of the United States

Yi LU [a], Long CHEN [a], Xueming LIU [b], Yuwen YANG [b, c], Wenyan XU [b, c], Chris WEBSTER [d], William C. SULLIVAN [e], Bin JIANG [b, c *]

[a] Department of Architecture and Civil Engineering, College of Engineering, City University of Hong Kong, Hong Kong SAR

[b] Virtual Reality Lab of Urban Environments and Human Health, HKUrbanLabs, The University of Hong Kong, Hong Kong SAR

[c] Division of Landscape Architecture, Department of Architecture, The University of Hong Kong, Hong Kong SAR

[d] HKUrbanLabs, Faculty of Architecture, The University of Hong Kong, Hong Kong SAR

[e] Smart, Healthy Communities initiative, University of Illinois at Urbana-Champaign, U.S.A.

Corresponding author:

[1*.] Postal Address: 614 Knowles Building, The University of Hong Kong, Pokfulam Road, Hong Kong SAR

Email: jiangbin@hku.hk

Tel: 852-55796958

Highlights

- The first study identified significant relationships between green spaces and the racial disparity of SARS-CoV-2 infection rates.

- A nationwide study of 135 most urbanized counties of the United States.

- The black-white racial disparity of SARS-CoV-2 infection rates was measured within each county.

- A higher ratio of green spaces is associated with a lower racial disparity of SARS-CoV-2 infection rates.

- Four green space factors are positively associated with a lower racial disparity of SARS-CoV-2 infection rates.

Abstract

There is striking racial disparity in the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection rates in the United States. We hypothesize that the disparity is significantly smaller in areas with a higher ratio of green spaces at the county level, because green spaces provide health benefits to people of different races through multiple pathways. We suggest that they are an equalizing salutogenic factor, modifying infection exposure. This study used the 135 most urbanized counties across the United States as sample sites. County level data on the SARS-CoV-2 infection rates of blacks and whites in each county were collected. The ratio of green spaces by land-cover type at the county level was calculated from satellite imagery. An ecological hierarchical regression analysis measured cross-sectional associations between racial disparity in infection rates and green spaces, after controlling for confounding factors. We found significantly higher infection rate among black individuals compared to white individuals. More importantly, a higher ratio of green spaces at the county level is significantly associated with a lower racial disparity in the SARS-CoV-2 infection rate. Further, we identified four green space factors that have significant negative associations with the racial disparity in SARS-CoV-2 infection rates, including open space in developed areas, forest, shrub and scrub, and grassland and herbaceous.

Keywords: racial disparity; health disparity; SARS-CoV-2; COVID-19; green spaces; pathway

Introduction

Racial disparity in health is a significant problem in many countries and can lead to social conflicts, economic crises, and loss of life[1, 2]. The black–white health disparity in the United States is a representative example in developed economies and is unsurprisingly manifest in the ongoing coronavirus disease 2019 (COVID-19) pandemic[3, 4, 5, 6]. COVID-19 results from infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and the SARS-CoV-2 infection rate in black individuals is significantly higher, proportionally, than that in white individuals[4, 7, 8]. Although studies of the COVID-19 pandemic are quickly accumulating, only a small portion have examined racial disparity in infection rates[9, 10, 11, 12]. Most of these studies focused on relationships between socioeconomic, demographic, or pre-existing chronic disease factors and racial disparity in SARS-CoV-2 infection rates [4, 13, 14]. None of the studies have directly examined effects of urban environmental factors. Evidence suggests that green spaces may have positive and independent effects on reducing the racial disparity in various long-term health outcomes [15, 16, 17, 18, 19, 20, 21]. Yet, to our knowledge, there is no evidence to support green space's effect on reducing racial disparity in SARS-CoV-2 infection rates. This lack of knowledge may mean missing opportunities to slow down the COVID-19 pandemic, and to moderate future epidemics by through urban greening. Our nationwide study is an initial ecological cross-sectional study aiming to establish a county-level relationship between black–white racial disparity in SARS-CoV-2 infection rates and the amount and type of green spaces in the United States.

A new trend: Significant environmental effects on racial disparity in health

Many studies have explored potential mechanisms that maintain and exacerbate general racial disparity in health outcomes to identify effective policies to address such inequities. Much of the racial disparity seems to be influenced by differences in socioeconomic and demographic factors[3, 4, 14]. Still, disparities remain significant after controlling for these factors. For example, the life expectancies of higher socioeconomic status (SES) blacks and whites are 7.1 and 6.8 years longer, respectively, than those of their lower SES counterparts[22, 23]. Life expectancy of blacks is at least three years less than that of whites with a comparable income level, at every level of income[23]. Racial disparity is also pronounced in birth outcomes; for example, the infant mortality rate for college-educated black women is 2.5 times higher than that for college-educated white women[23,

[24]. These studies suggest that persistent racial health disparities are due, in part, to other types of systemic disparities between races.

After controlling for SES factors, an under-studied reason for racial disparity in health outcomes is built environment disparity [25]. People of different races, especially blacks and whites in the USA, have lived in neighborhoods with distinct environmental qualities[26]. Neighborhoods that are the long-term home of more black people have, on average, fewer green spaces for physical activity, lower accessibility to healthy food, and lower traffic or crime-related safety[27, 28, 29, 30]. These findings have prompted studies in new directions. Researchers across disciplines have shifted from investigating exclusively social, economic, and educational interventions, and are now also investigating physical environmental interventions as a means to alleviate persistent racial-disparities[19, 29, 31].

Positive effects of green spaces on human health

The significantly positive effects that green spaces have on human health have drawn considerable attention from researchers, public health professionals, and governmental officers over the past decades. A growing number of studies have measured these effects at the regional[19, 32], municipal[33], neighborhood[34], and property-site[35] levels. Consensus has emerged that green spaces can influence human health through five major pathways[36], namely attention restoration[37], stress reduction[38, 39], enhanced social cohesion and social capital[40], increased physical activity[41], and supply of ecological products and services[42, 43, 44]. These pathways have been demonstrated across different geographic, social, and cultural contexts[45, 46, 47].

Green spaces reduce racial disparity in health: Circumstantial evidence and knowledge gaps

One line of research might be highly relevant to racial disparity in SARS-CoV-2 infection rates: green spaces can mitigate health disparities among populations with distinctively different SESs. A series of ground-breaking observational and survey studies conducted by Kuo and Sullivan (2001)[48, 49, 50], Sullivan et al. (2004)[51], and Taylor et al. (2002)[52] : They found that a greater amount of green space in public housing communities was significantly correlated with a higher level of self-discipline[52], a lower level of mental fatigue[50, 53], a higher level of social activity and cohesion[54], and a lower level of stress and aggressive behavior[55] in residents. Mitchell and Popham (2008)[19] published a nationwide survey-based study revealing that the association between income level

and all-cause mortality and circulatory disease mortality at the neighborhood level can be moderated by the amount of green space in a neighborhood. The study suggested that more exposure to green space can reduce the health disparity caused by income disparity. Lovasi et al. (2009)[21] found that the supply of public green spaces can reduce health disadvantages due to obesity and obesity-related illnesses in residents of low SES, by encouraging physical exercise. Branas et al. (2011, 2018)[56, 57] conducted citywide experimental studies, and found that the supply of green open space in low-income neighborhoods can significantly reduce crime, violence, stress, and fear, and increase daily exercise in those neighborhoods.

Many of these studies have examined the relationships between the supply of green spaces and health disparities in long-term health between populations with distinctively different SESs[58, 59]. While SES disparity is relevant to racial disparity, it is not fully representative of it[60]. Furthermore, the potential for exposure to green spaces to attenuate racial disparity in the risk of infection by airborne viruses, such as SARS-CoV-2, has not been investigated. This is a critical knowledge gap that may limit our opportunities to moderate the current pandemic and epidemics in the future.

Hypothesis

We hypothesized that the black–white racial disparity in the USA's SARS-CoV-2 infection rates is significantly less in urbanized counties with a higher ratio of various types of green spaces.

Methods

Study design

We compared the black–white disparity in SARS-CoV-2 infection rates in populations living in urbanized counties of the United States that have different amounts of green spaces, while adjusting for potential confounding factors of these counties. The infection data were retrieved on July 10, 2020. We adopted a within-subject (within-county) research design with a representative sample across the country. Black–white disparity was measured as the difference in the infection rate of blacks and whites in the same county. This allowed us to largely remove the bias caused by between-subject (between-county) factors that may lead to the uneven spread of SARS-CoV-2 in different counties, such as the distance to COVID-19 epicenters, climate, or local government policies. The study is cross-sectional in the sense that it does not analyze across time and ecological in the sense that it uses data aggregated at spatial unit of analysis (county).

Study areas

The United States is among the countries most severely impacted by COVID-19. It also has severe black–white racial inequalities in SARS-CoV-2 infection rates. Our study used counties as the basic unit of analysis, which are the fundamental administrative unit in the United States. There are a total of 3,142 counties or county-equivalent areas in the United States.

The most urbanized counties were chosen because we assumed that the racial inequality in the SARS-CoV-2 infection rates is more pronounced in denser urban environments. First, we identified a list of 314 large cities with a population ≥ 100,000 in 2019. Second, we identified a total of 229 counties containing or overlapping these large cities. Third, counties without infection data for blacks and whites were excluded. A total of 135 counties were chosen as our study areas. The total population in these counties was 132,350,027, which comprised 40.3% of the total population in the United States. We chose counties rather cities as the unit of analysis because the infection data for blacks and whites were often unavailable at the city level.

SARS-CoV-2 infection rates

The infection data were collected from the county health department portals on July 10, 2020. Only SARS-CoV-2 infections in whites and blacks were collected in this study. We calculated the SARS-CoV-2 infection rates for whites and blacks (numbers of cases per 100k) based on the total white and black populations in each county, respectively, which were retrieved from the 2019 census data[61]. The racial disparity in the SARS-CoV-2 infection rates was calculated as the difference between the infection rate in blacks and the infection rate in whites in the same county.

Green spaces

We used three datasets to assess green spaces, namely the National Land Cover Datasets in 2016 (NLCD 2016), the Tree Canopy Cover Datasets of the United States Forest Service (USFS), and NDVI.

The NLCD 2016 provides spatially explicit and reliable information on the national land-cover classification based on Landsat imagery at 30-meter resolution[62]. The overall agreement ranges from 71% to 97% between different land-cover classifications and reference data[62]. The NLCD 2016 has 16 types of land covers. We considered the following land-cover types with dominant natural elements: developed open space, deciduous forest, evergreen forest, mixed forest, shrub

and scrub, grassland and herbaceous, pasture and hay, cultivated crops, woody wetlands, and emergent herbaceous wetlands (Appendix Table 2). Deciduous/evergreen/mixed forest were combined into a new land-cover denoted "forest". The ratio of these green spaces over the total county area in each county was calculated.

The Tree Canopy Cover Dataset was derived from the NLCD by the USFS. It provides estimates of the percent tree canopy cover at 30-meter resolution. We aggregated the total tree-canopy cover estimates for each county and then divided it by the county area.

The NDVI is a widely used index that measures the quantity of green vegetation cover at the pixel level via remote sensing. We obtained the NDVI at a 30-meter resolution from the Google Earth Engine, which integrates Landsat 8 imagery in its cloud platform[63]. Four pixel-level average values were calculated for each county in each month from March to July in 2020, and we finally obtained the mean of four monthly NDVI values as a proxy for overall green spaces coverage during the research period.

However, we found that the land-cover dataset had high multicollinearity with two other measurements of green spaces. For instance, the NDVI and forest land-cover are strongly correlated (Pearson correlation coefficient: $r = 0.64$, $p < 0.001$), as are tree canopy and forest ($r = 0.85$, $p < 0.001$). Therefore, we only used measures from the land-cover dataset.

Confounding factors

Previous studies have shown that demographic and socioeconomic characteristics are important predictors of SARS-CoV-2 infection risk or racial disparity in this risk[1, 2, 64]. From the 2019 census data, we obtained the population density, the female population ratio, the difference in black–white population, the difference in black–white older adults, household size, housing value, the rate of high school graduates or higher, the rate of households with broadband, median household income, the poverty rate, healthcare receipts data, travel time to work, the employment rate, and the number of firms for our study areas (Appendix Table 1)[61, 65].

Pre-existing chronic disease is another indispensable dimension that might be connected with the racial disparity in SARS-CoV-2 infection rates[64]. Thus, county-level pre-existing chronic disease factors were collected from the 2016–2018 interactive heart disease and stroke data collated by the Centers for Disease Control and Prevention[66]. The pre-existing chronic disease

factors that we used were the coronary heart disease death rate, the heart failure death rate, the diagnosed diabetes rate, and the obesity rate for each county (Appendix Table 1).

All confounding factors were represented by proportional variables (e.g., per 100 k, density, ratio, and median value) or the difference between blacks and whites at the county level. Therefore, they were consistent with the dependent variable.

Statistical analysis

Three statistical analysis steps were performed. First, a paired *t*-test was used to examine whether there was a significant difference between SARS-CoV-2 infection rates in blacks and whites in the same county. Second, a variance inflation factor (VIF) test was used to remove potential multicollinearity among the independent variables. We adopted the threshold value of VIF = 4[67]. All of the factors with a VIF ≥ 4 were excluded from our models (Appendix Tables 1 & 2).

Third, hierarchical linear-regression models were used to examine the associations between the black–white difference in the SARS-CoV-2 infection rates and green space factors, while controlling for other confounding factors. The first model (Model 1) included the socioeconomic and demographic factors only. The second model (Model 2) additionally included pre-existing chronic disease factors. The last model (Model 3) additionally included green space factors, on the basis of Model 2.

All of the analyses were performed using R v4.0.2[68]. The model R^2 values, adjusted R^2 values, standardized coefficient (β) values, 95% confidence intervals, and *p*-values were reported.

Results

Results are presented in three sections. First, we present the incidence of SARS-CoV-2 infections in the sample of 135 most urbanized counties in the United States and examine the extent to which there are differences in infection rates among black and white populations. Second, we report demographic and green space characteristics of the sampled counties. Third, we employ hierarchical linear modeling to examine the extent to which, after controlling for confounding factors, the ratio of green spaces are associated with black-white disparities in SARS-CoV-2 infection rates. Finally, we report four green space factors that have significant negative associations with the racial disparity in SARS-CoV-2 infection rates.

Does a racial disparity in infection rates exist?

As of July 10, 2020, there were a total of 1,425,461 cases of SARS-CoV-2 infection in the 135 most urbanized counties in the United States, which accounted for 47% of the total cases of infection (3,038,325) in the United States (Figure 1). The 135 most urbanized counties were selected because they contain or overlap with 314 large cities with a population ≥ 100,000 in 2019 (see details in Methods). The county-level average infection rate for whites was 497 persons per 100,000 population, whereas the infection rate for blacks was approximately twice this (988 persons per 100,000 population) (Figure 1 & Table 1). The average black–white difference in the infection rate was 447 persons per 100,000 population. Whites had a higher infection rate than blacks in only 11 out of the 135 counties (Figure 2). As expected, a paired *t*-test revealed that the difference infection rates between blacks and whites is significant, with *t* (134) = 12.757 and *p* < 0.001 (Figure 3).

Figure 1. SARS-CoV-2 infection rates of blacks and whites in the 135 most urbanized counties of the United States. Pink column indicates infection rate for blacks; blue column indicates infection rate for whites. The height of columns indicates the magnitude of infection rate.

Figure 2. Racial disparity in the SARS-CoV-2 infection rates between blacks and whites in the 135 most urbanized counties of the United States. Cyan column indicates black infection rate is higher than white; purple column indicates white infection rate is higher than black. The height of columns indicates the magnitude of disparity.

Figure 3. The mean SARS-CoV-2 infection rate of blacks is approximately twice that of whites (988 vs. 497 per 100,000 population). Significant at $p < 0.001$ using a paired t-test. The error bar represents standard deviation.

Characteristics of the sampled counties

Table 1 shows that there are proportionally more whites than blacks in the study areas, which is consistent with the overall racial composition in the United States. The sample includes an average of 2.6 persons per household and a notably large range in median household income across counties. In terms of the green space factors, the sample includes a high quantity of developed open space and forest cover (Figure 4), and low levels of shrub and scrub, and grassland and herbaceous cover. Figure 4 shows the ratio of four types of green space in each county.

Table 1. Descriptive statistics for SARS-CoV-2 infection rates, socioeconomic and demographic factors, pre-existing chronic disease factors, and green space factors in the USA's 135 most urbanized counties.

Variable Categories	Variables	Min	Max	Mean	SD	Unit or Formula
Infection outcomes	Infection rate in blacks	91.6	2,514.8	987.9	583.7	Cases per 100k
	Infection rate in whites	29.1	2,486.2	496.8	336.2	Cases per 100k
	Difference in black–white infection rates	-367.9	1,874.2	491.1	447.3	Cases per 100k
Socioeconomic and demographic factors	Population density	27.8	19,625.8	1,003.2	2,356.6	Persons per square kilometer
	Female population ratio	0.479	0.533	0.511	0.010	Female/ total population
	Difference in black–white population	-0.878	0.481	-0.407	0.274	Black-to-white population ratio
	Difference in black–white older adults	-0.201	0.050	-0.078	0.041	Black older adults-to-white older adults population ratio
	Household size	2.1	3.3	2.6	0.3	Persons per household
	Households with broadband	0.666	0.917	0.818	0.056	Ratio
	Median household income	38,085	117,374	62,484	16,126	USD
	Healthcare receipts	222,956	2,129,326	771,836	310,816	1000 USD per 100k
	Number of firms	5,000	19,365	8,595	2,076	Number of firms per 100k
Pre-existing chronic disease factors	Coronary heart disease death rate	45.3	144.9	85.1	22.0	Cases per 100k
	Heart failure death rate	34.4	170.3	89.1	24.6	Cases per 100k
	Diagnosed diabetes rate	0.046	0.140	0.090	0.018	Cases/ total population
Green space factors	Developed open space	0.005	0.289	0.102	0.064	Developed open space area/ county area
	Forest	0.001	0.738	0.172	0.163	Forest area/ county area
	Shrub and scrub	0.000	0.829	0.082	0.177	Shrub and scrub area/ county area
	Grassland and herbaceous	0.001	0.498	0.061	0.093	Grassland and herbaceous area/ county area
	Pasture and hay	0.000	0.485	0.069	0.092	Pasture and hay area/ county area
	Cultivated crops	0.000	0.719	0.124	0.168	Cultivated crops area/ county area
	Woody wetlands	0.000	0.453	0.050	0.078	Woody wetlands area/ county area
	Emergent herbaceous wetlands	0.000	0.605	0.026	0.080	Emergent herbaceous wetlands area/ county area

Note: Min = minimum; Max = maximum; SD = standard deviation; Per 100k = 100,000 population; USD = United States Dollar.

Figure 4. Ratio of four types of green space to total county area, by county. a developed open space.
b forest. **c** shrub and scrub. **d** grassland and herbaceous.

Modeling association of green spaces and infection disparity

By fitting three hierarchical regression models, we identify three relevant associations (Table 2).
Model 1 shows that socioeconomic and demographic factors have a significant association with
racial disparity in the SARS-CoV-2 infection rate across the USA's 135 most urbanized counties
(adjusted R^2 of 0.12, $p = 0.003$). We also note that there is a significantly negative association
between household size and racial disparity in SARS-CoV-2 infection rates, such that as average
household size increases, infection disparity falls.

   After adding pre-existing chronic disease factors into Model 2, the explanatory power of Model
2 slightly decreases (adjusted $R^2 = 0.11$, $p = 0.007$). Model 2 shows that the combination of
socioeconomic, demographic, and pre-existing chronic disease factors is associated with racial
disparity. However, the overall statistical significance of Model 2 was not substantially better than
that of Model 1 ($p = 0.430$ for the sum-of-squares difference).

After adding green space factors into Model 3, the overall explanatory power of Model 3 increased by 18% (adjusted $R^2 = 0.29$, $p < 0.001$). Model 3 reveals that four green space factors are independently negatively associated with racial disparity. These are: land-cover ratio of developed open space; forest; shrub and scrub; and grassland and herbaceous. In addition, household size is negatively associated with SARS-CoV-2 infection rate disparity between races, while death from heart failure rate is positively associated with it.

Table 2. Three hierarchical linear regression results (N = 135).

Model Predictors	Model 1		Model 2		Model 3	
	β (95% CI)	p-value	β (95% CI)	p-value	β (95% CI)	p-value
Population density	-0.02 (-0.22, 0.18)	0.860	0.02 (-0.21, 0.24)	0.884	-0.14 (-0.38, 0.11)	0.279
Female population ratio	0.06 (-0.15, 0.27)	0.552	0.10 (-0.12, 0.31)	0.362	0.06 (-0.15, 0.28)	0.558
Difference in black–white population	-0.03 (-0.25, 0.19)	0.782	-0.03 (-0.25, 0.20)	0.812	0.03 (-0.20, 0.25)	0.801
Difference in black–white older adults	-0.12 (-0.31, 0.08)	0.232	-0.13 (-0.32, 0.07)	0.210	-0.17 (-0.36, 0.02)	0.079
Household size	-0.42 (-0.65, -0.19)	< 0.001***	-0.42 (-0.65, -0.18)	< 0.001***	-0.37 (-0.61, -0.14)	0.002**
Households with broadband	-0.15 (-0.42, 0.12)	0.276	-0.11 (-0.39, 0.18)	0.467	-0.07 (-0.34, 0.20)	0.611
Median household income	0.17 (-0.10, 0.44)	0.221	0.20 (-0.09, 0.49)	0.168	0.23 (-0.04, 0.50)	0.095
Healthcare receipts	-0.02 (-0.24, 0.20)	0.866	-0.04 (-0.26, 0.19)	0.751	-0.01 (-0.23, 0.22)	0.963
Number of firms	0.02 (-0.18, 0.21)	0.873	0.04 (-0.18, 0.25)	0.744	0.01 (-0.23, 0.25)	0.915
Coronary heart disease death rate			0.06 (-0.15, 0.27)	0.557	0.01 (-0.19, 0.21)	0.907
Heart failure death rate			0.14 (-0.06, 0.35)	0.175	0.26 (0.04, 0.48)	0.020*
Diagnosed diabetes rate			-0.04 (-0.29, 0.21)	0.760	-0.07 (-0.32, 0.18)	0.578
Developed open space					-0.31 (-0.56, -0.07)	0.011*
Forest					-0.31 (-0.53, -0.09)	0.006**
Shrub and scrub					-0.32 (-0.56, -0.07)	0.012*
Grassland and herbaceous					-0.42 (-0.62, -0.22)	< 0.001***
Pasture and hay					-0.12 (-0.30, 0.06)	0.180
Cultivated crops					0.01 (-0.23, 0.25)	0.947
Woody wetlands					0.01 (-0.18, 0.19)	0.963
Emergent herbaceous wetlands					-0.09 (-0.30, 0.13)	0.430
R^2/Adjusted R^2	0.178/0.119	0.003**	0.193/0.113	0.007**	0.294/0.288	< 0.001***
ANOVA F-statistic of R^2 change			0.927 (vs. Model 1)	0.430	4.739 (vs. Model 2)	< 0.001***

Note: * indicates $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$.

To improve our understanding of the associations between the four green spaces and racial disparity in SARS-CoV-2 infection rates, we plotted the independent effects of the four green spaces (using the effects package of the statistical software R[69]). Figure 5 clearly demonstrates

the negative relationships between the proportion of the four types of green space in counties and the racial disparity in SARS-CoV-2 infection rates.

Figure 5. Individual effects of the ratio of four green spaces on the black–white racial disparity in SARS-CoV-2 infection rates (in Model 3, with non-greenspace predictors fixed). a developed open space. **b** forest. **c** shrub and scrub. **d** grassland and herbaceous. Shaded areas represent the pointwise 95% confidence interval. Points represent partial residuals. Straight line represents the linear fitting of the effects. Dashed line represents the progressive fitting of the effects.

Discussion

Significant associations between socioeconomic, demographic, and pre-existing chronic disease factors and the racial disparity in SARS-CoV-2 infection rates have been reported by a number of ecological studies[3, 4, 5, 13, 14]. Some urban environmental factors, such as overcrowding housing conditions, living in senior living communities, living in high-density urban areas, and a long commute distance, are associated with a greater racial disparity in SARS-CoV-2 infection rates [70,

[71, 72, 73, 74]. However, no study examined the relationship between green spaces and racial disparity in SARS-CoV-2 infection rates.

In this study of the 135 urbanized counties in the United States, we found a large and statistically significant disparity in infection rates of the SARS-CoV-2 virus for black and white populations. Using hierarchical linear modelling, we found that the combination of socioeconomic, demographic, and pre-existing chronic disease conditions explained around 11% of variance in racial disparity. Adding green spaces explained an additional 18% of variance. Specifically, we identified four types of green space factors that are significantly associated with the racial disparity in SARS-CoV-2 infection: open space in developed areas, forest, shrub and scrub, and grassland and herbaceous.

In the paragraphs that follow, we consider possible mechanisms through which greater amounts of green spaces within a county might contribute to a reduction in racial disparity in SARS-CoV-2 infection rates. The plausible mechanisms are supported by theoretical and empirical evidence. We elaborate the contributions of our findings and identify questions for future research, We think that our study is the first to find that green space factors have significant independent effects on the racial disparity in COVID-19 infection rates, albeit using an ecological study design.

Interpretation of key findings

Variations in county-level open space in developed areas, forest, shrub and scrub, and grassland and herbaceous were all significantly negatively correlated with the size of discrepancy in black-white infection rates. In urban counties with more green spaces available, the racial disparity in SARS-CoV-2 infection rates was lower.

We propose that four mechanisms may account for this pattern. Green spaces deliver the following benefits: facilitate physical activity[33, 51, 53]; promote mental and social health[4748]; encourage social activities at a safe physical distance[75]; and improve air quality at the neighborhood and city levels[76]. All of these may have an influence on reducing racial disparities in infection rates.

First, areas with a large portion of green spaces provided free access to residents of all racial population and promoted physical activity before and during the pandemic[77, 78]. This may have enhanced residents' immune system regardless of race and moderated factors behind race-based infection disparities[79, 80, 81]. Physical activity conducted in nature provides additional benefits

compared with physical activity conducted in indoor environment[82, 83, 84]. As we know, black people tend to living in communities with fewer green spaces for physical activity[27, 28, 29, 30]. Therefore, black people who can access open green spaces may accrue health benefits from conducting physical activity in these green spaces, and therefore potentially have lower infection rate than black people who cannot access open green spaces.

Second, the presence of more green spaces may have enhanced mental and social health regardless race before and during the pandemic[55], providing multiple benefits that moderate risks that otherwise fall more heavily on black communities. It is feasible that disadvantaged groups and races may have heavier stress burdens, especially during the pandemic[18, 19, 20, 21]. Visual or physical contact with urban green spaces can reduce mental fatigue[53, 85], reduce mental stress[35, 86, 87, 88], and enhance self-discipline[52] at the individual level, which can promote immune system function and self-control behaviors in daily life. It can also reduce negative moods[89, 90] and verbal and behavioral aggressiveness[50], which can enhance trust[54, 91], collaboration[92]. This could promote a stronger collective stewardship of the living environment and previous studies found a stronger environmental stewardship can reduce environment-health disparities[16, 93].

Third, more green spaces can decrease the SARS-CoV-2 infection risk by improving air quality and decreasing exposure to air pollutants (e.g., $PM_{2.5}$) that are associated with higher SARS-CoV-2 infection rates[94, 95]. Black households tend to live in higher density, more polluted streets, compared with white households[27, 28, 29, 30]. The supply of green spaces would be of greater relative benefits to black populations than white populations. Studies have reported that air quality in high-density residential areas in inner cities is significantly worse than that in suburban and rural residential areas[96]. A greater supply of green spaces, especially in urban areas, may reduce this significant environmental disparity[17, 19], thereby contributing to a lower racial disparity in SARS-CoV-2 infection rates. We note, however, that relying on urban green spaces is not sufficient to create healthy air environments[97] for the whole urban area thus provision of large-scale green spaces beyond the urban area tis critical to achieving this goal[76].

Fourth, more green spaces can lower the risk of infection by providing all racial populations with ample areas for social activities, while allowing safe physical distance to be maintained[75]. Staying with the idea of unequal effects of green space increments, for economic and social reasons, black households in the USA have lower average levels of ownership of private open green space than whites, and live in more crowded neighborhoods[25]. Black populations are more likely than

white populations in the USA to socialize in crowded indoor spaces. Therefore, black people with access to open green spaces are more likely to socialize in outdoor spaces, than black people without access to such green spaces.

In sum, greater amounts of green spaces will have a stronger health effect on black households compared to white. This, we suggest, is a crucial mechanism explaining our regression findings. Other things being equal, a rise in the quantity or accessibility of green spaces in an area might be expected to have an equal effect on black and white[98, 99]. However, black households are less likely to access green spaces than white households. The marginal health (and other) benefits to black households resulting from more green spaces, will be greater than for whites.

Contributions and implications

This study makes three theoretical and practical contributions, which will influence future research, policymaking, and design practice.

First, to the best of our knowledge, this study is the first to measure whether and to what extent green spaces within and beyond developed urban areas are associated with racial disparities in rates of contagious disease infection. Although a few studies have identified some built environment factors, such as crowded living conditions, staying in senior living communities, and dense urban areas, are related to racial disparity in SARS-CoV-2 infection rates[70, 71, 73, 74], none have addressed green spaces.

Second, we adopted a within-subject research design with a standard spatial sampling unit across the country, giving a representative study for USA urban counties. In contrast with previous studies on COVID-19 that examined associations between SARS-CoV-2 infection rate or COVID-19 mortality rate across counties or cities[10, 100, 101], our study focused on comparing difference in SARS-CoV-2 infection rates among whites and blacks in the same county. By comparing racial disparity in infection rates within each county we obtain greater statistical validity, as this approach mitigates bias caused by uneven spread of infections across counties due to differences in national road network accessibility, airports and railway connectivity, governmental regulations, social norms, and quantity and quality of healthcare services.

Third, our study suggests that green space as a type of urban infrastructure[102] should be considered to be relevant interventions to reduce the racial disparity of infectious diseases with characteristics similar to the current pandemic. Providing an adequate supply of accessible and

well-designed green spaces in urban areas, and preserving and developing natural green spaces across county areas, is part of an epidemic and pandemic-resilience strategy for highly urbanized areas. Considering that urban and agricultural areas are rapidly encroaching on forest, grassland, and many other natural landscapes inside and/or outside cities worldwide[103, 104], it is crucial to maintain and increase efforts to preserve scarce urban green spaces. As well as achieving environmental goals, we show that this approach can also be expected to help deliver health and racial equality agenda.

Limitations and opportunities for future research

This study has some limitations, that point to opportunities for future research.

We collected SARS-CoV-2 infection data from the governmental website of each county in the United States. The quality of data is credible for each county, but the methods of collecting and sorting data may have varied across counties. Moreover, there are a total 229 counties containing or overlapping all cities with a population $\geq$ 100,000 in the United States. However, infection data were unavailable for 94 counties, leaving a set of high-quality data from 135 counties.

This study only examined the problem in the US at the county level. Although the racial disparity in SARS-CoV-2 infection rates in the United States might be a good example for countries with diverse racial populations[6, 9, 105], we suggest that this research should be duplicated in other countries, to fully determine the complex issue of generally disadvantaged sub-groups in relation to environmental exposure and pandemic dynamics.

This study provides a base-line analysis of green spaces and racial disparity in pandemic infections. It is limited by the normal constraints of a cross-sectional design and of an ecological design. Cross-sectional associations do not imply causality and we cannot avoid the possibility of ecological fallacy. There are possibilities that can be investigated in further studies. Nevertheless, findings of this study echo findings of many published studies that reported causal links between more green spaces and better health status and lower health disparities.

Lastly, a caveat can be raised with respect to possible ecological bias: The access to data in SARS-CoV-2 infection rates at the neighborhood scale was not available, which made us cannot analyze the distribution of green spaces for races at the neighborhood scale. It is possible an unequal allocation of green spaces could lead to races' unequal exposures to green spaces, which need to be further examined in a future study at a finer spatial scale. Two arguments remain

reasonable albeit the possibility of that bias: More green spaces lead to greater marginal health benefits to black households and then a smaller racial disparity in SARS-CoV-2 infection rates; Large patches of natural green spaces outside the city, such as forest and natural grassland and herbaceous , can create adjusting environmental quality for the whole urban area thus can reduce racial disparity in SARS-CoV-2 infection rates. Nevertheless, we call future studies should further examine the problem and proposed arguments.

Conclusion

This study is an initial effort to understand the relationships between environmental factors and racial disparity in SARS-CoV-2 infection rates. We employed hierarchical regression analysis to perform within-county comparisons of the infection rates of the two major racial groups. After controlling for potential confounding factors, we found that four green space factors have independent statistical power to interpret the racial disparity. The findings suggest that the supply of open green space in developed urban areas and natural green spaces across a county may help to reduce racial disparity in SARS-CoV-2 infection rates. The findings in this study tentatively support the potential for exposure to green spaces to attenuate racial disparity.[58]

Author Contributions

BJ proposed the research concept. BJ and YL developed the concept into a full research plan. LC, XML, YWY, WYX conducted the data collection and data analysis under YL's and BJ's supervision. YL, BJ and YWY conducted writing of the introduction, discussion, and conclusion. YL, LC, XML, YWY, WYX conducted writing of the methods and results. CW and WCS provided critical revisions for the introduction and discussion.

References

1. Der Ananian C, Winham DM, Thompson SV, Tisue ME. Perceptions of heart-healthy behaviors among African American Adults: A mixed methods study. *International journal of environmental research and public health* **15**, 2433 (2018).

2. Thomas SB*, et al.* Racial differences in the association between self-rated health status and objective clinical measures among participants in the BARI 2D trial. *American journal of public health* **100**, S269-S276 (2010).

3. Holmes L*, et al.* Black–White risk differentials in COVID-19 (SARS-COV2) transmission, mortality and case fatality in the United States: translational epidemiologic perspective and challenges. *International journal of environmental research and public health* **17**, 4322 (2020).

4. Figueroa JF, Wadhera RK, Lee D, Yeh RW, Sommers BD. Community-Level Factors Associated With Racial And Ethnic Disparities In COVID-19 Rates In Massachusetts: Study examines community-level factors associated with racial and ethnic disparities in COVID-19 rates in Massachusetts. *Health Affairs*, 10.1377/hlthaff. 2020.01040 (2020).

5. Gu T*, et al.* COVID-19 outcomes, risk factors and associations by race: a comprehensive analysis using electronic health records data in Michigan Medicine. *medRxiv*, (2020).

6. Kirby T. Evidence mounts on the disproportionate effect of COVID-19 on ethnic minorities. *The Lancet Respiratory Medicine* **8**, 547-548 (2020).

7. Murphy T*, et al.* State of Black America Unmasked.(National Urban League, accessed 16 Oct 2020); http://sobadev.iamempowered.com/sites/soba.iamempowered.com/files/NUL-SOBA-2020-ES-web.pdf.

8. Wrigley-Field E. US racial inequality may be as deadly as COVID-19. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 21854-21856 (2020).

9. Kullar R*, et al.* Racial disparity of coronavirus disease 2019 in African American Communities. *The Journal of infectious diseases* **222**, 890-893 (2020).

10. Abedi V*, et al.* Racial, economic, and health inequality and COVID-19 infection in the United States. *Journal of racial and ethnic health disparities*, 1-11 (2020).

11. Joseph NP*, et al.* Racial/ethnic disparities in disease severity on admission chest radiographs among patients admitted with confirmed COVID-19: a retrospective cohort study. *Radiology*, 202602 (2020).

12. Hooper MW, Nápoles AM, Pérez-Stable EJ. COVID-19 and racial/ethnic disparities. *Jama*, (2020).

13. Townsend MJ, Kyle TK, Stanford FC. Outcomes of COVID-19: disparities in obesity and by ethnicity/race.). Nature Publishing Group (2020).

14. Sarmadi M, Moghanddam VK, Dickerson AS, Martelletti L. Association of COVID-19 distribution with air quality, sociodemographic factors, and comorbidities: an ecological study of US states. *Air Quality, Atmosphere & Health*, 1-11 (2020).

15. Jennings V, Gaither CJ. Approaching environmental health disparities and green spaces: an ecosystem services perspective. *International journal of environmental research and public health* **12**, 1952-1968 (2015).

16. Jennings V, Johnson Gaither C, Gragg RS. Promoting environmental justice through urban green space access: A synopsis. *Environmental Justice* **5**, 1-7 (2012).

17. Wolch JR, Byrne J, Newell JP. Urban green space, public health, and environmental justice: The challenge of making cities 'just green enough'. *Landscape and urban planning* **125**, 234-244 (2014).

18. Payne-Sturges D, Gee GC. National environmental health measures for minority and low-income populations: tracking social disparities in environmental health. *Environmental Research* **102**, 154-171 (2006).

19. Mitchell R, Popham F. Effect of exposure to natural environment on health inequalities: an observational population study. *The Lancet* **372**, 1655-1660 (2008).

20. Dadvand P, *et al.* Green spaces and general health: roles of mental health status, social support, and physical activity. *Environment international* **91**, 161-167 (2016).

21. Lovasi GS, Hutson MA, Guerra M, Neckerman KM. Built environments and obesity in disadvantaged populations. *Epidemiologic reviews* **31**, 7-20 (2009).

22. Bucholz EM, Ma S, Normand S-LT, Krumholz HM. Race, socioeconomic status, and life expectancy after acute myocardial infarction. *Circulation* **132**, 1338-1346 (2015).

23. Braveman PA, Cubbin C, Egerter S, Williams DR, Pamuk E. Socioeconomic disparities in health in the United States: what the patterns tell us. *American journal of public health* **100**, S186-S196 (2010).

24. Singh GK, Stella MY. Infant mortality in the United States, 1915-2017: large social inequalities have persisted for over a century. *International Journal of Maternal and Child Health and AIDS* **8**, 19 (2019).

25. Popescu I, Duffy E, Mendelsohn J, Escarce JJ. Racial residential segregation, socioeconomic disparities, and the White-Black survival gap. *PLoS One* **13**, e0193222 (2018).

26. Massey DS. Residential segregation and neighborhood conditions in US metropolitan areas. *America becoming: Racial trends and their consequences* **1**, 391-434 (2001).

27. Wen M, Zhang X, Harris CD, Holt JB, Croft JB. Spatial disparities in the distribution of parks and green spaces in the USA. *Annals of Behavioral Medicine* **45**, S18-S27 (2013).

28. Phelan JC, Link BG. Is Racism a Fundamental Cause of Inequalities in Health? In: *Annual Review of Sociology*) (2015).

29. Williams DR, Sternthal M. Understanding Racial-ethnic Disparities in Health: Sociological Contributions. *J Health Soc Behav* **51**, S15-S27 (2010).

30. Williams DR. Miles to Go before We Sleep: Racial Inequities in Health. *J Health Soc Behav* **53**, 279-295 (2012).

31. Black JL, Macinko J. Neighborhoods and obesity. *Nutrition Reviews* **66**, 2-20 (2008).

32. Mitchell R, Popham F. Greenspace, urbanity and health: relationships in England. *J Epidemiol Community Health* **61**, 681-683 (2007).

33. Lu Y. Using Google Street View to investigate the association between street greenery and physical activity. *Landscape and Urban Planning*, (2018).

34. Lee M, Kim S, Ha M. Community greenness and neurobehavioral health in children and adolescents. *Science of The Total Environment* **672**, 381-388 (2019).

35. Jiang B, Chang C-Y, Sullivan WC. A dose of nature: Tree cover, stress reduction, and gender differences. *Landscape and Urban Planning* **132**, 26-36 (2014).

36. Jiang B, Zhang T, Sullivan WC. Healthy Cities: Mechanisms and Research Questions Regarding the Impacts of Urban Green Landscapes on Public Health and Well-being. *Landsc Archit Front* **3**, 24-35 (2015).

37. Kaplan S. The Restorative Benefits of Nature - toward an Integrative Framework. *Journal of Environmental Psychology* **15**, 169-182 (1995).

38. Ulrich R. View through a window may influence recovery from surgery. *Science* **224**, (1984).

39. Ulrich RS. NATURAL VERSUS URBAN SCENES - SOME PSYCHOPHYSIOLOGICAL EFFECTS. *Environ Behav* **13**, 523-556 (1981).

40. Holtan MT, Dieterlen SL, Sullivan WC. Social life under cover: tree canopy and scoia capital in Baltimore, Maryland. *Environment & Behavior* **46**, 1-24 (2014).

41. Cohen DA, McKenzie TL, Sehgal A, Williamson S, Golinelli D, Lurie N. Contribution of public parks to physical activity. *Am J Public Health* **97**, 509-514 (2007).

42. Bolund P, Hunhammar S. Ecosystem services in urban areas. *Ecol Econ* **29**, 293-301 (1999).

43. Yu K. Security patterns and surface model in landscape ecological planning. *Landscape and Urban Planning* **36**, 1-17 (1996).

44. Coutts C, Hahn M. Green Infrastructure, Ecosystem Services, and Human Health. *International Journal of Environmental Research and Public Health* **12**, 9768-9798 (2015).

45. Kuo M. How might contact with nature promote human health? Promising mechanisms and a possible central pathway. *Frontiers in Psychology* **6**, (2015).

46. Hartig T, Mitchell R, de Vries S, Frumkin H. Nature and Health. *Annual Review of Public Health* **35**, 207-228 (2014).

47. Bratman GN, Hamilton JP, Daily GC. The impacts of nature experience on human cognitive function and mental health. *Year in Ecology and Conservation Biology* **1249**, 118-136 (2012).

48. Kuo F, Sullivan CW. Environment and crime in the inner city: does vegetation reduce crime? *Environ Behav* **33**, 343-367 (2001).

49. Kuo FE. Coping with poverty - Impacts of environment and attention in the inner city. *Environ Behav* **33**, 5-34 (2001).

50.   Kuo FE, Sullivan WC. Aggression and violence in the inner city - Effects of environment via mental fatigue. *Environment and Behavior* **33**, 543-571 (2001).

51.   Sullivan WC, Kuo FE, DePooter SF. The fruit of urban nature - Vital neighborhood spaces. *Environment and Behavior* **36**, 678-700 (2004).

52.   Taylor AF, Kuo FE, Sullivan WC. Views of nature and self-discipline: Evidence from inner city children. *Journal of environmental psychology* **22**, 49-63 (2002).

53.   Jiang B, Schmillen R, Sullivan WC. How to waste a break: Using portable electronic devices substantially counteracts attention enhancement effects of green spaces. *Environment and Behavior* **51**, 1133-1160 (2019).

54.   Arnberger A, Eder R. The influence of green space on community attachment of urban and suburban residents. *Urban For Urban Green* **11**, 41-49 (2012).

55.   Beyer KM, Kaltenbach A, Szabo A, Bogar S, Nieto FJ, Malecki KM. Exposure to neighborhood green space and mental health: evidence from the survey of the health of Wisconsin. *International journal of environmental research and public health* **11**, 3453-3472 (2014).

56.   Branas CC*, et al.* Citywide cluster randomized trial to restore blighted vacant land and its effects on violence, crime, and fear. *Proceedings of the National Academy of Sciences*, (2018).

57.   Branas CC, Cheney RA, MacDonald JM, Tam VW, Jackson TD, Ten Have TR. A Difference-in-Differences Analysis of Health, Safety, and Greening Vacant Urban Space. *American Journal of Epidemiology* **174**, 1296-1306 (2011).

58.   Browning MH, Rigolon A. Do income, race and ethnicity, and sprawl influence the greenspace-human health link in city-level analyses? Findings from 496 cities in the United States. *International journal of environmental research and public health* **15**, 1541 (2018).

59.   Braubach M, Egorov A, Mudu P, Wolf T, Ward Thompson C, Martuzzi M. Effects of Urban Green Space on Environmental Health, Equity and Resilience.). Springer International Publishing (2017).

60.   Williams DR, Priest N, Anderson NB. Understanding associations among race, socioeconomic status, and health: Patterns and prospects. *Health Psychology* **35**, 407 (2016).

61.   United States Census Bureau. Annual County Resident Population Estimates by Age, Sex, Race, and Hispanic Origin: April 1, 2010 to July 1, 2019 (CC-EST2019-ALLDATA). (United States Census Bureau, accessed 16 Oct 2020); https://www.census.gov/data/tables/time-series/demo/popest/2010s-counties-detail.html.

62.   Yang L*, et al.* A new generation of the United States National Land Cover Database: Requirements, research priorities, design, and implementation strategies. *ISPRS Journal of Photogrammetry and Remote Sensing* **146**, 108-123 (2018).

63.   Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment* **202**, 18-27 (2017).

64.   Centers for Disease Control and Prevention. People with Certain Medical Conditions. (United States Census Bureau, accessed 16 Oct 2020); https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html.

65. United States Census Bureau. U.S. Census Bureau Releases 2014-2018 ACS 5-Year Estimates. (United States Census Bureau, accessed 16 Oct 2020); https://www.census.gov/programs-surveys/acs/news/updates/2019.html.

66. Centers for Disease Control and Prevention. Interactive Atlas of Heart Disease and Stroke Tables. (United States Census Bureau, accessed 16 Oct 2020); https://nccd.cdc.gov/DHDSPAtlas/Reports.aspx.

67. O'brien RM. A Caution Regarding Rules of Thumb for Variance Inflation Factors. *Quality & Quantity* **41**, 673-690 (2007).

68. Team RC. R: A language and environment for statistical computing. (Vienna, Austria: R Foundation for Statistical Computing, accessed 16 Oct 2020); http://www.R-project.org

69. Fox J, Weisberg S. An R Companion to Applied Regression, 3rd edition. Sage, Thousand Oaks CA. (2019).

70. Emeruwa UN, *et al.* Associations between built environment, neighborhood socioeconomic status, and SARS-CoV-2 infection among pregnant women in New York City. *JAMA* **324**, 390-392 (2020).

71. Megahed NA, Ghoneim EM. Antivirus-built environment: Lessons learned from Covid-19 pandemic. *Sustainable Cities and Society* **61**, 102350 (2020).

72. Jin X, *et al.* Neighborhood-Level Public Facilities and COVID-19 Transmission: A Nationwide Geospatial Study In China. *medRxiv* doi: https://doi.org/10.1101/2020.08.25.20181362 (2020). Preprint.

73. Nguyen QC, *et al.* Using 164 Million Google Street View Images to Derive Built Environment Predictors of COVID-19 Cases. *International journal of environmental research and public health* **17**, 6359 (2020).

74. Rozenfeld Y, *et al.* A model of disparities: risk factors associated with COVID-19 infection. *International journal for equity in health* **19**, 1-10 (2020).

75. Venter Z, Barton D, Gundersen V, Figari H, Nowell M. Urban nature in a time of crisis: recreational use of green space increases during the COVID-19 outbreak in Oslo, Norway. *Environmental Research Letters*, **15**, 10 (2020).

76. Doughty MR, Hammond GP. Sustainability and the built environment at and beyond the city scale. *Building and environment* **39**, 1223-1233 (2004).

77. Richardson E, Pearce J, Mitchell R, Day P, Kingham S. The association between green space and cause-specific mortality in urban New Zealand: an ecological analysis of green space utility. *BMC public health* **10**, 240 (2010).

78. Pearce JR, Maddison R. Do enhancements to the urban built environment improve physical activity levels among socially disadvantaged populations? *International journal for equity in health* **10**, 1-9 (2011).

79. Shi Y, *et al.* COVID-19 infection: the perspectives on immune responses. *Cell Death & Differentiation,* **27**, 1451–1454 (2020).

80. Rook GA. Regulation of the immune system by biodiversity from the natural environment: an ecosystem service essential to health. *Proceedings of the National Academy of Sciences* **110**, 18360-18367 (2013).

81.   Rook GA, Raison CL, Lowry CA. Microbial 'old friends', immunoregulation and socioeconomic status. *Clinical & Experimental Immunology* **177**, 1-12 (2014).

82.   Li Q. Effect of forest bathing trips on human immune function. *Environmental health and preventive medicine* **15**, 9-17 (2010).

83.   Li Q*, et al.* Forest bathing enhances human natural killer activity and expression of anti-cancer proteins. *International journal of immunopathology and pharmacology* **20**, 3-8 (2007).

84.   Lyu B*, et al.* Benefits of a three-day bamboo forest therapy session on the psychophysiology and immune system responses of male college students. *International Journal of Environmental Research and Public Health* **16**, 4991 (2019).

85.   Ohly H*, et al.* Attention Restoration Theory: A systematic review of the attention restoration potential of exposure to natural environments. *Journal of Toxicology and Environmental Health, Part B* **19**, 305-343 (2016).

86.   Ulrich RS, Simons RF, Losito BD, Fiorito E, Miles MA, Zelson M. Stress recovery during exposure to natural and urban environments. *Journal of environmental psychology* **11**, 201-230 (1991).

87.   Norwood MF, Lakhani A, Maujean A, Zeeman H, Creux O, Kendall E. Brain activity, underlying mood and the environment: A systematic review. *Journal of Environmental Psychology* **65**, 101321 (2019).

88.   Jiang X, Larsen L, Sullivan W. Connections-between Daily Greenness Exposure and Health Outcomes. *Int J Environ Res Public Health* **17**, (2020).

89.   Brooks AM, Ottley KM, Arbuthnott KD, Sevigny P. Nature-related mood effects: Season and type of nature contact. *Journal of Environmental Psychology* **54**, 91-102 (2017).

90.   Nisbet EK, Zelenski JM, Grandpierre Z. Mindfulness in nature enhances connectedness and mood. *Ecopsychology* **11**, 81-91 (2019).

91.   Jennings V, Bamkole O. The relationship between social cohesion and urban green space: An avenue for health promotion. *International journal of environmental research and public health* **16**, 452 (2019).

92.   Zoellner J, Zanko A, Price B, Bonner J, Hill JL. Exploring community gardens in a health disparate population: findings from a mixed methods pilot study. *Progress in community health partnerships: research, education, and action* **6**, 153-165 (2012).

93.   McOliver CA*, et al.* Community-based research as a mechanism to reduce environmental health disparities in american Indian and alaska native communities. *Int J Environ Res Public Health* **12**, 4076-4100 (2015).

94.   Wu X, Nethery RC, Sabath BM, Braun D, Dominici F. Exposure to air pollution and COVID-19 mortality in the United States. *medRxiv*, https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7277007/ (2020). Preprint.

95.   Yongjian Z, Jingu X, Fengming H, Liqing C. Association between short-term exposure to air pollution and COVID-19 infection: Evidence from China. *Science of the total environment*, 138704 (2020).

96.   Strosnider H, Kennedy C, Monti M, Yip F. Rural and Urban Differences in Air Quality, 2008–2012, and Community Drinking Water Quality, 2010–2015 — United States. *MMWR Surveill Summ* **66**, 1-10 (2017).

97.   Richardson EA, Mitchell R, Hartig T, De Vries S, Astell-Burt T, Frumkin H. Green cities and health: a question of scale? *J Epidemiol Community Health* **66**, 160-165 (2012).

98.   Webster C. Pricing accessibility: Urban morphology, design and missing markets. *Prog Plann* **73**, 77-111 (2010).

99.   Woo Y, Webster C. A Co-evolution of gated communities and local public goods. *Urban Stud* **51**, 2539-2554 (2014).

100.   Pan W, Miyazaki Y, Tsumura H, Miyazaki E, Yang W. Identification of county-level health factors associated with COVID-19 mortality in the United States. *The Journal of Biomedical Research* **1**, 1-9 (2020).

101.   Vahidy FS*, et al.* Racial and Ethnic Disparities in SARS-CoV-2 Pandemic: Analysis of a COVID-19 Observational Registry for a Diverse US Metropolitan Population. *medRxiv*, https://www.medrxiv.org/content/10.1101/2020.04.24.20073148v2 (2020). Preprint.

102.   Suppakittpaisarn P, Jiang X, Sullivan WC. Green Infrastructure, Green Stormwater Infrastructure, and Human Health: A Review. *Current Landscape Ecology Reports* **2**, 96-110 (2017).

103.   Miller MD. The impacts of Atlanta's urban sprawl on forest cover and fragmentation. *Applied geography (Sevenoaks)* **34**, 171-179 (2012).

104.   Hung K-LJ, Ascher JS, Holway DA. Urbanization-induced habitat fragmentation erodes multiple components of temporal diversity in a Southern California native bee assemblage. *PLoS One* **12**, e0184136-e0184136 (2017).

105.   Witteveen D. Sociodemographic inequality in exposure to COVID-19-induced economic hardship in the United Kingdom. *Research in Social Stratification and Mobility* **69**, 100551 (2020).

Appendix

Table 1. The definition for socioeconomic and demographic factors, and pre-existing chronic disease factors.

Variables	Item description (formula)	Official definition	VIF Test
Population density	Total population / county area	Population: 2019 county-level census population	Retained
Female population ratio	Female persons, percent	2019 county-level census female percent	Retained
Difference in black-white population	Black population / total population - white population / total population	Black: 2019 county-level census black population White: 2019 county-level census white population	Retained
Difference in black-white older adults	Older adults of black / black population - older adults of white / white population	Older adults: Age above 65 of the 2019 age group population statistics	Retained
Household size	Persons per household, 2014-2018	Persons per household, or average household size, is obtained by dividing the number of persons in households by the number of households (or householders).	Retained
Housing value	Median value of owner-occupied housing units, 2014-2018	Specified owner-occupied housing units--one-family houses on less than 10 acres without a business or medical office on the property.	Removed
Rate of high school graduate or higher	High school graduate or higher, percent of persons age 25 years+, 2014-2018	High school graduates include people whose highest degree was a high school diploma or its equivalent. These data include only persons 25 years old and over.	Removed
Rate of households with broadband	Households with a broadband Internet subscription, percent, 2014-2018	Households with a broadband Internet subscription：The Internet question asked if any member of the household accesses the Internet.	Retained
Median household income	Median household income (in 2018 dollars), 2014-2018	Income of households: This includes the income of the householder and all other individuals 15 years old and over in the household, whether they are related to the householder or not.	Retained
Poverty rate	Persons in poverty, percent (in 2019 census)	Census Bureau uses a set of money income thresholds that vary by family size and composition to determine who is in poverty. If a family's total income is less than the family's threshold, then that family and every individual in it is considered in poverty.	Removed
Healthcare receipts	Total health care and social assistance receipts/revenue, 2012 ($1,000) / per 100k	The sector includes both health care and social assistance because it is sometimes difficult to distinguish between the boundaries of these two activities.	Retained
Travel time to work	Mean travel time to work (minutes), workers age 16 years+, 2014-2018	Travel time to work refers to the total number of minutes that it usually took the person to get from home to work each day during the reference week.	Removed

Employment rate	Total employment, 2018 / per 100k	Paid employment consists of full- and part-time employees, including salaried officers and executives of corporations. Included are employees on paid sick leave, holidays, and vacations; not included are proprietors and partners of unincorporated businesses.	Removed
Number of firms	All firms, 2012 / per 100k	Firms equally male-/female-owned, equally minority-/nonminority-owned, and equally veteran-/nonveteran-owned are counted and tabulated as separate categories.	Retained
Coronary heart disease death rate	Coronary heart disease death rate per 100,000	All ages, all races/ethnicities, both genders, 2016-2018 (not spatial smoothed data)	Retained
Heart failure death rate	Heart failure death rate per 100,000	All ages, all races/ethnicities, both genders, 2016-2018 (not spatial smoothed data)	Retained
Diagnosed diabetes rate	Diagnosed diabetes percentage	Age-adjusted percentage, 20+, 2016	Retained
Obesity rate	Obesity percentage	Age-adjusted percentage, 20+. 2016	Removed

Note. The detail definition is from the United States Census and Heart disease and stroke data of Centers for Disease Control and Prevention

Table 2: Official definition of greenspace factors.

Classes	Official definition	VIF Test
Developed open space	Areas with a mixture of some constructed materials, but mostly vegetation in the form of lawn grasses. Impervious surfaces account for less than 20% of total cover. These areas most commonly include large-lot single-family housing units, parks, golf courses, and vegetation planted in developed settings for recreation, erosion control, or aesthetic purposes.	Retained
Deciduous forest	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. More than 75% of the tree species shed foliage simultaneously in response to seasonal change.	
Evergreen forest	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. More than 75% of the tree species maintain their leaves all year. Canopy is never without green foliage.	
Mixed forest	Areas dominated by trees generally greater than 5 meters tall, and greater than 20% of total vegetation cover. Neither deciduous nor evergreen species are greater than 75% of total tree cover.	
Forest	Combination of deciduous, evergreen, and mixed forest.	Retained
Shrub and scrub	Areas dominated by shrubs; less than 5 meters tall with shrub canopy typically greater than 20% of total vegetation. This class includes true shrubs, young trees in an early successional stage or trees stunted from environmental conditions.	Retained
Grassland and herbaceous	Areas dominated by gramanoid or herbaceous vegetation, generally greater than 80% of total vegetation. These areas are not subject to intensive management such as tilling, but can be utilized for grazing.	Retained
Pasture and hay	Areas of grasses, legumes, or grass-legume mixtures planted for livestock grazing or the production of seed or hay crops, typically on a perennial cycle. Pasture/hay vegetation accounts for greater than 20% of total vegetation.	Retained
Cultivated crops	Areas used for the production of annual crops, such as corn, soybeans, vegetables, tobacco, and cotton, and also perennial woody crops such as orchards and vineyards. Crop vegetation accounts for greater than 20% of total vegetation. This class also includes all land being actively tilled.	Retained
Woody wetlands	Areas where forest or shrubland vegetation accounts for greater than 20% of vegetative cover and the soil or substrate is periodically saturated with or covered with water.	Retained
Emergent herbaceous wetlands	Areas where perennial herbaceous vegetation accounts for greater than 80% of vegetative cover and the soil or substrate is periodically saturated with or covered with water.	Retained

Note: The definition of land classes is derived from The Multi-Resolution Land Characteristics (MRLC) consortium