The effect of body image dissatisfaction on goal-directed decision making in a population marked by negative appearance beliefs and disordered eating

Jakub Onysk[1][¶], Peggy Seriès[2][&][*]

[1] Computational and Biological Learning Laboratory, Department of Engineering, University of

Cambridge, Cambridge, UK

[2] Institute for Adaptive and Neural Computation, School of Informatics, University of Edinburgh,

Edinburgh, UK

* Corresponding author

E-mail: pseries@inf.ed.ac.uk (PS)

[¶] Writing (original draft), visualisation, investigation, formal analysis, methodology, code.

[&] Conceptualisation, Supervision, Review and editing, project administration.

Abstract

Eating disorders are associated with one of the highest mortality rates among all mental

disorders, yet there is very little research about them within the newly emerging and promising

field of computational psychiatry. As such, we focus on investigating a previously unexplored,

yet a core aspect of eating disorders – body image dissatisfaction. We continue a freshly opened

debate about model-based learning and its trade-off against model-free learning – a proxy for

goal-directed and habitual behaviour. We perform a behavioural study that utilises a two-step

decision-making task and a reinforcement learning model to understand the effect of body image

dissatisfaction on model-based learning in a population characterised by high scores of

disordered eating and negative appearance beliefs, as recruited using Prolific. We find a

significantly reduced model-based contribution in the body image dissatisfaction task condition

in the population of interest as compared to a healthy control.

34   **Introduction**

Eating disorders (ED) form a group of different conditions that, in a detrimental way

affects a person's relationship with food. This usually leads to physical and psychological

problems, which severely decrease the quality of life. In fact, these problems can result in many

deaths, as eating disorders are reported to have one of the highest, if not the highest, mortality

rate among all mental disorders (1–4).

Generally, ED involve heightened preoccupation with food, such as restriction, or

consumption of unusually large amounts of food, which in some cases is followed by

compensatory behaviours such as vomiting, use of laxatives, or overexercising (5). In extreme

cases, eating disorders result in death either due to severe malnutrition, morbid obesity or suicide

(4,6). Moreover, the recovery from ED can be as low as 24% even after 10 years (7), which in

combination with high mortality and a significant decrease in quality of life, calls for extended

research into the roots and treatments of ED.

In addition to eating related behaviours, one of the core symptoms of ED is body image

disturbance, which can be understood as a negative misrepresentation of one's body, body image

preoccupation, usually involving disgust, shame and dissatisfaction (8). We here aim to

investigate how body image dissatisfaction in eating disorders affects basic mechanisms of

decision making, namely, habitual (repeating of the same action as a response to a stimulus) and

goal-directed (intentional and deliberate decision-making) behaviour (9). On top of logistic

regression analysis of the behavioural data(10), the mechanism will be captured in a

computational, reinforcement learning (RL) framework.

Background

Eating disorders.

There are three main subtypes of eating disorders – Anorexia Nervosa (AN), Bulimia

Nervosa (BN) and Binge-Eating Disorder (BED) as in 5th edition of The Diagnostic and

Statistical Manual of Mental Disorders (5). All share low life satisfaction, greatly impaired

quality of life, and increased mortality or suicide risk, and sometimes body image issues. A brief

distinction between the types, but nowhere close to accurately describing the nuance and lived-

experience of each subtype, would characterise: AN by extreme restriction of food and pursuit of

weight loss; BN by undergoing recurrent binge eating episode followed by compensatory

behaviours; BED by the recurrence of uncontrollable binges, but usually without compensatory

behaviours (5). In addition, there exists an array of atypical eating disorders that do not fit the

description of the main (clinical) types (11), but nevertheless constitutes a majority of cases (12).

In this paper, we do not focus on any particular type, but rather aim to explore and quantify the

effect of body image dissatisfaction on decision making in broadly understood eating disorders

(see Methods).

Computational psychiatry.

One possible avenue to understand and help devise treatments for eating disorders,

alongside the traditional approach of psychotherapy, psychiatry and neuroimaging (13–15) is the

emerging field of computational psychiatry (CP).This framework is based in the assumption that

the brain's characteristic function is one of computation and information processing. As such, it

offers an understanding of mental illness whereby the differences and/or errors within these

computations may result in malfunctioning, maladaptive behaviours and mental states(16). In an

attempt to describe these mechanisms, some researchers have focused on multiple modes of

decision making that can be gauged with a decision-making task and quantified with a

computational model that captures individual and group differences (17,18). Mental disorders

such as schizophrenia, anxiety, and obsessive-compulsive disorder (OCD) have been linked to

various significant changes in decision-making processes (19–21). Moreover, employing

computational methods to understand mental disorders promises bridging the neurobiological

and psychiatric levels of descriptions of populations of interest, for example the role of

dopaminergic signals in reward learning and it implications in addiction (22). Such link can

potentially inform better and more advanced theories of mental disorders, as well as inspire new

treatment approaches, and promises improved early detection leading to prevention (23).

Computational psychiatry of eating disorders.

As mentioned above, depression, OCD, anxiety and schizophrenia have received a lot

attention in the field of CP, with very promising results and theories. Eating disorders have

received less attention, with only a couple papers in the last nine years (24–34). To illustrate the

difference a Google Scholar search for eating- disorder related papers:

("computational psychiatry" "OR" "computational" "OR" "reinforcement learning" "OR"

"reinforcement" "OR" "bayesian" "OR" "decision making" "OR" "decision-making")

"AND" ("anorexia" "OR" "anorexia nervosa" "OR" "bulimia nervosa" "OR" "binge" "OR"

"binge eating" OR" "eating" "OR" "bulimia") (35),

returns 11 results, while a search for schizophrenia related papers:

("computational psychiatry" "OR" "computational" "OR" "reinforcement learning" "OR"

"reinforcement" "OR" "bayesian" "OR" "decision making" "OR" "decision-making")

"AND" ("schizophrenia") (36),

returns 466 results.

102   The main focus has been on two main aspects of decision making in eating disorders.

103 One is related to how sensitive to punishments individuals with AN are, that is how well they

104 learn from negative feedback from the environment (in this case a decision-making task).

105 Unfortunately, the results prove to be quite contradictory (24,33).

106   The second main aspect of ED that has been of interest to computational psychiatrists is

107 the trade-off between goal-directed and habitual system utilisation in decision-making. As

108 mentioned in the introduction, goal-directed decision-making is related to acting in the

109 environment with a goal in mind, intentionally and deliberately. It is usually characterised by

110 forming an internal model of the environment that describes which states and actions will bring

111 about the best results over prolonged time. On the other hand, habitual decision-making is

112 associated with responding to stimuli in the environment in an automatic manner, without much

113 deliberation, usually repeating those actions that immediately yield the best results. In this case,

114 an agent does not create a model of the environment. As such, goal-directed behaviour utilises

115 model-based learning, which allows building an accurate model of the states and actions and

116 their associated values that takes into account a hidden probabilistic structure of the environment.

117 However, this kind of learning is computationally more demanding, using more resources to

118 support the process. The habitual behaviour employs model-free learning, which updates a

119 running score of possible states and actions, based on the last experience, without registering any

120 hidden structures in the environment. Such a process is computationally efficient as it relies

121 mainly on the memory of the last events (9).

122   It has been shown that healthy population employs both model-free and model-based

123 learning that trade off against each other (10). Several studies attempted to see how this trade-off

124 is different within eating disorders. The results are converging and suggest that reduced model

based learning is associated with high eating disorder questionnaire scores in a general

population (30). The reduction is also evident clinical AN and BED groups (27,34). Moreover, a

recent study by Foerde et al. (27) suggests even further reduced model-based learning

contribution in AN when the task is strictly food related as compared to a monetary task. This is

of particular significance for two reasons. Firstly, AN is characterised by extreme pursuit of

weight-loss, which can be understood as extreme goal-directed behaviour. As such, one would

expect to see increased model-based learning in AN, yet the results suggest otherwise – such

"goal-pursuit" of weight-loss in AN is actually habitual, almost compulsive. Secondly, the study

investigates a food-related task (in addition to monetary), showing that the goal-directed deficits

are domain-general rather than specifically due to the monetary nature of the task, which on its

own may be construed as not motivating enough for that population.

Problem Statement, Objectives and Hypothesis

Evidently, there is a need for more computational psychiatry research about eating

disorders, given the high mortality. In particular, the focus should be directed at the body image

disturbance aspect of the disorder, which has not been previously investigated

To that end, we aim to explore the effect of body image dissatisfaction on decision

making in generally defined eating disorders by implementing a two-step decision-making task

that captures both model-based and model-free contribution (10). The task is given to a

population that scores high on an eating disorder, and body image disturbance questionnaires.

For comparison, a healthy control group (HC) is also be recruited. We employ two task

conditions, one of which attempts to target the dissatisfaction with body image and its

manifestation in decision making. The introduction of body image dissatisfaction condition aims

to create a context in which ED finds themselves during everyday struggle - excessive worry

about body image and how to change it through dieting. Secondly, the condition mirrors the

food-related task of Foerde et al.(27), aiming to support the domain-general nature of the goal-

directed deficit.

We hypothesise that: 1) In the monetary (neutral) condition and as described by previous

studies, the group with ED will report a significantly decreased model-based learning as

compared to HC, and no difference in model-free learning as previously shown (27,30). 2)

Model-based learning will be further decreased in the body image dissatisfaction condition in the

group with ED. HC will not report any significant differences between conditions.

158   **Materials and Methods**

Participants

An online study was performed. Participants for the study were recruited using Prolific,

which is an online participant recruitment service used mainly for research and academic

purposes (37). The platform provides a pool of participants that are reliable and allows to custom

screen for different groups before recruitment.

Since two groups were needed – a healthy control (HC) and an eating disorder (ED)

group - we applied two separate sets of pre-screening criteria on Prolific (pre-set filters that are

available in the study designer). For HC we looked for people who:

a) "Have never gone on a diet in the past."

b) "Do no currently for at least one week restrict food intake to manage weight."

c) "Have no diagnosed mental health condition that is uncontrolled (by medication

170   or intervention) and which has a significant impact on your daily life/activities."

171   d) "Do not have or have not had a diagnosed, on-going mental health illness or

172   condition."

173   For ED, participants had to select that they have gone on a diet, as in criterion (a), and

174 that they restrict food intake to either lose or maintain weight, as in criterion (b). Moreover, to

175 ensure data of good quality both groups had additional criteria to meet:

176   (i)   age between 18-38

177   (ii)   have normal or corrected-to-normal vision

178   (iii)   female as an assigned sex at birth

179   (iv)   an approval rate of 98%

180   (v)   a minimum of 20 previous submissions on Prolific.

181   Participants first completed a range of questionnaires (sub-study 1) to be then further

selected to complete a decision-making task (sub-study 2). The questionnaires were hosted on

Qualtrics (an online survey software) (38), while the decision-making task was hosted partly on

Qualtrics and partly on Pavlovia (an online behavioural experiment platform) (39), designed

using PsychoPy3 v. 2020.1.2 software (40). For each sub-study, participants were paid at a rate

of £6.25 per hour. The questionnaire took on average 7 minutes to complete, while the task took

on average 27 minutes. Subjects were based all around the world. The study was approved

according to the University of Edinburgh's Informatics Research Ethics Process, with an RT

number 2019/48215.

190   After further selection based on questionnaires and application of exclusion criteria for

the decision-making task (see Supplementary material), 35 (ED group) and 32 (HC group)

participants were selected for data analysis.

Power analysis

A prior power analysis was performed to determine the sample size needed to detect a

significant difference (at the 5% level) between groups in the neutral condition. Following a

similarly designed study by Foerde et al. (27), for calculation 1, we aimed for a medium effect

size of d=0.75 (two-tailed t-test between two independent means), and the power $(1 - \beta) = 0.8$.

The sample size calculation using G*Power software (41) suggested recruiting 29 participants

per group. Moreover, we calculated (calculation 2) the sample size needed to detect the

difference between conditions in the ED group (two-tailed t-test between two dependent means).

Setting the effect size at d=0.75 and the power at $(1 - \beta) = 0.8$, resulted in n=16 for the ED

group.

Additional post hoc power analysis was performed as above. Using the results from this

study with 35 participants in the ED group and 32 in the HC group, calculation 1 revealed effect

size of d=0.88 and power $(1 - \beta) = 0.94$, while calculation 2 revealed effect size of d=1.04 and

power $(1 - \beta) = 0.99$.

Self-report questionnaires

In the first part of the study, participants answered questions about their age, gender,

weight, and height. They also completed three questionnaires - Eating Attitudes Test (EAT-26)

(42), Appearance Anxiety Inventory (AAI) (43) and The Obsessive–Compulsive Inventory (OCI-

R) (44) – to assess the spectrum of: an eating disorder, body image disturbance and obsessive-

compulsive behaviours, respectively.

In order to divide participants into HC and ED groups, cut-off points were applied for the

questionnaires as in the Table 1, chosen based on the literature. For EAT-26, it is suggested to use

a cut-off of 11 for subclinical populations (45) that are characterised as displaying disordered

eating behaviours that could warrant further clinical diagnosis. Hence, for HC we set a threshold

of ≤ 10 and for ED ≥ 14 to achieve stronger effects. For AAI, a few cut-off points have been

suggested: a score above 6 may suggest some issues with body image (46), while scores between

15-20 is reported in an appearance-concerned adult population, and a score above 20 indicates a

high-risk group for body image disturbance (47). We therefore chose as cut-off for ED a score $\geq$

14, whereas for HC a score of ≤ 10 was chosen. Lastly, an additional criterion for HC group was

applied so that they do not display worrying obsessive-compulsive behaviours as these have also

been associated with reduced model-based learning (30). We selected a cut-off point of ≤ 10 on

OCI-R questionnaire for HC, which is twice as low as the recommended optimal cut-off point

(44). Moreover, two attention checks were implemented in the questionnaires to filter out

participants not taking part in the study in good faith.

Table 1. Cut-off points for ED and HC on EAT-26, AAI, and OCI-R questionnaires

Measure	Group	
	HC	ED
EAT-26	≤ 10	≥ 14
AAI	≤ 10	≥ 14
OCI-R	≤ 10	any

Two-step decision making task

Selected participants were asked to complete a two-step decision-making task in order to

quantify model-based and model-free learning contribution (10). Before proceeding with the

task, participants were asked to select a body type that is most similar to their own from a diverse

range of body types in the S1 Fig in the Supporting Information. Furthermore, they were

instructed on how to complete the task by reading through an illustrated tutorial. After reading

the instructions, they completed 25 trials of the task as a practice.

Fig 1. The structure of the two-step decision-making task.

In the first stage (grey), a dashed arrow corresponds to a rare transition with the probability of

30%, while the bold arrow corresponds to a common transition with the probability of 70%. In

the second stage, the speckled arrow is associated with the probability of receiving a reward. The

duration of each stage is noted to the right (stage one, stage two, reward).

The task reproduces that of Daw et al. (10) but was conceived as a treasure game to make

it as engaging and easy to understand as possible (Fig 1). In the first stage (grey stage),

participants are presented with a choice between two stimuli - on the left, a ship against a sunset,

while on the right, a pirate ship near a lighthouse. They are asked to play a role of a treasure

hunter and decide which ship to board. Each ship can sail to two destinations - a blue island and

a purple island. The game has a hidden structure, which participants tend to learn with time, such

that the normal ship sails to the blue island (left) 70% of the time (common transition) and to the

purple island (right) 30% of the time (rare transition), while the pirate ship sails to the blue island

30% of the time and to the purple island 70% of the time.

Once the participant boards the chosen ship, they are taken to one of the islands (stage

two), where two chests await them. Here, they are asked to quickly select the chest to try their

chances at finding a treasure (a pirate coin). However, each chest is assigned a probability of

containing the coin, which evolves over time according to a Gaussian random walk with a

standard deviation $\sigma = 0.0275$, as in the Fig 2 (more details in the Supporting Information). One

of the strategies is to track, over time, which chest is the most favourable, i.e. has the highest

chance of yielding the coin. After the chest is opened and the participant receives the coin (or

not), they are taken back to the first stage to repeat the trial.

Fig 2. The sample evolution of the reward probability for each of the chest in stage 2.

The probabilities (in the order as in Fig 1) evolve over 150 trials, according to a Gaussian

random walk with σ=0.0275. One set of two evolutions always starts randomly in a range [0.58,

0.72], while the other set of two starts in [0.31, 0.45]. The values of probabilities are bounded in

[0.25, 0.75].

There are two conditions, 150 trials each. Each condition consists of two series of 75

trials, with a short break in between to counteract a possible loss of attention or tiredness. The

neutral (NT) condition is exactly like in the Fig 1, with a pirate coin as a reward (Fig 3A). The

other condition, the body image dissatisfaction (BID) condition, differs from the neutral in that

the reward is a pirate coin next to a body type that the participant selected as the most similar to

their own (Fig 3C). The aim of the BID condition is to gauge the effects of body image

dissatisfaction on decision-making.

Fig 3. Possible reward outcomes during the task.

In the neutral condition, participants could receive either (A) or (B), while in the BID condition

they could receive either (C) or (B). An empty box (B) indicates no reward. The body type was

selected by the participant before the task, more body type examples can be found in the S1 Fig.

For a balanced design half of the participants in each group completed the neutral

condition first, followed by the BID condition. The other half completed the task in a reverse

order. All results are then based on the average across two subgroups.

Model-based and model-free learning in the task.

The task allows to capture the contribution of model-based and model-free learning.

Thanks to its stage-like structure and probabilistic nature of we can easily distinguish between

two strategies a participant can have, usually a mix of the two.

For the purpose of this example, we focus on two extreme cases (30). On one hand, an

agent could utilise only model-free learning strategy. This means that they would make their

decision as to which ship and chest to select purely based on the reward they received (or lack

thereof), usually repeating the rewarded action, regardless of the fact the transition type. On the

other hand, a participant who uses only model-based strategy, takes into account the learnt

transition structure of the task, on top of the knowledge about previous rewards. Moreover, they

track the probability of receiving the reward to know which chest is the most favourable. In other

words, they chose the same ship if the previous trial had a common transition with a reward or a

rare transition and no reward.

Mixed-effects logistic regression of raw choice data

Before fitting RL model parameters to the data, a mixed-effects logistic regression

analysis of raw choice data was performed to quantify the model-based and mode-free learning

effects (10). The analysis focuses only on the choices made in the first stage of the task and how

these are influenced by received a reward and transition type in the previous trial. The regression

models the probability of repeating the same choice, p(stay) and how it's influenced by reward

(rewarded=1, unrewarded=-1) and transition (common=1, rare =-1), and their interaction. The

main effect of reward is interpreted as model-free learning contribution and the reward x

tranision interaction as model-based learning contribution. These effects were regressed

alongside with group, condition, and age (z-scored) as fixed effects as below:

glmer(stay ~ r * transition * (group * condition + age_z) + (r * transition + 1 |

sub), family = binomial),

The estimates were obtained with the lme4 package in R, using Bound Optimization by

Quadratic Approximation (bobyqa) with 1e5 functional evaluations (30).

Stay probabilities calculation (frequency-based)

As an additional measure, for each participant we calculated the following probabilities

p(stay|rewarded, common), p(stay|unrewarded, common), p(stay|rewarded, rare),

p(stay|unrewarded, rare) based on the frequency of repeated choice in each of the four (reward,

transition) cases. These were used for additional regression analyses and figures.

Model-based score calculation (frequency-based)

Yet another measure that helps to capture model-based learning contribution that is

independent of the reinforcement learning is the model-based score (MB score) (48). This is

calculated as follows:

MBscore = p(stay|rewarded, common) - p(stay|rewarded, rare) -

p(stay|unrewarded, common) + p(stay|unrewarded, rare)

Model-based and model free learning - an RL model

Finally, the full computational RL model, described and quantified below (27,30,49),

incorporates choice data from two stages of the task, For simplicity, we begin with the update

equation of the state-action value function, $Q^{II}(s_2, c_2)$ for the second stage (II) states and actions.

There are two possible stage two states s_2: 1 - the blue island; 2 - the purple island, such that

$s_2 \in \{1,2\}$. On each island, an agent can make two separate choices c_2: L - open the left chest; R

- open the right chest, such that $c_2 \in \{L, R\}$. Moreover, after opening the chest, the agent can

receive the reward, $r \in \{0,1\}$, where 0 corresponds to an empty chest, and 1 corresponds to the

pirate coin. At the start of the task all value functions are initialised at 0.5. On any trial, t, we

update the value function $Q^{II}(s_2, c_2)$ of the visited state and action taken as in the Eq. 1. The

subscript, t, indicates the current trial values, whereas $t + 1$ indicates the values at the following

trial. α is the learning rate.

$$Q^{II}_{t+1}(s_{2,t}, c_{2,t}) = (1 - \alpha)Q^{II}_t(s_{2,t}, c_{2,t}) + r_t \tag{1}$$

To calculate the probability of making a choice $c \in \{L, R\}$ at stage two, we use the

softmax function as in Eq. 2, with an inverse temperature parameter β_2 quantifying the influence

of the value function on making the choice.

$$P(c_{2,t} = c) = \frac{\exp\{\beta_2 Q^{II}_t(s_{2,t}, c)\}}{\sum_{c' \in \{L,R\}} \exp\{\beta_2 Q^{II}_t(s_{2,t}, c')\}} \tag{2}$$

In stage one (I), we directly see how model-free and model-based learning play their part.

Here, we have two sets of update equations. The first set, Eq. 3-5 is model-based, where the

agent uses their knowledge about the environment - which stage two state-action is the best as

well as the structure of the task (common vs. rare transition) to update the values of the ships in

stage one. In this case, the value functions for both ships are updated simultaneously, such that

the value of choosing a ship is the weighted sum (by transition probability) of maximums over

stage two actions values.

$$\mathbf{Q}^{MB}_t = [Q^{MB}_t(c_{1,t} = L), Q^{MB}_t(c_{1,t} = R)] \tag{3}$$

$$Q_t^{MB}(c_{1,t} = L) = 0.7 \times \max_{c_2}\{Q_t^{II}(s_2 = 1, c_2)\} + 0.3 \times \max_{c_2}\{Q_t^{II}(s_2 = 2, c_2)\} \qquad (4)$$

$$Q_t^{MB}(c_{1,t} = R) = 0.7 \times \max_{c_2}\{Q_t^{II}(s_2 = 2, c_2)\} + 0.3 \times \max_{c_2}\{Q_t^{II}(s_2 = 1, c_2)\} \qquad (5)$$

The second, model-free update in Eq. 6, updates the values of the ships based only on the reward

received at the end of the trial.

$$Q_{t+1}^{MF}(c_{1,t}) = (1 - \alpha)Q_t^{MF}(c_{1,t}) + r_t \qquad (6)$$

Model-free and model-based contributions are joined together in a weighted value function over

two ship choices, Q^I, as in Eq. 7.

$$Q_t^I(c) = \beta_{MB}Q_t^{MB}(c) + \beta_{MF}Q_t^{MF}(c) + \rho I(c = c_{1,t-1}) \qquad (7)$$

The contribution of each system is captured with β_{MB} (model-based) and β_{MF} (model-free)

parameters. An additional indicator $I(c = c_{1,t-1})$ tells if the choice made on the current trial is

repeated as in the previous one, with a parameter ρ describing how much switching or staying is

done regardless of the feedback. Finally, the probability of choosing either ship is calculated as

in Eq. 8, analogously to the second stage.

$$P(c_{1,t} = c) = \frac{\exp\{Q_t^I(c)\}}{\sum_{c' \in \{L,R\}} \exp\{Q_t^I(c')\}} \qquad (8)$$

The model has a total of five parameters: $\beta_{MB}, \beta_{MF}, \beta_2, \alpha, \rho$.

Moreover, for each subject, a relative model-based fractional reliance was also calculated,

treated as a new parameter (27), which allows to capture the trade-off between model-free and

model-based learning:

$$w = \frac{\beta_{MB}}{\beta_{MB} + \beta_{MF}}$$

Model fitting

Model parameters for each group and condition were estimated using hierarchical

Bayesian approaches, which provides the best test-retest reliability for this particular model, and

follows the procedure as in Brown et al. (50). The estimation was performed using RStan

package (v.2.21.0) (51) in R (v. 4.0.2) based on Markov Chain Monte Carlo techniques (No-U-

Turn Hamiltonian Monte Carlo). Each parameter was estimated with a mean, scale, and

individual error estimates. The learning rate α was constrained to (0,1) with an inverse logit

transformation, while the means of β_{MB}, β_{MF}, were bounded below by 0. We used weakly

informative and uninformative priors for the means: $\alpha \sim N(0,2.5), \beta_{MB} \sim N(0,100), \beta_{MF} \sim$

$N(0,100), \beta_2 \sim N(0,100), \rho \sim N(0,100)$. For the scales of each parameter, we used

Cauchy(0,2.5) distribution, all constrained to be greater than 0. Individual error terms were all

$N(0,1)$, where for β_{MB}, β_{MF} these were constrained to be greater than 0. For each of the four

chains we ran 2000 samples (after discarding 2000 warm-up ones). Chains were inspected for

convergence and their $\hat{R}$ values were all around 1 (below 1.1). The mean value (across chains) of

each parameter for each participant were used in the analysis.

Mixed-effects linear regression of parameter estimates

The estimates for each parameter were analysed using mixed-effects linear regression to

compare them between groups and conditions, with group, condition and age (z-scored) as fixed

effects per subject:

lmer(parameter ~ group * condition + age_z + (1 | sub))

Parameter recovery

To quantify the reliability of the parameter estimates, parameter recovery was performed after

data collection (see Supporting Information). For the collected data, the model and fitting

procedure described above provide fair to excellent reliability, with average (across parameters,

groups and conditions) parameter recovery Pearson correlation coefficient (PCC), $\rho = 0.834$.

Results

Demographic and questionnaires summary

The screening procedures resulted in recruiting two significantly different groups -

healthy control (HC) and eating disorder (ED). The summarised information, along with two-

sample t-tests, can be found in Table 2. The groups are mainly characterised by average total

scores on the three questionnaires, with scores significantly different between groups across all

questionnaires (p<0.001). As the groups also significantly differ in age, the variable (z-scored)

was included as a covariate in the regression analysis.

Table 2. Summary of demographic information and questionnaire scores in each group.

Measure	HC (n=32) Mean ±SD	ED (n=35) Mean ±SD	t value	p value
EAT-26	2.94 ± 2.72	25.83 ± 10.67	-11.78	**<0.001***
AAI	4.59 ± 2.87	23.94 ± 7.26	-14.10	**<0.001***
OCI-R	5.5 ± 3.59	22.83 ± 11.09	-8.44	**<0.001***
BMI (kg/m²)	21.83 ± 4.27	26.07 ± 6.21	-3.01	**0.004***
Age	26.38 ± 4.61	30.57 ± 4.45	-3.79	**<0.001***

This includes means and standard deviations (SD) of EAT-26, AAI, OCI-R scores, age, and BMI,

as well as t- and p- values of the two-sample t-tests.

Task performance - rewards and reaction times

The performance in the task was analysed independent of the reinforcement learning

model. Average characteristics were calculated for each group (detailed results in S3 Table).

These measures include: total reward in the neutral and BID condition, and the total reward after

completing the full task, as well as mean reaction times (RT) during the neutral, BID, and across

both conditions. There were no differences in the above measures between groups.

Collected raw choice data analysis

Mixed-effects logistic regression analysis – collected data.

To assess the model-based and model-free contribution during the task, raw choice data

was analysed with mixed-effects logistic regression of the probability of repeating the same

choice in stage one (Table 3). HC group showed a significant contribution of both model-free

system as indicated by the reward effect (p-value<0.001), and model-based system as indicated

by the reward x transition interaction effect (p-value=0.018). According to our hypothesis, we

would expect a significantly lower estimate for the reward × transition × groupED effect to

detect a difference between groups in the neutral condition; as well as a significantly lower

estimate for the reward × transition × groupED × conditionBID effect to detect a difference

between condition in the ED group. However, the analysis revealed no differences between

groups or conditions in terms of model-based and model-free learning. The estimated

probabilities (frequency-based) for each case in each group and condition are depicted in Fig 4.

Fig 4. Stay probabilities in the collected data. (A) Neutral condition. (B) BID condition.

Table 3. Random effects logistic regression for probability of staying (collected data).

Effects	Estimate	SE	z value	p value
Intercept (HC, conditionNT)	1.68	0.24	6.86	**<0.001***
reward	0.92	0.33	2.77	**0.006***
transition	-0.57	0.16	-3.68	**<0.001***
groupED	-0.46	0.35	-1.33	0.183
conditionBID	-0.21	0.13	-1.56	0.120
age_z	0.18	0.17	1.07	0.285
reward × transition	0.84	0.36	2.37	**0.018***
groupED × conditionBID	-0.07	0.18	-0.36	0.718
reward × groupED	-0.19	0.47	-0.40	0.689
reward × conditionBID	-0.03	0.22	-0.13	0.900
reward × age_z	-0.05	0.22	-0.21	0.833
transition × groupED	0.26	0.21	1.23	0.219
transition × conditionBID	0.09	0.16	0.57	0.569
transition × age_z	-0.05	0.09	-0.57	0.566
reward × groupED × conditionBID	0.21	0.30	0.72	0.473
transition × groupED × conditionBID	0.11	0.22	0.51	0.612
reward × transition × groupED	-0.43	0.50	-0.87	0.387
reward × transition × conditionBID	0.05	0.27	0.21	0.837
reward × transition × age_z	0.00	0.23	0.00	0.999

Effects	Estimate	SE	z value	p value
reward × transition × groupED × conditionBID	-0.30	0.36	-0.85	0.397

Note. Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

To check whether model-based and model-free learning overall manifests itself across the

two populations together, group and condition were removed as fixed effects from the regression

analysis (Table 4). The simpler model revealed significant contribution of both learning systems

regardless of the group and condition (Reward effect p-value<0.001, Reward x Transition effect

p-value=0.008). Moreover, there was a significant effect of transition (p-value <0.001)

Table 4. Random effects logistic regression for probability of staying (collected data).

Effects	Estimate	SE	z value	p value
Intercept	1.31	0.15	8.45	**<0.001***
r	0.87	0.20	4.30	**<0.001***
transition	-0.36	0.09	-4.07	**<0.001***
age_z	0.08	0.16	0.48	0.628
r:transition	0.56	0.21	2.64	**0.008***
r:age_z	-0.07	0.20	-0.34	0.732
transition:age_z	0.02	0.09	0.18	0.857
r:transition:age_z	-0.13	0.21	-0.60	0.552

Note. Age (z-scored) is treated as fixed-effect covariates per subject.

Frequency based-probability regression – collected data.

Furthermore, we estimated the probabilities of staying after common and rewarded trials

using the frequency calculation and used them in mixed-effects linear regression analysis (Table

5). This simpler comparison revealed reduced probability of staying after common, rewarded

trials between groups in the neutral condition, which was just short of significance (p-value =

0.054), indicating a potentially weaker model-based learning capacity in ED. There were no

differences between conditions in either group.

Table 5. Random effects for probability of staying based on collected data (frequency

estimate).

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	0.93	0.03	35.79	**<0.001***
groupED	-0.07	0.04	-1.96	0.054
conditionBID	-0.01	0.01	-0.46	0.649
age_z	0.00	0.02	-0.09	0.930
groupED:conditionBID	-0.01	0.02	-0.50	0.622

Note. Age is treated as fixed-effect covariates per subject.

MB score analysis – collected raw choice data

Lastly, we calculated MB scores in the collected data (Fig *5*). These were regressed

against group and condition variables (Table 6). There were no differences in MB scores between

groups and conditions.

Fig 5. MB scores per group and condition in collected data.

Table 6. Random effects linear regression for MB score (collected data).

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	0.12	0.04	2.80	**0.006***
groupED	-0.06	0.06	-0.93	0.354
conditionBID	0.01	0.04	0.20	0.845
age_z	-0.01	0.03	-0.20	0.843
groupED:conditionBID	-0.04	0.06	-0.69	0.490

RL parameter estimates - model-based and model-free learning

Following the negative results from the logistic regression, an attempt to capture the

group and condition differences was made using a more sensitive RL model, taking into account

incremental learning over many trials. Model parameters were fit as described in the Methods. A

comparison of average parameter estimates with standard errors (SE) of $\beta_{MB}, \beta_{MF,}, w$ between

groups and conditions can be found in Fig 6.

Fig 6. Estimated model parameters.

Mean $\pm$ standard error (SE) of the estimated model parameters in HC (blue) and ED (red) during

each condition (neutral and BID). (A) β_{MB}, (B) β_{MF}, (C) w.

The results from the mixed effects linesar regression model for each parameter, showing

a significant difference between groups and conditions can be found in Tables 7-11.

Table 7. Mixed effects linear regression analysis of model-based learning parameter β_{MB}

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	1.42	0.17	8.23	**<0.001***
ED group	-0.61	0.25	-2.46	**0.016***
BID condition	0.22	0.11	1.92	0.060
Age	-0.09	0.12	-0.75	0.454
ED group × BID condition	-0.49	0.16	-3.08	**0.003***

481   Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

483   **Table 8. Mixed effects linear regression analysis of model-free learning parameter β_{MF}**

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	1.56	0.14	10.94	**<0.001***
ED group	-0.70	0.21	-3.81	**<0.001***
BID condition	-0.32	0.14	-2.62	**0.011***
*Age	0.02	0.09	0.30	0.764
ED group × BID condition	0.52	0.20	1.94	0.057

484   Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

486   **Table 9. Mixed effects linear regression analysis of relative model-based learning**

487   **parameter w**

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	0.51	0.04	14.10	**<0.001***
ED group	0.05	0.05	0.98	0.329
BID condition	0.04	0.03	1.23	0.223
Age	0.01	0.02	0.22	0.825
ED group × BID condition	-0.15	0.04	-3.58	**<0.001***

488   Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

490   **Table 10. Mixed effects linear regression analysis of learning rate parameter α**

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	0.61	0.05	12.60	**<0.001***
ED group	-0.09	0.07	-1.33	0.188
BID condition	0.15	0.05	3.02	**0.004***

Effects	Estimate	SE	t value	p value
Age	0.02	0.03	0.50	0.619
ED group × BID condition	-0.10	0.07	-1.43	0.159

491 Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

493 **Table 11. Mixed effects linear regression analysis of inverse temperature parameter β_2**

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	1.63	0.12	13.06	**<0.001***
ED group	-0.60	0.18	-3.34	**0.001**
BID condition	-0.03	0.11	-0.25	0.802
Age	0.10	0.08	1.23	0.224
ED group × BID condition	0.09	0.16	0.56	0.576

Group, condition, age (z-scored) are treated as fixed-effect covariates per subject.

As hypothesised the model-based contribution (as quantified with β_{MB} parameter) in the

neutral condition is decreased in ED as compared to HC (Table 7; p-value = 0.016). Moreover,

there is a further reduction in model-based learning in the BID condition in ED (p-value =

0.003), which is not present in the HC group (p-value = 0.060). However, looking at the w

parameter (Table 9), that captures the trade-off between model-free and model-based learning,

the reduction in model-based contribution is only present in the BID condition in ED group (p-

value <0.001), while there is no difference between groups in the neutral condition (p-value =

0.329).

Surprisingly, model-free learning is significantly reduced in the neutral condition in the

ED group (compared to HC p-value <0.001, Table 8), which then increases (short of

significance) in the BID condition (p-value = 0.057). Furthermore, model-free learning is slightly

attenuated in the BID condition in HC (p-values = 0.011).

In addition to model-based and model-free learning, there was an increase in the learning

rate between conditions for HC (p-value = 0.004; Table 10), as well as a reduction in the

β_2 inverse temperature parameter estimate in the ED group in the neutral condition (p-value =

0.001; Table 11), potentially indicating a more exploratory choice strategy in the second stage.

Simulated raw choice data analysis

Mixed-effects logistic regression analysis – simulated data.

In an attempt to check whether the RL model is even able to capture the contribution of

model-based and model-free system in the recruited populations, ensuring meaningful and

interpretable results, we used the individually estimated parameters to simulate fake choice data,

which was then analysed used mixed-effects logistic regression, looking for the effects of reward

and its interaction with transition (27). In the model with the interaction between group and

condition as a variable (Table 12), we detected model-free and model-based contribution in the

neutral condition in HC (p-value=0.013, p-value<0.001, respectively). Moreover, the model also

detected reduced model-based contribution in the ED group as compared to HC group in the

neutral condition (p-value=0.013). There were no differences between conditions in either group.

The estimated probabilities (frequency-based) for each case in each group and condition are

depicted in Fig 7.

Fig 7. Stay probabilities in the simulated data. (A) Neutral condition. (B) BID condition.

Table 12. Random effects logistic regression for probability of staying (simulated data).

Effects	Estimate	SE	z value	p value
Intercept (HC, conditionNT)	1.64	0.20	8.16	**<0.001***
reward	0.52	0.21	2.48	**0.013***
transition	-0.42	0.13	-3.20	**0.001***
groupED	-0.36	0.29	-1.23	0.217
conditionBID	-0.27	0.14	-2.00	**0.045***
age_z	0.09	0.14	0.68	0.496
reward × transition	0.82	0.23	3.63	**<0.001***
groupED × conditionBID	0.14	0.19	0.76	0.446
reward × groupED	0.06	0.29	0.22	0.826
reward × conditionBID	0.26	0.21	1.25	0.213
reward × age_z	0.01	0.13	0.12	0.907
transition × groupED	0.24	0.18	1.31	0.190
transition × conditionBID	0.12	0.16	0.77	0.444
transition × age_z	0.00	0.07	-0.05	0.961
reward × groupED × conditionBID	-0.35	0.29	-1.23	0.220
transition × groupED × conditionBID	-0.02	0.22	-0.09	0.929
reward × transition × groupED	-0.77	0.31	-2.49	**0.013***

Effects	Estimate	SE	z value	p value
reward × transition × conditionBID	-0.22	0.25	-0.87	0.382
reward × transition × age_z	-0.06	0.13	-0.44	0.660
reward × transition × groupED × conditionBID	0.26	0.34	0.77	0.443

Note. Group, condition, age are treated as fixed-effect covariates per subject.

Frequency based-probability regression – simulated data.

Furthermore, comparing the simulated probabilities of staying after common and

rewarded trials using the frequency calculation in random effects linear regression (Table 13), we

again found a weaker model-based learning capacity in ED (p-value = 0.033). There were no

differences between conditions in either group.

Table 13. Random effects linear regression for probability of staying (frequency-based;

simulated data).

Effects	Estimate	SE	t value	p value
Intercept (HC, neutral)	0.91	0.03	35.61	**<0.001***
groupED	-0.08	0.04	-2.17	**0.033***
conditionBID	-0.01	0.02	-0.46	0.649
age_z	0.00	0.02	-0.25	0.803
groupED:conditionBID	0.00	0.03	0.04	0.968

Note. Age is treated as fixed-effect covariates per subject.

MB score analysis – simulated raw choice data

Lastly, we calculated MB scores in the simulated data (Fig 8). These were regressed

against group and condition variables (

Table **14**). The simulated data captured reduced MB scores in the ED group in the neutral

condition (p-value = 0.040).

Fig 8. MB scores per group and condition. (A) collected and (B) simulated data.

Table 14. Random effects linear regression for MB score (simulated data).

Effects	Estimate	SE	t value	p value
Intercept (HC, conditionNT)	0.10	0.03	3.42	**<0.001***
groupED	-0.09	0.04	-2.08	**0.040***
conditionBID	-0.02	0.03	-0.73	0.471
age_z	-0.01	0.02	-0.39	0.699
groupED:conditionBID	0.02	0.04	0.53	0.598

Correlations of Δ*w* with other covariates

To check how the difference in model-based learning between condition correlates with

questionnaire scores and demographics (age, EAT-26, AAI, and OCI-R scores all z-scored), we

introduced Δ*w* that captures the difference in *w* between conditions:

$$\Delta w = w_{BID} - w_{NT}$$

when negative, this would indicate the BID condition had a reducing effect on model-based

learning.

The correlation plots with Pearson correlation coefficient (PCC), ρ, and p-values for the

hypothesis of no relationship between Δw and the covariates can be found in Fig 9. We found

significant negative correlations with EAT-26 and AAI scores and OCI-R scores as in Table 15

(ρ=-0.269, ρ=-0.363, ρ=-0.324 with p-value=0.028, p-value=0.003, p-value=0.007).

Fig 9. Correlation of EAT-26, AAI, OCI-R and age measurements (z-scored) with Δw.

Correlation is across both groups. Pearson's correlation coefficient (ρ) and p-values are included.

Table 15. PCC, ρ, between Δw and the covariates across HC and ED together.

	Covariates			
	EAT-26	**AAI**	**OCI-R**	**Age**
ρ	-0.269	-0.363	-0.324	0.029
p-value	**0.028**	**0.003***	**0.007***	0.817

568   P-values for the hypothesis of no relationship between the Δw and covariates (EAT-26, AAI,

569   OCI-R scores and age - all z-scored) are included.

571   **Discussion**

The current study focused on the missing element in computational psychiatry research

on eating disorders - body image dissatisfaction. In particular, the reduced utilisation of goal-

directed behaviour was explored.

First of all, the findings from a two-step decision-making task show a significantly

smaller contribution of model-based learning in a population characterised by high body image

dissatisfaction and disordered eating as compared to the HC group. We find a further model-

based learning reduction in this population when the task aims at capturing the additional effect

of body image dissatisfaction (as implemented in the BID condition). We did not find such an

effect in the healthy control. Additionally, the reduction in model-based learning between groups

and between conditions in the ED group was not associated with age.

Furthermore, we found a significant correlation of the difference between the measures of

model-based learning in the neutral and BID condition with EAT-26 and AAI scores. It was also

correlated with OCI-R scores, which may suggest that the BID condition is not able to capture

the difference between the effect of the body issues related and OCD-like behaviours on model-

based learning. Nevertheless, this is an important finding that extends previous results that

correlate reduced model-based learning in one general set of eating disorders and compulsive

disorders (30). Despite the exploratory nature of this study, the results are promising in that they

can point further research to better understand the mechanisms of eating disorders and associated

with it body image dissatisfaction and how these manifest in decision-making.

Moreover, we replicated previous findings that suggest a decreased model-based

learning, during a neutral condition of the two-step decision-making task, in groups characterised

by disordered eating and compulsive behaviours (27,30,34). The results confirm the set

hypothesis of decreased model-based learning between groups and between conditions in ED.

However, the simultaneous decrease in model-free learning in the HC group is surprising and

will deserve further investigation. This result is also contrary to previous studies that find no

difference in model-free learning between groups (27,30). One potential reason for the difference

of results is the high heterogeneity in the estimates of model-free learning parameter both in ED

and HC group, which may have skewed the results. In fact, HC group, in general, has more

heterogeneous estimates of parameters than ED. This may be due to the HC group being in fact

composed of different subgroups, possibly even along the eating disorder continuum, for

example if they were not entirely sincere when answering the questionnaires. Alternatively, ED

may exhibit a generally reduced learning capability (see below for further discussion).

Lastly, the logistic regression of the raw choice data showed the two populations employ

both learning systems. However, to detect the differences between groups and conditions we

used a more sensitive RL model analysis that takes into account the incremental learning from

many trials as well as stage two choices, which are not part of the logistic regression model. The

group difference in model-based learning RL parameter in the neutral condition was further

confirmed in the logistic regression of the simulated data using RL parameter estimates.

Implications

First of all, as far as we know, this is the first study that looked at the body image

dissatisfaction from a computational perspective. Our study complies with a novel RDoC

framework (52) that aims to investigate and understand mental disorders by integrating many

levels of information such as behaviour and self-report. Such a framework helps overcome the

downsides of a classical psychiatric approach, namely, comorbidity of disorders, heterogeneity,

and arbitrary definition of what constitutes a disorder (53). Our results are based on a rigorous

and quantified behavioural paradigm that addresses an underlying mechanism of the disorder –

the effect body image dissatisfaction on model-based learning.

The fact that body image dissatisfaction is associated with decreased contribution of

model-based learning provides some support for the mechanism of extreme habitual body

preoccupation. The fear of gaining weight and associated with it goal of either losing or not

gaining weight set as a goal, evolves over time into a rigid and habitual body checking. When an

individual's body type is displayed onto a screen it may act as a trigger for the habitual behaviour

of body comparison and concern. As such, a body-preoccupied state leads to significant

reduction in the model-based capabilities that could be allocated towards task completion in a

more goal-directed manner, as is done in the neutral condition. Our result adds to the discussion

reopened by Foerde et al. (27) about domain-general/specific deficits in goal-directed learning. It

provides support for the view that goal-directed control impairments are not specific to purely

monetary tasks but suggests that it could be aggravated in conditions that trigger body image

preoccupations.

The high heterogeneity of parameter estimates (model-based/free) in the healthy control

may suggest that some healthy controls share some traits with the ED group that explains

reduced model-based learning. This might be related to traits that we have not directly explored

in our study but that have been associated with deficits in model-based learning, e.g. alcohol

addiction, or impulsivity (30), despite partially controlling for the effect of compulsivity on

model-based learning (30) by excluding participants with high OCI-R scores in the healthy

control group. Alternatively, it is possible that some participants from a healthy control group do

in fact have body image preoccupation issues. A potential factor in group misclassification could

be a wide-spread and widely accepted societal preoccupation with dieting, looks as well as the

phenomenon of body shaming that could render the recruitment of a 'healthy' population

difficult (54,55). This could manifest in participants as selecting the option of "no past diet

experience" in the screening stage despite 'clean eating' they might engage in as a widely

accepted 'health standard', which is actually emotionally distressing and linked with functional

impairments (56). As such, the heterogeneity in the parameter estimates in the healthy control

may highlight the blurred boundary between health and dieting, and indeed call into question the

possibility of a truly healthy control group within this field of research.

Limitations

We employed the most reliable fitting method for this particular task and model (50), as

well as applying strict exclusion criteria to participants' behavioural data and questionnaire

attention checks to arrive at these initial promising results. Nevertheless, it would be beneficial to

perform a similar study on a larger online population and/or in the lab on a clinical group to

strengthen the evidence for the hypothesis of this paper and test its replicability. This should also

further address any possible concerns related to the varied nationalities of the participants.

A possible limitation presents itself in the selection of body types/silhouettes by the

participants. These may be seen as overly expressive and caricature-like, failing to accurately

capture their body type, which may cast doubt on the source of the reduction in model-based

learning. As such, we are currently working on a paradigm that offers a more rigorously defined

range of body types for the participants to select from, where a selected image is compared to the

participant's basic body parameters (e.g. weight, height), capturing the discrepancy between

reality and perception. That being said, the results of this study offer an interesting point for

further examination: the significant change in goal-directed decision-making exclusively in the

ED group, caused by the mere presence of said silhouettes.

An alternative explanation of the results may expose potential issues with the design of

the BID condition. Since the silhouette is displayed with the reward, this may induce aversive

behaviour (due to high-body dissatisfaction), whereby learning from the rewards is reduced,

which results in lower values of inverse temperate parameters both for model-free and model-

based learning. Though a valid concern, which could be more concretely verified by re-running

the study using a different design, we believe that the reduction of the fractional model-based

reliance parameter (w) between conditions may still serve as promising evidence for the reduced

model-based learning. The reasoning being that the w parameter captures the trade-off between

the two systems (relative measurement) regardless of the absolute reduced learning from the

rewards. Moreover, we did not detect any differences in total reward received between

conditions in the ED group. A different interpretation then arises, which could indicate that the

ED group may employ a slightly different strategy where they rely more on action exploration

rather than exploitation to perform well in the task (19).

Furthermore, we are aware of the not so excellent parameter recovery, especially in the

ED group in the BID condition. Having informally tested various models that allow for

constraining of the β_{MB}, β_{MF} parameters to be greater than 0, the only way to improve the

parameter recovery would be to increase the sample size.

Kool et al. (57) suggest that the structure of the two step decision making task used here

does not accurately estimate the trade-off between model-based and model-free learning. The

authors propose certain modifications to the task to increase that accuracy such as changing the

drift rate in the Gaussian random walk of reward probabilities, reducing the number of stage two

choices to one per state, or introducing a deterministic transition structure. However, it has been

shown that in some cases of slight variations to the task structure, the reinforcement learning

model will not be able to distinguish between model-based and model-free actions as efficiently

as in the standard task (58), while the task in the current form has proven to yield reliable and

consistent results (27,30).

An alternative explanation to the reduction of model-based learning in the BID condition

in the ED group can be attributed to a stress response rather than an effect that is strictly body

image related. Otto et al. (59) showed that as a result of a stress response (submerging a hand in

ice cold water) model-based learning noticed a decreased contribution to decision-making. Given

that body image preoccupation can be characterised as a distressing experience, the result from

our study could be potentially explained by a generic stress response. However, to verify this

alternative explanation would be beyond the scope of this exploratory study, where for example

an cortisol-measurement/imaging study could be performed to identify which cortisol/neural

response is associated with reduced model-based learning in the BID condition and compare it to

a hypothesised, stress-related response(60–62).

Lastly, there are a couple of steps that could be taken in order to further explore the

mechanisms and effects of body image dissatisfaction on decision-making in eating disorders. A

natural step is to perform a very similar study as here, on a clinical population and use brain

imaging such as fMRI to try and correlate the behavioural-computational changes in the ED

group with neural signatures to provide a neurobiological basis of body image dissatisfaction

effects on decision-making. Ideally, in the future, a more whole understanding of eating disorders

and accompanying body image issues could help devise new and more effective treatment

options.

709   **Conclusions**

710   Given the high mortality of eating disorders, this study expands the sparse field of

computational psychiatry of eating disorders that so far has focused on the general perception of

reward (24,33), and quantification of model-based learning in a neutral setting (30,34) or in

relation to food choices (27). Since one of the prevalent aspects of eating disorders is body image

preoccupation, we explored its effect on model-based learning in comparison with a healthy

control. The results from the online study on a population characterised by high scores on eating

disorder and body image dissatisfaction questionnaires show a significantly negative effect of

body image dissatisfaction on model-based learning that is not present in the healthy control.

This finding offers additional insight into the mechanisms of the disorder and the effect that the

core element of the disorder, such as body image dissatisfaction, has on decision-making.

Directed by the above result, further inquiry can be performed into more nuanced treatment

strategies that could help to break from rigid habits and strengthen model-based capabilities of

patients related to the perception of their body through well-tailored therapeutic activities.

References

1.   Arcelus J, Mitchell AJ, Wales J, Nielsen S. Mortality Rates in Patients With Anorexia
Nervosa and Other Eating Disorders: A Meta-analysis of 36 Studies. Arch Gen Psychiatry.
2011;68(7):724–31.

2.   Fichter MM, Quadflieg N. Mortality in eating disorders - results of a large prospective
clinical longitudinal study. Int J Eat Disord. 2016;49(4):391–401.

3.   Godier LR, Park RJ. Compulsivity in anorexia nervosa: a transdiagnostic concept. Front
Psychol. 2014;5.

4.   Smink FRE, van Hoeken D, Hoek HW. Epidemiology of Eating Disorders: Incidence,
Prevalence and Mortality Rates. Curr Psychiatry Rep. 2012;14(4):406–14.

5.   American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders
(DSM-5®). American Psychiatric Pub; 2013. 1520 p.

6.   Morris J, Twaddle S. Anorexia nervosa. BMJ. 2007;334(7599):894–8.

7.   Berkman ND, Lohr KN, Bulik CM. Outcomes of eating disorders: A systematic review of
the literature. International Journal of Eating Disorders. 2007;40(4):293–309.

8.   Henn AT, Taube CO, Vocks S, Hartmann AS. Body Image as Well as Eating Disorder and
Body Dysmorphic Disorder Symptoms in Heterosexual, Homosexual, and Bisexual
Women. Front Psychiatry. 2019;10:531.

9.   Dolan RJ, Dayan P. Goals and Habits in the Brain. Neuron. 2013;80(2):312–25.

10.   Daw ND, Gershman SJ, Seymour B, Dayan P, Dolan RJ. Model-Based Influences on
Humans' Choices and Striatal Prediction Errors. Neuron. 2011;69(6):1204–15.

11.   Garcia F, Delavenne H, Déchelotte P. Atypical eating disorders: a review. NDS. 2011;67.

12.   Fairburn CG, Bohn K. Eating disorder NOS (EDNOS): an example of the troublesome "not
otherwise specified" (NOS) category in DSM-IV. Behav Res Ther. 2005;43(6):691–701.

13.   Kessler RM, Hutson PH, Herman BK, Potenza MN. The neurobiological basis of binge-
eating disorder. Neuroscience & Biobehavioral Reviews. 2016;63:223–38.

14.   Linardon J, Kothe EJ, Fuller-Tyszkiewicz M. Efficacy of psychotherapy for bulimia
nervosa and binge-eating disorder on self-esteem improvement: Meta-analysis. European
Eating Disorders Review. 2019;27(2):109–23.

15.   Simon JJ, Skunde M, Walther S, Bendszus M, Herzog W, Friederich H-C. Neural signature
of food reward processing in bulimic-type eating disorders. Soc Cogn Affect Neurosci.
2016;11(9):1393–401.

16.   Huys QJM, Browning M, Paulus MP, Frank MJ. Advances in the computational
756   understanding of mental illness. Neuropsychopharmacology. 2021;46(1):3–19.

757   17.   Huys QJM, Moutoussis M, Williams J. Are computational models of any use to psychiatry?
758   Neural Networks. 2011;24(6):544–51.

759   18.   Seriès P, editor. Computational psychiatry: a primer. Cambridge, Massachusetts: The MIT
760   Press; 2020.

761   19.   Raymond JG, Steele JD, Seriès P. Modeling Trait Anxiety: From Computational Processes
762   to Personality. Front Psychiatry. 2017;8.

763   20.   Rocha FF da, Alvarenga NB, Malloy-Diniz L, Corrêa H. Decision-making impairment in
764   obsessive-compulsive disorder as measured by the Iowa Gambling Task. Arq Neuro-
765   Psiquiatr. 2011;69(4):642–7.

766   21.   Valton V, Romaniuk L, Douglas Steele J, Lawrie S, Seriès P. Comprehensive review:
767   Computational modelling of schizophrenia. Neuroscience & Biobehavioral Reviews.
768   2017;83:631–46.

769   22.   Huys QJM, Tobler PN, Hasler G, Flagel SB. Chapter 3 - The role of learning-related
770   dopamine signals in addiction vulnerability. In: Diana M, Di Chiara G, Spano P, editors.
771   Progress in Brain Research. Elsevier; 2014. p. 31–77. (Dopamine; vol. 211).

772   23.   Huys QJ, Pizzagalli DA, Bogdan R, Dayan P. Mapping anhedonia onto reinforcement
773   learning: a behavioural meta-analysis. Biol Mood Anxiety Disord. 2013 Dec;3(1):12.

774   24.   Bernardoni F, Geisler D, King JA, Javadi A-H, Ritschel F, Murr J, et al. Altered Medial
775   Frontal Feedback Learning Signals in Anorexia Nervosa. Biological Psychiatry.
776   2018;83(3):235–43.

777   25.   Chan TWS, Ahn W-Y, Bates JE, Busemeyer JR, Guillaume S, Redgrave GW, et al.
778   Differential impairments underlying decision making in anorexia nervosa and bulimia
779   nervosa: A cognitive modeling analysis: Decision-Maing In Anorexia And Bulimia. Int J
780   Eat Disord. 2014;47(2):157–67.

781   26.   DeGuzman M, Shott ME, Yang TT, Riederer J, Frank GKW. Association of Elevated
782   Reward Prediction Error Response With Weight Gain in Adolescent Anorexia Nervosa. AJP.
783   2017;174(6):557–65.

784   27.   Foerde K, Daw ND, Rufin T, Walsh BT, Shohamy D, Steinglass JE. Deficient Goal-
785   Directed Control in a Population Characterized by Extreme Goal Pursuit. Journal of
786   Cognitive Neuroscience. 2020;1–19.

787   28.   Frank GKW, Reynolds JR, Shott ME, O'Reilly RC. Altered Temporal Difference Learning
788   in Bulimia Nervosa. Biological Psychiatry. 2011;70(8):728–35.

29.   Giannunzio V, Degortes D, Tenconi E, Collantoni E, Solmi M, Santonastaso P, et al.
Decision-making impairment in anorexia nervosa: New insights into the role of age and
decision-making style. European Eating Disorders Review. 2018;26(4):302–14.

30.   Gillan CM, Kosinski M, Whelan R, Phelps EA, Daw ND. Characterizing a psychiatric
symptom dimension related to deficits in goal-directed control. eLife. 2016;5:e11305.

31.   Patzelt EH, Kool W, Millner AJ, Gershman SJ. Incentives Boost Model-Based Control
Across a Range of Severity on Several Psychiatric Constructs. Biological Psychiatry.
2019;85(5):425–33.

32.   Reiter AMF, Heinze H-J, Schlagenhauf F, Deserno L. Impaired Flexible Reward-Based
Decision-Making in Binge Eating Disorder: Evidence from Computational Modeling and
Functional Neuroimaging. Neuropsychopharmacol. 2017;42(3):628–37.

33.   Verharen JPH, Danner UN, Schröder S, Aarts E, van Elburg AA, Adan RAH. Insensitivity
to Losses: A Core Feature in Patients With Anorexia Nervosa? Biological Psychiatry:
Cognitive Neuroscience and Neuroimaging. 2019;1–9.

34.   Voon V, Derbyshire K, Rück C, Irvine MA, Worbe Y, Enander J, et al. Disorders of
compulsivity: a common bias towards learning habits. Mol Psychiatry. 2015;20(3):345–52.

35.   Google Scholar. Computational Psychiatry of Eating Disorders Search - Google Scholar
[Internet]. [cited 2021 Sep 12]. Available from:
https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=%28%22computational+psy
chiatry%22+%22OR%22+%22computational%22+%22OR%22+%22reinforcement+learni
ng%22+%22OR%22+%22reinforcement%22+%22OR%22+%22bayesian%22+%22OR%2
2+%22decision+making%22+%22OR%22+%22decision-
making%22%29+%22AND%22+%28%22anorexia%22+%22OR%22+%22anorexia+nervo
sa%22+%22OR%22+%E2%80%9Cbulimia+nervosa%22+%22OR%22+%22binge%22+%
22OR%22+%22binge+eating%E2%80%9D+OR%22+%22eating%22+%22OR%22+%22b
ulimia%22%29&btnG=

36.   Google Scholar. Computational Psychiatry of Schizophrenia Search - Google Scholar
[Internet]. [cited 2021 Sep 12]. Available from:
https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=%28%22computational+psy
chiatry%22+%22OR%22+%22computational%22+%22OR%22+%22reinforcement+learni
ng%22+%22OR%22+%22reinforcement%22+%22OR%22+%22bayesian%22+%22OR%2
2+%22decision+making%22+%22OR%22+%22decision-
making%22%29+%22AND%22+%28%E2%80%9Cschizophrenia%E2%80%9D%29&btn
G=

37.   Prolific. Prolific | Online participant recruitment for surveys and market research [Internet].
[cited 2020 Aug 4]. Available from: https://www.prolific.co/

38.   Qualtrics. Qualtrics - Leading Experience Management & Survey Software [Internet].
Qualtrics. [cited 2020 Aug 5]. Available from: https://www.qualtrics.com/uk/

39.   Pavlovia. Pavlovia [Internet]. [cited 2020 Aug 5]. Available from: https://pavlovia.org/

40.   PsychoPy. Home — PsychoPy v3.0 [Internet]. [cited 2020 Apr 14]. Available from:
829   https://www.psychopy.org/

830   41.   Faul F, Erdfelder E, Lang A-G, Buchner A. G*Power 3: A flexible statistical power analysis
831   program for the social, behavioral, and biomedical sciences. Behavior Research Methods.
832   2007;39(2):175–91.

833   42.   Garner DM, Olmsted MP, Bohr Y, Garfinkel PE. The Eating Attitudes Test: psychometric
834   features and clinical correlates. Psychological Medicine. 1982;12(4):871–8.

835   43.   Veale D, Eshkevari E, Kanakam N, Ellison N, Costa A, Werner T. The Appearance Anxiety
836   Inventory: Validation of a Process Measure in the Treatment of Body Dysmorphic Disorder.
837   Behavioural and Cognitive Psychotherapy. 2014;42(5):605–16.

838   44.   Foa EB, Huppert JD, Leiberg S, Langner R, Kichic R, Hajcak G, et al. The Obsessive-
839   Compulsive Inventory: Development and validation of a short version. Psychological
840   Assessment. 2002;14(4):485–96.

841   45.   Orbitello B, Ciano R, Corsaro M, Rocco PL, Taboga C, Tonutti L, et al. The EAT-26 as
842   screening instrument for clinical nutrition unit attenders. International Journal of Obesity.
843   2006;30(6):977–81.

844   46.   Bjureberg J, Enander J, Andersson E, Ivanov VZ, Rück C, Fernández de la Cruz L. Sudden
845   Gains in Internet-Based Cognitive Behavior Therapy for Body Dysmorphic Disorder.
846   Behavior Therapy. 2019;

847   47.   Mastro S, Zimmer-Gembeck MJ, Webb HJ, Farrell L, Waters A. Young adolescents'
848   appearance anxiety and body dysmorphic symptoms: Social problems, self-perceptions and
849   comorbidities. Journal of Obsessive-Compulsive and Related Disorders. 2016;8:50–5.

850   48.   Chen H, Mojtahedzadeh N, Belanger MJ, Nebe S, Kuitunen-Paul S, Sebold M, et al.
851   Model-Based and Model-Free Control Predicts Alcohol Consumption Developmental
852   Trajectory in Young Adults: A 3-Year Prospective Study. Biological Psychiatry.
853   2021;89(10):980–9.

854   49.   Decker JH, Otto AR, Daw ND, Hartley CA. From Creatures of Habit to Goal-Directed
855   Learners: Tracking the Developmental Emergence of Model-Based Reinforcement
856   Learning. Psychol Sci. 2016;27(6):848–58.

857   50.   Brown VM, Chen J, Gillan CM, Price RB. Improving the Reliability of Computational
858   Analyses: Model-Based Planning and Its Relationship With Compulsivity. Biological
859   Psychiatry: Cognitive Neuroscience and Neuroimaging. 2020;

860   51.   Stan Development Team. RStan: the R interface to Stan [Internet]. 2020. Available from:
861   http://mc-stan.org/

52. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, et al. Research Domain
Criteria (RDoC): Toward a New Classification Framework for Research on Mental
Disorders. AJP. 2010;167(7):748–51.

53. NIMH. NIMH | About RDoC [Internet]. [cited 2020 Aug 16]. Available from:
https://www.nimh.nih.gov/research/research-funded-by-nimh/rdoc/about-rdoc.shtml

54. Kar P. Partha Kar: Dieting and body shaming. BMJ [Internet]. 2019 [cited 2020 Aug
16];364. Available from: https://www.bmj.com/content/364/bmj.l1222

55. Sujoldzić A, De Lucia A. A cross-cultural study of adolescents--BMI, body image and
psychological well-being. Coll Antropol. 2007;31(1):123–30.

56. Ambwani S, Shippe M, Gao Z, Austin SB. Is #cleaneating a healthy or harmful dietary
strategy? Perceptions of clean eating and associations with disordered eating among young
adults. Journal of Eating Disorders. 2019;7(1):17.

57. Kool W, Cushman FA, Gershman SJ. When Does Model-Based Control Pay Off? O'Reilly
JX, editor. PLoS Comput Biol. 2016;12(8):e1005090.

58. Silva CF da, Hare TA. A note on the analysis of two-stage task results: How changes in task
structure affect what model-free and model-based strategies predict about the effects of
reward and transition on the stay probability. PLOS ONE. 2018;13(4):e0195328.

59. Otto AR, Raio CM, Chiang A, Phelps EA, Daw ND. Working-memory capacity protects
model-based learning from stress. Proceedings of the National Academy of Sciences.
2013;110(52):20941–6.

60. Alexi J, Dommisse K, Cleary D, Palermo R, Kloth N, Bell J. An Assessment of Computer-
Generated Stimuli for Use in Studies of Body Size Estimation and Bias. Frontiers in
Psychology. 2019;10:2390.

61. Ralph-Nearman C, Filik R. Development and validation of new figural scales for female
body dissatisfaction assessment on two dimensions: thin-ideal and muscularity-ideal. BMC
Public Health. 2020;20(1):1114.

62. Hudson GM, Lu Y, Zhang X, Hahn J, Zabal JE, Latif F, et al. The Development of a BMI-
Guided Shape Morphing Technique and the Effects of an Individualized Figure Rating
Scale on Self-Perception of Body Size. European Journal of Investigation in Health,
Psychology and Education. 2020;10(2):579–94.