Locus coeruleus integrity is related to tau burden and memory loss

in autosomal-dominant Alzheimer's disease

Authors:

Martin J. Dahl*[1,2], Mara Mather[2], Markus Werkle-Bergner[1], Briana L. Kennedy[2,3], Samuel Guzman[4], Kyle Hurth[4], Carol A. Miller[4], Yuchuan Qiao[5], Yonggang Shi[5], Helena C. Chui[6], & John M. Ringman[6]

Affiliations:

[1]Center for Lifespan Psychology, Max Planck Institute for Human Development, 14195 Berlin, Germany

[2]Davis School of Gerontology, University of Southern California, 90089 Los Angeles, CA, USA

[3]School of Psychological Science, University of Western Australia, Perth, Australia

[4]Department of Pathology, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

[5]Laboratory of Neuro Imaging (LONI), USC Stevens Neuroimaging and Informatics Institute, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

[6]Department of Neurology, Keck School of Medicine, University of Southern California, 90033 Los Angeles, CA, USA

* Corresponding author:

MJD (dahl@mpib-berlin.mpg.de)

Abstract

Abnormally phosphorylated tau, an indicator of Alzheimer's disease, accumulates in the
first decades of life in the locus coeruleus (LC), the brain's main noradrenaline supply. However,
technical challenges in reliable in-vivo assessments have impeded research into the role of the
LC in Alzheimer's disease.

We studied participants with or known to be at-risk for mutations in genes causing
autosomal-dominant Alzheimer's disease (ADAD) of early onset, providing a unique window
into the pathogenesis of Alzheimer's largely disentangled from age-related factors. Using high-
resolution MRI and tau PET, we revealed lower rostral LC integrity in symptomatic participants.
LC integrity was associated with individual differences in tau burden and memory decline. Post-
mortem analyses in a separate set of carriers of the same mutation confirmed substantial
neuronal loss in the LC.

Our findings link LC degeneration to tau burden and memory in Alzheimer's and
highlight a role of the noradrenergic system in this neurodegenerative disease.

Keywords: Locus coeruleus; autosomal-dominant Alzheimer's disease; noradrenaline; norepinephrine;
tau pathology*;* neurodegeneration; A431E

1. Introduction

Alzheimer's disease (AD) is presently an incurable neurodegenerative disease leading

to dementia with the number of worldwide cases predicted to triple over the next thirty years

(Alzheimer's Disease International, 2019; Canter et al., 2016; World Health Organization,

2004). Post-mortem studies indicate that abnormally phosphorylated tau, an indicator of AD,

begins to appear early in life in the locus coeruleus (LC), the primary source of cortical

noradrenaline (Braak et al., 2011; Ehrenberg et al., 2017; Poe et al., 2020; Stratmann et al.,

2016; Theofilas et al., 2015). With increasing age, abnormal tau appears in a characteristic

topographical sequence in noradrenergic projection targets like the mediotemporal lobe

(referred to as Braak stages, a classification system of the progression of tau spread; Braak et

al., 2011; Chalermpalanupap, Weinshenker, & Rorabaugh, 2017; Stratmann et al., 2016),

suggesting pathology slowly spreads from the LC to its target regions (Braak & Del Tredici,

2016). Consistent with this possibility, using an animal model, Gosh and colleagues

demonstrated that injecting hyperphosphorylated human tau into the rodent LC leads to its

slow spread to other brainstem nuclei and eventually to cortical regions (Ghosh et al., 2019).

Tau deposition is strongly linked to both neural and cognitive decline in Alzheimer's

(Hanseeuw et al., 2019; Jagust, 2018; La Joie et al., 2020). Tau pathology in the LC increases

linearly with the progression of Braak stages while noradrenergic neurons first decrease in

size and then degenerate (Ehrenberg et al., 2017; Kelly et al., 2017; Theofilas et al., 2017).

Indeed, a meta-analysis of twenty-four post-mortem investigations revealed substantial cell

loss in the LC of Alzheimer's patients relative to controls (mean d = 2.28; 95% CI = 2.06–

2.51; Lyness, Zarow, & Chui, 2003), and reduced neuron counts are evident already at

prodromal stages of the disease (Arendt et al., 2015; Kelly et al., 2017). Topographically,

noradrenergic neurodegeneration is most pronounced in rostral and middle segments of the

nucleus that project to the hippocampus and other memory-relevant areas (Ehrenberg et al.,

2017; Lyness et al., 2003). LC degeneration, however, does not only constitute a consequence

of Alzheimer's pathology but also may contribute to its disease development (Marien et al.,

2004; Mather & Harley, 2016; Satoh & Iijima, 2019; Weinshenker, 2018). Studies with

genetically modified animals indicate that abnormally phosphorylated tau in the LC leads to

dysfunctional noradrenergic neuromodulation in memory-relevant brain areas (Ghosh et al.,

2019; Rorabaugh et al., 2017; Weinshenker, 2018) and that tau pathology and LC

degeneration synergistically aggravate neural and behavioral deterioration (Chalermpalanupap

et al., 2018; also see Jacobs, Riphagen, Ramakers, & Verhey, 2019). Specifically,

experimentally decreased noradrenaline levels have been associated with increased tau and

amyloid-β deposition, another hallmark of AD (Chalermpalanupap et al., 2018; Heneka et al.,

2010). Insights into the role of noradrenergic neurodegeneration in the progression of AD are

thus of high clinical significance (Grinberg & Heinsen, 2017), yet *in-vivo* human data are

scarce.

Studies in humans have been impeded by methodological challenges in non-invasive

assessments of the LC due to its small size and its location deep within the brainstem (for

discussions, see Astafiev, Snyder, Shulman, & Corbetta, 2010; Keren, Lozar, Harris, Morgan,

& Eckert, 2009). Advances in high-resolution brainstem magnetic resonance imaging (MRI)

may help the field overcome these hurdles (Betts, Kirilina, et al., 2019; Liu et al., 2017; Sun

et al., 2020). Over the last fifteen years, several MRI sequences have been developed that

reveal the LC as a hyperintense cluster of voxels bordering the fourth ventricle (e.g., Betts,

Cardenas-Blanco, Kanowski, Jessen, & Düzel, 2017; Nakane, Nihashi, Kawai, & Naganawa,

2008; Priovoulos et al., 2017; Sasaki et al., 2006). Keren and colleagues (2015) validated a

MR-based LC imaging sequence by first scanning human post-mortem samples at ultra-high

field strength and afterwards performing histological analyses of the same samples. They

demonstrated that the hyperintensities observed on MRI scans closely co-localize with

noradrenergic cells as identified by tyrosine hydroxylase (the rate-limiting enzyme of

catecholamine synthesis) staining, opening the door for non-invasive LC assessments (Keren

et al., 2015; also see Cassidy et al., 2019 for a postmortem validation in dopaminergic

structures and Betts, Kirilina, et al., 2019 for a discussion of potential contrast mechanisms).

Probing the utility of LC MRI to detect pathology-related changes in the noradrenergic

system *in vivo*, several studies compared patients diagnosed with AD to healthy controls

(Betts, Cardenas-Blanco, et al., 2019; Dordevic et al., 2017; Miyoshi et al., 2013; Takahashi

et al., 2015). In general agreement with earlier post-mortem investigations (Lyness et al.,

2003), the majority of studies observed lower LC MR-intensity, a proxy for the density of

noradrenergic cells (Keren et al., 2015), in the patient groups (but see Miyoshi et al., 2013).

However, all of these studies investigated the by far most common, late-onset type of AD

(>65 years; LOAD). LOAD is considered a heterogeneous, age-related disease that

encompasses not only tau and amyloid pathology but also the aggregation of other proteins

such as α-synuclein, the effects of cerebrovascular ischemia, and other processes (Jagust,

2018; Van Cauwenberghe et al., 2016). Given that LOAD incidence increases exponentially

with age and so is highly correlated with age (Brookmeyer et al., 2007) and large lifespan

studies suggest spatially confined age differences in LC integrity in healthy older adults (Dahl

et al., 2019; Liu et al., 2019), unambiguously distinguishing disease- from age-related

changes on this basis is challenging (Lindenberger et al., 2011). Focusing instead on forms of

AD that begin early in life allows the possibility of distinguishing age- from disease-related

changes in LC integrity (cf. Jacobs, Becker, et al., 2019).

Autosomal dominant Alzheimer's (ADAD) is a rare inherited disease caused by

mutations in genes associated with the generation or accumulation of amyloid-β (Bertram et

al., 2010; Tanzi & Bertram, 2005; Van Cauwenberghe et al., 2016). In contrast to LOAD, it

constitutes a more "pure" tau and amyloid pathology in which cerebrovascular- or other age-

related neuropathological changes are less frequent (Jagust, 2018; Ringman et al., 2016; Spina

et al., 2021) (but see Lee et al., 2016). In ADAD, symptoms develop early in life with a

relatively predictable age at onset (Ryman et al., 2014), thus providing a unique window into

the pathogenesis of Alzheimer's (Bateman et al., 2012; Ringman, 2005). Whether or not there

is MR-identifiable LC degeneration in patients diagnosed with Alzheimer's dementia early in

life is still an open question. Importantly, answering this question could help shed light on

whether LC involvement is a core feature of AD or whether it should be considered more

indicative of a non-Alzheimer's process generally associated with aging, such as the process

outlined in the "primary age-related tauopathy" framework (Crary et al., 2014).

While previous post-mortem research suggests a topographical pattern of

noradrenergic neurodegeneration within the LC in LOAD (Ehrenberg et al., 2017; Lyness et

al., 2003), most current *in-vivo* (MRI) studies lack the spatial specificity to draw comparable

conclusions. That is, they report disease-related differences in MR-indexed LC integrity

(Dordevic et al., 2017; Takahashi et al., 2015) without exploring potential topographical

patterns therein, precluding direct comparisons (but see Betts, Cardenas-Blanco, et al., 2019).

Investigations in healthy participants successfully applied LC MRI to map the spatial extent

of the central noradrenergic nucleus in standard space (Betts et al., 2017; Dahl et al., 2019;

Keren et al., 2009; Liu et al., 2019; Tona et al., 2017; Ye et al., 2021), paving the way for

topographical cross-study comparisons. However, estimates of locus coeruleus' dimensions

and localizations show a sizeable variance across publications (range of agreement: 48–94%;

cf. Dahl et al., 2019; Liu et al., 2019; Ye et al., 2021; or even 1–40% using a different

methodology, Mäki-Marttunen & Espeseth, 2020). Such lack of consensus considerably limits

the replicability of findings and impedes scientific progress.

Thus, the present study pursued two main goals: First, we aimed at improving the

reliability and validity of MR-based LC detection. Leveraging a meta-analytical approach, we

aggregated across previously published maps of the LC (Betts et al., 2017; Dahl et al., 2019;

Keren et al., 2009; Liu et al., 2019; Tona et al., 2017; Ye et al., 2021) to derive a biologically

plausible volume of interest (meta mask) that shows high agreement across investigations.

Second, we aimed at determining whether MR-indexed LC integrity can serve as marker for

noradrenergic degeneration in early-onset ADAD. To this end, we applied the newly

generated meta mask to an independent clinical sample to semi-automatically extract MR-

indexed LC integrity across the rostrocaudal extent of the nucleus (cf. Dahl et al., 2019). We

specifically focused on a sample of participants with or known to be at-risk for rare mutations

in genes causing ADAD (Presenilin-1 protein gene [*PSEN1*]; Amyloid-β Precursor Protein

gene [*APP];* Goate et al., 1991; Murrell et al., 2006; Yescas et al., 2006). In addition, using an

independent sample of banked post-mortem neuropathological specimens, we characterized

changes occurring in the LC in persons dying with the A431E mutation in *PSEN1* relative to

controls. We hypothesized that middle to rostral MR-indexed LC integrity would be lower in

living symptomatic participants relative to healthy matched controls, corroborating earlier

post-mortem findings (in LOAD; Lyness et al., 2003). Similarly, post-mortem LC specimens

were hypothesized to show signs of neurodegeneration in ADAD relative to controls. Finally,

we predicted MR-indexed LC integrity would be associated with cortical tau burden

(Chalermpalanupap et al., 2018), as assessed using positron emission tomography (PET), and

attention and memory performance (Dahl et al., 2019; Elman et al., 2021; Liu et al., 2020).

2. Material and methods

2.1.1. Aggregating across published locus coeruleus maps to derive a high confidence

138   **meta mask**

139   Previous mapping studies used a variety of LC-sensitive MRI sequences (Turbo Spin

Echo [TSE; Dahl et al., 2019; Keren et al., 2009; Tona et al., 2017]; Magnetization Transfer

[MT; Liu et al., 2019; Ye et al., 2021]; Fast Low Angle Shot [FLASH; Betts et al., 2017]) at

different field strength (7 Tesla: Ye et al., 2021; 3 Tesla: all other studies) to visualize the

nucleus. Most investigations were conducted across younger and older adults or in lifespan

samples (but see Tona et al., 2017 and Ye et al., 2021 for studies in younger and older adults,

respectively). After data collection, individual brainstem scans were transformed into a group

or standard space in which a manual (Betts et al., 2017; Liu et al., 2019; Tona et al., 2017) or

semi-automatic (Dahl et al., 2019; Keren et al., 2009; Ye et al., 2021) approach was taken to

segment the LC from surrounding tissue. While some earlier reports (Dahl et al., 2019; Liu et

al., 2019; Ye et al., 2021) include a comparison of a subset of the published masks, to date

there is no systematic evaluation of their agreement. More importantly, no previous study

attempted to resolve the apparent disagreement in LC dimensions and localizations across

publications.

153   In keeping with earlier analyses (Dahl et al., 2019; Liu et al., 2019; Ye et al., 2021),

154 the LC masks noted in Table 1 were used for cross-study comparisons. Before relating

155 different masks to one-another, all volumes were binarized and moved to a common space

156 (MNI-ICBM 152 linear space, 0.5 mm resolution), if necessary. In standard space, first an

157 unthresholded aggregate mask was generated using the following formula:

158
$$LC_{\text{aggregate}} = \frac{1}{n}\sum_{i=1}^{n}(v)$$

159   Whereby n denotes the number of masks included in the analyses ($n = 6$) and v

denotes the value for a given voxel (either zero [non-LC tissue] or one [LC]). That is, at every

voxel the individual binarized masks were averaged, resulting in an aggregate mask

($LC_{aggregate}$; see Figure 1). The generated volume, $LC_{aggregate}$, showed a value range between 0

and 1 (complete agreement that a voxel does not/does belong to the LC, respectively) in steps

of $\frac{1}{n}$. Please note that we decided against weighing the contribution of the individual masks to

the $LC_{aggregate}$ by sample size as two masks (Dahl et al., 2019; Liu et al., 2019) account for

more than 80% of the total number of participants (see Table 1). Hence, the aggregate would

be strongly biased in favor of a subset of masks that are themselves highly spatially congruent

(agreement 94%; Liu et al., 2019).

169   In a second step, the aggregate mask was pruned by removing low-agreement voxels

170   until a majority vote was achieved. That is, only areas with agreement values > 50% were

171   retained (i.e., more than $\frac{1}{2}n$ masks voted for a voxel to belong to the LC). Essentially, the

172   described mask generation process is comparable to label fusion with majority voting

173   (Rohlfing et al., 2004; Sabuncu et al., 2010; Wang et al., 2013) .

174   Finally, to obtain a biologically more plausible volume, the pruned aggregate mask

was smoothed with a 0.1 mm full width at half maximum (FWHM) kernel using SPM12

(Penny et al., 2007) in Matlab (The MathWorks Inc., Natick, MA, USA). Voxels exceeding a

threshold of 0.05 were considered part of the LC in the meta mask ($LC_{metaMask}$; see Figure 1).

The resulting mask had a rostrocaudal extent of 14 mm and a volume of 74 mm³ (592 voxels),

corresponding well with previous post-mortem findings (Fernandes et al., 2012; Tona et al.,

2017). We share the generated meta mask with the scientific community via osf.io/sf2ky/

Table 1: Locus coeruleus masks included in comparisons

Publication	MRI acquisition	Sample	Locus coeruleus segmentaion	Mask source	Mask version
Betts et al., 2017	3 T; FLASH	N = 82; YA; OA	Manual	https://doi.org/10.1016/j.neuroimage.2017.09.042	--
Dahl et al., 2019	3 T; TSE	N = 294; YA; OA	Semi-automatic	https://www.mpib-berlin.mpg.de/LC-Map	--
Keren et al., 2009	3 T; TSE	N = 44; YA–OA	Semi-automatic	http://www.eckertlab.org/LC	1 SD
Liu et al., 2019	3 T; MT	N = 605; YA–OA	Manual	https://osf.io/r2bwk	--
Tona et al. 2017	3 T; TSE	N = 17; YA	Manual	http://www.nitrc.org/projects/prob_lc_3t	[1]
Ye et al., 2021	7 T; MT	N = 53; OA	Semi-automatic	https://www.nitrc.org/projects/lc_7t_prob	5% prob.[2]

Note: [1] *The masks provided by Tona and colleagues were averaged across scan sessions and*
segmentations. [2]The mask by Ye and colleagues was linearly and non-linearly co-registered
from MNI-ICBM 152 non-linear 2009b to MNI-ICBM 152 linear space (0.5 mm resolution)
using Advanced Normalization Tools (antsRegistration). MNI-ICBM 152 linear was chosen
as target space as all but one masks share this space. For co-registration, the standard brains
of the respective spaces were aligned and the transformations were applied to the mask file
with a nearest neighbor interpolation. The transformations from MNI 2009b non-lin ⇔ MNI
linear are shared via osf.io/sf2ky/; T, Tesla; FLASH, Fast Long Angle Shot; TSE, Turbo Spin
Echo; MT, Magnetization Transfer; YA, Younger Adults; OA, Older Adults; YA–OA, lifespan
sample including younger, middle-aged and older adults; SD, standard deviation; prob.,
probability.

2.1.2 Evaluation of previously published locus coeruleus masks

Once a high-agreement volume of interest ($LC_{metaMask}$) was established, published

locus coeruleus masks were compared regarding their specificity and sensitivity. That is, we

evaluated (1) how many voxels of a given mask fell within the $LC_{metaMask}$ relative to the total

number of voxels in that mask and (2) how many voxels of the $LC_{metaMask}$ were included in

each mask (relative to the total number of voxels in the $LC_{metaMask}$). The mean of the quality

metrics (specificity, sensitivity) was taken as indicator for the accuracy of a mask.

2.2. Application of locus coeruleus meta mask to an independent clinical sample

2.2.1. Study design and participants

To determine whether the $LC_{metaMask}$ can detect clinically significant differences in LC

integrity, eighteen participants with or known to be at-risk for mutations in genes associated

with ADAD (*PSEN1* or *APP*) were investigated (34.7 ± 10.1 years [mean ± standard

deviation]; 9♀; see Table 2; data collection was interrupted by the COVID-19 pandemic).

Participants' consent was obtained according to the Declaration of Helsinki. The study was

approved by the ethical committee of the University of Southern California. Data were

acquired as part of a study employing the Human Connectome Protocol (HCP) in genetic

subtypes of AD (NIH U01AG051218, PI: Ringman; for more details, see

adrc.usc.edu/scientists-researchers/ [Estudio de la enfermaded de Alzheimer en Jalisciences;

EEAJ]). Genotyping confirmed mutations in nine of the eighteen participants (the A431E

substitution in *PSEN1*, n = 6; the I180F substitution in *PSEN1*, n = 1; and the V717I

substitution in *APP*, n = 2; Goate et al., 1991; Murrell et al., 2006; Yescas et al., 2006), of

which four were symptomatic. We defined symptomatology based on the Washington

University Clinical Dementia Rating Scale [CDR] (see Table 2). Symptomatic participants

showed CDR values greater than zero, while all participants with a value of zero were

considered cognitively normal (n = 14).

Participants underwent 3 T MRI, flortaucipir PET, and cognitive testing. LC MR-

intensity, a non-invasive proxy for neuronal density (Keren et al., 2015), was semi-

automatically extracted from high-resolution brainstem scans (Dahl et al., 2019). LC ratios—

that is, a ratio of peak LC intensity standardized to a pontine reference (Betts, Kirilina, et al.,

2019; Liu et al., 2017)—were computed across the rostrocaudal extent of the nucleus.

Standard uptake value ratios (SUVR) were calculated from partial volume corrected

flortaucipir PET data using cerebellar gray matter as the reference based on the PETSurfer

tool from FreeSurfer (Greve et al., 2014). SUVR images were mapped to the cortical surface

which was parcellated into 36 regions of interest (Desikan et al., 2006). Univariate

associations between LC ratios, clinical variables, cognitive status, and tau burden were

evaluated using non-parametric Wilcoxon rank sum tests and bootstrapped Spearman's

correlations. For Wilcoxon rank sum tests, effect sizes (r) are calculated by dividing the test

statistic (Z) by the square root of the sample size (n; (Rosenthal, 1991)). The multivariate

pattern between LC ratios, cortical tau burden and cognitive decline was evaluated using a

partial least squares correlation (see below; Krishnan, Williams, McIntosh, & Abdi, 2011;

McIntosh & Lobaugh, 2004). For bivariate correlations and partial least squares correlations,

we provide the rho values as effect size measure. Two-tailed statistical tests were used with an

alpha level of $p < 0.05$. The applied statistical tests did not include covariates. Participants

with missing data were excluded from the respective analyses (see below). We did not

conduct a formal power calculation, given that there was no available prior evidence on

the studied phenomena.

Table 2: Descriptive statistics for imaging subgroups

	ADAD-control	ADAD-presymptomatic	ADAD-symptomatic
n	9	5	4
Mutation	0/9	5/5	4/4
Mutation type		*PSEN1* (A431E; 3/5) *PSEN1* (I180F; 1/5) *APP* (V717I; 1/4)	*PSEN1* (A431E; 3/4) *APP* (V717I; 1/4)
Diagnosis	0/9	0/5	4/4
CASI	95.3	95.0	74.9
(std, range)	(1.8; 92–97.7)	(4.2; 90–99)	(11.8; 62.8–86.8)
CDR	0	0	0.75
(std, range)	(0, 0)	(0, 0)	(0.3; 0.5–1)
Age	33.4	30	43.5
(std, range)	(12.2; 19–58)	(4.5; 22–33)	(1.3; 42–45)
Gender	5 male 4 female	1 male 4 female	3 male 1 female

Note: ADAD, autosomal-dominant Alzheimer's disease; ADAD-control: control participants
not carrying the mutation, but genetically related to mutation carriers; ADAD-
presymptomatic: presymptomatic mutation carriers (Washington University Clinical
Dementia Rating Scale [CDR] scores of 0); ADAD-symptomatic: symptomatic mutation
carriers; std. standard deviation; CASI, Cognitive Abilities Screening Instrument (Teng et al.,
1994); *PSEN1*: Presenilin-1 protein (gene); *APP*: Amyloid-β Precursor Protein (gene)
250

2.2.2. Assessment of imaging data

Structural MRI data were collected employing a 3 T Siemens Prisma scanner with a

32-channel head coil following a Human Connectome Project imaging approach (HCP; cf.

humanconnectome.org/). PET data were acquired on a 64-slice Biograph PET/CT Siemens

scanner at the USC PET Imaging Sciences Center. Only those sequences used in the current

analyses are described below.

A three-dimensional (3D) T1-weighted magnetization prepared gradient-echo

(MPRAGE) sequence with a duration of 6:38 min and the following parameters was applied:

repetition time (TR) = 2400 ms; echo time (TE) = 2.22 ms; inversion time (TI) = 1000 ms;

flip angle = 8°; bandwidth = 220 Hz/Px; dimensions = 208 × 300 × 320; isometric voxel size

= 0.8 mm^3. Based on this whole-brain (MPRAGE) sequence, a high-resolution, two-

dimensional (2D) T1-weighted TSE sequence was aligned perpendicularly to the plane of the

brainstem. Acquisition of the TSE sequence took 1:53 min, and the following parameters

were used: TR = 750 ms; TE = 10 ms; flip angle = 120°; bandwidth = 285 Hz/Px; dimensions

= 512 × 512 × 11; anisometric voxel size = 0.43 × 0.43 × 3.5 mm^3 (mimicking the elongated

shape of the locus coeruleus). Each TSE scan consisted of eleven axial slices.

PET data were collected within the period of 90–120 min after the injection of the

compound 18F-AV-1451 (total dose: 344 MBq). Low dose computerized tomography (CT)

scans were acquired prior to the PET scans for attenuation correction. The dynamic PET scans

included six frames of five-minute duration, with the dimensions of 168 × 168 × 56 and a

voxel size of 1.5 × 1.5 × 4 mm^3. The six dynamic scans were rigidly aligned to the first one

and averaged to create a static image for SUVR-based analysis.

2.2.3. Semi-automatic locus coeruleus intensity assessment using meta mask

Leveraging the generated $LC_{metaMask}$ (see above), locus coeruleus intensity was semi-

automatically extracted using a previously validated pipeline (Dahl et al., 2019; also see Ye et

al., 2021 for a comparable approach). In short, individual whole brain and brainstem scans

were iteratively aligned across participants using a template-based procedure implemented in

Advanced Normalization Tools (version 2.1; ANTs; Avants et al., 2011; Avants, Tustison, &

Song, 2009), and subsequently transformed to standard space (MNI-ICBM 152 linear, 0.5

mm; see Dahl et al., 2019 for a step-by-step description of the standardization). In standard

space, individual brainstem scans were masked using the high-confidence $LC_{metaMask}$ to

remove non-LC tissue (using SPM12 in Matlab). To allow inter-participant comparisons of

LC data, a normalization of the arbitrarily scaled MR-intensity values is required (Betts,

Kirilina, et al., 2019). Thus, brainstem scans were additionally masked using a reference

volume of interest positioned in the central pontine white matter, as previously suggested (Ye

et al., 2021; dimensions: 4 × 4 mm in plane, following the $LC_{metaMask}$ with a constant distance

of 8.5 mm [y-dimension] along the rostrocaudal axis; see Figure 3). Within the masked

brainstem scans, we then fully automatically searched for voxels of brightest intensity in the

LC and reference regions. Next, individual, spatially-resolved LC intensity ratios were

computed for each slice using the following formula (Dahl et al., 2019; Liu et al., 2017):

293
$$LC_{Ratio} = \frac{\max(LC) - \max(Ref)}{\max(Ref)}$$

Where max(*LC*) denotes the peak intensity for a given slice in the LC volume of interest and

max(*Ref*) indicates the peak intensity in the reference region. After extraction, ratios of the left

and right hemisphere were averaged for further analyses to obtain more stable intensity

estimates. Finally, the peak ratio across the rostrocaudal LC extent was calculated as proxy for

overall locus coeruleus integrity.

2.2.4. Assessment of cortical tau burden

To evaluate the association of MR-indexed LC integrity to hallmarks of AD, we

calculated cortical tau burden from flortaucipir PET data.

First, each participant's whole brain MR data (MPRAGE) was preprocessed using a

HCP pipeline (version 3.27; Makropoulos et al., 2018). Next, the cortical surface was

reconstructed and parcellated into thirty-six anatomical regions of interest using Freesurfer

6.0 (Dale et al., 1999; Desikan et al., 2006). Flortaucipir PET images were then co-registered

within participants to whole brain (MPRAGE) native space. The Muller-Gartner (MG)

method was used for partial volume correction of the PET data, as implemented in PETSurfer

(Greve et al., 2016). Standard uptake value ratios (SUVR) were calculated using cerebellar

gray matter as reference region. Finally, partial-volume corrected SUVR images were mapped

to the cortical surface and parcellated into thirty-six regions of interest (Desikan et al., 2006).

For further analyses, each region's SUVR data were averaged across hemispheres to obtain

more stable estimates of cortical tau burden. Tau PET data were not available for two

participants.

2.2.5. Association of locus coeruleus intensity and cortical tau burden

We employed a multivariate statistical approach to assess the relation between MR-

indexed locus coeruleus integrity and tau pathology across the cortex. Specifically, using a

Partial Least Squares Correlation (PLSC; Krishnan et al., 2011; McIntosh & Lobaugh, 2004)

we estimated a latent tau pathology profile that is maximally related to LC integrity (cf.

Keresztes et al., 2017; Muehlroth et al., 2020).

First, a between-person Pearson's correlation matrix (R) was computed between the

overall LC ratios (i.e., a vector Y_{LC} with $n_{\text{Participants}} \times 1$ [overall intensity ratios]) and the tau

SUVR data (i.e., a matrix X_{Tau} with $n_{\text{Participants}} \times 36$ [regions of interest]). This correlation

matrix ($R_{LC,Tau} = Y_{LC}^{\text{T}} X_{Tau}$) was then decomposed into three matrices using a singular value

decomposition (SVD; SVD($R_{LC,Tau}$) = USV^{T}). Whereby U refers to a left singular vector of LC

weights, V^{T} denotes a right singular matrix of regional tau weights, and S is a diagonal matrix

of singular values. Based on this matrix decomposition, a single latent variable was extracted.

The latent variable optimally expresses (in a least squares sense) the pattern of interindividual

differences in regional tau burden that shares the largest amount of variance with

interindividual differences in overall LC intensity ratios. The statistical strength of the

extracted pattern (i.e., the latent variable) was evaluated using a permutation test (by

randomly re-ordering the observations in X_{Tau} while leaving Y_{LC} unchanged and re-calculating

the SVD; $n_{\text{Permutations}} = 10,000$). Subsequently, the reliability of the contribution (i.e., weights

[V^{T}]) of individual cortical regions, that is, the columns in X_{Tau}, to this latent variable was

determined using a bootstrapping procedure ($n_{\text{Bootraps}} = 10,000$). A ratio of the region-specific

weights (V^{T}) and their corresponding bootstrapped standard errors provided bootstrap ratios

(BSR) that can be interpreted akin to Z-scores. Finally, multiplying participants' regional

pattern of tau pathology (X_{Tau}) with each regions' contribution to the latent variable (i.e., its

weight, V^{T}) yielded a summary measure, reflecting participants' LC-related tau pathology.

2.2.6. Cognitive assessments

Participants completed a series of standardized neuropsychological tests to assess

memory and attention. The cognitive assessment included screening instruments (the

Montreal Cognitive Assessment [MoCA, Nasreddine et al., 2005]; the Cognitive Abilities

Screening Instrument [CASI; Teng et al., 1994], and the Washington University Clinical

Dementia Rating Scale [CDR; measure: sum of boxes; cf. O'Bryant et al., 2008]). In addition,

attention (Digit-Symbol-Substitution Test [DSST; Wechsler, 1981]; Digit Span [measure:

backward report; Wechsler, 1997], Trail Making Test [measure: part B; Tombaugh, 2004]) and

memory performance were assessed (the Spanish English Verbal Learning Test [SEVLT;

measures: average performance over learning trials and delayed recall; González, Mungas, &

Haan, 2002], a learning and memory test akin to the Rey Auditory Verbal Learning Test

[RAVLT] in which Spanish and English versions were simultaneously developed; Benson

Complex Figure Test [measure: delayed recall; cf. Possin, Laluz, Alcantar, Miller, & Kramer,

2011]; and the Craft Story Test [delayed recall; Craft et al., 1996]). When available, for

further analyses individual test scores were transformed to Z-scores using published

normative data. Trail Making Test data were not available for one participant.

A partial least squares correlation (PLSC; Krishnan et al., 2011; McIntosh & Lobaugh,

2004) was applied to capture the multivariate association between MR-indexed LC integrity

and cognitive performance, using the same methods as described above.

2.3.1 Neuropathological assessment

To characterize the pathology underlying the LC signal on MRI in ADAD, we

examined the LC in an independent sample of four persons dying with the A431E mutation in

PSEN1 in comparison to five age-matched neurologically normal controls. In short, paraffin-

embedded sections of the pons were cut and stained with hematoxyin and eosin (H&E), glial

fibrillary acid protein (GFAP), AT8 antibodies for neurofibrillary pathology, 4G8 antibodies

for amyloid-β, and anti-alpha synuclein antibodies for Lewy Body pathology. Qualitative

comparisons were made between patients and controls.

2.3.2. Histopathology methods

After 1-month fixation, tissue areas were dissected and paraffin-embedded and serially

sectioned into 5 µm-thick sections using a microtome (Microm International Waldorf,

Germany). Brain sections were counterstained with H&E and reagents were purchased from

Sigma-Aldrich (St. Louis, MO, USA). Sections adjacent to H&E were immunostained using

the Leica Bond RXm™ automated staining processor (Leica Biosystems, Buffalo Grove, IL,

USA). Tissue sections were stained using the Bond Polymer Refine AFP (Red) Detection

System for all Abs except GFAP Ab and NeuN Ab which used Bond Polymer Refine DAB

Detection System (Leica Biosystems). Antigen retrieval was performed with Epitope Retrieval

Solution 1 (ER1, pH 6) for 20 min except for GFAP Ab which was for 10 min. Sections were

then incubated with amyloid-beta (4G8, BioLegend, SIG #800701, 1:4000) for 30 min, TAR

DNA binding protein (TDP-43, Sigma, T11705; 1:1250) for 15 min, phosphor-tau (AT8,

MN1020, Thermo Scientific, USA; 1:1000 for 15 min; GFAP Rb pAb, Chemicon, Temecula

AB5804, CA, USA; 1;2500) for 15 min; (NeuN, AbCAM, ab104224, USA; 1:800) for 30m. a-

Synuclein Rb pAb, Chemicon, AB5038, 1:2000 for 15 min.

Staining with 4-repeat isoform tau (RD4, Cosmo Bio Co. LTD. Inspiration of Life

Science, Cat# TIP-4RT-P01; 1:3000) and 3-repeat isoform tau (RD3, 05-803, Millipore, USA;

1:5000) required manual pretreatment outside autostainer. Sections heated in a pressure cooker

at 125° in citrate buffer (CC2, pH 6) for 90 min and treated with 88% Formic Acid for 10 min.

Sections were then loaded in autostainer with Bond Polymer Refine Red Detection System for

15 min. Images were obtained with a Nikon Eclipse E400 brightfield microscope (Nikon

Instruments Inc, Melville, NY, USA). All stains were performed at Children's Hospital of Los

Angeles.

2.4. Data and code availability

The here described LC meta mask as well as pontine reference mask is available for

download on an Open Science Framework repository (osf.io/sf2ky/). All sources for

previously published masks are listed in Table 1. For access to raw imaging and clinical data,

please refer to the HCP (https://nda.nih.gov/ccf). The customized code used for these analyses

is available from the corresponding authors upon request.

3.   Results

3.1.1. Aggregating across published locus coeruleus maps yields high confidence meta

mask

Previously published masks of the LC (Betts et al., 2017; Dahl et al., 2019; Keren et al.,

2009; Liu et al., 2019; Tona et al., 2017; Ye et al., 2021) were averaged voxel-per-voxel to

generate an unthresholded overlay (*LC_{aggregate}*). The *LC_{aggregate}* demonstrated a core of voxels

that, across publications, were consistently judged to belong to the LC (see dark-blue areas in

Figure 1), surrounded by a cloud of ambiguous, low-agreement voxels (see yellow areas in

Figure 1). Topographically, consensus was lowest in most rostral and caudal areas (see inlay in

Figure 1).

Pruning low-agreement voxels yielded a high-confidence LC volume of interest

(*LC_{metaMask;}* available for download via: osf.io/sf2ky/) that matches the dimensions reported in

post-mortem investigations (Fernandes et al., 2012) and allows assessment of the accuracy of

previously published masks.

417
Figure 1. Aggregating across published locus coeruleus (LC) maps yields high confidence meta mask.
Left. Unthresholded voxel-by-voxel average across published LC masks. Areas with voxels that are
consistently judged to belong to the LC are shown in dark blue while low agreement areas are shown in
yellow. **Inlay**. Agreement across masks (averaged across voxels in-plane) is lowest in most rostral and
caudal regions. **Middle**. Pruning low agreement areas isolates a core of high confidence voxels. **Right**.
Smoothing and filtering the thresholded overlay yields a biologically plausible LC volume of interest
(rostrocaudal extent of 14 mm; volume of 74 mm³ [592 voxels]), corresponding well with earlier post-
mortem reports (Fernandes et al., 2012). Sl. #, slice number in MNI 152 lin (0.5 mm) space; The LC
meta mask is available for download via: osf.io/sf2ky/).

3.1.2. Accuracy of previously published locus coeruleus masks

Once a high-confidence LC volume of interest ($LC_{metaMask}$) was generated, published

masks were evaluated regarding their specificity and sensitivity. That is, we assessed (1) how

many voxels judged by a given mask as belonging to the locus coeruleus were part of the

$LC_{metaMask}$, and (2) what percentage of the $LC_{metaMask}$ was included in each individual mask.

The mean across specificity and sensitivity was taken as accuracy measure to rank the masks

(see Figure 2).

The masks that were computed based on the largest samples (cf. Table 1; Dahl et al.,

2019; Liu et al., 2019) demonstrated the highest accuracy. Concerning specificity and

sensitivity, the two masks appeared almost as mirror images. That is, the volume of interest

published by Liu and colleagues evinced a high sensitivity at the cost of a relatively lower

specificity (i.e., it included also non-LC voxels), while the situation was reversed for the mask

by Dahl and colleagues (in line with it mapping peak LC coordinates; Dahl et al., 2019).

Notably, the two masks themselves show a high agreement (94%; Liu et al., 2019). Ye and

colleagues tested a smaller sample, albeit with a higher field strength (7 T) to generate their

volume of interest which appears to strike a balance between sensitivity and specificity. None

of the tested masks, however, exceeded an accuracy rating of $\frac{2}{3}$, indicating that, by

aggregating across publications, the $LC_{metaMask}$ forms a volume of interest that conveys

valuable information (i.e., is not redundant with previous masks).

Figure 2. Accuracy of previously published locus coeruleus (LC) masks. Binarized masks in standard space (MNI 152, lin; top row) are ranked based on their accuracy (mean across specificity and sensitivity). Specificity expresses the percentage of a given mask that is included in the *LCmetaMask* whereas sensitivity denotes the percentage of the *LCmetaMask* that is part of each mask. For access to the individual masks, please refer to Table 1. Note that the relative position of the individual masks (top row) corresponds to their position in MNI 152 linear space; x/y/z axes are not displayed for clarity.

3.2.1. Locus coeruleus meta mask captures high intensity voxels in independent clinical sample

We aligned and pooled across brainstem scans of participants with or known to be at-risk for ADAD mutations (cf. Dahl et al., 2019). On the group level, we observed a cluster of hyperintense voxels bordering the lateral floor of the fourth ventricle (Betts, Kirilina, et al., 2019). This LC-related hyperintensity was accurately captured by the $LC_{metaMask}$ while excluding more medial, non-LC related high intensity voxels (see Figure 3). Taken together, this indicates that the $LC_{metaMask}$ can be applied to independent datasets to reliably extract MR-indexed integrity across the rostrocaudal axis.

Figure 3. LC-related hyperintensity is accurately captured by meta mask. **Upper row**. Axial and coronal view of brainstem template overlaid on standard MNI brain. **Inlays**. The area surrounding the fourth ventricle is depicted with (1) a greyscale colormap and (2) a black-to-white-to-red colormap to highlight hyperintense areas. (3) LC-related hyperintensities closely overlap with the LC meta mask (blue overlay; white overlay, pontine reference region). **Lower row**. 3-D representation of (**a**) the LC meta mask, (**b**) above-average intensity voxels, and (**c**) their overlap from a front and back view. Sl. #, slice number in MNI 152 linear (0.5 mm) space; LC, locus coeruleus. Right hemisphere is plotted on the left.

3.2.2. Lower MR-indexed locus coeruleus integrity in autosomal-dominant Alzheimer's disease

Comparing LC intensity ratios—an *in-vivo* proxy for the integrity for the structure (Keren et al., 2015)—across cognitively normal and symptomatic participants, we observed reliably lower intensity in the patient group (Wilcoxon rank sum test, $Z = -2.177$; $p = 0.03$; effect size (r) = -0.513; see Figure 4; mean (SD)$_{symptomatic}$ = 0.123 (0.051); mean (SD)$_{cog.\ normal}$ = 0.204 (0.05)). This difference was most pronounced in middle–rostral segments of the nucleus that project to the mediotemporal lobe (Wilcoxon rank sum test of group differences in rostral vs. caudal segment; $Z = -2.816$; $p = 0.005$; effect size (r) = -0.532; see Figure 4). Among carriers of ADAD mutations, closer proximity to the mutation-specific median age of dementia diagnosis—termed adjusted age—was associated with lower LC ratios (mean correlation coefficient across 1,000,000 bootstraps: *rho* = -0.671; $p = 0.024$; see Figure 4). Taken together, corroborating earlier post-mortem work (in LOAD; Lyness et al., 2003), we observed lower MR-indexed LC integrity in middle–rostral segments of the nucleus, potentially indicating noradrenergic neurodegeneration.

Figure 4. Lower LC intensity ratios in autosomal-dominant Alzheimer's disease (ADAD). Cognitively normal (n = 14) participants are compared against mutation carriers that show cognitive impairments based on the Washington University Clinical Dementia Rating Scale (i.e., CDR > 0; n = 4). **a.** Symptomatic participants show lower overall LC intensity ratios relative to cognitively normal controls ($Z = -2.177$; $p = 0.03$). **b.** Group differences are most

pronounced in middle–rostral segments of the LC (plotted on the y-axis; $Z = –2.816$; $p =$

0.005; shaded areas indicate ± 1 standard error of the mean [SEM]). **c**. Closer proximity to the

mutation-specific median age of dementia diagnosis (adjusted age) is associated with lower

overall LC ratios (mean correlation coefficient across 1,000,000 bootstraps: *rho* = –0.671; *p* =

0.024). Note that in panel **c** only confirmed mutation carriers (n = 9) with and without

cognitive symptoms are included (black and red markers, respectively). **d**. LC-MRI of

cognitively normal and symptomatic participants (four per group) were standardized and

averaged. The crosshair highlights lower LC intensity in symptomatic participants. LC, locus

coeruleus; Cog. normal, cognitively unimpaired participants.

3.2.3. MR-indexed locus coeruleus integrity is associated with cortical tau burden

Previous animal research suggests that noradrenergic neurodegeneration may

exacerbate neural decline and contribute to Alzheimer's pathogenesis (Chalermpalanupap et

al., 2018; Rorabaugh et al., 2017). Thus, we next tested whether LC intensity would be

associated with cortical tau burden, a hallmark of AD.

Across cortical regions (Desikan et al., 2006), symptomatic participants demonstrated

higher flortaucipir SUVR (see Figure 5). Importantly, leveraging a multivariate statistical

approach (partial least squares correlation [PLSC]; Krishnan et al., 2011), we revealed a

topographical pattern of LC-related tau pathology ($p = 0.037$; see Figure 5). That is, we

extracted a latent variable (latent PLS score) that optimally captures the multivariate

association between participants' LC intensity and regional tau burden ($r = 0.54$ [95%

confidence interval (CI): 0.167, 0.806]; lower LC intensity was associated with higher

flortaucipir SUVR). Especially tau pathology in occipito-temporo-parietal regions contributed

to this latent variable, as indicated by reliable bootstrap ratios (BSR; < -3; see Figure 5; BSR

can be interpreted akin to Z-values; please note that we opted for a conservative threshold [–

3] due to the large number of cortical ROI). For associations of LC intensity and regional tau

burden, see Table S1 in the supplementary information. In sum, in line with research in

genetically modified animals (Chalermpalanupap et al., 2018; Rorabaugh et al., 2017), we

observed a prominent association between in-vivo proxies of LC integrity and tau pathology.

Figure 5. Lower LC intensity is associated with tau burden in posterior brain regions. **a.** Symptomatic participants show higher flortaucipir standardized uptake value ratios (SUVR) across most cortical regions relative to cognitively normal controls (missing PET data for two participants [dark blue bars]). **b.** Contribution of individual cortical regions of interest to a partial least squares (PLS) latent variable indexing LC-related tau pathology (Bootstrap ratios < −3 are considered reliable). **c.** Scatter plot depicting the relation between latent PLS scores and LC intensity ($p = 0.037$; $r = 0.54$ [95% CI: 0.167, 0.806]). LC, locus coeruleus; ADAD, autosomal-dominant Alzheimer's disease.

3.2.4. MR-indexed locus coeruleus integrity is associated with cognitive deficits

Finally, we set out to test the behavioral relevance of our *in-vivo* proxy of LC integrity. Noradrenaline release from the LC is strongly implicated in the modulation of attentional and mnemonic processing (Berridge & Waterhouse, 2003; Dahl et al., 2020; T.-H. Lee et al., 2018; Sara, 2009). Accordingly, recent imaging studies reported positive associations between LC intensity and cognition in healthy aging (Dahl et al., 2019; Hämmerer et al., 2018; Liu et al., 2020). However, whether this relation extends to ADAD of young onset is currently unknown.

Across a range of cognitive tasks, we detected worse cognitive abilities and higher dementia symptoms in the ADAD group relative to controls (see Figure 6). Reliable group differences (Wilcoxon rank sum test; $Z < −1.96 \mid Z > 1.96$) were observed for screening instruments (the Montreal Cognitive Assessment [MoCA, Nasreddine et al., 2005]; the Cognitive Abilities Screening Instrument [CASI; Teng et al., 1994], and the Washington

University Clinical Dementia Rating Scale [CDR; measure: sum of boxes; cf. O'Bryant et al.,

2008]). In addition, symptomatic participants performed significantly worse on the Spanish

English Verbal Learning Test (SEVLT; measures: average performance over learning trials

and delayed recall; González, Mungas, & Haan, 2002), in line with reports of verbal learning

tests to convey information about participants' current and future cognitive status (Albert et

al., 2001; Belleville et al., 2017; Moradi et al., 2017; Schoenberg et al., 2006).

We applied a partial least squares correlation (PLSC) to isolate the pattern of cognitive

impairment reliably linked to MR-indexed LC integrity. Our analyses isolated a single latent

variable ($p = 0.016$) that optimally expresses the multivariate association between

participants' LC intensity and cognitive performance ($r = 0.728$ [95% CI: 0.408, 0.888]). All

cognitive tasks that were sensitive to distinguish symptomatic from cognitively normal

participants contributed reliably to this latent variable (bootstrap ratios [BSR]< –1.96 | BSR >

1.96; see Figure 6). To conclude, across a range of neuropsychological tests, higher MR-

indexed LC integrity was observed in participants with unimpaired cognitive performance.

Figure 6. Lower LC intensity is associated with cognitive deficits. **a**. Symptomatic
participants show lower cognitive performance as indicated from screening instruments for
mild cognitive impairment and dementia (Clinical Dementia Rating Scale, sum of boxes
[CDR]; Cognitive Abilities Screening Instrument [CASI]; Montreal Cognitive Assessment
[MoCA]) as well as the Spanish English Verbal learning task (SEVLT). **b**. Contribution of
individual neuropsychological tests to a partial least squares (PLS) latent variable indexing
LC-related cognitive deficits (Bootstrap ratios < –1.96 and > 1.96 are considered reliable). **c**.
Scatter plot depicting the relation between latent PLS scores and LC intensity ($p = 0.016$; $r =$
0.728 [95% CI: 0.408, 0.888]). LC, locus coeruleus; ADAD, autosomal-dominant
Alzheimer's disease. For references to the cognitive tasks, please refer to the main text.

3.3.   Neuropathological findings in the locus coeruleus

Descriptions of the neuropathological changes occurring in the brainstem in ADAD

are scarce. To characterize the pathology underlying the LC signal on MRI in ADAD, LC

histopathology was compared between four carriers of the A431E mutation in *PSEN1* (aged

39–59 years, 2 ♀), including one person homozygous for the A431E mutation (Parker et al.,

2019), and five neurologically normal controls (aged 37–58 years, 2 ♀). All A431E mutation

carriers demonstrated severe loss of LC neurons and background astrogliosis by GFAP

relative to controls. The remaining LC neurons from the A431E mutant carriers demonstrated

a severe loss of neuromelanin and reactivity (see Figure 7). AT8 staining showed neuritic

changes in all A431E mutation carriers with three of four showing neurofibrillary tangle

formation. On staining with 4G8, three mutation carriers showed diffuse extracellular neuritic

and coarse grained or compact plaques (see Figure 8). Two of five controls showed rare

coarse grained extracellular plaques. Cotton-wool amyloid plaques are large ball-like

extracellular plaques that displace nearby structures and are commonly seen in autosomal

dominant forms of Alzheimer's disease. The A431E homozygote case has a higher density of

these plaques throughout the brain and brainstem relative to the heterozygote cases. Alpha

synuclein staining did not reveal any Lewy body pathology in any samples. Immunostaining

using antibodies against 4-repeat tau (RD4) and 3-repeat-tau (RD3) showed positivity for both

isoforms.

Figure 7. Neuropathological findings in the locus coeruleus of a 45–50 year old woman
heterozygous for the A431E mutation in *PSEN1*, a 35–40 year old man homozygous for the
A431E mutation in PSEN1, and a normal control (45–50 year old). Hematoxylin and Eosin

(H&E) staining shows profound drop-out of neurons in the *PSEN1* mutation carriers
compared to control. Neurons within carriers lack neuromelanin and show loss of pigment;
whereas, neurons within control show robust pigmentation and neuromelanin expression.
Immunostained slides with anti-tau (AT8) antibodies using red chromagen demonstrate
neurofibrillary tangles within neurons and the presence of many tau positive threads in
PSEN1 mutation carriers with only a very rare single neurofibrillary tangle observed in
normal control (lower portion of image).

Table 3: Neuropathological assessment of amyloid β deposits (A), staging of neurofibrillary
tangles (B), and scoring of neuritic plaques (C)[*], as well as Braak and Braak[**] staging for
specimen of mutation carriers

Case	A	B	C	Braak & Braak
Case 1024	3	3	3	VI
Case 412[1]	3	3	3	VI
Case 838[2]	3	3	3	VI
Case 697	3	3	3	VI

Note: [1] Staging performed in 1996; [2] No official score available for CASE 838, provided
staging indicates an estimate by the pathologist.
[*](Montine et al., 2012); [**](Braak et al., 2011)

Figure 8. Neuropathological findings in the locus coeruleus of a 45–50 year old woman
heterozygous for the A431E mutation in PSEN1, a 35–40 year old man homozygous for the
A431E mutation, and a normal control (45–50 years old). GFAP highlights the background
reactive astrogliosis surrounding the locus coeruleus with the homozygous case demonstrating
the most reactive background and increased number of astrocytes with cell body and process
hypertrophy. The amyloid stain (4G8) highlights the extracellular amyloid plaques in the
locus coeruleus with the heterozygote case demonstrating diffuse and compact amyloid
plaques. The homozygote case demonstrates large cotton-wool amyloid plaques. The control
does not have any amyloid plaques in the area. The alpha-synuclein did not reveal Lewy body
pathology.

4.   Discussion

Animal and post-mortem research indicates that the brain's noradrenergic system

plays a central role in the pathogenesis of Alzheimer's disease (Braak et al., 2011;

Weinshenker, 2018). *In vivo* investigations, however, have long been hindered by challenges

in reliable non-invasive LC assessments (Astafiev et al., 2010; Keren et al., 2009). While

initial intriguing *in vivo* studies in late-onset Alzheimer's are underway (LOAD; e.g., Betts,

Cardenas-Blanco, et al., 2019; Jessen et al., 2018), unequivocally distinguishing age- from

disease-related LC alterations is difficult on this basis (Dahl et al., 2019; Liu et al., 2019).

Focusing instead on types of dementia developing earlier in life provides a unique window

into the mechanisms driving Alzheimer's development (ADAD; Bateman et al., 2012; Jacobs,

Becker, et al., 2019; Ringman, 2005).

Leveraging a meta-analytical approach, we first synthesized LC localizations and

dimensions across previously published studies to improve the reliability and validity of MR-

based LC detection. In a second step, we applied this newly generated volume of interest to

determine whether MR-indexed LC integrity can serve as marker for noradrenergic

degeneration in ADAD and verified this pathology using post-mortem samples.

Our analyses isolated a cluster of high-confidence voxels that (1) is consistently

identified as LC across published studies and (2) corresponds well to post-mortem reports

(Fernandes et al., 2012). Next, exploiting this refined spatial characterization we semi-

automatically extracted intensity data in an independent sample of participants with or known

to be at-risk for mutations in genes associated with ADAD. We revealed lower MR-indexed

LC integrity in symptomatic mutation carriers. Group differences were most pronounced in

middle to rostral segments of the LC (Ehrenberg et al., 2017; Lyness et al., 2003) and scaled

with proximity to the mutation-specific median age of dementia diagnosis (Ryman et al.,

2014). Beyond that, lower overall LC intensity was linked to higher tau burden in posterior

brain regions (Chalermpalanupap et al., 2018) and worse cognitive performance across

several neuropsychological tests (Dahl et al., 2019; Liu et al., 2020). Taken together, our

finding of diminished LC integrity in ADAD suggests a prominent role of the noradrenergic

system in this neurodegenerative disease.

4.1. Aggregating across published locus coeruleus maps yields high confidence meta

mask

Imaging of small brainstem structures that are not visible using conventional MR

sequences is notoriously challenging (for discussions, see Astafiev et al., 2010; Keren et al.,

2009) and warrants a close alignment of individual scans (for recommendations, see Betts,

Kirilina, et al., 2019; Liu et al., 2017). Here we used Advanced Normalization Tools (ANTs;

Avants et al., 2011, 2009) for brainstem normalization. ANTs is becoming increasingly

recognized as a powerful tool for accurate coregistrations – it has won several internationally

recognized medical image processing challenges (e.g., Klein et al., 2009 for a full list see

github.com/ANTsX/ANTs) and was used in all of the currently best ranked LC mapping

studies (3/3, [relative to 1/3 of the remaining studies]; Dahl et al., 2019; Liu et al., 2019; Ye et

al., 2021).

Besides normalization, LC mapping studies employing larger sample sizes and a

higher MR-field strength evinced a higher accuracy, as indicated by spatial congruence with

the meta mask (Dahl et al., 2019; Liu et al., 2019; Ye et al., 2021). We hope that by sharing an

unbiased, biologically-plausible LC volume of interest that is based on more than 1,000

participants and all commonly used sequence types (FSE, MT, FLASH; see Table 1) we assist

future functional and structural imaging studies in classifying brainstem effects (this LC meta

mask is available for download via osf.io/sf2ky/).

Moreover, the LC meta mask can be combined with a template-based semi-automatic

LC intensity assessment to systematically compare LC integrity in patients and controls (for a

step-by-step description of the procedure, see (Dahl et al., 2019); for an application, see e.g.,

(Gallant et al., 2021)). In short, LC-MRI are standardized to MNI space and masked with the

LC volume of interest. After masking, the relative intensity within the LC (i.e., standardized

with respect to a reference area) can be compared across groups.

4.2.1. Lower MR-indexed locus coeruleus integrity in autosomal-dominant

Alzheimer's disease

Applying the newly generated LC meta mask to an independent clinical sample, we

demonstrate its utility for accurately extracting intensity information across the rostrocaudal

extent of the nucleus. We observed lower middle–rostral LC intensity in symptomatic carriers

of ADAD-causing mutations. This finding is consistent with post-mortem reports of

substantial noradrenergic neurodegeneration in LOAD (Lyness et al., 2003). Some previous

post-mortem studies in ADAD also noted disproportionate depigmentation and neuronal loss

in the LC (Janssen et al., 2000; Snider et al., 2005) which we verified in four persons dying

with the A431E mutation in *PSEN1*. *In vivo* imaging studies are in general agreement with the

histological findings and indicate lower LC contrast in LOAD patients (Betts, Cardenas-

Blanco, et al., 2019; Dordevic et al., 2017; Takahashi et al., 2015). To date there is only a

single imaging study focusing on early-onset Alzheimer's (ADAD; Jacobs, Becker, et al.,

2019), albeit in non-symptomatic mutation carriers (i.e., at a preclinical stage). Our

observation that LC intensity scaled with proximity of dementia onset (adjusted age) across

non-symptomatic and symptomatic mutation carriers matches a negative age association in

mutation carriers reported by Jacobs and colleagues (2019) and might indicate a progressive

deterioration. In conjunction with prior post-mortem validations of LC MRI (Keren et al.,

2015; Cassidy et al., 2019) and decreased cell counts in LOAD (Lyness et al., 2003), our post-

mortem validation of decreased cell counts in the LC in ADAD are indicative of

noradrenergic neurodegeneration in ADAD.

4.2.2.   MR-indexed locus coeruleus integrity is associated with cortical tau burden

Leveraging a multivariate statistical approach, we further revealed a pattern of LC-

related tau pathology that was most pronounced in posterior brain regions (cf. Jacobs, Becker,

et al., 2019). This observation supports the notion that loss of noradrenergic neurons may

exacerbate the severity of other Alzheimer's hallmarks, like tau pathology (Chalermpalanupap

et al., 2018). The LC is one of the first brain regions in which aberrant tau can be detected at

relatively early ages (i.e., the first decades of life; Braak et al., 2011). The accumulation of

abnormal tau has been hypothesized to shift noradrenergic cells to a mode of hyperactivity

which in turn promotes tau release and spread along LC' widespread axonal pathways

(Chalermpalanupap et al., 2017; Weinshenker, 2018). While tau gradually propagates through

cortical regions (categorized as Braak stages), noradrenergic neurons first shrink in size and

then degenerate (Ehrenberg et al., 2017; Kelly et al., 2017; Theofilas et al., 2017). The frank

decline of noradrenergic cells is preceded by a period of dysfunctional neurotransmission in

LC's terminal regions like the mediotemporal lobe (Weinshenker, 2018). Thus, the negative

association between *in vivo* proxies for LC integrity and tau burden may signify a more

advanced Alzheimer's disease stage including noradrenergic neurodegeneration and wide-

spread tau pathology.

4.2.3.   MR-indexed locus coeruleus integrity is associated with cognitive deficits

Finally, we found evidence for an association between MR-indexed LC integrity and

cognitive performance across several neuropsychological tests. Noradrenergic

neuromodulation has been implicated in a range of cognitive functions, prominently including

attention and memory (Berridge & Waterhouse, 2003; Bouret & Sara, 2005; Corbetta et al.,

2008; Mather et al., 2016; Sara, 2009). In particular, mediated via β-adrenoceptors,

noradrenaline release promotes long-term-potentiation in the hippocampus—a key

determinant of synaptic plasticity and memory (O'Dell et al., 2015). Recent optogenetic

research moreover indicates a causal role of LC activity in memory acquisition and

consolidation, potentially supported by co-release of dopamine (Duszkiewicz et al., 2019;

Takeuchi et al., 2016; Uematsu et al., 2017; Wagatsuma et al., 2018). Consistent with these

observations in animals, recent *in vivo* human research demonstrated an association between

MR-indexed LC integrity and memory performance in healthy aging (Dahl et al., 2019;

Elman et al., 2021; Hämmerer et al., 2018; Liu et al., 2020). Extending evidence from healthy

populations, we here confirm the behavioral relevance of MR-indexed LC integrity in

tracking cognitive decline in a clinical sample. Neuropsychological tests that proved sensitive

to distinguish patients from controls were reliably linked to LC integrity (González et al.,

2002; Nasreddine et al., 2005; O'Bryant et al., 2008; Teng et al., 1994). In line with reports

indicating verbal learning tests as effective early markers of Alzheimer's pathology (Albert et

al., 2001; Belleville et al., 2017; Moradi et al., 2017; Schoenberg et al., 2006), verbal learning

and memory showed the closest link to LC integrity (cf. Dahl et al., 2019, for analogous

results in later life). Taken together, our findings suggest that noradrenergic neurotransmission

supports memory performance whereas dysfunctions therein are associated with Alzheimer's

related cognitive decline.

4.3. Limitations and conclusions

Some limitations should be noted. First, the contrast mechanisms underlying LC-

MRI constitute an active area of research (for a discussion, see Betts, Kirilina, et al., 2019)

and future studies may help determine what precise physiological processes are reflected in

MR-contrast differences. Second, as a cross-sectional observational study, it is not possible to

conclude a causative relationship between loss of LC integrity and cognitive symptoms and

deposition of abnormal tau in persons with ADAD. Finally, with a prevalence of about 5 cases

per 100,000 persons at risk, ADAD is a rare disease (Campion et al., 1999) accounting for less

than 1% of all Alzheimer's cases (Bateman et al., 2011). All but one ADAD mutation carriers

in the current study had the same mutation in *PSEN1* (A431E) which has specific features

(e.g., co-occurring spastic paraparesis and possibly atypical posterior-predominant tau

deposition) that may limit generalizability of the current findings. While providing insights

into the pathogenesis of Alzheimer's largely disentangled from factors associated with aging,

the presented findings need to be corroborated by longitudinal evidence in larger samples

(Lindenberger et al., 2011).

763   In conclusion, we applied a meta-analytical approach to advance the reliability and

validity of MR-based LC detection. Using non-invasive *in vivo* proxies, we revealed lower

LC integrity in symptomatic carriers of ADAD-causing mutations. Moreover, LC integrity

was strongly associated with cortical tau pathology and memory decline. In summary, this

provides support for a prominent role of the noradrenergic system in Alzheimer's disease

(Mather & Harley, 2016; Weinshenker, 2018).

5.   Acknowledgements

The authors thank all study participants for their contribution.
771
772 **6.   Competing interests**

773 The authors declare no competing financial interests.
774

7.   Funding

MJD is recipient of a stipend from the GA. Lienert-Foundation and was supported by a
stipend from the Davis School of Gerontology, University of Southern California. MM's work
was supported by National Institutes of Health (NIH) grant R01AG025340, an Alexander von
Humboldt fellowship, and a Max Planck Sabbatical Award. MW-B received support from the
German Research Foundation (DFG, WE 4269/5-1) and the Jacobs Foundation (Early Career
Research Fellowship 2017–2019). BLK was supported by a NIH grant F32AG057162. YS
was supported by NIH grants RF1AG064584 and RF1AG056573. SG, KH, and CCM were
supported by P30AG066530, JMR was supported by NIH grants U01AG051218,
R01AG062007, and P30AG066530. The funders had no role in study design, data collection
and analysis, decision to publish, or preparation of the manuscript. We thank Dr. Abhay
Sagare for help with genotyping.
787

8. References

Albert, M. S., Moss, M. B., Tanzi, R., & Jones, K. (2001). Preclinical prediction of AD using

neuropsychological tests. *Journal of the International Neuropsychological Society*, *7*(5),

631–639. https://doi.org/10.1017/S1355617701755105

Alzheimer's Disease International. (2019). World Alzheimer Report 2019: Attitudes to

Dementia. *Alzheimer's Disease International: London*.

Arendt, T., Brückner, M. K., Morawski, M., Jäger, C., & Gertz, H. J. (2015). Early neurone

loss in Alzheimer's disease: cortical or subcortical? *Acta Neuropathologica*

Communications, *3*, 10. https://doi.org/10.1186/s40478-015-0187-1

Astafiev, S. V., Snyder, A. Z., Shulman, G. L., & Corbetta, M. (2010). Comment on

"Modafinil shifts human locus coeruleus to low-tonic, high-phasic activity during

functional MRI" and "Homeostatic sleep pressure and responses to sustained attention in

the suprachiasmatic area." *Science*, *328*(5976), 309.

https://doi.org/10.1126/science.1177200

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A

reproducible evaluation of ANTs similarity metric performance in brain image

registration. *NeuroImage*, *54*(3), 2033–2044.

https://doi.org/10.1016/j.neuroimage.2010.09.025

Avants, B. B., Tustison, N., & Song, G. (2009). Advanced Normalization Tools: V1.0. *Insight*

Journal, *2*. https://doi.org/http://hdl.handle.net/10380/3113

Bateman, R. J., Aisen, P. S., De Strooper, B., Fox, N. C., Lemere, C. A., Ringman, J. M.,

Salloway, S., Sperling, R. A., Windisch, M., & Xiong, C. (2011). Autosomal-dominant

Alzheimer's disease: A review and proposal for the prevention of Alzheimer's disease. In

Alzheimer's Research and Therapy (Vol. 2, Issue 6, p. 1). BioMed Central.

https://doi.org/10.1186/alzrt59

Bateman, R. J., Xiong, C., Benzinger, T. L. S., Fagan, A. M., Goate, A., Fox, N. C., Marcus,

D. S., Cairns, N. J., Xie, X., Blazey, T. M., Holtzman, D. M., Santacruz, A., Buckles, V.,

Oliver, A., Moulder, K., Aisen, P. S., Ghetti, B., Klunk, W. E., McDade, E., … Morris, J.

C. (2012). Clinical and biomarker changes in dominantly inherited Alzheimer's disease.

New England Journal of Medicine, *367*(9), 795–804.

https://doi.org/10.1056/NEJMoa1202753

Belleville, S., Fouquet, C., Hudon, C., Zomahoun, H. T. V., & Croteau, J. (2017).

Neuropsychological measures that predict progression from mild cognitive impairment

to Alzheimer's type dementia in older adults: A systematic review and meta-analysis.

Neuropsychology Review, *27*(4), 328–353. https://doi.org/10.1007/s11065-017-9361-5

Berridge, C. W., & Waterhouse, B. D. (2003). The locus coeruleus-noradrenergic system:

Modulation of behavioral state and state-dependent cognitive processes. *Brain Research*

Reviews, *42*(1), 33–84. https://doi.org/10.1016/S0165-0173(03)00143-7

Bertram, L., Lill, C. M., & Tanzi, R. E. (2010). The genetics of alzheimer disease: Back to the

future. In *Neuron* (Vol. 68, Issue 2, pp. 270–281). Neuron.

https://doi.org/10.1016/j.neuron.2010.10.013

Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Jessen, F., & Düzel, E. (2017). In vivo MRI

assessment of the human locus coeruleus along its rostrocaudal extent in young and older

adults. *NeuroImage*, *163*, 150–159. https://doi.org/10.1016/j.neuroimage.2017.09.042

Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Spottke, A., Teipel, S. J., Kilimann, I.,

Jessen, F., & Düzel, E. (2019). Locus coeruleus MRI contrast is reduced in Alzheimer's

disease dementia and correlates with CSF Aβ levels. *Alzheimer's & Dementia:*

Diagnosis, Assessment & Disease Monitoring, *11*, 281–285.

https://doi.org/10.1016/j.dadm.2019.02.001

Betts, M. J., Kirilina, E., Otaduy, M. C. G., Ivanov, D., Acosta-Cabronero, J., Callaghan, M.

F., Lambert, C., Cardenas-Blanco, A., Pine, K., Passamonti, L., Loane, C., Keuken, M.

C., Trujillo, P., Lüsebrink, F., Mattern, H., Liu, K. Y., Priovoulos, N., Fliessbach, K.,

Dahl, M. J., … Hämmerer, D. (2019). Locus coeruleus imaging as a biomarker for

noradrenergic dysfunction in neurodegenerative diseases. *Brain*, *142*(9), 2558–2571.

https://doi.org/10.1093/brain/awz193

Bouret, S., & Sara, S. J. (2005). Network reset: A simplified overarching theory of locus

coeruleus noradrenaline function. *Trends in Neurosciences*, *28*(11), 574–582.

https://doi.org/10.1016/j.tins.2005.09.002

Braak, H., & Del Tredici, K. (2016). Potential Pathways of Abnormal Tau and α-Synuclein

Dissemination in Sporadic Alzheimer's and Parkinson's Diseases. *Cold Spring Harbor*

Perspectives in Biology, *8*(11), a023630.

https://doi.org/10.1101/CSHPERSPECT.A023630

Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the pathologic

process in Alzheimer disease: Age categories from 1 to 100 years. *Journal of*

Neuropathology and Experimental Neurology, *70*(11), 960–969.

https://doi.org/10.1097/NEN.0b013e318232a379

Brookmeyer, R., Johnson, E., Ziegler-Graham, K., & Arrighi, H. M. (2007). Forecasting the

global burden of Alzheimer's disease. *Alzheimer's & Dementia*, *3*(3), 186–191.

https://doi.org/10.1016/J.JALZ.2007.04.381

Campion, D., Dumanchin, C., Hannequin, D., Dubois, B., Belliard, S., Puel, M., Thomas-

Anterion, C., Michon, A., Martin, C., Charbonnier, F., Raux, G., Camuzat, A., Penet, C.,

Mesnage, V., Martinez, M., Clerget-Darpoux, F., Brice, A., & Frebourg, T. (1999). Early-

onset autosomal dominant Alzheimer disease: Prevalence, genetic heterogeneity, and

mutation spectrum. *American Journal of Human Genetics*, *65*(3), 664–670.

https://doi.org/10.1086/302553

Canter, R. G., Penney, J., & Tsai, L. H. (2016). The road to restoring neural circuits for the

treatment of Alzheimer's disease. In *Nature* (Vol. 539, Issue 7628, pp. 187–196). Nature

Publishing Group. https://doi.org/10.1038/nature20412

Cassidy, C. M., Zucca, F. A., Girgis, R. R., Baker, S. C., Weinstein, J. J., Sharp, M. E., Bellei,

C., Valmadre, A., Vanegas, N., Kegeles, L. S., Brucato, G., Jung Kang, U., Sulzer, D.,

Zecca, L., Abi-Dargham, A., & Horga, G. (2019). Neuromelanin-sensitive MRI as a

noninvasive proxy measure of dopamine function in the human brain. *Proceedings of the*

National Academy of Sciences of the United States of America, *116*(11), 5108–5117.

https://doi.org/10.1073/pnas.1807983116

Chalermpalanupap, T., Schroeder, J. P., Rorabaugh, J. M., Liles, L. C., Lah, J. J., Levey, A. I.,

& Weinshenker, D. (2018). Locus coeruleus ablation exacerbates cognitive deficits,

neuropathology, and lethality in P301S tau transgenic mice. *The Journal of*

Neuroscience : The Official Journal of the Society for Neuroscience, *38*(1), 74–92.

https://doi.org/10.1523/JNEUROSCI.1483-17.2017

Chalermpalanupap, T., Weinshenker, D., & Rorabaugh, J. M. (2017). Down but not out: The

consequences of pretangle tau in the locus coeruleus. *Neural Plasticity*, *2017*, 7829507.

https://doi.org/10.1155/2017/7829507

Corbetta, M., Patel, G., & Shulman, G. L. (2008). The reorienting system of the human brain:

From environment to theory of mind. *Neuron*, *58*(3), 306–324.

https://doi.org/10.1016/j.neuron.2008.04.017

Craft, S., Newcomer, J., Kanne, S., Dagogo-Jack, S., Cryer, P., Sheline, Y., Luby, J., Dagogo-

Jack, A., & Alderson, A. (1996). Memory improvement following induced

hyperinsulinemia in Alzheimer's disease. *Neurobiology of Aging*, *17*(1), 123–130.

https://doi.org/10.1016/0197-4580(95)02002-0

Crary, J. F., Trojanowski, J. Q., Schneider, J. A., Abisambra, J. F., Abner, E. L., Alafuzoff, I.,

Arnold, S. E., Attems, J., Beach, T. G., Bigio, E. H., Cairns, N. J., Dickson, D. W.,

Gearing, M., Grinberg, L. T., Hof, P. R., Hyman, B. T., Jellinger, K., Jicha, G. A.,

Kovacs, G. G., … Nelson, P. T. (2014). Primary age-related tauopathy (PART): a

common pathology associated with human aging. *Acta Neuropathologica*, *128*(6), 755–

766. https://doi.org/10.1007/s00401-014-1349-0

Dahl, M. J., Mather, M., Düzel, S., Bodammer, N. C., Lindenberger, U., Kühn, S., & Werkle-

Bergner, M. (2019). Rostral locus coeruleus integrity is associated with better memory

performance in older adults. *Nature Human Behaviour*, *3*(11), 1203–1214.

https://doi.org/10.1038/s41562-019-0715-2

Dahl, M. J., Mather, M., Sander, M. C., & Werkle-Bergner, M. (2020). Noradrenergic

responsiveness supports selective attention across the adult lifespan. *Journal of*

Neuroscience, *40*(22), 4372–4390. https://doi.org/10.1523/JNEUROSCI.0398-19.2020

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I.

Segmentation and surface reconstruction. *NeuroImage*, *9*(2), 179–194.

https://doi.org/10.1006/nimg.1998.0395

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,

R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006).

An automated labeling system for subdividing the human cerebral cortex on MRI scans

into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Dordevic, M., Müller-Fotti, A., Müller, P., Schmicker, M., Kaufmann, J., & Müller, N. G.

(2017). Optimal Cut-Off Value for Locus Coeruleus-to-Pons Intensity Ratio as Clinical

Biomarker for Alzheimer's Disease: A Pilot Study. *Journal of Alzheimer's Disease*

Reports, *1*(1), 159–167. https://doi.org/10.3233/adr-170021

Duszkiewicz, A. J., McNamara, C. G., Takeuchi, T., & Genzel, L. (2019). Novelty and

Dopaminergic Modulation of Memory Persistence: A Tale of Two Systems. *Trends in*

Neurosciences, *42*(2), 102–114. https://doi.org/10.1016/J.TINS.2018.10.002

Ehrenberg, A. J., Nguy, A. K., Theofilas, P., Dunlop, S., Suemoto, C. K., Di Lorenzo Alho, A.

T., Leite, R. P., Diehl Rodriguez, R., Mejia, M. B., Rüb, U., Farfel, J. M., de Lucena

Ferretti-Rebustini, R. E., Nascimento, C. F., Nitrini, R., Pasquallucci, C. A., Jacob-Filho,

W., Miller, B., Seeley, W. W., Heinsen, H., & Grinberg, L. T. (2017). Quantifying the

accretion of hyperphosphorylated tau in the locus coeruleus and dorsal raphe nucleus:

The pathological building blocks of early Alzheimer's disease. *Neuropathology and*

Applied Neurobiology, *43*(5), 393–408. https://doi.org/10.1111/nan.12387

Elman, J. A., Puckett, O. K., Beck, A., Panizzon, M. S., Sanderson-Cimino, M. E., Gustavson,

D. E., Lyons, M. J., Franz, C. E., & Kremen, W. S. (2021). MRI-assessed locus coeruleus

integrity is heritable and associated with cognition, Alzheimer's risk, and sleep-wake

disturbance. *Alzheimer's & Dementia*, *17*(6), 1017– 1025.

https://doi.org/10.1002/alz.044862

Fernandes, P., Regala, J., Correia, F., & Gonçalves-Ferreira, A. J. (2012). The human locus

coeruleus 3-D stereotactic anatomy. *Surgical and Radiologic Anatomy*, *34*(10), 879–885.

https://doi.org/10.1007/s00276-012-0979-y

Gallant, S. N., Kennedy, B. L., Bachman, S. L., Huang, R., Lee, T.-H., & Mather, M. (2021).

Behavioral and fMRI evidence that arousal enhances bottom-up attention and memory

selectivity in young but not older adults. *BioRxiv*, 2021.07.02.450802.

https://doi.org/10.1101/2021.07.02.450802

Ghosh, A., Torraville, S. E., Mukherjee, B., Walling, S. G., Martin, G. M., Harley, C. W., &

Yuan, Q. (2019). An experimental model of Braak's pretangle proposal for the origin of

Alzheimer's disease: The role of locus coeruleus in early symptom development.

Alzheimer's Research and Therapy, *11*(1), 59. https://doi.org/10.1186/s13195-019-0511-

2

Goate, A., Chartier-Harlin, M. C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L.,

Haynes, A., Irving, N., James, L., Mant, R., Newton, P., Rooke, K., Roques, P., Talbot,

C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., … Hardy, J. (1991).

Segregation of a missense mutation in the amyloid precursor protein gene with familial

Alzheimer's disease. *Nature*, *349*(6311), 704–706. https://doi.org/10.1038/349704a0

González, H. M., Mungas, D., & Haan, M. N. (2002). A verbal learning and memory test for

English- and Spanish-speaking older Mexican-American adults. *Clinical*

Neuropsychologist, *16*(4), 439–451. https://doi.org/10.1076/clin.16.4.439.13908

Greve, D. N., Salat, D. H., Bowen, S. L., Izquierdo-Garcia, D., Schultz, A. P., Catana, C.,

Becker, J. A., Svarer, C., Knudsen, G. M., Sperling, R. A., & Johnson, K. A. (2016).

Different partial volume correction methods lead to different conclusions: An 18F-FDG-

PET study of aging. *NeuroImage*, *132*, 334–343.

https://doi.org/10.1016/j.neuroimage.2016.02.042

Greve, D. N., Svarer, C., Fisher, P. M., Feng, L., Hansen, A. E., Baare, W., Rosen, B., Fischl,

B., & Knudsen, G. M. (2014). Cortical surface-based analysis reduces bias and variance

in kinetic modeling of brain PET data. *NeuroImage*, *92*, 225–236.

https://doi.org/10.1016/j.neuroimage.2013.12.021

Grinberg, L. T., & Heinsen, H. (2017). Light at the beginning of the tunnel? Investigating

early mechanistic changes in Alzheimer's disease. *Brain*, *140*(11), 2770–2773.

https://doi.org/10.1093/brain/awx261

Hämmerer, D., Callaghan, M. F., Hopkins, A., Kosciessa, J., Betts, M., Cardenas-Blanco, A.,

Kanowski, M., Weiskopf, N., Dayan, P., Dolan, R. J., & Düzel, E. (2018). Locus

coeruleus integrity in old age is selectively related to memories linked with salient

negative events. *Proceedings of the National Academy of Sciences of the United States of*

America, *115*, 2228–2233. https://doi.org/10.1073/pnas.1712268115

Hanseeuw, B. J., Betensky, R. A., Jacobs, H. I. L., Schultz, A. P., Sepulcre, J., Becker, J. A.,

Cosio, D. M. O., Farrell, M., Quiroz, Y. T., Mormino, E. C., Buckley, R. F., Papp, K. V.,

Amariglio, R. A., Dewachter, I., Ivanoiu, A., Huijbers, W., Hedden, T., Marshall, G. A.,

Chhatwal, J. P., … Johnson, K. (2019). Association of amyloid and tau with cognition in

preclinical Alzheimer disease. *JAMA Neurology*, Advance online publication.

https://doi.org/10.1001/jamaneurol.2019.1424

Heneka, M. T., Nadrigny, F., Regen, T., Martinez-Hernandez, A., Dumitrescu-Ozimek, L.,

Terwel, D., Jardanhazi-Kurutz, D., Walter, J., Kirchhoff, F., Hanisch, U. K., & Kummer,

M. P. (2010). Locus ceruleus controls Alzheimer's disease pathology by modulating

microglial functions through norepinephrine. *Proceedings of the National Academy of*

Sciences of the United States of America, *107*(13), 6058–6063.

https://doi.org/10.1073/pnas.0909586107

Jacobs, H. I. L., Becker, A., Sperling, R. A., Guzman-Velez, E., Baena, A., Uquillas, F.

d'Oleire, Artola, A., Pardilla-Delgado, E., Gatchel, J. R., Reiman, E. M., Johnson, K. A.,

Lopera, F., & Quiroz, Y. T. (2019). P2-425: Locus coeruleus intensity is associated with

early amyloid and tau pathology in preclinical autosomal dominant Alzheimer's disease.

Alzheimer's & Dementia, *15*, P774–P775. https://doi.org/10.1016/j.jalz.2019.06.2832

Jacobs, H. I. L., Riphagen, J. M., Ramakers, I. H. G. B., & Verhey, F. R. J. (2019).

Alzheimer's disease pathology: Pathways between central norepinephrine activity,

memory, and neuropsychiatric symptoms. *Molecular Psychiatry*, Advance online

publication. https://doi.org/10.1038/s41380-019-0437-x

Jagust, W. (2018). Imaging the evolution and pathophysiology of Alzheimer disease. *Nature*

Reviews Neuroscience, *19*, 687–700. https://doi.org/10.1038/s41583-018-0067-3

Janssen, J. C., Hall, M., Fox, N. C., Harvey, R. J., Beck, J., Dickinson, A., Campbell, T.,

Collinge, J., Lantos, P. L., Cipolotti, L., Stevens, J. M., & Rossor, M. N. (2000).

Alzheimer's disease due to an intronic presenilin-1 (PSEN1 intron 4) mutation. A

clinicopathological study. *Brain*, *123*(5), 894–907.

https://doi.org/10.1093/brain/123.5.894

Jessen, F., Spottke, A., Boecker, H., Brosseron, F., Buerger, K., Catak, C., Fliessbach, K.,

Franke, C., Fuentes, M., Heneka, M. T., Janowitz, D., Kilimann, I., Laske, C., Menne, F.,

Nestor, P., Peters, O., Priller, J., Pross, V., Ramirez, A., … Düzel, E. (2018). Design and

first baseline data of the DZNE multicenter observational study on predementia

Alzheimer's disease (DELCODE). *Alzheimer's Research & Therapy*, *10*, 15.

https://doi.org/10.1186/s13195-017-0314-2

Kelly, S. C., He, B., Perez, S. E., Ginsberg, S. D., Mufson, E. J., & Counts, S. E. (2017).

Locus coeruleus cellular and molecular pathology during the progression of Alzheimer's

disease. *Acta Neuropathologica Communications*, *5*(1), 8.

https://doi.org/10.1186/s40478-017-0411-2

Keren, N. I., Lozar, C. T., Harris, K. C., Morgan, P. S., & Eckert, M. A. (2009). In vivo

mapping of the human locus coeruleus. *NeuroImage*, *47*(4), 1261–1267.

https://doi.org/10.1016/j.neuroimage.2009.06.012

Keren, N. I., Taheri, S., Vazey, E. M., Morgan, P. S., Granholm, A. C. E., Aston-Jones, G., &

Eckert, M. A. (2015). Histologic validation of locus coeruleus MRI contrast in post-

mortem tissue. *NeuroImage*, *113*, 235–245.

https://doi.org/10.1016/j.neuroimage.2015.03.020

Keresztes, A., Bender, A. R., Bodammer, N. C., Lindenberger, U., Shing, Y. L., & Werkle-

Bergner, M. (2017). Hippocampal maturity promotes memory distinctiveness in

childhood and adolescence. *Proceedings of the National Academy of Sciences of the*

United States of America, *114*(34), 9212–9217. https://doi.org/10.1073/pnas.1710654114

Klein, A., Andersson, J., Ardekani, B. A., Ashburner, J., Avants, B., Chiang, M.-C.,

Christensen, G. E., Collins, D. L., Gee, J., Hellier, P., Song, J. H., Jenkinson, M., Lepage,

C., Rueckert, D., Thompson, P., Vercauteren, T., Woods, R. P., Mann, J. J., & Parsey, R.

V. (2009). Evaluation of 14 nonlinear deformation algorithms applied to human brain

MRI registration. *NeuroImage*, *46*(3), 786–802.

https://doi.org/10.1016/J.NEUROIMAGE.2008.12.037

Krishnan, A., Williams, L. J., McIntosh, A. R., & Abdi, H. (2011). Partial Least Squares (PLS)

methods for neuroimaging: A tutorial and review. *NeuroImage*, *56*(2), 455–475.

https://doi.org/10.1016/j.neuroimage.2010.07.034

La Joie, R., Visani, A. V., Baker, S. L., Brown, J. A., Bourakova, V., Cha, J., Chaudhary, K.,

Edwards, L., Iaccarino, L., Janabi, M., Lesman-Segev, O. H., Miller, Z. A., Perry, D. C.,

O'Neil, J. P., Pham, J., Rojas, J. C., Rosen, H. J., Seeley, W. W., Tsai, R. M., …

Rabinovici, G. D. (2020). Prospective longitudinal atrophy in Alzheimer's disease

correlates with the intensity and topography of baseline tau-PET. *Science Translational*

Medicine, *12*(524), eaau5732. https://doi.org/10.1126/scitranslmed.aau5732

Lee, S., Viqar, F., Zimmerman, M. E., Narkhede, A., Tosto, G., Benzinger, T. L. S., Marcus,

D. S., Fagan, A. M., Goate, A., Fox, N. C., Cairns, N. J., Holtzman, D. M., Buckles, V.,

Ghetti, B., McDade, E., Martins, R. N., Saykin, A. J., Masters, C. L., Ringman, J. M., …

Network, for the D. I. A. (2016). White matter hyperintensities are a core feature of

Alzheimer's disease: Evidence from the dominantly inherited Alzheimer network.

Annals of Neurology, *79*(6), 929–939. https://doi.org/https://doi.org/10.1002/ana.24647

Lee, T.-H., Greening, S. G., Ueno, T., Clewett, D., Ponzio, A., Sakaki, M., & Mather, M.

(2018). Arousal increases neural gain via the locus coeruleus–noradrenaline system in

younger adults but not in older adults. *Nature Human Behaviour*, *2*(5), 356–366.

https://doi.org/10.1038/s41562-018-0344-1

Lindenberger, U., von Oertzen, T., Ghisletta, P., & Hertzog, C. (2011). Cross-sectional age

variance extraction: What's change got to do with it? *Psychology and Aging*, *26*(1), 34–

47. https://doi.org/10.1037/a0020525

Liu, K. Y., Acosta-Cabronero, J., Cardenas-Blanco, A., Loane, C., Berry, A. J., Betts, M. J.,

Kievit, R. A., Henson, R. N., Düzel, E., Howard, R., & Hämmerer, D. (2019). In vivo

visualization of age-related differences in the locus coeruleus. *Neurobiology of Aging*,

74, 101–111. https://doi.org/10.1016/j.neurobiolaging.2018.10.014

Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Howard, R., &

Hämmerer, D. (2020). Noradrenergic-dependent functions are associated with age-

related locus coeruleus signal intensity differences. *Nature Communications*, *11*(1),

1712. https://doi.org/10.1038/s41467-020-15410-w

Liu, K. Y., Marijatta, F., Hämmerer, D., Acosta-Cabronero, J., Düzel, E., & Howard, R. J.

(2017). Magnetic resonance imaging of the human locus coeruleus: A systematic review.

Neuroscience and Biobehavioral Reviews, *83*, 325–355.

https://doi.org/10.1016/j.neubiorev.2017.10.023

Lyness, S. A., Zarow, C., & Chui, H. C. (2003). Neuron loss in key cholinergic and aminergic

nuclei in Alzheimer disease: A meta-analysis. In *Neurobiology of Aging* (Vol. 24, Issue 1,

pp. 1–23). Elsevier. https://doi.org/10.1016/S0197-4580(02)00057-X

Mäki-Marttunen, V., & Espeseth, T. (2020). Uncovering the locus coeruleus: comparison of

localization methods for functional analysis. *NeuroImage*, 117409.

https://doi.org/10.1016/j.neuroimage.2020.117409

Makropoulos, A., Robinson, E. C., Schuh, A., Wright, R., Fitzgibbon, S., Bozek, J., Counsell,

S. J., Steinweg, J., Vecchiato, K., Passerat-Palmbach, J., Lenz, G., Mortari, F., Tenev, T.,

Duff, E. P., Bastiani, M., Cordero-Grande, L., Hughes, E., Tusor, N., Tournier, J. D., …

Rueckert, D. (2018). The developing human connectome project: A minimal processing

pipeline for neonatal cortical surface reconstruction. *NeuroImage*, *173*, 88–112.

https://doi.org/10.1016/j.neuroimage.2018.01.054

Marien, M. R., Colpaert, F. C., & Rosenquist, A. C. (2004). Noradrenergic mechanisms in

neurodegenerative diseases: A theory. *Brain Research Reviews*, *45*(1), 38–78.

https://doi.org/10.1016/j.brainresrev.2004.02.002

Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). Norepinephrine ignites local

hotspots of neuronal excitation: How arousal amplifies selectivity in perception and

memory. *Behavioral and Brain Sciences*, *39*, e200.

https://doi.org/10.1017/S0140525X15000667

Mather, M., & Harley, C. W. (2016). The locus coeruleus: Essential for maintaining cognitive

function and the aging brain. *Trends in Cognitive Sciences*, *20*(3), 214–226.

https://doi.org/10.1016/j.tics.2016.01.001

McIntosh, A. R., & Lobaugh, N. J. (2004). Partial least squares analysis of neuroimaging data:

Applications and advances. *NeuroImage*, *23*(SUPPL. 1).

https://doi.org/10.1016/j.neuroimage.2004.07.020

Miyoshi, F., Ogawa, T., Kitao, S. I., Kitayama, M., Shinohara, Y., Takasugi, M., Fujii, S., &

Kaminou, T. (2013). Evaluation of Parkinson disease and Alzheimer disease with the use

of neuromelanin MR imaging and123I-metaiodobenzylguanidine scintigraphy. *American*

Journal of Neuroradiology, *34*(11), 2113–2118. https://doi.org/10.3174/ajnr.A3567

Montine, T. J., Phelps, C. H., Beach, T. G., Bigio, E. H., Cairns, N. J., Dickson, D. W.,

Duyckaerts, C., Frosch, M. P., Masliah, E., Mirra, S. S., Nelson, P. T., Schneider, J. A.,

Thal, D. R., Trojanowski, J. Q., Vinters, H. V, & Hyman, B. T. (2012). National Institute

on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of

Alzheimer's disease: a practical approach. *Acta Neuropathologica*, *123*(1), 1–11.

https://doi.org/10.1007/s00401-011-0910-3

Moradi, E., Hallikainen, I., Hänninen, T., & Tohka, J. (2017). Rey's Auditory Verbal Learning

Test scores can be predicted from whole brain MRI in Alzheimer's disease. *NeuroImage:*

Clinical, *13*, 415–427. https://doi.org/10.1016/j.nicl.2016.12.011

Muehlroth, B. E., Sander, M. C., Fandakova, Y., Grandy, T. H., Rasch, B., Lee Shing, Y., &

Werkle-Bergner, M. (2020). Memory quality modulates the effect of aging on memory

consolidation during sleep: Reduced maintenance but intact gain. *NeuroImage*, *209*.

https://doi.org/10.1016/j.neuroimage.2019.116490

Murrell, J., Ghetti, B., Cochran, E., Macias-Islas, M. A., Medina, L., Varpetian, A.,

Cummings, J. L., Mendez, M. F., Kawas, C., Chui, H., & Ringman, J. M. (2006). The

A431E mutation in PSEN1 causing Familial Alzheimer's Disease originating in Jalisco

State, Mexico: An additional fifteen families. *Neurogenetics*, *7*(4), 277–279.

https://doi.org/10.1007/s10048-006-0053-1

Nakane, T., Nihashi, T., Kawai, H., & Naganawa, S. (2008). Visualization of neuromelanin in

the Substantia nigra and locus ceruleus at 1.5T using a 3D-gradient echo sequence with

magnetization transfer contrast. *Magnetic Resonance in Medical Sciences : MRMS : An*

Official Journal of Japan Society of Magnetic Resonance in Medicine, *7*(4), 205–210.

http://www.ncbi.nlm.nih.gov/pubmed/19110515

Nasreddine, Z. S., Phillips, N. A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I.,

Cummings, J. L., & Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA:

A brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics*

Society, *53*(4), 695–699. https://doi.org/10.1111/j.1532-5415.2005.53221.x

O'Bryant, S. E., Waring, S. C., Cullum, C. M., Hall, J., Lacritz, L., Massman, P. J., Lupo, P.

J., Reisch, J. S., & Doody, R. (2008). Staging dementia using clinical dementia rating

scale sum of boxes scores: A Texas Alzheimer's research consortium study. *Archives of*

Neurology, *65*(8), 1091–1095. https://doi.org/10.1001/archneur.65.8.1091

O'Dell, T. J., Connor, S. A., Guglietta, R., & Nguyen, P. V. (2015). β-Adrenergic receptor

signaling and modulation of long-term potentiation in the mammalian hippocampus.

Learning & Memory, *22*(9), 461–471. https://doi.org/10.1101/lm.031088.113

Parker, J., Mozaffar, T., Messmore, A., Deignan, J. L., Kimonis, V. E., & Ringman, J. M.

(2019). Homozygosity for the A431E mutation in PSEN1 presenting with a relatively

aggressive phenotype. *Neuroscience Letters*, *699*, 195–198.

https://doi.org/10.1016/j.neulet.2019.01.047

Penny, W., Friston, K., Ashburner, J., Kiebel, S., & Nichols, T. (2007). Statistical Parametric

Mapping: The Analysis of Functional Brain Images. In *Statistical Parametric Mapping:*

The Analysis of Functional Brain Images. Elsevier Ltd. https://doi.org/10.1016/B978-0-

12-372560-8.X5000-1

Poe, G. R., Foote, S., Eschenko, O., Johansen, J. P., Bouret, S., Aston-Jones, G., Harley, C.

W., Manahan-Vaughan, D., Weinshenker, D., Valentino, R., Berridge, C., Chandler, D. J.,

Waterhouse, B., & Sara, S. J. (2020). Locus coeruleus: a new look at the blue spot.

Nature Reviews Neuroscience, *21*(11), 644–659. https://doi.org/10.1038/s41583-020-

0360-9

Possin, K. L., Laluz, V. R., Alcantar, O. Z., Miller, B. L., & Kramer, J. H. (2011). Distinct

neuroanatomical substrates and cognitive mechanisms of figure copy performance in

Alzheimer's disease and behavioral variant frontotemporal dementia. *Neuropsychologia*,

49(1), 43–48. https://doi.org/10.1016/j.neuropsychologia.2010.10.026

Priovoulos, N., Jacobs, H. I. L., Ivanov, D., Uludag, K., Verhey, F. R. J., & Poser, B. A.

(2017). High-resolution in vivo imaging of human locus coeruleus by magnetization

transfer MRI at 3T and 7T. *NeuroImage*, *168*, 127–136.

https://doi.org/10.1016/j.neuroimage.2017.07.045

Ringman, J. M. (2005). What the study of persons at risk for familial Alzheimer's disease can

tell us about the earliest stages of the disorder: A review. *Journal of Geriatric Psychiatry*

and Neurology, *18*(4), 228–233. https://doi.org/10.1177/0891988705281878

Ringman, J. M., Monsell, S., Ng, D. W., Zhou, Y., Nguyen, A., Coppola, G., Van Berlo, V.,

Mendez, M. F., Tung, S., Weintraub, S., Mesulam, M.-M., Bigio, E. H., Gitelman, D. R.,

Fisher-Hubbard, A. O., Albin, R. L., & Vinters, H. V. (2016). Neuropathology of

Autosomal Dominant Alzheimer Disease in the National Alzheimer Coordinating Center

Database. *Journal of Neuropathology & Experimental Neurology*, *75*(3), 284–290.

https://doi.org/10.1093/jnen/nlv028

Rohlfing, T., Brandt, R., Menzel, R., & Maurer, C. R. (2004). Evaluation of atlas selection

strategies for atlas-based image segmentation with application to confocal microscopy

images of bee brains. *NeuroImage*, *21*(4), 1428–1442.

https://doi.org/10.1016/j.neuroimage.2003.11.010

Rorabaugh, J. M., Chalermpalanupap, T., Botz-Zapp, C. A., Fu, V. M., Lembeck, N. A.,

Cohen, R. M., & Weinshenker, D. (2017). Chemogenetic locus coeruleus activation

restores reversal learning in a rat model of Alzheimer's disease. *Brain*, *140*(11), 3023–

3038. https://doi.org/10.1093/brain/awx232

Rosenthal, R. (1991). *Meta-Analytic Procedures for Social Research*. SAGE Publications, Inc.

https://doi.org/10.4135/9781412984997

Ryman, D. C., Acosta-Baena, N., Aisen, P. S., Bird, T., Danek, A., Fox, N. C., Goate, A.,

Frommelt, P., Ghetti, B., Langbaum, J. B. S., Lopera, F., Martins, R., Masters, C. L.,

Mayeux, R. P., McDade, E., Moreno, S., Reiman, E. M., Ringman, J. M., Salloway, S.,

… Bateman, R. J. (2014). Symptom onset in autosomal dominant Alzheimer disease: A

systematic review and meta-analysis. *Neurology*, *83*(3), 253–260.

https://doi.org/10.1212/WNL.0000000000000596

Sabuncu, M. R., Yeo, B. T. T., Van Leemput, K., Fischl, B., & Golland, P. (2010). A

generative model for image segmentation based on label fusion. *IEEE Transactions on*

Medical Imaging, *29*(10), 1714–1729. https://doi.org/10.1109/TMI.2010.2050897

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature*

Reviews Neuroscience, *10*(3), 211–223. https://doi.org/10.1038/nrn2573

Sasaki, M., Shibata, E., Tohyama, K., Takahashi, J., Otsuka, K., Tsuchiya, K., Takahashi, S.,

Ehara, S., Terayama, Y., & Sakai, A. (2006). Neuromelanin magnetic resonance imaging

of locus ceruleus and substantia nigra in Parkinson's disease. *NeuroReport*, *17*(11),

1215–1218. https://doi.org/10.1097/01.wnr.0000227984.84927.a7

Satoh, A., & Iijima, K. M. (2019). Roles of tau pathology in the locus coeruleus (LC) in age-

associated pathophysiology and Alzheimer's disease pathogenesis: Potential strategies to

protect the LC against aging. *Brain Research*, *1702*, 17–28.

https://doi.org/10.1016/J.BRAINRES.2017.12.027

Schoenberg, M. R., Dawson, K. A., Duff, K., Patton, D., Scott, J. G., & Adams, R. L. (2006).

Test performance and classification statistics for the Rey Auditory Verbal Learning Test

in selected clinical samples. *Archives of Clinical Neuropsychology*, *21*(7), 693–703.

https://doi.org/10.1016/j.acn.2006.06.010

Snider, B. J., Norton, J., Coats, M. A., Chakraverty, S., Hou, C. E., Jervis, R., Lendon, C. L.,
Goate, A. M., McKeel, D. W., & Morris, J. C. (2005). Novel presenilin 1 mutation
(S170F) causing Alzheimer disease with lewy bodies in the third decade of life. *Archives
of Neurology*, *62*(12), 1821–1830. https://doi.org/10.1001/archneur.62.12.1821

Spina, S., La Joie, R., Petersen, C., Nolan, A. L., Cuevas, D., Cosme, C., Hepker, M., Hwang,
J.-H., Miller, Z. A., Huang, E. J., Karydas, A. M., Grant, H., Boxer, A. L., Gorno-
Tempini, M. L., Rosen, H. J., Kramer, J. H., Miller, B. L., Seeley, W. W., Rabinovici, G.
D., & Grinberg, L. T. (2021). Comorbid neuropathological diagnoses in early versus late-
onset Alzheimer's disease. *Brain : A Journal of Neurology*, *144*(7), 2186–2198.
https://doi.org/10.1093/brain/awab099

Stratmann, K., Heinsen, H., Korf, H. W., Del Turco, D., Ghebremedhin, E., Seidel, K.,
Bouzrou, M., Grinberg, L. T., Bohl, J., Wharton, S. B., Den Dunnen, W., & Rüb, U.
(2016). Precortical Phase of Alzheimer's Disease (AD)-Related Tau Cytoskeletal
Pathology. *Brain Pathology*, *26*(3), 371–386. https://doi.org/10.1111/bpa.12289

Sun, W., Tang, Y., Qiao, Y., Ge, X., Mather, M., Ringman, J. M., & Shi, Y. (2020). A
probabilistic atlas of locus coeruleus pathways to transentorhinal cortex for connectome
imaging in Alzheimer's disease. *NeuroImage*, *223*, 117301.
https://doi.org/10.1016/j.neuroimage.2020.117301

Takahashi, J., Shibata, T., Sasaki, M., Kudo, M., Yanezawa, H., Obara, S., Kudo, K., Ito, K.,
Yamashita, F., & Terayama, Y. (2015). Detection of changes in the locus coeruleus in
patients with mild cognitive impairment and Alzheimer's disease: High-resolution fast
spin-echo T1-weighted imaging. *Geriatrics and Gerontology International*, *15*(3), 334–
340. https://doi.org/10.1111/ggi.12280

Takeuchi, T., Duszkiewicz, A. J., Sonneborn, A., Spooner, P. A., Yamasaki, M., Watanabe, M.,
Smith, C. C., Fernández, G., Deisseroth, K., Greene, R. W., & Morris, R. G. M. (2016).

Locus coeruleus and dopaminergic consolidation of everyday memory. *Nature*, *537*(7620), 357–362. https://doi.org/10.1038/nature19325

Tanzi, R. E., & Bertram, L. (2005). Twenty years of the Alzheimer's disease amyloid hypothesis: A genetic perspective. In *Cell* (Vol. 120, Issue 4, pp. 545–555). Cell Press. https://doi.org/10.1016/j.cell.2005.02.008

Teng, E. L., Hasegawa, K., Homma, A., Imai, Y., Larson, E., Graves, A., Sugimoto, K., Yamaguchi, T., Sasaki, H., Chiu, D., & White, L. R. (1994). The Cognitive Abilities Screening Instrument (CASI): A Practical Test for Cross-Cultural Epidemiological Studies of Dementia. *International Psychogeriatrics*, *6*(1), 45–58. https://doi.org/10.1017/s1041610294001602

Theofilas, P., Dunlop, S., Heinsen, H., & Grinberg, L. T. (2015). Turning on the light within: Subcortical nuclei of the isodentritic core and their role in Alzheimer's disease pathogenesis. *Journal of Alzheimer's Disease*, *46*(1), 17–34. https://doi.org/10.3233/JAD-142682

Theofilas, P., Ehrenberg, A. J., Dunlop, S., Di Lorenzo Alho, A. T., Nguy, A., Leite, R. E. P., Rodriguez, R. D., Mejia, M. B., Suemoto, C. K., Ferretti-Rebustini, R. E. D. L., Polichiso, L., Nascimento, C. F., Seeley, W. W., Nitrini, R., Pasqualucci, C. A., Jacob Filho, W., Rueb, U., Neuhaus, J., Heinsen, H., & Grinberg, L. T. (2017). Locus coeruleus volume and cell population changes during Alzheimer's disease progression: A stereological study in human postmortem brains with potential implication for early-stage biomarker discovery. *Alzheimer's and Dementia*, *13*(3), 236–246. https://doi.org/10.1016/j.jalz.2016.06.2362

Tombaugh, T. (2004). Trail Making Test A and B: Normative data stratified by age and education. *Archives of Clinical Neuropsychology*, *19*(2), 203–214. https://doi.org/10.1016/S0887-6177(03)00039-8

Tona, K. D., Keuken, M. C., de Rover, M., Lakke, E., Forstmann, B. U., Nieuwenhuis, S., &

van Osch, M. J. P. (2017). In vivo visualization of the locus coeruleus in humans:

Quantifying the test–retest reliability. *Brain Structure and Function*, *222*(9), 4203–4217.

https://doi.org/10.1007/s00429-017-1464-5

Uematsu, A., Tan, B. Z., Ycu, E. A., Cuevas, J. S., Koivumaa, J., Junyent, F., Kremer, E. J.,

Witten, I. B., Deisseroth, K., & Johansen, J. P. (2017). Modular organization of the

brainstem noradrenaline system coordinates opposing learning states. *Nature*

Neuroscience, *20*(11), 1602–1611. https://doi.org/10.1038/nn.4642

Van Cauwenberghe, C., Van Broeckhoven, C., & Sleegers, K. (2016). The genetic landscape

of Alzheimer disease: Clinical implications and perspectives. In *Genetics in Medicine*

(Vol. 18, Issue 5, pp. 421–430). Nature Publishing Group.

https://doi.org/10.1038/gim.2015.117

Wagatsuma, A., Okuyama, T., Sun, C., Smith, L. M., Abe, K., & Tonegawa, S. (2018). Locus

coeruleus input to hippocampal CA3 drives single-trial learning of a novel context.

Proceedings of the National Academy of Sciences of the United States of America,

115(2), E310–E316. https://doi.org/10.1073/pnas.1714082115

Wang, H., Suh, J. W., Das, S. R., Pluta, J. B., Craige, C., & Yushkevich, P. A. (2013). Multi-

atlas segmentation with joint label fusion. *IEEE Transactions on Pattern Analysis and*

Machine Intelligence, *35*(3), 611–623. https://doi.org/10.1109/TPAMI.2012.143

Wechsler, D. (1981). *WAIS-R manual: Wechsler Adult Intelligence Scale-Revised*.

Psychological Corporation.

Wechsler, D. (1997). *WAIS-III : administration and scoring manual : Wechsler Adult*

Intelligence Scale (3rd ed.). Psychological Corporation.

Weinshenker, D. (2018). Long road to ruin: Noradrenergic dysfunction in neurodegenerative

disease. *Trends in Neurosciences*, *41*(4), 211–223.

https://doi.org/10.1016/j.tins.2018.01.010

World Health Organization. (2004). *ICD-10 : international statistical classification of*

diseases and related health problems : tenth revision (2nd ed). World Health

Organization.

Ye, R., Rua, C., O'Callaghan, C., Jones, P. S., Hezemans, F. H., Kaalund, S. S., Tsvetanov, K.

A., Rodgers, C. T., Williams, G., Passamonti, L., & Rowe, J. B. (2021). An in vivo

probabilistic atlas of the human locus coeruleus at ultra-high field. *NeuroImage*, *225*,

117487. https://doi.org/10.1016/j.neuroimage.2020.117487

Yescas, P., Huertas-Vazquez, A., Villarreal-Molina, M. T., Rasmussen, A., Tusié-Luna, M. T.,

López, M., Canizales-Quinteros, S., & Alonso, M. E. (2006). Founder effect for the

Ala431Glu mutation of the presenilin 1 gene causing early-onset Alzheimer's disease in

Mexican families. *Neurogenetics*, *7*(3), 195–200. https://doi.org/10.1007/s10048-006-

0043-3

1267

1268

1269 **9. Supplementary information**

1270

1271 **Table S1:** Associations of locus coeruleus intensity and regional tau burden

Region of interest[1]	*rho*	*p*
bankssts	-0.688	0.004
transversetemporal	-0.656	0.007
postcentral	-0.641	0.009
inferiorparietal	-0.641	0.009
inferiortemporal	-0.638	0.009
cuneus	-0.624	0.012
fusiform	-0.609	0.014
parstriangularis	-0.582	0.02
pericalcarine	-0.553	0.029
posteriorcingulate	-0.544	0.032
precuneus	-0.532	0.036
caudalmiddlefrontal	-0.526	0.038
entorhinal	-0.524	0.04
lingual	-0.509	0.046
middletemporal	-0.509	0.046
isthmuscingulate	-0.503	0.049
temporalpole	-0.497	0.052
lateraloccipital	-0.494	0.054
caudalanteriorcingulate	-0.488	0.057
rostralanteriorcingulate	-0.476	0.064
superiorparietal	-0.459	0.076
paracentral	-0.459	0.076
parahippocampal	-0.456	0.078
insula	-0.441	0.089
lateralorbitofrontal	-0.435	0.094
parsorbitalis	-0.418	0.109
frontalpole	-0.409	0.117
parsopercularis	-0.403	0.123
precentral	-0.382	0.145
medialorbitofrontal	-0.376	0.151
superiortemporal	-0.365	0.165
superiorfrontal	-0.312	0.239
rostralmiddlefrontal	-0.282	0.288
supramarginal	-0.232	0.385

1272 *Note*: [1]Parcellation based on Desikan et al., (2006).
1273 Tau data was averaged across hemispheres before analysis (see *Material and methods*).
1274 *Rho* and *p* values are based on non-parametric Spearman's correlations (n = 16).
1275

1276 **Table S2:** Associations of locus coeruleus intensity and cognitive performance

Cognitive task	rho	p
Craft Story	0.62	0.008
Complex Figure Delay	0.589	0.013
Trails B	0.498	0.042
Number Span Backward	0.459	0.064
SEVLT_total (trial 1–5)	0.447	0.072
CASI	0.286	0.265
SEVLT long delay	0.006	0.981
MoCA	-0.02	0.94
Digit Symbol	-0.16	0.539
CDR	-0.405	0.107

1277 *Note*: *Rho* and *p* values are based on non-parametric Spearman's correlations (n = 17). See
1278 *Material and methods* for a description of the cognitive tasks.
1279