Valuation system connectivity is correlated with poly-drug use in young adults

Kavinash Loganathan[1*], Jinglei Lv[2,3,4], Vanessa Cropley[3], Andrew Zalesky[3,4], Eric Tatt Wei Ho[1,5],

1 Centre for Intelligent Signal & Imaging Research, Universiti Teknologi PETRONAS, Perak, Malaysia.

2 School of Biomedical Engineering & Brain and Mind Centre, The University of Sydney, Sydney, Australia

3 Melbourne Neuropsychiatry Centre, Department of Psychiatry, University of Melbourne & Melbourne Health, Melbourne Australia

4 Department of Biomedical Engineering, University of Melbourne, Melbourne, Australia

5 Department of Electrical & Electronics Engineering, Universiti Teknologi PETRONAS, Perak, Malaysia

*Corresponding author:

Kavinash Loganathan

Centre for Intelligent Signal & Imaging Research

Universiti Teknologi PETRONAS

32610 Bandar Seri Iskandar, Perak, Malaysia.

Tel: +6 05 368 7893

Fax: +6 05 365 7443

Email: kavinash@gmail.com

Abstract

Poly-drug consumption contributes to fatal overdose in more than half of all poly-drug users. Analyzing decision-making networks may give insight into the motivations behind poly-drug use. We correlated average functional connectivity of the valuation system (VS), executive control system (ECS) and valuation-control complex (VCC) in a large population sample (n=992) with drug use behaviour. VS connectivity is correlated with sedative use, ECS connectivity is separately correlated with hallucinogens and opiates. Network connectivity is also correlated with drug use via two-way interactions with other substances including alcohol and tobacco. These preliminary findings can contribute to our understanding of the common combinations of substance co-use and associated neural patterns.

Main Text

The pursuit of a desired internal state is known to drive drug-seeking and -taking behaviour (Hellberg et al., 2019). For example, an individual in pain may consume sedating substances such as marijuana or sedatives for relief (Choi et al., 2017; Shurman et al., 2010), while individuals looking to experience enhanced levels of euphoria may choose stimulants or cocaine (Breiter et al., 1997; Klega and Keehbauch, 2018). Poly-drug users mix-and-match different combinations in an attempt to increase the efficacy (Scott et al., 2007) or attenuate drawbacks (Fisk et al., 2011) of a particular substance. This tendency to co-use multiple drugs can cause a fatal overdose (Su et al., 2019). Many rehabilitating patients having a history of co-using more than one drug (Su et al., 2019; Wang et al., 2018) thereby requiring different treatment strategies (Hiebler-Ragger and Unterrainer, 2019; Weigl et al., 2017) due to higher levels of risky behaviour and psychopathology, decreased cognitive functioning as well as poorer treatment outcome and engagement (Wang et al., 2017).

Addictive drugs are thought to disrupt the brains' reward system (Koob and Le Moal, 2008). These substances are consumed to achieve desired internal states (e.g. euphoria or pain relief), with regions of the Valuation and Executive Control System responsible for weighing drug value in relation to these states (Loganathan and Ho, 2021). Given that valuation of drugs changes based on internal and external factors as well as the increased complexity of the addiction phenotype among polyu-drug users, this study will focus on the interaction between Valuation and Executive Control System connectivity and poly-drug use in young adults. Value, the currency of choice, fluctuates based on internal states (e.g. satiety, deprivation) as well as external circumstances (e.g. socioeconomic factors, health status) (Compton et al., 2007; Davis and Johnson, 2008; Helie et al., 2017). The process of assigning value to different options may be regulated by the valuation and executive control systems of the brain (van den Bos and McClure, 2013). The valuation system (VS, comprised of the

ventromedial prefrontal cortex, posterior cingulate cortex, amygdala and nucleus accumbens) may be involved in estimating the incentive value of different options (van den Bos and McClure, 2013) and is hypothesized to interface between pleasure and pain (Kringelbach, 2005). The executive control system (ECS, consisting of the lateral prefrontal cortex, dorsal anterior cingulate cortex and posterior parietal cortex), guides behaviour in pursuit of long-term goals (van den Bos and McClure, 2013). The ECS may assist in refining VS-encoded values by incorporating elements of future-thinking and careful deliberation (van den Bos and McClure, 2013). Addiction is thought to be associated with an imbalance in valuation, as drug-related stimuli are attributed greater salience compared to non-drug stimuli (Helie et al., 2017). The VS and ECS are linked with substance use behaviour, both as a network (Zhai et al., 2015) and as individual regions (Bedi et al., 2015). Normally, the interaction between VS and ECS is balanced, modulating reward processing by assigning values to various reward and non-reward options to facilitate optimal decision-making (Dalley et al., 2011; Xie et al., 2014; Zhai et al., 2015). During drug dependence the valuation system activation appears not to be counteracted by the control system which has since surrendered its regulatory abilities, tilting this once-mutually-reinforcing system towards the instrumental pursuit of drugs (Xie et al., 2014; Zhai et al., 2015).

Addictive drugs operate via different pathways in the brain (Leurquin-Sterk et al., 2018; Solinas et al., 2019). Taken together with the tendency to exclude poly-drug users from individual substance use neuroimaging studies (Zou et al., 2015), these limitations have restricted the development of a unifying concept underpinning drug use, further complicating efforts to develop focused treatment plans targeted at poly-drug users (for a review, see Wang *et al* (2017)). Given that poly-drug use contributes to more than 50% of fatal overdoses (Coffin et al., 2003), interrogating the interactions between different drugs is a necessary first step in unravelling the complex psychopathology underpinning drug addiction and what is

required to treat it. This study aims to provide preliminary evidence linking functional

connectivity of the VS and ECS with poly-substance use to infer the common combinations

of substance co-use and associated neural patterns. This is done by correlating the

connectivity of the VS, ECS and the valuation-control complex (VCC, a hybrid network

comprised of VS and ECS regions) with poly-drug use in the Human Connectome Project

Healthy Young Adult dataset using multivariate generalized linear models. We hypothesize

that functional connectivity within the VS, ECS and VCC would all be correlated with

alcohol, tobacco, opiates, cocaine, stimulants, sedatives and marijuana use. We further

hypothesize that in some cases, connectivity may be correlated with substance use via a

paired interaction with another drug (eg VS connectivity paired with sedative use may be

correlated with cocaine use).

Functional MRI data from the Human Connectome Project (HCP, Smith *et al.*, 2013)

was sourced for healthy adults of both genders (n = 992, 463 males, 529 females; age range =

22-37). Recruitment procedures, inclusion/exclusion criteria and pre-processing steps are

described elsewhere (Glasser et al., 2013; Smith et al., 2013a). Additional steps such as

global signal regression are deemed too aggressive for use with HCP imaging data and are

not implemented as it has been known to introduce artifactual changes in correlations

patterns, fundamentally altering interregional correlations. These changes are thought to be

dependent on the underlying true interregional correlation pattern (Anderson et al., 2011;

Glasser et al., 2013; Saad et al., 2012; Smith et al., 2013b). The functional scans used in this

study have already been cleaned of structured noise via joint use of independent component

analysis (ICA) and FMRIB's ICA-based X-noisifier (FIX) to remove non-neural spatio-

temporal components from each highpass-filtered fMRI scan. Initial head motion correction

is performed by registering fMRI data to a distorted gradient echo EPI single-band reference

image. For each subject, rigid body head motion was estimated using FSL's MCFLIRT

routine to derive a motion transformation matrix for each time point. Each transform is described by 6 motion parameters consisting of three translations and three rotations, which is then condensed to form a vector of framewise displacement by summing the absolute values of the differentials of the 6 parameters (Power et al., 2012). As a part of the artefactual cleanup process, 24-counfound timeseries derived from the motion estimation (comprised of the 6 rigid body-parameter timeseries, their backwards-looking temporal derivatives, plus all 12 resulting regressors squared) had temporal highpass filtering applied to them and then regressed out of the data aggressively (WU-Minn, 2017). The HCP pipeline ultimately resulted in the removal of spatial and temporal artefacts without removing potentially useful data, particularly relevant for resting-state fMRI data that is particularly sensitive to artefactual corruption across voxels (Smith et al., 2013a).

Only subjects with all four repeated resting-state fMRI sessions (first and second scan sessions with left-right and right-left phase encoding directions), were used for this study. Preparation of masks used in this study has been detailed elsewhere (Loganathan et al., 2020). Briefly, the VS, ECS and VCC were delineated using the binary masks that combined regions of interest (ROIs) from both the Desikan-Killiany (Desikan et al., 2006) and Destrieux (Destrieux et al., 2010) parcellations. All anatomical labels were extracted and merged using the FMRIB Software Library (Smith *et al.*, 2004, https://fsl.fmrib.ox.ac.uk/fsl/). The complete list of all anatomical ROIs used to delineate the VS and ECS are shown in Table Supplementary 1, along with their central coordinates given in Montreal Neurological Institute (MNI) space.

The following substance use measures were obtained from the Human Connectome Project Restricted Access Data (Table 1). These substance use and frequency measures were collected using two approaches: a detailed questionnaire prepared specifically for the Human Connectome Project (for short-term alcohol and tobacco use), followed by the Semi-

Structured Assessment for the Genetics of Alcoholism (SSAGA) for long-term alcohol,

tobacco and illicit drug use. Additionally, parental use history, age, gender and framewise

displacement (FD) measures were included as demographic variable.

Table 1: List of drug-based dependent variables and independent variables comprised of both substance use measures, as well as demographic and family use history variables.

Dependent Variables	Independent Variables		
Cocaine Use Hallucinogen Use Opiate Use Sedative Use Stimulant Use Marijuana Use	Average Brain Network Connectivity	Substance Use Measures	Demographics and Family Use Variables
	Valuation System (VS) Executive Control System (ECS) Valuation-Control Complex (VCC)	Total Drinks 7 Days Drink Frequency 7 Days Drinks/Day 12 Months Drink Frequency 12 Months Total Tobacco Use 7 Days Tobacco Use Frequency 7 Days Cocaine Use Hallucinogen Use Opiate Use Sedative Use Stimulant Use Marijuana Use	Age Gender Father Use History Mother Use History Average Framewise Displacement

The rsfMRI signal was averaged over all voxels comprising each ROI (node) of the

VS (12 nodes), ECS (20 nodes) and VCC (32 nodes), yielding a regionally averaged signal

for each node. The Pearson correlation coefficient in the regionally averaged signals were

then computed between all pairs of nodes, resulting in a 12 x 12 (VS), 20 x 20 (ECS) and 32

x 32 (VCC) connectivity matrix for each of the four scan runs (Rest 1-LR, Rest 1-RL, Rest 2-

LR, Rest 2-RL). This computation was repeated for each subject. The four connectivity

matrices computed from the four rsfMRI sessions for each subject were then averaged to

improve the signal-to-noise ratio, yielding a single connectivity matrix for each of the three

systems for every subject (see Supplementary Results). These connectivity matrices were

then averaged per subject, such that for each of the VS, ECS and VCC, each subject's

connectivity would be represented by a single value. Generalized linear models (GLM) were then computed to correlate the average connectivity of VS, ECS and VCC with each subject's substance use profile. All dependent variables, independent variables and connectivity measures are standardized to z-scores with mean of 0 and standard deviation of 1. Regression coefficients are therefore representative of SD change in outcome per unit SD change in predictor and may be interpreted as partial correlation coefficients (Bring, 1994). Throughout this study, a total of 288 tests were performed (16 independent variables X 3 brain networks X 6 dependent variables). Each dependent variable was tested against 15 independent variables for all 3 networks (VS, ECS, VCC) producing a single model per dependent variable per network. For the purposes of illustration, let us assume one wishes to model Cocaine use (dependent variable) and its correlation with Valuation System average connectivity (brain network). The 16 independent variables would therefore comprise of Total Drinks 7 Days, Drink Frequency 7 Days, Drinks/Day 12 Months, Drink Frequency 12 Months, Total Tobacco Use 7 Days, Tobacco Use Frequency 7 Days, Hallucinogen, Opiate, Sedatives, Stimulants, Marijuana, Age, Gender, Fathers' Use History, Mothers' Use History, Average Framewise Displacement). Multiple comparisons across all models were controlled using the False Discovery Rate using the p-values attributed to each component of each model. p-values < 0.05 post-correction are deemed significant.

Correlation between substance use measures is shown in Table 2. The most highly correlated pairing of substances are opiates and sedatives ($r^2 = 0.725$), followed by cocaine and hallucinogens ($r^2 = 0.614$). Marijuana use was not strongly correlated with other substances, except with hallucinogens ($r^2 = 0.559$). Alcohol (Total Drinks 7 Days, Drink Frequency 7 Days, Drinks/Day 12 Months and Drink Frequency 12 Months) and tobacco measures (Total Tobacco 7 Days, Tobacco Frequency 7 Days) did not show strong correlation with illicit drug use measures. Stronger correlations were reported between

alcohol and Marijuana measures as well as tobacco and Marijuana measures, but even these r^2 values were no higher than 0.326 and 0.336 respectively. Correlations between resting-state functional connectivity and substance use measures are shown in Table 3. Network connectivity of all three systems is weakly correlated with substance use measures, with no r^2 value higher than 0.1.

Table 2: Correlation matrix of substance use measures. Intercepts highlighted in grey.

	Total Drinks 7 Days	Drink Frequency 7 Days	Drinks/Day 12 Months	Drink Frequency 12 Months	Total Tobacco 7 Days	Tobacco Frequency 7 Days	Cocaine	Hallucinogens	Opiates	Sedatives	Stimulants	Marijuana
Total Drinks 7 Days	1.000											
Drink Frequency 7 Days	0.999	1.000										
Drinks/Day 12 Months	0.282	0.282	1.000									
Drink Frequency 12 Months	-0.410	-0.410	-0.157	1.000								
Total Tobacco 7 Days	0.018	0.018	0.195	-0.034	1.000							
Tobacco Frequency 7 Days	0.077	0.077	0.208	-0.051	0.803	1.000						
Cocaine	0.119	0.119	0.197	-0.051	0.181	0.189	1.000					
Hallucinogens	0.148	0.148	0.209	-0.088	0.142	0.167	0.614	1.000				
Opiates	0.103	0.103	0.151	-0.028	0.193	0.220	0.545	0.578	1.000			
Sedatives	0.048	0.048	0.169	0.020	0.194	0.227	0.581	0.536	0.725	1.000		
Stimulants	0.029	0.029	0.220	0.000	0.097	0.165	0.544	0.480	0.535	0.570	1.000	
Marijuana	0.241	0.241	0.326	-0.131	0.270	0.336	0.436	0.559	0.444	0.411	0.401	1.000

Table 3: Correlation between VS, ECS and VCC connectivity and substance use measures

	VS	ECS	VCC
Cocaine	0.052	0.004	0.022
Hallucinogens	0.093	0.054	0.073
Opiates	0.080	0.058	0.072
Sedatives	0.058	0.024	0.040
Stimulants	0.042	0.005	0.017
MJ	0.058	0.039	0.051

Average connectivity of VS, ECS and VCC is correlated, either independently or via a two-way interaction with substance use (Table 4). VS connectivity is correlated with Sedative use (Estimate=0.04, t=2.3, p=0.03) while ECS connectivity is correlated with Hallucinogen (Estimate=0.08, t=2.4, p=0.025) and Opiate (Estimate=0.05, t=2.3, p=0.03) use. Cocaine use is correlated with VS, ECS and VCC connectivity via two-way interactions with Total Tobacco 7 Days, Hallucinogen use, Opiate use, Sedative use and Fathers' Use History. Opiate use is correlated with VS (Estimate=0.07, t=3.3, p=0.002), ECS (Estimate=0.08, t=3.6, p=0.0008) and VCC (Estimate=0.1, t=4.4, p<0.00001) connectivity via two-way interaction with Total Tobacco 7 Days. Sedative use is correlated with VS (Estimate=0.04, t=2.6, p=0.017), ECS (Estimate=0.04, t=2.2, p=0.04) and VCC (Estimate=0.04, t=2.3, p=0.03) connectivity via two-way interaction with Drinks/Day 12 Months. Stimulant use is correlated with VS (Estimate=-0.07, t=-2.9 p=0.007), ECS (Estimate=-0.25, t=-7.2, p<0.00001) and VCC (Estimate=-0.16, t=-4.8 p<0.00001) connectivity via two-way interaction with Opiate use. Hallucinogen use is correlated with VS (Estimate=-0.06, t=-2.6, p=0.015) and VCC (Estimate=-0.05, t=7.8, p=0.04) connectivity via two-way interaction with Total Tobacco 7 Days. Opiate use is correlated with ECS (Estimate=0.19, t=5.8, p<0.00001) and VCC (Estimate=0.09, t=3.6, p=0.0009) connectivity via two-way interaction with Hallucinogen use. Sedative use is correlated with ECS (Estimate=0.05, t=2.45, p=0.02) and VCC (Estimate=0.08, t=3.6, p=0.0006) connectivity via two-way interaction with Tobacco Frequency 7 Days. Sedative use is also correlated with ECS (Estimate=0.19, t=7.96 p<0.00001) and VCC (Estimate=0.18, t=8.3, p<0.00001) connectivity via two-way interaction with Opiate use. Sedative use is correlated with VS (Estimate=0.09, t=3.17 p=0.003) and ECS (Estimate=-0.08, t=-2.86, p=0.008) use via two-way interaction with Cocaine use.

Table 4: VS, ECS and VCC connectivity is correlated with drug use, both independently (highlighted) and as part of a two-way interaction with other substances. Effect sizes of each GLM are represented by r^2 values. Estimates displayed are standardized partial coefficients for each component of the GLM. *p*-values shown here are FDR-corrected. Exact p-values are displayed where possible. If the p-value is too small (less than 0.00001), then it is displayed as p<0.00001. Father, Fathers' Use History; Mother, Mothers' Use History; E, Estimate

Dependents	VS	ECS	VCC
Cocaine	$r^2 = 0.7227$ VS – Total Tobacco 7 Days (E=0.07, t=2.92 p=0.007) VS – Tobacco Frequency 7 Days (E=0.08, t=2.53 p=0.02) VS – Hallucinogens (E=0.12, t=3.79 p=0.00038) VS – Opiates (E=-0.2, t=-6.1, p<0.00001) VS – Sedatives (E=-0.06, t=-2.83 p=0.0085) VS – Father (E=-0.07, t=-3.3 p=0.0021)	$r^2 = 0.7208$ ECS – Total Tobacco 7 Days (E=0.06, t=2.84 p=0.08) ECS – Hallucinogens (E=0.17, t=4.65 p=0.00001) ECS – Opiates (E=-0.21, t=-5.82, p<0.00001) ECS – Sedatives (E=-0.06, t=-2.66 p=0.01) ECS – Father (E=-0.06, t=-3.3, p=0.002)	$r^2 = 0.7198$ VCC – Total Tobacco 7 Days (E=0.07, t=3.0 p=0.005) VCC – Hallucinogens (E=0.18, t=5.4 p<0.00001) VCC – Opiates (E=-0.20, t=-5.7, p<0.00001) VCC – Sedatives (E=-0.05, t=-2.22, p=0.04) VCC – Father (E=-0.06, t=-3.0, p=0.006)
Hallucinogens	$r^2 = 0.6707$ VS – Total Tobacco 7 Days (E=-0.06, t=-2.6, p=0.015) VS – Tobacco Frequency 7 Days (E=0.14, t=4.8 p<0.00001) VS – Cocaine (E=-0.21, t=-6.5 p<0.00001) VS – Opiates (E=0.22, t=7.15, p<0.00001) VS – Stimulants (E=0.006, t=2.4, p<0.02)	$r^2 = 0.7856$ ECS (E=0.08, t=2.4, p=0.025) ECS – Tobacco Frequency 7 Days (E=0.03, t=2.37 p=0.03) ECS – Gender (E=-0.1, t=-2.24, p=0.03)	$r^2 = 0.6612$ VCC – Total Tobacco 7 Days (E=-0.05, t=-2.2 p=0.04) VCC – Opiates (E= 0.19, t=7.8 p<0.00001)
Opiates	$r^2 = 0.7056$ VS – Drink Frequency 12 Months (E=-0.05, t=0.19, p=0.03) VS – Total Tobacco 7 Days (E=0.07, t=3.3, p=0.002) VS – Tobacco Frequency 7 Days (E=0.1, t=5.9, p<0.00001)	$r^2 = 0.7137$ ECS (E=0.05, t=2.3, p=0.03) ECS – Total Tobacco 7 Days (E=0.08, t=3.6, p=0.0008) ECS – Hallucinogens (E=0.19, t=5.8, p<0.00001) ECS – Sedatives (E=-0.07, t=-3.1, p=0.005)	$r^2 = 0.6996$ VCC – Drinks/Day 12 Months (E=-0.04, t=-2.3 p=0.024) VCC – Total Tobacco 7 Days (E=0.1, t=4.4, p<0.00001) VCC – Hallucinogens (E=0.09, t=3.6, p=0.0009)
Sedatives	$r^2 = 0.7797$ VS (E=0.04, t=2.3, p=0.03) VS – Drinks/Day 12 Months (E=0.04, t=2.6, p=0.017) VS – Cocaine (E=0.09, t=3.17 p=0.003) VS – Hallucinogens (E=-0.11, t=-4.42 p<0.00001) VS – Opiates (E=0.11, t=5.7 p<0.00001)	$r^2 = 0.806$ ECS – Drinks/Day 12 Months (E=0.04, t=2.2, p=0.04) ECS – Tobacco Frequency 7 Days (E=0.05, t=2.45, p=0.02) ECS – Cocaine (E=-0.08, t=-2.86, p=0.008) ECS – Opiates (E=0.19, t=7.96 p<0.00001)	$r^2 = 0.8083$ VCC – Drinks/Day 12 Months (E=0.04, t=2.3, p=0.03) VCC – Tobacco Frequency 7 Days (E=0.08, t=3.6, p=0.0006) VCC – Hallucinogens (E=-0.14, t=-4.7, p<0.00001) VCC – Opiates (E=0.18, t=8.3, p<0.00001)
Stimulants	$r^2 = 0.6265$ VS – Drink Frequency 12 Months (E=-0.05, t=-2.5 p=0.022) VS – Opiates (E=-0.07, t=-2.9 p=0.007) VS – Father (E=-0.07, t=-2.4 p=0.025)	$r^2 = 0.6392$ ECS – Tobacco Frequency 7 Days (E=0.08, t=3.0 p=0.005) ECS – Hallucinogens (E=-0.1, t=-3.2 p=0.0033) ECS – Opiates (E = -0.25, t=-7.2, p<0.00001)	$r^2 = 0.6288$ VCC – Tobacco Frequency 7 Days (E=0.06, t=2.9, p=0.04) VCC – Hallucinogens (E=-0.09, t=-3.1 p=0.0035) VCC – Opiates (E=-0.16, t=-4.8, p<0.00001)
Marijuana	$r^2 = 0.4715$ VS – Tobacco Frequency 7 Days (E=-0.5, t=-2.5, p=0.019) VS – Stimulants (E=0.7, t=2.62, p=0.015)	$r^2 = 0.4703$ ECS – Drink Frequency 12 Months (E=0.73, t=3.5 p=0.00094)	$r^2 = 0.4732$ VCC – Drink Frequency 12 Months (E=0.7, t=3.4, p=0.0018) VCC – Opiates (E=-0.8, t=-3.0, p=0.005) VCC – Sedatives (E=0.6, t=2.5, p=0.018)

Research suggests that substance use may stem in part from a disrupted decision-making system stemming from aberrant valuation of drugs (Zhai et al., 2015). The value of drugs varies between individuals, driven by hedonic sensations in some or pain relief in others (Hogarth, 2020). These underlying motivations increase the value of drugs and bias choice towards their use (Hogarth and Field, 2020). Additionally, connections both within and between the Valuation and Executive Control System act as dependency risk and resilience markers (Ersche et al., 2020). Here, we correlated the average network connectivity of the Valuation and Executive Control Systems, as well as the Valuation-Control Complex with drug use in the brains of healthy young adults. Common co-use patterns and their network correlations can thus be inferred based on trends observed in this study. Understanding the needs of one who takes addictive substances with respect to a desired internal state (e.g., euphoria or numbness) represents the first step to improving existing forms of interventions, with greater focus on individual needs that are being met via substance use.

Correlations are observed between combinations of addictive substances, for example Sedatives and Cocaine, Opiates and Sedatives. Highly correlated pairs of drugs reflect popular combinations used to achieve a particular state (euphoria, numbness etc) (Floyd et al., 2010; Laursen et al., 2016). Thus, two-way interactions involving substance use measures and average network connectivity of choice networks may signify the high value placed on these combinations to achieve said state. Stimulating and sedating substances are commonly co-used due to their synergistic interactions. Stimulant-type substances (cocaine, stimulants like amphetamine) energize the user, producing longer-lasting euphoria while off-setting sedating effects (Hernández-López et al., 2002). Sedative-type substances (alcohol, tobacco, marijuana, pain-killing sedatives, heroin) calm the user, ameliorating unpleasant side-effects of stimulant use such as anxiety and aggression (Fisk et al., 2011).

Tobacco and alcohol (Drink Frequency, Total Drinks and Drinks/Day) use measures form two-way interactions with VS, ECS and VCC average network connectivity for all six substances tested as dependent variables (Cocaine, Hallucinogens, Opiates, Sedatives, Stimulants, Marijuana). Tobacco and alcohol both possess sedating qualities (Fultz et al., 2021; Kishore, 2014; Walker, 1980) and are considered gateway substances that pave the way towards the development of illicit drug consumption (Kandel and Kandel, 2015), underpinning poly-substance use (Anderson et al., 2018; Jongenelis et al., 2019; Moss et al., 2014). Stimulants complement alcohol to produce longer-lasting euphoria while off-setting the sedating qualities of alcohol (Hernández-López et al., 2002), reducing the anxiety and agitation (Fisk et al., 2011; Knackstedt and Ettenberg, 2005). Alcohol and sedating substances (opiates, sedatives, marijuana) are often co-used for pain management (Downey et al., 2013; Hood et al., 2020; Ronen et al., 2010; Wilson et al., 2020), albeit at the expense of cognitive control over reward-driven behaviour (Crean et al., 2011; Müller-Oehring et al., 2013). Similarly, tobacco use sensitizes the reward centres of the brain making one susceptible to other drugs (Kandel and Kandel, 2015). Tobacco is thought to encourage cocaine and opiate use (Cross et al., 2017; Mello et al., 1980; Stark and Campbell, 1993) by stimulating the reward centres of the brain (Cross et al., 2017; Nuechterlein et al., 2016). The findings of the current study support existing evidence regarding the involvement of alcohol and tobacco in poly-drug use, as these two substances form two-way interactions with average connectivity of VS, ECS and VCC for all illicit drugs (Cocaine, Hallucinogens, Opiates, Sedatives, Stimulants, Marijuana). Given the "gateway" nature of these two substances (Kandel and Kandel, 2015), particularly among young adults (Laursen et al., 2016; McKetin et al., 2014), it is theorized that these participants may have started using alcohol and/or tobacco to achieve an internal state of euphoria or numbness, but began to incorporate illicit drugs to boost the strength of the desired sensation.

Two-way interactions between all three networks and Opiates in Cocaine use could be indicative of "speedball", a cocaine-heroin mixture that induces enhanced feelings of euphoria (Lacy et al., 2014), while reducing the aggression and anxiety (Fisk et al., 2011). A similar effect may underpin other two-way interactions between all three networks and sedating substances (Opiates, Sedatives) with Stimulants and vice versa. Studies revealed ECS involvement in regulation of attentional processing of external cues (Sarter et al., 2001) including auditory (Napoli et al., 2021; Plakke et al., 2013) and visual stimuli (Geliebter et al., 2016). Both stimulant and cocaine use are capable of enhancing the senses (Devonshire et al., 2007; Oken et al., 2006) and inducing euphoria (Bellone et al., 2020; Dlugos et al., 2011) by increasing activation within the ECS (Jan et al., 2014) and VS (Crane et al., 2018) respectively.

Co-use of sedating substances are commonly used for pain management (Peele, 2016) especially when a single substance is found to be inadequate for the purpose (Wilson et al., 2020). The VS acts as the middle ground for processing both pleasure and pain (Leknes and Tracey, 2008). Connections between the VS and ECS regions is observed in patients experiencing chronic pain (Jiang et al., 2016). In the current study, VS, ECS and VCC each form a two-way interaction with alcohol in Sedatives use, suggesting a pain-killing combination. Two-way interactions involving VS, ECS and VCC connectivity and sedating substances (Marijuana, Sedatives) when paired with other sedating substances (such as alcohol) could indicate these combinations of substances are being used for pain management.

This study is not without limitations. There are various types of stimulants, sedatives and hallucinogens, some of which have been manufactured to have pharmacological properties of other abusable substances (Klega and Keehbauch, 2018). Given that each drug has its own mechanism of action that targets specific receptors or pathways, information

regarding the specific types of drugs used may help shed additional light on the relationships between network connectivity and poly-drug use measures. Additionally, global signal regression was not performed in this study. The fMRI scans used in this study have been minimally pre-processed by the HCP team to ensure minimal information loss (Glasser et al., 2013). Global signal regression is deemed too aggressive for use with HCP imaging data and never implemented. This method may introduce artifactual changes in correlations patterns, fundamentally altering interregional correlations. These changes are thought to be dependent on the underlying true interregional correlation pattern (Anderson et al., 2011; Saad et al., 2012).

In conclusion, we correlated the average network connectivity of the Valuation and Executive Control Systems, as well as the Valuation-Control Complex with drug use in the brains of healthy young adults. Common co-use patterns and their network correlations can thus be inferred based on trends observed in this study. Two-way interactions involving substance use measures and average network connectivity of choice networks may signify the high value placed on these combinations to achieve said state. Particularly, tobacco and alcohol use measures form interactions with VS, ECS and VCC average network connectivity across all forms of illicit substances (Cocaine, Hallucinogens, Opiates, Sedatives, Stimulants and Marijuana), suggesting that participants may have started using alcohol and/or tobacco initially but began to incorporate illicit drugs to enhance their experience. Additionally, two-way interactions involving sedating substances (e.g. Opiates) with connectivity of all 3 networks is observed in stimulant-type substances (e.g. Cocaine) and vice-versa, suggesting a combination aimed at increasing feelings of euphoria while dampening anxiety and aggression. Finally, connectivity of all three networks each form a two-way interaction with alcohol among sedating substances, suggesting a combination for pain management. These results provide early evidence linking functional connectivity of the VS and ECS with poly-

substance use, enabling us to infer common combinations of substance co-use and associated neural patterns.

Acknowledgements: Kavinash Loganathan performed analysis, interpreted results and prepared this manuscript, Jinglei Lv was involved in experimental design, assisted in analysis and interpretation of results, Eric Tatt Wei Ho performed analysis and interpreted results, Vanessa Cropley assisted with brain mask generation and provided technical feedback, Andrew Zalesky was involved in experimental design, provided technical feedback and assisted with result interpretation.

Funding: Kavinash Loganathan is grateful to IBRO-APRC and Yayasan Universiti Teknologi PETRONAS (YUTP) for funding to undertake this study. Eric Tatt Wei Ho gratefully acknowledges the Ministry of Higher Education, Malaysia for funding through the Higher Institution Center of Excellence (HI-CoE) program awarded to the Center for Intelligent Signal & Imaging Research, Universiti Teknologi PETRONAS and the YUTP-Fundamental Research Grant award. Vanessa Cropley was supported by an Australian National Health and Medical Research Council (NHMRC) Investigator Grant (1177370).

References

Anderson, J.S., Druzgal, T.J., Lopez-Larson, M., Jeong, E.-K., Desai, K., Yurgelun-Todd, D., 2011. Network anticorrelations, global regression, and phase-shifted soft tissue correction. Hum. Brain Mapp. 32, 919–934. https://doi.org/10.1002/hbm.21079

Anderson, R.E., Hruska, B., Boros, A.P., Richardson, C.J., Delahanty, D.L., 2018. Patterns of co-occurring addictions, posttraumatic stress disorder, and major depressive disorder in detoxification treatment seekers: Implications for improving detoxification treatment outcomes. J. Subst. Abuse Treat. 86, 45–51. https://doi.org/10.1016/j.jsat.2017.12.009

Bedi, G., Lindquist, M.A., Haney, M., 2015. An fMRI-Based Neural Signature of Decisions to Smoke Cannabis. Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol. 40, 2657–2665. https://doi.org/10.1038/npp.2015.135

Bellone, C., Loureiro, M., Lüscher, C., 2020. Drug-Evoked Synaptic Plasticity of Excitatory Transmission in the Ventral Tegmental Area. Cold Spring Harb. Perspect. Med. https://doi.org/10.1101/cshperspect.a039701

Breiter, H.C., Gollub, R.L., Weisskoff, R.M., Kennedy, D.N., Makris, N., Berke, J.D., Goodman, J.M., Kantor, H.L., Gastfriend, D.R., Riorden, J.P., Mathew, R.T., Rosen, B.R., Hyman, S.E., 1997. Acute effects of cocaine on human brain activity and emotion. Neuron 19, 591–611.

Bring, J., 1994. How to Standardize Regression Coefficients. Am. Stat. 48, 209–213. https://doi.org/10.1080/00031305.1994.10476059

Choi, N.G., DiNitto, D.M., Marti, C.N., Choi, B.Y., 2017. Association between Nonmedical Marijuana and Pain Reliever Uses among Individuals Aged 50. J. Psychoactive Drugs 49, 267–278. https://doi.org/10.1080/02791072.2017.1342153

Ciric, R., Rosen, A.F.G., Erus, G., Cieslak, M., Adebimpe, A., Cook, P.A., Bassett, D.S., Davatzikos, C., Wolf, D.H., Satterthwaite, T.D., 2018. Mitigating head motion artifact in functional connectivity MRI. Nat. Protoc. 13, 2801–2826. https://doi.org/10.1038/s41596-018-0065-y

Coffin, P.O., Galea, S., Ahern, J., Leon, A.C., Vlahov, D., Tardiff, K., 2003. Opiates, cocaine and alcohol combinations in accidental drug overdose deaths in New York City, 1990–98. Addiction 98, 739–747.

Compton, W.M., Thomas, Y.F., Stinson, F.S., Grant, B.F., 2007. Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions. Arch. Gen. Psychiatry 64, 566–576. https://doi.org/10.1001/archpsyc.64.5.566

Crane, N.A., Gorka, S.M., Weafer, J., Langenecker, S.A., de Wit, H., Phan, K.L., 2018. Neural activation to monetary reward is associated with amphetamine reward sensitivity. Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol. 43, 1738–1744. https://doi.org/10.1038/s41386-018-0042-8

Crean, R.D., Crane, N.A., Mason, B.J., 2011. An Evidence-Based Review of Acute and Long-Term Effects of Cannabis Use on Executive Cognitive Functions. J. Addict. Med. 5.

Cross, S.J., Lotfipour, S., Leslie, F.M., 2017. Mechanisms and genetic factors underlying co-use of nicotine and alcohol or other drugs of abuse. Am. J. Drug Alcohol Abuse 43, 171–185. https://doi.org/10.1080/00952990.2016.1209512

Dalley, J.W., Everitt, B.J., Robbins, T.W., 2011. Impulsivity, compulsivity, and top-down cognitive control. Neuron 69, 680–694. https://doi.org/10.1016/j.neuron.2011.01.020

Davis, W.R., Johnson, B.D., 2008. Prescription opioid use, misuse, and diversion among street drug users in New York City. Drug Alcohol Depend. 92, 267–276. https://doi.org/10.1016/j.drugalcdep.2007.08.008

Desikan, R.S., Ségonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L., Dale, A.M., Maguire, R.P., Hyman, B.T., Albert, M.S., Killiany, R.J., 2006. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage 31, 968–980. https://doi.org/10.1016/j.neuroimage.2006.01.021

Destrieux, C., Fischl, B., Dale, A., Halgren, E., 2010. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. Neuroimage 53, 1–15. https://doi.org/https://doi.org/10.1016/j.neuroimage.2010.06.010

Devonshire, I.M., Mayhew, J.E.W., Overton, P.G., 2007. Cocaine preferentially enhances sensory processing in the upper layers of the primary sensory cortex. Neuroscience 146, 841–851. https://doi.org/10.1016/j.neuroscience.2007.01.070

Dlugos, A.M., Hamidovic, A., Hodgkinson, C., Shen, P.H., Goldman, D., Palmer, A.A., de Wit, H., 2011. OPRM1 gene variants modulate amphetamine-induced euphoria in humans. Genes. Brain. Behav. 10, 199–209. https://doi.org/10.1111/j.1601-183X.2010.00655.x

Downey, L.A., King, R., Papafotiou, K., Swann, P., Ogden, E., Boorman, M., Stough, C., 2013. The effects of cannabis and alcohol on simulated driving: Influences of dose and experience. Accid. Anal. Prev. 50, 879–886. https://doi.org/10.1016/j.aap.2012.07.016

Ersche, K.D., Meng, C., Ziauddeen, H., Stochl, J., Williams, G.B., Bullmore, E.T., Robbins, T.W., 2020. Brain networks underlying vulnerability and resilience to drug addiction. Proc. Natl. Acad. Sci. U. S. A. 117, 15253–15261.

https://doi.org/10.1073/pnas.2002509117

Fisk, J.E., Murphy, P.N., Montgomery, C., Hadjiefthyvoulou, F., 2011. Modelling the
adverse effects associated with ecstasy use. Addiction 106, 798–805.
https://doi.org/10.1111/j.1360-0443.2010.03272.x

Floyd, L.J., Hedden, S., Lawson, A., Salama, C., Moleko, A.G., Latimer, W., 2010. The
association between poly-substance use, coping, and sex trade among black South
African substance users. Subst. Use Misuse 45, 1971–1987.
https://doi.org/10.3109/10826081003767635

Fultz, E.K., Coelho, M.A., Lieberman, D., Jimenez-Chavez, C.L., Bryant, C.D., Szumlinski,
K.K., 2021. Hnrnph1 is a novel regulator of alcohol reward. Drug Alcohol Depend. 220,
108518.

Geliebter, A., Benson, L., Pantazatos, S.P., Hirsch, J., Carnell, S., 2016. Greater anterior
cingulate activation and connectivity in response to visual and auditory high-calorie
food cues in binge eating: Preliminary findings. Appetite 96, 195–202.
https://doi.org/10.1016/j.appet.2015.08.009

Givens, C.J., 2016. Adverse drug reactions associated with antipsychotics, antidepressants,
mood stabilizers, and stimulants. Nurs. Clin. 51, 309–321.

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L.,
Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M.,
Consortium, W.-M.H.C.P., 2013. The minimal preprocessing pipelines for the Human
Connectome Project. Neuroimage 80, 105–124.
https://doi.org/10.1016/j.neuroimage.2013.04.127

Helie, S., Shamloo, F., Novak, K., Foti, D., 2017. The roles of valuation and reward

processing in cognitive function and psychiatric disorders. Ann. N. Y. Acad. Sci. 1395, 33–48. https://doi.org/10.1111/nyas.13327

Hellberg, S.N., Russell, T.I., Robinson, M.J.F., 2019. Cued for risk: Evidence for an incentive sensitization framework to explain the interplay between stress and anxiety, substance abuse, and reward uncertainty in disordered gambling behavior. Cogn. Affect. Behav. Neurosci. 19, 737–758. https://doi.org/10.3758/s13415-018-00662-3

Hernández-López, C., Farré, M., Roset, P.N., Menoyo, E., Pizarro, N., Ortuño, J., Torrens, M., Camí, J., de La Torre, R., 2002. 3,4-Methylenedioxymethamphetamine (ecstasy) and alcohol interactions in humans: psychomotor performance, subjective effects, and pharmacokinetics. J. Pharmacol. Exp. Ther. 300, 236–244. https://doi.org/10.1124/jpet.300.1.236

Hiebler-Ragger, M., Unterrainer, H.-F., 2019. The Role of Attachment in Poly-Drug Use Disorder: An Overview of the Literature, Recent Findings and Clinical Implications. Front. psychiatry 10, 579. https://doi.org/10.3389/fpsyt.2019.00579

Hogarth, L., 2020. Addiction is driven by excessive goal-directed drug choice under negative affect: translational critique of habit and compulsion theory. Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol. 45, 720–735. https://doi.org/10.1038/s41386-020-0600-8

Hogarth, L., Field, M., 2020. Relative expected value of drugs versus competing rewards underpins vulnerability to and recovery from addiction. Behav. Brain Res. 112815. https://doi.org/https://doi.org/10.1016/j.bbr.2020.112815

Hood, L.E., Leyrer-Jackson, J.M., Olive, M.F., 2020. Pharmacotherapeutic management of co-morbid alcohol and opioid use. Expert Opin. Pharmacother. 21, 823–839. https://doi.org/10.1080/14656566.2020.1732349

Jan, R.K., Lin, J.C., McLaren, D.G., Kirk, I.J., Kydd, R.R., Russell, B.R., 2014. The effects of methylphenidate on cognitive control in active methamphetamine dependence using functional magnetic resonance imaging. Front. Psychiatry 5, 20.

Jiang, Y., Oathes, D., Hush, J., Darnall, B., Charvat, M., Mackey, S., Etkin, A., 2016. Perturbed connectivity of the amygdala and its subregions with the central executive and default mode networks in chronic pain. Pain 157, 1970–1978. https://doi.org/10.1097/j.pain.0000000000000606

Jongenelis, M., Pettigrew, S., Lawrence, D., Rikkers, W., 2019. Factors Associated with Poly Drug Use in Adolescents. Prev. Sci. 20, 695–704. https://doi.org/10.1007/s11121-019-00993-8

Kandel, D., Kandel, E., 2015. The Gateway Hypothesis of substance abuse: developmental, biological and societal perspectives. Acta Paediatr. 104, 130–137. https://doi.org/10.1111/apa.12851

Kishore, K., 2014. Monograph of tobacco (Nicotiana tabacum). Indian J. Drugs 2, 5–23.

Klega, A.E., Keehbauch, J.T., 2018. Stimulant and Designer Drug Use: Primary Care Management. Am. Fam. Physician 98, 85–92.

Knackstedt, L.A., Ettenberg, A., 2005. Ethanol consumption reduces the adverse consequences of self-administered intravenous cocaine in rats. Psychopharmacology (Berl). 178, 143–150. https://doi.org/10.1007/s00213-004-1996-2

Koob, G.F., Le Moal, M., 2008. Addiction and the brain antireward system. Annu. Rev. Psychol. 59, 29–53. https://doi.org/10.1146/annurev.psych.59.103006.093548

Kringelbach, M.L., 2005. The human orbitofrontal cortex: linking reward to hedonic experience. Nat. Rev. Neurosci. 6, 691–702. https://doi.org/10.1038/nrn1747

Lacy, R.T., Strickland, J.C., Brophy, M.K., Witte, M.A., Smith, M.A., 2014. Exercise decreases speedball self-administration. Life Sci. 114, 86–92. https://doi.org/10.1016/j.lfs.2014.08.005

Laursen, H.R., Henningsson, S., Macoveanu, J., Jernigan, T.L., Siebner, H.R., Holst, K.K., Skimminge, A., Knudsen, G.M., Ramsoy, T.Z., Erritzoe, D., 2016. Serotonergic neurotransmission in emotional processing: New evidence from long-term recreational poly-drug ecstasy use. J. Psychopharmacol. 30, 1296–1304. https://doi.org/10.1177/0269881116662633

Leknes, S., Tracey, I., 2008. A common neurobiology for pain and pleasure. Nat. Rev. Neurosci. 9, 314–320. https://doi.org/10.1038/nrn2333

Leurquin-Sterk, G., Ceccarini, J., Crunelle, C.L., Weerasekera, A., de Laat, B., Himmelreich, U., Bormans, G., Van Laere, K., 2018. Cerebral dopaminergic and glutamatergic transmission relate to different subjective responses of acute alcohol intake: an in vivo multimodal imaging study. Addict. Biol. 23, 931–944. https://doi.org/10.1111/adb.12542

Loganathan, K., Ho, E.T.W., 2021. Value, drug addiction and the brain. Addict. Behav. 116, 106816. https://doi.org/10.1016/j.addbeh.2021.106816

Loganathan, K., Lv, J., Cropley, V., Ho, E.T.W., Zalesky, A., 2020. Associations between delay discounting and connectivity of the valuation-control system in healthy young adults. Neuroscience. https://doi.org/https://doi.org/10.1016/j.neuroscience.2020.11.026

McKetin, R., Chalmers, J., Sunderland, M., Bright, D.A., 2014. Recreational drug use and binge drinking: stimulant but not cannabis intoxication is associated with excessive alcohol consumption. Drug Alcohol Rev. 33, 436–445. https://doi.org/10.1111/dar.12147

Mello, N.K., Mendelson, J.H., Sellers, M.L., Kuehnle, J.C., 1980. Effects of heroin self-administration on cigarette smoking. Psychopharmacology (Berl). 67, 45–52. https://doi.org/10.1007/BF00427594

Moss, H.B., Chen, C.M., Yi, H., 2014. Early adolescent patterns of alcohol, cigarettes, and marijuana polysubstance use and young adult substance use outcomes in a nationally representative sample. Drug Alcohol Depend. 136, 51–62. https://doi.org/https://doi.org/10.1016/j.drugalcdep.2013.12.011

Müller-Oehring, E.M., Jung, Y.-C., Sullivan, E. V, Hawkes, W.C., Pfefferbaum, A., Schulte, T., 2013. Midbrain-Driven Emotion and Reward Processing in Alcoholism. Neuropsychopharmacology 38, 1844–1853. https://doi.org/10.1038/npp.2013.102

Napoli, J.L., Camalier, C.R., Brown, A.-L., Jacobs, J., Mishkin, M.M., Averbeck, B.B., 2021. Correlates of Auditory Decision-Making in Prefrontal, Auditory, and Basal Lateral Amygdala Cortical Areas. J. Neurosci. 41, 1301–1316. https://doi.org/10.1523/JNEUROSCI.2217-20.2020

Nuechterlein, E.B., Ni, L., Domino, E.F., Zubieta, J.-K., 2016. Nicotine-specific and non-specific effects of cigarette smoking on endogenous opioid mechanisms. Prog. Neuropsychopharmacol. Biol. Psychiatry 69, 69–77. https://doi.org/10.1016/j.pnpbp.2016.04.006

Oken, B.S., Salinsky, M.C., Elsas, S.M., 2006. Vigilance, alertness, or sustained attention: physiological basis and measurement. Clin. Neurophysiol. 117, 1885–1901. https://doi.org/10.1016/j.clinph.2006.01.017

Peele, S., 2016. People Control Their Addictions: No matter how much the "chronic" brain disease model of addiction indicates otherwise, we know that people can quit addictions - with special reference to harm reduction and mindfulness. Addict. Behav. reports 4,

97–101. https://doi.org/10.1016/j.abrep.2016.05.003

Plakke, B., Ng, C.-W., Poremba, A., 2013. Neural correlates of auditory recognition memory in primate lateral prefrontal cortex. Neuroscience 244, 62–76. https://doi.org/10.1016/j.neuroscience.2013.04.002

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. Neuroimage 59, 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Ronen, A., Chassidim, H.S., Gershon, P., Parmet, Y., Rabinovich, A., Bar-Hamburger, R., Cassuto, Y., Shinar, D., 2010. The effect of alcohol, THC and their combination on perceived effects, willingness to drive and performance of driving and non-driving tasks. Accid. Anal. Prev. 42, 1855–1865. https://doi.org/10.1016/j.aap.2010.05.006

Saad, Z.S., Gotts, S.J., Murphy, K., Chen, G., Jo, H.J., Martin, A., Cox, R.W., 2012. Trouble at Rest: How Correlation Patterns and Group Differences Become Distorted After Global Signal Regression. Brain Connect. 2, 25–32. https://doi.org/10.1089/brain.2012.0080

Sarter, M., Givens, B., Bruno, J.P., 2001. The cognitive neuroscience of sustained attention: where top-down meets bottom-up. Brain Res. Brain Res. Rev. 35, 146–160. https://doi.org/10.1016/s0165-0173(01)00044-3

Scott, D.J., Domino, E.F., Heitzeg, M.M., Koeppe, R.A., Ni, L., Guthrie, S., Zubieta, J.-K., 2007. Smoking modulation of mu-opioid and dopamine D2 receptor-mediated neurotransmission in humans. Neuropsychopharmacology 32, 450–457. https://doi.org/10.1038/sj.npp.1301238

Shurman, J., Koob, G.F., Gutstein, H.B., 2010. Opioids, pain, the brain, and hyperkatifeia: a

framework for the rational use of opioids for pain. Pain Med. 11, 1092–1098. https://doi.org/10.1111/j.1526-4637.2010.00881.x

Smith, S.M., Beckmann, C.F., Andersson, J., Auerbach, E.J., Bijsterbosch, J., Douaud, G., Duff, E., Feinberg, D.A., Griffanti, L., Harms, M.P., Kelly, M., Laumann, T., Miller, K.L., Moeller, S., Petersen, S., Power, J., Salimi-Khorshidi, G., Snyder, A.Z., Vu, A.T., Woolrich, M.W., Xu, J., Yacoub, E., Ugurbil, K., Van Essen, D.C., Glasser, M.F., 2013a. Resting-state fMRI in the Human Connectome Project. Neuroimage 80, 144–168. https://doi.org/10.1016/j.neuroimage.2013.05.039

Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg, H., Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niazy, R.K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances in functional and structural MR image analysis and implementation as FSL. Neuroimage 23 Suppl 1, S208-19. https://doi.org/10.1016/j.neuroimage.2004.07.051

Smith, S.M., Vidaurre, D., Beckmann, C.F., Glasser, M.F., Jenkinson, M., Miller, K.L., Nichols, T.E., Robinson, E.C., Salimi-Khorshidi, G., Woolrich, M.W., Barch, D.M., Ugurbil, K., Van Essen, D.C., 2013b. Functional connectomics from resting-state fMRI. Trends Cogn. Sci. 17, 666–682. https://doi.org/10.1016/j.tics.2013.09.016

Solinas, M., Belujon, P., Fernagut, P.O., Jaber, M., Thiriet, N., 2019. Dopamine and addiction: what have we learned from 40 years of research. J. Neural Transm. 126, 481–516. https://doi.org/10.1007/s00702-018-1957-2

Stark, M.J., Campbell, B.K., 1993. Cigarette smoking and methadone dose levels. Am. J. Drug Alcohol Abuse 19, 209–217. https://doi.org/10.3109/00952999309002681

Su, S., Fairley, C.K., Mao, L., Medland, N.A., Jing, J., Cheng, F., Zhang, L., 2019. Estimates of the national trend of drugs use during 2000-2030 in China: A population-based

mathematical model. Addict. Behav. 93, 65–71.

https://doi.org/10.1016/j.addbeh.2019.01.022

van den Bos, W., McClure, S.M., 2013. Towards a general model of temporal discounting. J.

Exp. Anal. Behav. 99, 58–73. https://doi.org/10.1002/jeab.6

Walker, R.B., 1980. Medical aspects of tobacco smoking and the anti-tobacco movement in

Britain in the nineteenth century. Med. Hist. 24, 391–402.

Wang, L., Min, J.E., Krebs, E., Evans, E., Huang, D., Liu, L., Hser, Y.-I., Nosyk, B., 2017.

Polydrug use and its association with drug treatment outcomes among primary heroin,

methamphetamine, and cocaine users. Int. J. Drug Policy 49, 32–40.

Wang, T., Ma, J., Wang, R., Liu, Z., Shi, J., Lu, L., Bao, Y., 2018. Poly-Drug Use of

Prescription Medicine among People with Opioid Use Disorder in China: A Systematic

Review and Meta-Analysis. Subst. Use Misuse 53, 1117–1127.

https://doi.org/10.1080/10826084.2017.1400066

Weigl, M., Anzenberger, J., Grabenhofer-Eggerth, A., Horvath, I., Schmutterer, I., Strizek, J.,

Tanios, A., 2017. Bericht zur Drogensituation 2017. Gesundheit Österreich.

Wilson, T.D., Barry, K.L., Maust, D.T., Blow, F.C., 2020. Association between relationship

quality and concurrent alcohol use and sedative-tranquilizer misuse in middle and later

life. Aging Ment. Health 1–5. https://doi.org/10.1080/13607863.2020.1727850

WU-Minn, H.C.P., 2017. 1200 subjects data release reference manual. URL https//www.

humanconnectome. org.

Xie, C., Shao, Y., Ma, L., Zhai, T., Ye, E., Fu, L., Bi, G., Chen, G., Cohen, A., Li, W., Chen,

G., Yang, Z., Li, S.-J., 2014. Imbalanced functional link between valuation networks in

abstinent heroin-dependent subjects. Mol. Psychiatry.

https://doi.org/10.1038/mp.2012.169

Zhai, T., Shao, Y., Chen, Gang, Ye, E., Ma, L., Wang, L., Lei, Y., Chen, Guangyu, Li, W.,
Zou, F., Jin, X., Li, S.-J., Yang, Z., 2015. Nature of functional links in valuation
networks differentiates impulsive behaviors between abstinent heroin-dependent
subjects and nondrug-using subjects. Neuroimage 115, 76–84.
https://doi.org/10.1016/j.neuroimage.2015.04.060

Zou, F., Wu, X., Zhai, T., Lei, Y., Shao, Y., Jin, X., Tan, S., Wu, B., Wang, L., Yang, Z.,
2015. Abnormal resting-state functional connectivity of the nucleus accumbens in multi-
year abstinent heroin addicts. J. Neurosci. Res. 93, 1693–1702.
https://doi.org/10.1002/jnr.23608

Supplementary Methods

The following table shows the list of regions of interest (ROIs) and their center of gravity (in mm) used in the construction of the valuation, executive control and valuation-control complex masks.

Supplementary Table 1: List of ROIs and their corresponding center of gravity (in mm) used to assemble the Valuation and Control system brain masks. LH, left hemisphere; RH, right hemisphere; OFC, orbitofrontal cortex; PC, posterior cingulate; FG, frontal gyrus; MC, middle cingulate; lat, lateral; medial, med; vent, ventral; dors, dorsal; sup, superior; inf, inferior; ros, rostral; caud, caudal; mid, middle; ant, anterior; pos, posterior; opercularis, pars opercularis; triangularis, pars triangularis; orbitalis, pars orbitalis

System	Functional region	ROI	x	y	z
Valuation	vmPFC	LH lat OFC	-23.83	30.67	-17.77
		RH lat OFC	23.00	31.98	-18.46
		LH med OFC	-6.80	33.49	-16.02
		RH med OFC	6.32	37.38	-16.73
	PCC	LH dors PC	-4.07	-39.89	29.19
		RH dors PC	5.33	-36.50	32.92
		LH vent PC	-7.19	-47.48	8.64
		RH vent PC	8.55	-49.22	7.49
	Amygdala	LH amygdala	-23.18	-4.93	-19.56
		RH amygdala	24.20	-3.75	-19.71
	Accumbens	LH accumbens	-8.79	11.65	-6.46
		RH accumbens	9.53	12.09	-7.23

Control	PPC	LH inf parietal	-41.27	-68.97	34.93
		RH inf parietal	47.51	-60.80	33.32
		LH sup parietal	-20.92	-66.90	48.70
		RH sup parietal	24.23	-61.60	52.12
	dlPFC	LH sup FG	-9.79	27.31	43.89
		RH sup FG	12.23	30.84	43.45
		LH ros mid FG	-31.16	48.05	17.58
		RH ros mid FG	34.05	46.79	17.17
		LH caud mid FG	-35.85	11.95	47.34
		RH cau mid FG	38.06	11.21	48.12
	vlPFC	LH opercularis	-47.86	18.01	13.92
		RH opercularis	48.58	15.55	11.08
		LH orbitalis	-42.11	38.11	-13.59
		RH orbitalis	44.08	39.89	-12.09
		LH triangularis	-46.47	35.21	1.73
		RH triangularis	49.08	33.38	6.56
	dACC	LH ant MC	-7.19	12.19	34.99
		RH ant MC	7.96	15.22	36.09
		LH pos MC	-7.50	-14.65	40.42
		RH pos MC	7.98	-8.77	42.24

Supplementary Results

Supplementary Table 2: Descriptive statistics of the variables featured in this study.

Variable	Type	Mean	Standard Deviation	Range
Age	Continuous	28.73	3.71	15
Cognition scores	Continuous	114.08	20.25	93.32
Total Drinks 7 Days	Continuous	4.89	7.08	55
Number of Days Drank 7 Days	Continuous	1.68	1.78	7
Total Tobacco 7 Days	Continuous	7.4	23.06	175
Number of Days Any Tobacco 7 Days	Continuous	0.98	2.28	7
SSAGA Alcohol 12 Drinks/Day	Ordinal	2.28	1.57	
SSAGA Alcohol 12 Drink Frequency	Ordinal	4.29	1.54	
Cocaine	Ordinal	0.22	1.16	
Hallucinogens	Ordinal	0.27	0.9	
Opiates	Ordinal	0.23	0.92	
Sedatives	Ordinal	0.17	0.82	
Stimulants	Ordinal	0.21	0.88	
MJ	Ordinal	1.41	1.69	
Father's History of Drug Use	Binary	0.13	0.34	
Mother's History of Drug Use	Binary	0.03	0.18	

A multicollinearity test was performed among all substance use measures (Supplementary Table 3). Results do not indicate the presence of multicollinearity among these variables. Among the substance use measures, no condition index scored more than 15, while no variance inflation factor (VIF) readings are above 10.

Supplementary Table 3: Multicollinearity test between substance use measures

	s-value	Condition Index	VIF
Total Drinks 7 Days	2.272	1.000	2.948
Drink Frequency 7 Days	1.432	1.587	2.629
Drinks/Day 12 Months	1.161	1.958	2.704
Drink Frequency 12 Months	0.958	2.372	2.297
Total Tobacco 7 Days	0.731	3.108	2.819
Tobacco Frequency 7 Days	0.701	3.243	2.915
Cocaine	0.646	3.516	2.025
Hallucinogens	0.552	4.117	2.155
Opiates	0.502	4.525	2.491
Sedatives	0.477	4.764	2.532
Stimulants	0.412	5.518	1.781
Marijuana	0.353	6.434	1.785

The following figures show average VS, ECS and VCC connectivity in the brains of Human Connectome Project Healthy Young Adult participants used in this study. Connectivity within the VS was higher between the regions of the vmPFC and PCC (Supplementary Figure 1A). Connectivity within the ECS was higher between regions of the PPC, DLPFC and VLPFC (Supplementary Figure 1B). In the VCC, connectivity between the ECS and VS regions was higher between the PPC-DLPFC and the vmPFC-PCC as well as between the lateral OFC and VLPFC (Supplementary Figure 2).

Supplementary Figure 1: Valuation system component regions (upper) and resting-state functional connectivity matrix (lower) in the human brain (A). Control System component regions (upper) and resting-state functional connectivity matrix (lower) in the human brain (B).

Supplementary Figure 2: Valuation system (yellow-orange) and control system (green) component regions in the human brain. The cortical regions (vmPFC, PCC) of the valuation system are labelled yellow, while the subcortical regions (NA, amygdala) are labelled orange). The valuation-control complex is a combination of the valuation and control systems (A). Resting-state connectivity matrix of the valuation-control complex. Connections within the valuation system are shown in yellow, the control system in green. PPC-DLFPC connections with vmPFC-PCC are highlighted in red, VLPFC-lateral OFC connections are highlighted in orange (B).

Given that head motion both affects functional connectivity measures (Ciric et al., 2018) and is affected by substance use (Givens, 2016), average framewise displacement values were correlated with both average functional connectivity of VS, ECS and VCC (Supplementary Table 4) as well as substance use measures (Supplementary Table 5). Strong correlations were not observed between average framewise displacement with either average connectivity or substance use measures with only tobacco measures (Total Tobacco 7 Days and Tobacco Frequency 7 Days) registering r values higher than 0.1.

Supplementary Table 4: Correlations between average framewise displacement for each subject and average functional connectivity of the VS, ECS and VCC

Network	Correlation (r)	p-value
Valuation System	-0.010	0.3765
Executive Control System	0.090	0.0023
Valuation-Control Complex	0.080	0.0059

Supplementary Table 5: Correlations between average framewise displacement for each subject and substance use measures.

Substance Use Measure	Correlation (r)	p-value
Total Drinks 7 Days	-0.019	0.275
Drink Frequency 7 Days	-0.019	0.275
Drinks/Day 12 Months	0.090	0.0023
Drink Frequency 12 Months	0.053	0.0476
Total Tobacco 7 Days	0.187	<0.0001
Tobacco Frequency 7 Days	0.150	<0.0001
Cocaine	0.040	0.1041
Hallucinogens	0.029	0.1808
Opiates	0.053	0.0476
Sedatives	0.062	0.0255
Stimulants	-0.0039	0.4512
Marijuana	0.0798	0.006

Supplementary Table 6: GLM equations and their corresponding r²- and p-values (Valuation System). To enable comparison of effect sizes across relationships, all predictor and outcome variables were standardized to Z-scores with a mean of 0 and an SD of 1. Regression coefficients are therefore representative of SD change in outcome per unit SD change in predictor and may be interpreted as partial correlation coefficients. "7" (e.g. Total Tobacco 7, Drink Frequency 7) indicates 7 days; "12" (e.g. Drinks/Day 12) indicates 12 months.

Equation	r^2	p-value
ocaine ~ 0.19443 + 0.068862(VS – Total Tobacco 7) + 0.075595(VS – Tobacco Frequency 7) + 0.11683(VS - Hallucinogens) - 0.19901(VS - Opiates) – 0.05976(VS - Sedatives) - 0.07448(VS - Father) 0.059324(Age – Total Drinks 7) + 0.16187(Age – Tobacco Frequency 7) + 0.14899(Age – Hallucinogens) + 0.06162(Total Drinks 7 – Drinks/Day 12) + 0.10987(Total Drinks 7 – Tobacco Frequency) + 0.11095(Total Drinks 7 – Hallucinogen) – 0.31816(Total Drinks 7 – Hallucinogens) – 0.31816(Total Drinks 7 – Stimulants) + 0.092877(Total Drinks 7 – Marijuana) - 0.1899(Drink Frequency 7 – obacco Frequency 7) – 0.1584(Drink Frequency 7 – Hallucinogens) + 0.36753(Drink Frequency 7 - Sedatives) + 0.16412(Drinks/Day 12 – Total Tobacco 7) + 0.035364(Drink Frequency 7 – Total obacco 7) + 0.093752(Drink Frequency 7 – Tobacco Frequency 7) - 0.19016 (Drink Frequency 7 – Sedatives) + 0.10236(Total Tobacco 7 – Tobacco Frequency 7) – 0.078794(Total Tobacco 7 – allucinogens) + 0.090749(Total Tobacco 7 – Sedatives) – 0.51708(Total Tobacco 7 – Mother) + 0.083325(Tobacco Frequency 7 – Sedatives) + 0.21259(Tobacco Frequency 7 – Father) - .6422(Tobacco Frequency 7 – Mother) - 0.030834(Hallucinogens – Opiates) + 0.71706(Hallucinogens – Sedatives) – 0.32437(Hallucinogens – Father) - 2.2652(Hallucinogens – Mother) + .28473(Hallucinogens – Gender) – 0.10987(Opiates – Sedatives) - 0.1689(Opiates – Marijuana) + 1.2938(Opiates – Mother) + 0.24486(Opiates – Gender) – 0.054253(Sedatives – Marijuana) – .59964(Sedatives – Gender) + 0.11409(Stimulants – Mother)	0.7227	2.96e-230
allucinogens ~ -1.6301 – 0.06369(VS – Total Tobacco 7) + 0.13897(VS – Tobacco Frequency 7) - 0.22626 (VS – Cocaine) + 0.21878(VS – Opiates) + 0.0062508(VS – Stimulants) + 0.05203(Age – rinks/Day 12) + 0.20312(Age – Cocaine) – 0.12568(Age – Opiates) + 0.10987(Total Drinks 7 – Cocaine) – 0.065655(Total Drinks 7 – Marijuana) – 0.1457(Drink Frequency 7 – Cocaine) + .0087609(Drink Frequency 7 – Sedatives) + 0.141(Drinks/Day 12 – Opiates) – 0.094957(Drinks/Day 12 – Sedatives) + 0.10969(Drink Frequency 12 – Tobacco Frequency 7) + 0.070751(Drink requency 12 – Opiates) – 0.012169(Drink Frequency 12 – Stimulants) – 0.091478(Total Tobacco 7 – Opiates) – 0.066949(Total Tobacco 7 – Sedatives) + 0.062218(Tobacco Frequency 7 – Marijuana) – .081708(Cocaine – Sedatives) + 0.11838(Cocaine – Marijuana) + 0.4122(Cocaine – Father) + 0.42785(Cocaine – Gender) – 0.15891(Opiates – Father) + 4.9378(Opiates – Mother) + 0.013022(Sedatives Stimulants) – 3.465(Sedatives – Mother) – 0.26538(Sedatives – Gender) + 0.014303(Stimulants – Gender)	0.6707	1.98e-208
piates ~ -0.077541 – 0.047223 (VS – Drink Frequency 7) + 0.07354(VS – Total Tobacco 7) + 0.11462(VS – Tobacco Frequency 7) – 0.036665(VS – Father) + 0.065478(Age – Hallucinogens) – .17539(Age – Sedatives) + 0.042794(Total Drinks 7 – Cocaine) + 0.19854(Total Drinks – Cocaine) – 0.19573(Drink Frequency 7 – Cocaine) + 0.065957(Drink Frequency 7 – Hallucinogens) – .061074(Drinks/Day 12 – Total Drinks 7) – 0.063834(Drinks/Day 12 – Cocaine) + 0.083089(Drinks/Day 12 – Hallucinogens) – 0.14065(Drinks/Day 12 – Hallucinogens) – 0.071936(Drink Frequency 2 – Cocaine) – 0.12058(Drink Frequency 12 – Hallucinogens) + 0.16286(Drink Frequency 12 – Hallucinogens) – 0.047018(Total Tobacco 7 – Tobacco Frequency 7) + 0.14016(Total Tobacco 7 – ocaine) – 0.082762(Tobacco Frequency 12 – Hallucinogens) + 0.04842(Tobacco Frequency 12 – Sedatives) – 2.2096(Cocaine – Mother) – 0.19427(Cocaine – Gender) + 0.0052223(Hallucinogens – timulants) – 0.13244(Hallucinogens – Marijuana) + 3.216(Sedatives – Mother) + 0.31369(Stimulants – Gender)	0.7056	1.65e-228
edatives ~ 0.022372 + 0.044131(VS – Drinks/Day 12) + 0.05633(VS – Tobacco Frequency 7) + 0.087239(VS – Cocaine) – 0.11234(VS – Hallucinogens) + 0.1094(VS – Opiates) + 0.035095(Age – rink Frequency 7) – 0.13456(Age – Cocaine) + 0.18204 (Age – Opiates) – 0.038749(Total Drinks 7 – Drink Frequency 7) – 0.29372(Total Drinks 7 – Hallucinogens) + 0.37067(Total Drinks 7 – piates) – 0.95341(Total Drinks 7 – Mother) – 0.062641(Drink Frequency 7 – Tobacco Frequency 7) + 0.089163(Tobacco Frequency 7 – Hallucinogens) – 0.40739(Drink Frequency 7 – Opiate) + .034425(Drinks/Day 12 – Drink Frequency 12) + 0.040748(Drinks/Day 12 – Total Tobacco 7) – 0.1544(Drinks/Day 12 – Tobacco Frequency 7) – 0.095024(Drinks/Day 12 – Cocaine) + .41269(Drinks/Day 12 – Hallucinogens) – 0.21047(Drinks/Day 12 – Sedatives) – 0.16845(Drink Frequency 12 – Tobacco Frequency 7) + 0.1052(Total Tobacco 7 – Tobacco Frequency 7) – .078702(Total Tobacco 7 – Sedatives) – 0.030747(Total Tobacco 7 – Father) + 0.050822(Tobacco Frequency 7 – Cocaine) – 0.12095(Tobacco Frequency 7 – Hallucinogens) + 0.058513(Tobacco requency 7 – Opiates) + 5.4137(Tobacco Frequency 7 – Mother) – 0.24899(Tobacco Frequency 7 – Gender) + 0.10356(Cocaine – Hallucinogens) – 0.097371(Cocaine – Opiates) – 0.012427(Cocaine – timulants) + 0.1893(Cocaine – Gender) + 0.031576(Hallucinogens – Opiates) + 0.01372(Hallucinogens – Stimulants) + 0.26187(Hallucinogens – Marijuana) + 0.25412(Hallucinogens – Gender) – .0076042(Opiates – Stimulants) – 0.12105(Opiates – Marijuana) – 3.9793(Opiates – Mother) – 0.58812(Marijuana – Mother)	0.7797	1.24e-293
timulants ~ 0.088602 – 0.050192(VS – Drink Frequency 12) – 0.0724(VS – Opiates) – 0.060044(VS – Father) + 0.13787(Age – Opiates) + 0.078607(Total Drinks 7 – Drinks/Day 12) – 0.22526(Total rinks 7 – Drink Frequency 12) + 0.3232(Total Drinks 7 – Opiates) + 0.0088373(Total Drinks 7 – Stimulants) – 0.8904(Total Drinks 7 – Mother) + 0.15143(Drink Frequency 7 – Tobacco Frequency 7) – .36282(Drink Frequency 7 – Opiates) + 0.29091(Drinks/Day 12 – Drink Frequency 7) – 0.14723(Drinks/Day 12 – Opiates) + 0.22343(Drink Frequency 12 – Hallucinogens) – 0.066402(Total Tobacco 7 Cocaine) – 0.099145(Total Tobacco 7 – Opiates) + 0.038182(Tobacco Frequency 7 – Cocaine) – 0.065418(Tobacco Frequency 7 – Opiates) – 0.15984(Tobacco Frequency 7 – Marijuana) – .6715(Tobacco Frequency – Mother) – 0.3597(Tobacco Frequency – Gender) – 0.032032(Cocaine – Hallucinogens) + 0.081362(Hallucinogens – Father) + 2.3972(Hallucinogens – Mother) + .23292(Opiates – Gender) + 0.0064412(Stimulants – Father) – 0.81659(Father – Mother) + 0.15155(Father – Gender)	0.6265	6.35e-180
Marijuana ~ 8.3068 – 0.50742(VS – Tobacco Frequency 7) + 0.70711(VS – Stimulants) – 0.5338(Age – Total Tobacco 7) – 0.50275(Total Drinks 7 – Cocaine) – 1.0985(Drink Frequency 7 – Drinks/Day 2) – 1.0304(Drinks/Day 12 – Drink Frequency 12) + 0.38691(Drink Frequency 12 – Cocaine) + 0.82055(Drink Frequency 12 – Gender) – 0.49909(Total Tobacco 7 – Stimulants) – 0.80167(Opiates – ather) – 1.0505(Sedatives – Gender)	0.4715	4.31e-120

Supplementary Table 7: GLM equations and their corresponding r²- and p-values (Executive Control System). To enable comparison of effect sizes across relationships, all predictor and outcome variables were standardized to Z☐scores with a mean of 0 and an SD of 1. Regression coefficients are therefore representative of SD change in outcome per unit SD change in predictor and may be interpreted as partial correlation coefficients. "7" (e.g. Total Tobacco 7, Drink Frequency 7) indicates 7 days; "12" (e.g. Drinks/Day 12) indicates 12 months.

Equation	r^2	p-value
ocaine ~ 0.20424 + 0.061485(ECS – Total Tobacco 7) + 0.17194(ECS – Hallucinogens) – 0.2171(ECS – Opiates) – 0.059511(ECS – Sedatives) – 0.059615(ECS – Father) + 0.059518(Age – Total Drinks 7) + 0.15265(Age – Drink Frequency 12) + 0.18801 (Age – Hallucinogens) + 0.069751(Total Drinks 7 – Drinks/Day 12) + 0.13252(Total Drinks 7 – Drink Frequency 12) + 0.084688(Total Drinks 7 – Hallucinogens) – 0.31281 (Total Drinks 7 – Sedatives) + 0.0064776(Total Drinks – Stimulants) + 0.086285(Total Drinks 7 – Marijuana) – 0.18884(Drink Frequency 7 – Total Tobacco 7) – .13914(Total Drinks 7 – Hallucinogens) + 0.36538(Total Drinks 7 – Sedatives) – 0.044428(Drinks/Day 12 – Total Tobacco 7) + 0.1906(Drinks/Day 12 – Hallucinogens) + 0.13746(Drinks/Day 12 – edatives) + 0.04789(Drink Frequency 12 – Total Tobacco 7) + 0.14771(Drink Frequency 12 – Tobacco Frequency 7) – 0.19084(Drink Frequency 12 – Sedatives) – 0.0051556(Drink Frequency 12 – timulants) + 0.078983(Total Tobacco 7 – Tobacco Frequency 7) – 0.054534(Total Tobacco 7 – Hallucinogens) + 0.085068(Total Tobacco 7 – Sedatives) – 0.57584(Total Tobacco 7 – Mother) + .076395(Tobacco Frequency 7 – Sedatives) + 0.19187(Tobacco Frequency 7 – Father) – 3.7973(Tobacco Frequency 7 – Mother) – 0.025156(Hallucinogens – Opiates) + 0.07289(Hallucinogens – edatives) – 0.31752(Hallucinogens – Father) – 2.9559(Hallucinogens – Mother) + 0.16725(Hallucinogens – Gender) – 0.11995(Sedatives – Opiates) – 0.15598(Sedatives – Marijuana) + .2751(Sedatives – Mother) + 0.3902(Sedatives – Gender) – 0.057548(Opiates – Marijuana) – 0.59073(Sedatives – Gender) + 0.12277(Stimulants – Mother) + 0.008566(Stimulants – Gender)	0.7208	2.8e-228
allucinogens ~ 0.20424 + 0.036806(ECS – Drink Frequency 7) + 0.050288(ECS – Tobacco Frequency 7) – 0.039922 (ECS – Marijuana) – 0.096729(ECS – Gender) + 0.13239(Age – Cocaine) - .088736(Age – Sedatives) + 0.006427(Age – Stimulants) + 0.11837(Total Drinks 7 – Cocaine) – 0.056919(Total Drinks 7 – Cocaine) + 0.053899(Drink Frequency 7 – Drinks/Day 12) – 0.12396(Drink Frequency 12 – Cocaine) + 0.064681(Drink Frequency 12 – Sedatives) – 0.053216(Drink Frequency 7 – Marijuana) + 0.083314(Drinks/Day 12 – Cocaine) – 0.11902 (Drinks/Day 12 – Opiates) – .059315(Drinks/Day 12 – Sedatives) + 0.14191(Drink Frequency 12 – Tobacco Frequency 7) + 0.07462(Drink Frequency 12 – Opiates) – 0.011127(Drink Frequency 12 – Stimulants) – 0.082258(Drink requency 12 – Gender) – 0.069254(Total Tobacco 7 – Tobacco Frequency 7) – 0.070195(Total Tobacco 7 – Sedatives) + 0.034384(Tobacco Frequency 7 – Cocaine) + 0.064465(Tobacco Frequency 7 – Marijuana) – 2.3105(Tobacco Frequency 7 – Mother) – 0.12545(Cocaine – Sedatives) + 0.0083335(Cocaine – Stimulants) + 0.11842(Cocaine – Marijuana) + 0.22351(Cocaine – Father) - .020697(Opiates – Stimulants) + 0.066316(Opiates – Father) + 5.9728(Opiates – Father) + 0.58778(Opiates – Gender) + 0.025602(Sedatives – Stimulants) – 0.078412(Sedatives – Father) – .9675(Sedative – Mother) – 0.4959(Sedatives – Gender) – 0.073917(Stimulants – Mother) + 0.019456(Stimulants – Gender)	0.7856	3.58e-207
piates ~ 0.047756 + 0.075083(ECS – Total Tobacco 7) + 0.18812(ECS – Hallucinogens) – 0.071123(ECS – Sedatives) + 0.21161 (Age – Hallucinogens) – 0.2254(Age – Sedatives) + 0.037573(Total Drinks 7 – Drink Frequency 7) + 0.03691(Total Drinks 7 – Total Tobacco 7) – 0.033757(Total Drinks 7 – Tobacco Frequency 7) + 0.19832(Total Drinks 7 – Cocaine) + 0.042771(Total Drinks 7 – Marijuana) – 0.1885(Drink Frequency 7 – Cocaine) - 0.047404(Drinks/Day 12 – Total Tobacco 7) – 0.069289(Drinks/Day 12 – Cocaine) + 0.13164(Drinks/Day 12 – Hallucinogens) - .14536(Drinks/Day 12 – Sedatives) – 0.077845(Drink Frequency 12 – Cocaine) – 0.11868(Drink Frequency 12 – Hallucinogens) + 0.21566(Drink Frequency 12 – Sedatives) + 0.1148(Total Tobacco 7 – ocaine) + 0.034124(Tobacco Frequency 7 – Cocaine) – 0.086218(Tobacco Frequency – Hallucinogens) + 0.04198(Tobacco Frequency – Sedatives) – 0.016679(Tobacco Frequency – Stimulants) + .6212(Tobacco Frequency 7 – Mother) + 0.26391(Tobacco Frequency 7 – Gender) – 1.9114(Cocaine – Gender) + 0.017671(Hallucinogens – Stimulants) – 0.13732(Hallucinogens – Marijuana) - .713(Hallucinogens – Mother) - 0.45154(Hallucinogens – Gender) + 4.9326(Sedatives – Mother) + 0.50246(Sedatives – Gender) – 0.0042135(Stimulants – Marijuana)	0.7137	3.65e-230
edatives ~ 0.063295 + 0.037386(ECS – Drinks/Day 12) + 0.052648(ECS – Tobacco Frequency) – 0.07494(ECS – Cocaine) + 0.18558(ECS – Opiate) – 0.18269(Age – Hallucinogens) + 0.20098(Age – edatives) – 0.041226(Total Drinks 7 – Drink Frequency 7) – 0.056293(Total Drinks 7 – Tobacco Frequency 7) – 0.20718(Total Drinks 7 – Cocaine) – 0.11854(Total Drinks – Hallucinogens) + .41456(Total Drinks 7 – Opiates) – 0.56674(Total Drinks – Mother) + 0.15499(Tobacco Frequency 7 – Hallucinogens) – 0.44184(Drink Frequency 7 – Opiates) + 0.040378(Drinks/Day 12 – Drink requency 12) + 0.032768(Drinks/Day 12 – Total Tobacco 7) – 0.1174(Drinks/Day 12 – Tobacco Frequency 7) + 0.40788(Drinks/Day 12 – Cocaine) – 0.064984(Drinks/Day 12 – Hallucinogens) – .23925(Drinks/Day 12 – Sedatives) – 0.16679(Drink Frequency 12 – Tobacco Frequency 7) + 0.079715(Drink Frequency 12 – Cocaine) + 0.090469(Total Tobacco 7 – Tobacco Frequency 7) – 0.08084 Total Tobacco 7 – Opiates) – 0.12458(Tobacco Frequency 7 – Opiates) + 0.065791(Tobacco Frequency 7 + Hallucinogens + 0.065053(Tobacco Frequency 7 – Sedatives) – 0.07299(Tobacco Frequency – Father) + 5.6013(Tobacco Frequency 7 – Mother) – 0.25343(Tobacco Frequency 7 – Gender) + 0.12583(Cocaine – Hallucinogens) + 0.013536(Cocaine – Stimulants) + 0.30761(Cocaine – Marijuana) 0.24058(Cocaine – Gender) – 0.12079(Hallucinogens – Opiates) - 0.017375(Hallucinogens – Stimulants) + 0.10847(Hallucinogens – Father) + 0.25037(Hallucinogens – Gender) – 0.16537(Sedatives – Marijuana) – 5.0012(Opiates – Mother) – 0.16684(Opiates – Gender) – 0.56159(Marijuana – Mother)	0.806	5.55e-302
timulants ~ 0.11131 + 0.078438(ECS – Tobacco Frequency 7) – 0.098128(ECS – Hallucinogens) – 0.25233(ECS – Opiates) – 0.039931(Age – Drinks/Day 12) – 0.24174(Total Drinks 7 – Tobacco requency 7) + 0.26766(Total Drinks 7 – Opiates) + 0.0078777(Total Drinks 7 – Stimulants) – 0.70687(Total Drinks 7 – Mother) + 0.1713(Drink Frequency 7 – Tobacco Frequency 7) – 0.26031(Drink requency 7 – Opiates) + 0.38852(Drinks/Day 12 – Tobacco Frequency 7) + 0.1064(Drinks/Day 12 – Hallucinogens) – 0.2372(Drinks/Day 12 – Opiates) + 0.27017(Drink Frequency 12 – Opiates) – .091(Total Tobacco 7 – Tobacco Frequency 7) – 0.03657(Total Tobacco 7 – Marijuana) + 0.035504(Tobacco Frequency 7 – Hallucinogens) – 0.056698(Tobacco Frequency 7 – Opiates) – .099466(Tobacco Frequency 7 – Marijuana) – 4.6699(Tobacco Frequency 7 – Mother) + 0.49516(Tobacco Frequency – Gender) – 0.028789(Cocaine – Hallucinogens) – 1.4531(Cocaine – Mother) – .042229(Hallucinogens – Opiates) + 2.8089(Hallucinogens – Mother) + 2.1681(Opiates – Mother) + 0.5833(Opiates – Gender) + 0.0088012(Sedatives – Father) – 0.65884 (Father – Mother) + 0.10843 Father – Gender)	0.6392	1.12e-185
Marijuana ~ 8.1708 + 0.7277(ECS – Drink Frequency 12) + 0.45798(ECS - Marijuana) - 0.44289(Age – Total Tobacco 7) - 0.54216(Total Drinks 7 - Cocaine) - 1.0871(Drink Frequency 7 – Drinks/Day 2) - 0.86162(Drinks/Day 12 – Drink Frequency 12) - 0.40667(Total Tobacco 7 - Sedatives) - 0.62233(Hallucinogens - Father)	0.4703	3.18e-123

Supplementary Table 8: GLM equations and their corresponding r^2- and p-values (Valuation-Control Complex). To enable comparison of effect sizes across relationships, all predictor and outcome variables were standardized to Z-scores with a mean of 0 and an SD of 1. Regression coefficients are therefore representative of SD change in outcome per unit SD change in predictor and may be interpreted as partial correlation coefficients. "7" (e.g. Total Tobacco 7, Drink Frequency 7) indicates 7 days; "12" (e.g. Drinks/Day 12) indicates 12 months.

Equation	r^2	p-value
Cocaine ~ 0.21801 + 0.070669(VCC – Total Tobacco 7) + 0.18494(VCC – Hallucinogens) – 0.19881(VCC – Opiates) -0.048278(VCC – Sedatives) – 0.0050319(VCC – Stimulants) – 0.056556(VCC – Father) + 0.057609(Age – Total Drinks 7) + 0.14546(Age – Tobacco Frequency 7) + 0.1923(Age – Hallucinogens) + 0.074442(Total Drinks 7 – Drinks/Day 12) + 0.1322(Total Drinks 7 – Tobacco Frequency 7) + 0.091828(Total Drinks – Hallucinogens) – 0.31607(Total Drinks 7 – Sedatives) + 0.0084922(Total Drinks 7 – Stimulants) + 0.09164(Total Drinks 7 – Marijuana) – 0.19438(Drink Frequency 7 – Tobacco Frequency 7) – 0.15694(Tobacco Frequency 7 + Hallucinogens) + 0.38012(Tobacco Frequency 7 – Sedatives) – 0.054669(Drinks/Day 12 – Total Tobacco 7) + .1659(Drinks/Day 12 – Hallucinogens) + 0.12783(Drinks/Day 12 – Sedatives) + 0.14028(Drink Frequency 12 – Tobacco Frequency 7) – 0.19896(Drink Frequency 12 – Sedatives) + 0.090095(Total Tobacco 7 – Tobacco Frequency 7) – 0.084401(Total Tobacco 7 – Hallucinogens) + 0.088186(Total Tobacco 7 – Sedatives) + 0.066431(Tobacco Frequency 7 – Sedatives) + 0.20564(Tobacco Frequency – Father) – 3.4857(Tobacco Frequency 7 – Mother) + 0.061953(Hallucinogens – Sedatives) – 0.31718(Hallucinogens – Father) - 3.3182(Hallucinogens – Mother) + 0.21742(Hallucinogens – Gender) - .11906(Opiates – Sedatives) – 0.1508(Opiates – Marijuana) + 2.1964(Opiates – Mother) + 0.27352(Opiates – Mother) + 0.27352(Opiates – Gender) – 0.067314(Opiates – Marijuana) - 0.611(Opiates – Gender) + 0.11112(Stimulants – Mother) + 0.012633(Stimulants – Gender)	0.7198	1.95e-229
Hallucinogens ~ -0.2302 – 0.048987(VCC – Total Tobacco 7) + 0.18501(VCC – Opiates) + 0.054542(Age – Drinks/Day 12) + 0.18734(Age – Cocaine) + 0.047491(Total Drinks 7 – Drink Frequency 7) – 0.13593(Total Drinks 7 + Cocaine) – 0.045186(Total Drinks 7 – Opiates) – 0.058856(Total Drinks 7 – Marijuana) – 0.058856(Total Drinks 7 – Marijuana) – 0.1669(Drink Frequency 7 – Cocaine) + .060532(Drink Frequency 7 – Opiates) – 0.12282(Drinks/Day 12 – Opiates) + 0.1384(Drink Frequency 12 – Tobacco Frequency 7) – 0.010106(Drink Frequency 12 – Stimulants) – 0.056535(Total Tobacco 7 – Opiates) – 0.073854(Total Tobacco 7 – Sedatives) – 0.10552(Cocaine – Sedatives) + 0.11938(Cocaine – Marijuana) + 0.26035(Cocaine – Father) + 0.15462(Cocaine – Gender) + .017296(Sedatives – Stimulants) – 0.044036(Sedatives – Father) + 0.016868(Stimulants – Gender)	0.6612	5.64e-204
Opiates ~ -0.13254 – 0.045838(VCC – Drinks/Day 12) + 0.096573(VCC – Total Tobacco 7) + 0.08931(VCC – Hallucinogens) 0.063258(Age – Cocaine) – 0.21705(Age – Sedatives) + 0.037971(Total Drinks 7 – Drink Frequency 7) + 0.18642(Total Drinks 7 – Cocaine) – 0.17706(Drink Frequency 7 – Cocaine) - 0.0482(Drinks/Day 12 – Total Tobacco 7) -0.11275(Drinks/Day 12 – Opiate) - .056763(Drink Frequency 12 – Cocaine) – 0.18749(Drink Frequency 12 – Hallucinogens) + 0.20665(Drink Frequency 12 – Sedatives) + 0.10117(Total Tobacco 7 – Cocaine) – 0.056695(Tobacco Frequency 7 – Hallucinogens) + 0.047575(Tobacco Frequency 7 – Sedatives) + 0.1865(Tobacco Frequency 7 – Gender) – 2.2217(Cocaine – Mother) – 0.20784(Cocaine – Gender) – .094492(Hallucinogens – Marijuana) – 0.21276(Hallucinogens – Gender) + 3.3288(Sedatives – Mother) + 0.38016(Sedatives – Gender)	0.6996	3.94e-227
Sedatives ~ 0.024389 + 0.039047(VCC – Drinks/Day 12) + 0.082159(VCC – Tobacco Frequency 7) - 0.13514(VCC – Opiates) + 0.18292(VCC – Sedatives) – 0.16846(Age – Cocaine) + 0.18293(Age – Sedatives) – 0.042658(Total Drinks 7 – Drink Frequency 7) – 0.05001(Total Drinks 7 – Tobacco Frequency 7) – 0.1456(Total Drinks 7 – Cocaine) – 0.20886(Total Drinks 7 – Hallucinogens) + .43029(Total Drinks 7 – Opiate) - 0.68629(Total Drinks 7 – Mother) + 0.18042(Drink Frequency 7 – Cocaine) – 0.46683(Drink Frequency 7 – Opiates) + 0.041472(Drinks/Day 12 – Drink Frequency 12) + 0.031309(Drinks/Day 12 – Total Tobacco 7) – 0.10872(Drinks/Day 12 – Tobacco Frequency 7) – 0.062916 (Drinks/Day 12 – Cocaine) + 0.38104(Drinks/Day 12 – Hallucinogens) - 0.24011(Drink Frequency 12 – Tobacco Frequency 7) + 0.07902(Drink Frequency 12 – Hallucinogens) + 0.085114(Total Tobacco 7 – Tobacco Frequency 7) – 0.087971 (Total Tobacco 7 – Opiates) + 0.06194(Tobacco Frequency 7 – Cocaine) – 0.12195(Tobacco Frequency 7 – Hallucinogens) + 0.069098(Tobacco Frequency 7 – Sedatives) – 0.069737(Tobacco Frequency 7 – Father) + 5.4962(Tobacco Frequency 7 – Mother) – 0.2812(Tobacco Frequency – Gender) + 0.11883(Cocaine – Hallucinogens) – 0.11238(Cocaine – Opiates) – 0.014053(Cocaine – Stimulants) + 0.09909(Cocaine – Father) + 0.24428(Cocaine – Gender) + 0.015054(Hallucinogens – Stimulants) + 0.30791(Hallucinogens – Marijuana) + 1.3966(Hallucinogens – Mother) + 0.35179(Hallucinogens – Gender) - 0.0058883(Opiates – Stimulants) - .16606(Opiates – Marijuana) – 5.6098(Opiates – Mother) - 0.11617(Opiates – Gender) – 0.56871(Marijuana – Mother)	0.8083	5.7e-303
Stimulants ~ 0.12215 + 0.062193(VCC – Tobacco Frequency 7) – 0.089502(VCC – Opiates) – 0.16196(VCC – Sedatives) – 0.22405(Total Drinks 7 – Tobacco Frequency 7) + 0.25054(Total Drinks 7 – Opiates) + 0.0077219(Total Drinks 7 – Sedatives) – 1.0664(Total Drinks 7 – Mother) + 0.15457(Drink Frequency 7 – Tobacco Frequency 7) – 0.25852(Tobacco Frequency – Sedatives) + .33811(Drinks/Day 12 – Tobacco Frequency 7) – 0.13356(Drinks/Day 12 – Opiates) + 0.31323(Drink Frequency 12 – Hallucinogens) – 0.074914(Drink Frequency 12 – Hallucinogens) – 0.07229 (Total Tobacco 7 – Tobacco Frequency 7) – 0.044806(Total Tobacco 7 – Cocaine) – 0.03702(Total Tobacco 7 – Marijuana) – 0.048082(Tobacco Frequency 7 – Opiates) + 0.013621(Tobacco Frequency 7 – Marijuana) – 3.9026(Tobacco Frequency 7 – Mother) – 0.55796(Tobacco Frequency 7 – Gender) + 3.2924(Hallucinogens – Mother) – 0.016408(Opiates – Sedatives) + 0.45508(Opiates – Gender) + 0.0076174(Stimulants – Father) – 0.57193(Father – Mother) + 0.11931(Father – Gender)	0.6288	252e-182
Marijuana ~ 8.2017 + 0.69869(VCC – Drink Frequency 12) - 0.78946(VCC - Opiates) + 0.62535(VCC - Marijuana) - 0.62702(Age - Opiates) - 0.55758(Total Drinks 7 - Cocaine) - 1.0602(Drink Frequency 7 – Drinks/Day 12) - 0.84237(Drinks/Day 12 – Drink Frequency 12) - 0.37699(Total Tobacco 7 - Sedatives) - 0.82425(Hallucinogens – Father)	0.4732	4.32e-123

Supplementary Table 9: Beta coefficients for all dependent variables used in the Valuation System (VS) GLM analysis.

Cocaine	
-0.036798	VS average connectivity
-0.0040128	Age
0.053252	Total Drinks 7 Days
0.093316	Drink Frequency 7 Days
-0.029292	Drinks/Day 12 Months
-0.0020316	Drink Frequency 12 Months
0.0016601	Total Tobacco 7 Days
0.015079	Tobacco Frequency 7 Days
0.31723	Hallucinogens
0.051472	Opiates
0.22316	Sedatives
0.20316	Stimulants
0.0035696	Marijuana
-0.049317	Father's Use History
0.050529	Mother's Use History
0.021832	Gender

Hallucinogens	
-0.25552	VS average connectivity
0.039147	Age
0.056704	Total Drinks 7 Days
0.092725	Drink Frequency 7 Days
-0.061582	Drinks/Day 12 Months
-0.019646	Drink Frequency 12 Months
-0.017365	Total Tobacco 7 Days
-0.075815	Tobacco Frequency 7 Days
0.30166	Cocaine
0.20515	Opiates
0.073069	Sedatives
0.048425	Stimulants
0.033036	Marijuana
-0.02577	Father's Use History
0.018055	Mother's Use History
0.097249	Gender

Opiates	
-0.11587	VS average connectivity
0.023706	Age
-0.015377	Total Drinks 7 Days
0.15301	Drink Frequency 7 Days
-0.066566	Drinks/Day 12 Months
-0.081359	Drink Frequency 12 Months
0.012002	Total Tobacco 7 Days
0.032058	Tobacco Frequency 7 Days
0.041498	Cocaine
0.17394	Hallucinogens
0.49963	Sedatives
0.10283	Stimulants
0.0074145	Marijuana
-0.031873	Father's Use History
-0.062263	Mother's Use History
0.012615	Gender

Sedatives	
-0.00090637	VS average connectivity
0.0051333	Age
-0.017298	Total Drinks 7 Days
-0.11711	Drink Frequency 7 Days
0.050438	Drinks/Day 12 Months
0.043198	Drink Frequency 12 Months
0.025806	Total Tobacco 7 Days
0.053711	Tobacco Frequency 7 Days
0.18596	Cocaine
0.064031	Hallucinogens
0.51642	Opiates
0.17293	Stimulants
0.00015496	Marijuana
0.059432	Father's Use History
0.0060936	Mother's Use History
-0.087953	Gender

Stimulants	
0.096479	VS average connectivity
-0.023519	Age
-0.040469	Total Drinks 7 Days
0.037787	Drink Frequency 7 Days
-0.10873	Drinks/Day 12 Months
0.097314	Drink Frequency 12 Months
0.01813	Total Tobacco 7 Days
-0.02136	Tobacco Frequency 7 Days
0.23899	Cocaine
0.059906	Hallucinogens
0.15004	Opiates
0.24412	Sedatives
0.010415	Marijuana
-0.0036764	Father's Use History
0.016632	Mother's Use History
-0.013619	Gender

Marijuana	
7.0978	VS average connectivity
-0.13754	Age
-0.25428	Total Drinks 7 Days
-0.32861	Drink Frequency 7 Days
1.3669	Drinks/Day 12 Months
1.2826	Drink Frequency 12 Months
-0.18135	Total Tobacco 7 Days
1.8076	Tobacco Frequency 7 Days
0.33282	Cocaine
3.2392	Hallucinogens
0.85748	Opiates
0.017338	Sedatives
0.82553	Stimulants
0.50759	Father's Use History
0.2939	Gender

Supplementary Table 10: Beta coefficients for all dependent variables used in the Executive Control System (ECS) GLM analysis.

Cocaine	
-0.054278	ECS average connectivity
-0.030908	Age
0.049862	Total Drinks 7 Days
0.091928	Drink Frequency 7 Days
-0.029366	Drinks/Day 12 Months
-0.0009592	Drink Frequency 12 Months
0.0010022	Total Tobacco 7 Days
0.014475	Tobacco Frequency 7 Days
0.3177	Hallucinogens
0.053248	Opiates
0.22275	Sedatives
0.20171	Stims
0.0034632	Marijuana
-0.049623	Father's Use History
0.051625	Mother's Use History
0.038353	Gender
Hallucinogens	
-0.24993	ECS average connectivity
0.024356	Age
0.053158	Total Drinks 7 Days
0.091153	Drink Frequency 7 Days
-0.058269	Drinks/Day 12 Months
-0.018429	Drink Frequency 12 Months
-0.015766	Total Tobacco 7 Days
-0.077546	Tobacco Frequency 7 Days
0.30301	Cocaine
0.20567	Opiates
0.073608	Sedatives
0.048304	Stimulants
0.033086	Marijuana
-0.026739	Father's Use History
0.017635	Mother's Use History
0.096801	Gender
Opiates	
-0.099627	ECS average connectivity
0.035655	Age
-0.014834	Total Drinks 7 Days
0.15283	Drink Frequency 7 Days
-0.064595	Drinks/Day 12 Months
-0.081385	Drink Frequency 12 Months
0.013329	Total Tobacco 7 Days
0.031595	Tobacco Frequency 7 Days
0.042935	Cocaine
0.17388	Hallucinogens
0.4993	Sedatives
0.10346	Stimulants
0.0074785	Marijuana
-0.063282	Mother's Use History
8.081e-05	Gender
Sedatives	

-0.002682	ECS average connectivity
-0.0011138	Age
-0.018334	Total Drinks 7 Days
-0.11756	Drink Frequency 7 Days
0.050922	Drinks/Day 12 Months
0.043545	Drink Frequency 12 Months
0.025946	Total Tobacco 7 Days
0.05339	Tobacco Frequency 7 Days
0.18588	Cocaine
0.064405	Hallucinogens
0.51675	Opiates
0.17273	Stims
0.00013921	Marijuana
0.059243	Father's Use History
0.0062048	Mother's Use History
-0.085539	Gender
Stimulants	
0.081758	ECS average connectivity
-0.033177	Age
-0.040674	Total Drinks 7 Days
0.037855	Drink Frequency 7 Days
-0.11055	Drinks/Day 12 Months
0.097284	Drink Frequency 12 Months
0.016823	Total Tobacco 7 Days
-0.020869	Tobacco Frequency 7 Days
0.23746	Cocaine
0.059621	Hallucinogens
0.15105	Opiates
0.24367	Sedatives
0.010344	Marijuana
-0.0034245	Father's Use History
0.017622	Mother's Use History
-0.0024692	Gender
Marijuana	
6.9903	ECS average connectivity
-0.2248	Age
-0.2595	Total Drinks 7 Days
-0.32945	Drink Frequency 7 Days
1.3554	Drinks/Day 12 Months
1.2835	Drink Frequency 12 Months
-0.18936	Total Tobacco 7 Days
0.32319	Cocaine
3.2374	Hallucinogens
0.015567	Sedatives
0.81998	Stimulants
0.50829	Father's Use History

Supplementary Table 11: Beta coefficients for all dependent variables used in the Valuation-Control Complex (VCC) GLM analysis.

Cocaine	
-0.049962	VCC Average Connectivity
-0.025465	Age
0.050458	Total Drinks 7 Days
0.091885	Drink Frequency 7 Days
-0.028744	Drinks/Day 12 Months
-0.0011179	Drink Frequency 12 Months
0.0014423	Total Tobacco 7 Days
0.014491	Tobacco Frequency 7 Days
0.3181	Hallucinogens
0.052813	Opiates
0.22304	Sedatives
0.20204	Stimulants
0.0034975	Marijuana
-0.049828	Father's Use History
0.051446	Mother's Use History
0.033915	Gender

Hallucinogens	
-0.24443	VCC Average Connectivity
0.035679	Age
0.054761	Total Drinks 7 Days
0.092129	Drink Frequency 7 Days
-0.059251	Drinks/Day 12 Months
-0.018987	Drink Frequency 12 Months
-0.016038	Total Tobacco 7 Days
-0.076975	Tobacco Frequency 7 Days
0.30293	Cocaine
0.20484	Opiates
0.0733	Sedatives
0.048699	Stimulants
0.033072	Marijuana
-0.02614	Father's Use History
0.091916	Gender

Opiates	
-0.10261	VCC Average Connectivity
0.033092	Age
-0.014998	Total Drinks 7 Days
0.15322	Drink Frequency 7 Days
-0.065477	Drinks/Day 12 Months
-0.081394	Drink Frequency 12 Months
0.01284	Total Tobacco 7 Days
0.03172	Tobacco Frequency 7 Days
0.042579	Cocaine
0.17342	Hallucinogens
0.49921	Sedatives
0.10336	Stimulants
0.0074543	Marijuana
-0.063216	Mother's Use History
0.0032496	Gender

Sedatives	
-0.00047632	VCC Average Connectivity
0.0027925	Age
-0.017812	Total Drinks 7 Days
-0.11732	Drink Frequency 7 Days
0.050815	Drinks/Day 12 Months
0.043372	Drink Frequency 12 Months
0.025975	Total Tobacco 7 Days
0.053506	Tobacco Frequency 7 Days
0.18604	Hallucinogens
0.064204	Opiates
0.5165	Sedatives
0.17289	Stimulants
0.00015173	Marijuana
0.059336	Father's Use History
0.0060493	Mother's Use History
-0.087633	Gender

Stimulants	
0.083756	VCC Average Connectivity
-0.040712	Total Drinks 7 Days
0.037457	Drink Frequency 7 Days
-0.10971	Drinks/Day 12 Months
0.097337	Drink Frequency 12 Months
0.017278	Total Tobacco 7 Days
-0.021013	Tobacco Frequency 7 Days
0.23776	Hallucinogens
0.060179	Opiates
0.15087	Sedatives
0.24392	Stimulants
0.010365	Marijuana
-0.0037528	Father's Use History
0.017598	Mother's Use History
-0.0046949	Gender

Marijuana	
7.021	VCC Average Connectivity
-0.18956	Age
-0.25597	Total Drinks 7 Days
-0.33041	Drink Frequency 7 Days
1.3605	Drinks/Day 12 Months
1.2828	Drink Frequency 12 Months
-0.18623	Total Tobacco 7 Days
0.32632	Cocaine
3.2403	Hallucinogens
0.86268	Opiates
0.016972	Sedatives
0.82179	Stimulants
0.50696	Father's Use History

GLM Outputs

This section contains GLM outputs for all models analyzed in this study. The data shown here can be analyzed in conjunction with data from Table 4 in the main text as well as Supplementary Tables 6-11. FDR, False Discovery Rate

Valuation System

Cocaine

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.19443	0.089555	2.1711	0.04075
x1 (VS)	-0.02103	0.019787	-1.0628	0.32531
x2 (Age)	0.095394	0.02011	4.7435	7.64E-06
x3 (Total Drinks 7 Days)	-0.071648	0.059659	-1.201	0.26474
x4 (Drink Frequency 7 Days)	0.038705	0.038097	1.0159	0.34706
x5 (Drinks/Day 12 Months)	0.07269	0.022665	3.2071	0.00278
x6 (Drink Frequency 12 Months)	-0.02034	0.018867	-1.0781	0.31842
x7 (Total Tobacco 7 Days)	0.12467	0.057777	2.1578	0.04193
x8 (Tobacco Frequency 7 Days)	0.72958	0.090899	8.0262	3.32E-14
x9 (Hallucinogens)	0.22728	0.099522	2.2837	0.03187
x10 (Opiates)	0.15184	0.11115	1.3662	0.2012
x11 (Sedatives)	0.34883	0.048851	7.1407	1.37E-11
x12 (Stimulants)	-0.01502	0.0079137	-1.8979	0.07241
x13 (Marijuana)	-0.024416	0.01955	-1.2489	0.24464
x14 (Father)	-0.045062	0.021248	-2.1208	0.04539
x15 (Mother)	-1.7305	0.49376	-3.5047	0.00103
x16_1 (Gender)	0.0032325	0.041932	0.077088	0.94646
x1:x7	0.068862	0.023587	2.9195	0.00653
x1:x8	0.075595	0.029889	2.5292	0.01865
x1:x9	0.11683	0.030854	3.7866	0.00038
x1:x10	-0.19901	0.032633	-6.0985	7.99E-09
x1:x11	-0.059796	0.021154	-2.8267	0.00848
x1:x14	-0.07448	0.022633	-3.2908	0.0021
x2:x3	0.059324	0.023459	2.5288	0.01865
x2:x8	0.16187	0.029347	5.5156	1.86E-07
x2:x9	0.14899	0.02793	5.3342	4.68E-07
x3:x5	0.06162	0.024432	2.5221	0.01886
x3:x8	0.10987	0.038972	2.8192	0.00866
x3:x9	0.11095	0.040208	2.7594	0.01033
x3:x11	-0.31816	0.03554	-8.9522	0
x3:x12	0.009552	0.0024814	3.8495	0.0003
x3:x13	0.092877	0.022332	4.1589	9.25E-05
x4:x8	-0.1899	0.047031	-4.0377	0.00015
x4:x9	-0.1584	0.044303	-3.5754	0.00081
x4:x11	0.36753	0.044691	8.2238	8.35E-15
x5:x7	-0.04644	0.019066	-2.4358	0
x5:x9	0.14006	0.034529	4.0564	0.00014
x5:x11	0.16412	0.029924	5.4847	2.19E-07
x6:x7	0.035634	0.018024	1.977	0.06165
x6:x8	0.093752	0.025732	3.6434	0.00063
x6:x11	-0.19016	0.029731	-6.3962	1.42E-09
x7:x8	0.10236	0.022425	4.5645	1.68E-05
x7:x9	-0.078784	0.021791	-3.6154	0.0007
x7:x11	0.090749	0.018833	4.8187	5.46E-06
x7:x15	-0.51708	0.29644	-1.7443	0.09828
x8:x11	0.083325	0.014053	5.9292	2.04E-08
x8:x14	0.21259	0.025334	8.3914	3.27E-15
x8:x15	-3.6422	0.53523	-6.8048	1.16E-10
x9:x10	-0.030834	0.013474	-2.2883	0.03154
x9:x11	0.071706	0.017284	4.1486	9.64E-05
x9:x14	-0.32437	0.03888	-8.3428	3.27E-15

x9:x15	-2.2652	0.50995	-4.4419	2.84E-05
x9:x16_1	0.28473	0.076179	3.7377	0.00046
x10:x11	-0.10987	0.014158	-7.7603	2.01E-13
x10:x13	-0.1689	0.022263	-7.5865	6.57E-13
x10:x15	1.2938	0.52907	2.4455	0.02237
10:x16_1	0.24486	0.079641	3.0746	0.00418
x11:x13	-0.054253	0.025505	-2.1272	0.04482
11:x16_1	-0.59964	0.059227	-10.125	0
x12:x15	0.11409	0.044495	2.564	0.0171

Hallucinogens

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.16301	0.073223	-2.2262	0.0361
x1(VS)	-0.0060003	0.027468	-0.21845	0.85355
x2 (Age)	0.063478	0.021268	2.9846	0.00538
x3 (Total Drinks 7 Days)	0.016162	0.041173	0.39255	0.73685
x4 (Drink Frequency 7 Days)	-0.037504	0.034784	-1.0782	0.31842
X5 (Drinks/Day 12 Months)	-0.042398	0.023267	-1.8223	0.08416
x6 (Drink Frequency 12 Months)	0.057581	0.023303	2.471	0.02123
x7 (Total Tobacco 7 Days)	-0.078312	0.022684	-3.4523	0.00123
x8 (Tobacco Frequency 7 Days)	0.20352	0.033153	6.1386	6.37E-09
x9 (Cocaine)	0.17129	0.046585	3.6769	0.00057
x10 (Opiates)	-0.6347	0.10444	-6.0771	8.96E-09
x11 (Sedatives)	0.72223	0.12231	5.905	2.32E-08
x12 (Stimulants)	0.018758	0.0037142	5.0503	1.84E-06
x13 (Marijuana)	-0.0058552	0.021002	-0.2788	0.80937
x14 (Father)	0.065674	0.022569	2.9099	0.00668
x15 (Mother)	-0.10345	0.3671	-0.2818	0.80798
x16_1 (Gender)	0.0091284	0.057403	0.15902	0.89549
x1:x7	-0.06369	0.024232	-2.6284	0.01459
x1:x8	0.13897	0.028756	4.8328	5.11E-06
x1:x9	-0.22626	0.034568	-6.5455	5.66E-10
x1:x10	0.21878	0.030616	7.146	1.31E-11
x1:x12	0.0062508	0.002594	2.4097	0.0244
x2:x5	0.05203	0.019513	2.6664	0.01329
x2:x9	0.20312	0.032651	6.2209	4.00E-09
x2:x10	-0.12568	0.037407	-3.3598	0.00169
x3:x9	0.10987	0.018551	5.923	2.10E-08
x3:x13	-0.065655	0.024772	-2.6504	0.01376
x4:x9	-0.1457	0.031356	-4.6466	1.18E-05
x4:x11	0.087609	0.022483	3.8967	0.00025
x5:x10	-0.141	0.030708	-4.5917	1.50E-05
x5:x11	-0.094957	0.025718	-3.6922	0.00054
x6:x8	0.10969	0.023653	4.6376	1.22E-05
x6:x10	0.070751	0.028022	2.5248	0.01874
x6:x12	-0.012169	0.0021495	-5.6614	8.57E-08
x7:x10	-4.4688	0.020475	-0.091498	2.54E-05
x7:x11	-0.066949	0.0181	-3.6989	0.00053
x8:x13	0.062218	0.02816	2.2095	0.03722
x9:x11	-0.081708	0.013456	-6.0724	9.15E-09
x9:x13	0.11838	0.031475	3.7609	0.00042
x9:x14	0.4122	0.044501	9.2628	0
x9:x16_1	0.42785	0.063802	6.7059	2.09E-10
x10:x14	-0.15891	0.028499	-5.5759	1.35E-07
x10:x15	4.9378	0.64754	7.6254	5.10E-13
x11:x12	0.013022	0.0021101	6.1713	5.28E-09
x11:x15	-3.465	0.60398	-5.737	5.78E-08
x11:x16_1	-0.26538	0.065512	-4.051	0.00014
X12:x16_1	0.014303	0.0049602	2.8836	0.00722

Opiates

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.077541	0.070267	-1.1035	0.30701
x1 (VS)	0.040826	0.020083	2.0329	0.05481
x2 (Age)	-0.038028	0.020823	-1.8263	0.08354
x3 (Total Drinks 7 Days)	0.026743	0.040899	0.65387	0.55642
x4 (Drink Frequency 7 Days)	-0.010397	0.033645	-0.30903	0.79001
x5 (Drinks/Day 12 Months)	-0.065305	0.022145	-2.9489	0.00599
x6 (Drink Frequency 7 Days)	-0.0084335	0.018997	-0.44393	0.69878
x7 (Total Tobacco 7 Days)	0.041498	0.022255	1.8647	0.07748
x8 (Tobacco Frequency 7 Days)	0.062363	0.033979	1.8354	0.08211
x9 (Cocaine)	0.53118	0.097458	5.4503	2.59E-07
x10 (Hallucinogens)	0.41164	0.050822	8.0995	2.00E-14
x11 (Sedatives)	-0.52993	0.09063	-5.8472	3.18E-08
x12 (Stimulants)	0.0087431	0.0026062	3.3547	0.00172
x13 (Marijuana)	-0.046655	0.019967	-2.3367	0.02829
x14 (Father)	-0.016033	0.020014	-0.80108	0.46662
x15 (Mother)	-0.013956	0.35628	-0.039172	0.97223
x16_1 (Gender)	-0.00019884	0.04351	-0.00457	0.99635
x1:x5	-0.047223	0.02008	-2.3517	0.0275
x1:x7	0.07354	0.022599	3.2541	0.00237
x1:x8	0.11462	0.019541	5.8657	2.90E-08
x1:x14	-0.036665	0.021236	-1.7265	0.10179
x2:x10	0.065478	0.032658	2.005	0.05848
x2:x11	-0.17539	0.032013	-5.4786	2.24E-07
x3:x7	0.042794	0.018475	2.3163	0.02965
x3:x9	0.19854	0.022039	9.0082	0
x4:x9	-0.19573	0.033408	-5.8589	2.98E-08
x4:x10	0.065957	0.032399	2.0358	0.05451
x5:x7	-0.061074	0.020429	-2.9896	0.0053
x5:x9	-0.063834	0.028204	-2.2633	0.03322
x5:x10	0.083089	0.02691	3.0876	0.00404
x5:x11	-0.14065	0.026189	-5.3707	3.86E-07
x6:x9	-0.071936	0.024398	-2.9484	0.00599
x6:x10	-0.12058	0.031964	-3.7725	0.0004
x6:x11	0.16286	0.024471	6.6553	2.89E-10
x7:x8	-0.047018	0.018819	-2.4985	0.02
x7:x9	0.14016	0.023093	6.0696	9.25E-09
x8:x10	-0.082762	0.011907	-6.9506	4.69E-11
x8:x11	0.04842	0.010479	4.6206	1.32E-05
x9:x15	-2.2096	0.4736	-4.6655	1.09E-05
x9:x16_1	-0.19427	0.063385	-3.0649	0.00428
x10:x12	0.0052223	0.0025444	2.0524	0.0527
x10:x13	-0.13244	0.020595	-6.4307	1.15E-09
x11:x15	3.216	0.41204	7.8051	1.48E-13
x11:x16_1	0.31369	0.059312	5.2888	5.84E-07

Sedatives

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.022372	0.062735	0.35661	0.75943
x1 (VS)	0.04005	0.01716	2.3339	0.02845
x2 (Age)	0.0014991	0.017759	0.084415	0.94466
x3 (Total Drinks 7 Days)	0.20616	0.074978	2.7496	0.0106
x4 (Drink Frequency 7 Days)	-0.059464	0.029751	-1.9987	0.05903
x5 (Drinks/Day 12 Months)	0.00073822	0.019948	0.037007	0.97281
x6 (Drink Frequency 12 Months)	0.0039006	0.018343	0.21265	0.85716
x7 (Total Tobacco 7 Days)	0.022957	0.019201	1.1956	0.26677
x8 (Tobacco Frequency 7 Days)	-0.65301	0.10563	-6.1818	5.03E-09
x9 (Cocaine)	0.20559	0.082675	2.4867	0.02052
x10 (Hallucinogens)	0.10711	0.056903	1.8824	0.07478
x11 (Opiates)	0.94797	0.093351	10.155	0
x12 (Stimulants)	-0.0012394	0.0022654	-0.54707	0.62541
x13 (Marijuana)	0.17369	0.044953	3.8638	0.00029
x14 (Father)	0.0062953	0.018357	0.34294	0.76735
x15 (Mother)	0.29914	0.31024	0.96421	0.37293
x16_1 (Gender)	0.0035812	0.037342	0.095903	0.93814
x1:x5	0.044131	0.017117	2.5782	0.01658
x1:x8	0.05633	0.027295	2.0637	0.05137
x1:x9	0.087239	0.027482	3.1744	0.00309
x1:x10	-0.11234	0.025444	-4.415	3.20E-05
x1:x11	0.1094	0.01925	5.6831	7.78E-08
x2:x4	0.035095	0.01753	2.0019	0.05882
x2:x9	-0.13456	0.024858	-5.4131	3.14E-07
x2:x11	0.18204	0.030242	6.0193	1.21E-08
x3:x4	-0.038749	0.018143	-2.1358	0.04415
x3:x10	-0.29372	0.024593	-11.943	0
x3:x11	0.37067	0.032158	11.527	0
x3:x15	-0.95341	0.31225	-3.0533	0.0044
x4:x8	-0.062641	0.025853	-2.423	0.02363
x4:x10	0.089163	0.029154	3.0583	0.00437
x4:x11	-0.40739	0.04155	-9.8047	0
x5:x6	0.034425	0.01492	2.3074	0.03026
x5:x7	0.040748	0.016035	2.5413	0.01814
x5:x8	-0.1544	0.025616	-6.0276	1.16E-08
x5:x9	-0.095024	0.025905	-3.6681	0.00058
x5:x10	0.41269	0.029598	13.943	0
x5:x11	-0.21047	0.027074	-7.7739	1.84E-13
x6:x8	-0.16845	0.025648	-6.5679	4.98E-10
x7:x8	0.1052	0.017689	5.9472	1.85E-08
x7:x11	-0.078702	0.017486	-4.5008	2.20E-05
x7:x14	-0.030747	0.012369	-2.4859	0.02053
x8:x9	0.050822	0.012278	4.1392	0.0001
x8:x10	-0.12095	0.013251	-9.1279	0
x8:x11	0.058513	0.012748	4.59	1.51E-05
x8:x15	5.4137	0.50968	10.622	0
x8:x16_1	-0.24899	0.059585	-4.1788	8.56E-05
x9:x10	0.10356	0.01592	6.5047	7.31E-10
x9:x11	-0.097371	0.018421	-5.2859	5.90E-07
x9:x12	-0.012427	0.0032082	-3.8735	0.00028
x9:x16_1	0.1893	0.060695	3.1188	0.00367
x10:x11	0.031576	0.011918	2.6494	0.01377
x10:x12	0.01372	0.002781	4.9333	3.18E-06
x10:x13	0.26187	0.025894	10.113	0
x10:x16_1	0.25412	0.054138	4.694	9.55E-06
x11:x12	-0.0076042	0.0033078	-2.2989	0.03084
x11:x13	-0.12105	0.023165	-5.2257	7.85E-07
x11:x15	-3.9793	0.48393	-8.2228	8.35E-15
x13:x15	-0.58812	0.23422	-2.511	0.01935

Stimulants

	Estimate	Standard Error	t-stat	FDR-corrected p-values
x1 (VS)	-0.015797	0.02277	-0.69377	0.53072
x2 (Age)	-0.0021483	0.023112	-0.092952	0.93938
x3 (Total Drinks 7 Days)	0.04537	0.09465	0.47934	0.6735
x4 (Drink Frequency 7 Days)	-0.049917	0.044039	-1.1335	0.29366
x5 (Drinks/Day 12 Months)	0.12911	0.02559	5.0453	1.88E-06
x6 (Drink Frequency 12 Months)	0.046473	0.021799	2.1318	0.04444
x7 (Total Tobacco 7 Days)	-0.03839	0.024451	-1.5701	0.13849
x8 (Tobacco Frequency 7 Days)	1.014	0.10643	9.5275	0
x9 (Cocaine)	0.088885	0.034089	2.6074	0.01532
x10 (Hallucinogens)	-0.11112	0.086958	-1.2779	0.23327
x11 (Opiates)	0.30046	0.058197	5.1627	1.07E-06
x12 (Sedatives)	0.0076386	0.0030082	2.5393	0.0182
x13 (Marijuana)	-0.02888	0.022981	-1.2567	0.24168
x14 (Father)	0.064528	0.063226	1.0206	0.3451
x15 (Mother)	-0.6984	0.4068	-1.7168	0.10376
x16_1 (Gender)	0.0095395	0.049015	0.19463	0.8706
x1:x6	-0.050192	0.020432	-2.4566	0.02193
x1:x11	-0.0724	0.024844	-2.9142	0.00662
x1:x14	-0.060044	0.025055	-2.3965	0.02503
x2:x11	0.13787	0.032584	4.2312	6.95E-05
x3:x5	0.078607	0.027167	2.8935	0.00701
x3:x8	-0.22526	0.029655	-7.596	6.15E-13
x3:x11	0.3232	0.029161	11.083	0
x3:x12	0.0088373	0.0026012	3.3974	0.0015
x3:x15	-0.8904	0.37182	-2.3947	0.0251
x4:x8	0.15143	0.03608	4.1971	7.97E-05
x4:x11	-0.36282	0.038143	-9.5122	0
x5:x8	0.29091	0.036896	7.8848	8.46E-14
x5:x11	-0.14723	0.03378	-4.3586	4.06E-05
x6:x10	0.22343	0.027907	8.0062	3.66E-14
x7:x9	-0.066402	0.022901	-2.8995	0.00689
x7:x11	-0.099145	0.020526	-4.8303	5.16E-06
x8:x9	0.038182	0.014379	2.6554	0.01359
x8:x11	-0.065418	0.011601	-5.6388	9.64E-08
x8:x13	-0.15984	0.029059	-5.5007	2.01E-07
x8:x15	-3.6715	0.53941	-6.8066	1.14E-10
x8:x16_1	-0.3597	0.069834	-5.1508	1.12E-06
x9:x10	-0.032032	0.013357	-2.3982	0.02496
x10:x14	0.081362	0.029542	2.7541	0.01047
x10:x15	2.3972	0.47992	4.9951	2.39E-06
x11:x16_1	0.23292	0.058028	4.0138	0.00016
x12:x14	0.0064412	0.0024247	2.6565	0.01359
x14:x15	-0.81659	0.26801	-3.0469	0.00449
x14:x16_1	0.15155	0.049576	3.0569	0.00437

Marijuana

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	8.3068	0.34772	23.89	0
x1 (VS)	-0.019352	0.24243	-0.079828	0.94646
x2 (Age)	-0.18552	0.24165	-0.76773	0.48626
x3 (Total Drinks 7 Days)	0.78284	0.51027	1.5342	0.14825
x4 (Drink Frequency 7 Days)	0.45268	0.4113	1.1006	0.30801
x5 (Drinks/Day 12 Months)	0.63854	0.28214	2.2632	0.03322
x6 (Drink Frequency 12 Months)	-1.1296	0.34158	-3.307	0.002
x7 (Total Tobacco 7 Days)	1.9231	0.24825	7.7466	2.14E-13
x8 (Tobacco Frequency 7 Days)	0.52491	0.30152	1.7408	0.09886
x9 (Cocaine)	4.0786	0.34962	11.666	0
x10 (Hallucinogens)	1.1093	0.34118	3.2512	0.00239
x11 (Opiates)	0.47346	0.42039	1.1262	0.29672
x12 (Sedatives)	0.84531	0.29437	2.8716	0.00748
x13 (Stimulants)	0.64646	0.23954	2.6987	0.01217
x14 (Father)	0.25295	0.23984	1.0547	0.32907
x16_1 (Gender)	0.061843	0.51184	0.12082	0.92141
x1:x8	-0.50742	0.20172	-2.5154	0.01917
x1:x13	0.70711	0.27039	2.6152	0.01504
x2:x7	-0.5338	0.24006	-2.2236	0.03613
x3:x9	-0.50275	0.14888	-3.377	0.0016
x4:x5	-1.0985	0.27824	-3.9481	0.00021
x5:x6	-1.0304	0.21604	-4.7696	6.80E-06
x6:x9	0.38691	0.18561	2.0846	0.04913
x6:x16_1	0.82055	0.42794	1.9174	0.06945
x7:x12	-0.49909	0.1745	-2.8601	0.00772
x10:x14	-0.80167	0.26792	-2.9922	0.00528
x11:x16_1	-1.0505	0.44694	-2.3505	0.02754

Executive Control System

Cocaine

	Estimate	Standard Error	t-stat	FDR-corrected p-values
x1 (ECS)	-0.031471	0.020638	-1.5249	0.15056
x2 (Age)	0.098818	0.020121	4.9112	3.54E-06
x3 (Total Drinks 7 Days)	-0.072847	0.061224	-1.1898	0.269
x4 (Drink Frequency 7 Days)	0.049544	0.038299	1.2936	0.22726
x5 (Drinks/Day 12 Months)	0.071731	0.023154	3.098	0.00393
x6 (Drink Frequency 12 Months)	0.01366	0.02488	0.54902	0.62478
x7 (Total Tobacco 7 Days)	0.11902	0.058074	2.0494	0.05301
x8 (Tobacco Frequency 7 Days)	0.76604	0.092808	8.254	7.52E-15
x9 (Hallucinogens)	0.43617	0.11062	3.9431	0.00021
x10 (Opiates)	-0.14972	0.11575	-1.2935	0.22726
x11 (Sedatives)	0.37474	0.04997	7.4993	1.20E-12
x12 (Stimulants)	-0.021634	0.0084403	-2.5632	0.0171
x13 (Marijuana)	-0.026332	0.019564	-1.346	0.20838
x14 (Father)	-0.02156	0.021297	-1.0123	0.34812
x15 (Mother)	-1.6654	0.49743	-3.348	0.00176
x16_1 (Gender)	-0.031925	0.055064	-0.57979	0.60467
x1:x7	0.061485	0.021668	2.8376	0.00824
x1:x9	0.17194	0.036951	4.6534	1.15E-05
x1:x10	-0.2171	0.037325	-5.8165	3.77E-08
x1:x11	-0.059511	0.022342	-2.6637	0.01337
x1:x14	-0.059614	0.018062	-3.3005	0.00204
x2:x3	0.059518	0.023665	2.515	0.01917
x2:x8	0.15265	0.029622	5.1532	1.12E-06
x2:x9	0.18801	0.029605	6.3507	1.86E-09
x3:x5	0.069751	0.024554	2.8407	0.00819
x3:x8	0.13252	0.04014	3.3016	0.00203
x3:x9	0.084688	0.041412	2.045	0.05349
x3:x11	-0.31281	0.0366	-8.5467	0
x3:x12	0.0064776	0.0029109	2.2253	0.0361
x3:x13	0.086285	0.022464	3.841	0.00031
x4:x8	-0.18884	0.047314	-3.9912	0.00018
x4:x9	-0.13914	0.045219	-3.0769	0.00416
x4:x11	0.36538	0.045181	8.087	2.24E-14
x5:x7	-0.044428	0.01896	-2.3432	0.02789
x5:x9	0.1906	0.037116	5.1354	1.22E-06
x5:x11	0.13746	0.028916	4.7539	7.32E-06
x6:x7	0.04789	0.018668	2.5653	0.0171
x6:x8	0.14771	0.028293	5.2207	8.03E-07
x6:x11	-0.19084	0.030374	-6.2829	2.78E-09
x6:x12	-0.0051556	0.0026829	-1.9217	0.06889
x7:x8	0.078983	0.022407	3.5248	0.00096
x7:x9	-0.054534	0.02224	-2.452	0.02217
x7:x11	0.085068	0.018651	4.5609	1.69E-05
x7:x15	-0.57584	0.29704	-1.9386	0.06676
x8:x11	0.076395	0.014183	5.3865	3.61E-07
x8:x14	0.19187	0.023467	8.1763	1.23E-14
x8:x15	-3.7973	0.54146	-7.013	3.19E-11
x9:x10	-0.025156	0.013487	-1.8652	0.07748
x9:x11	0.07289	0.017395	4.1904	8.19E-05
x9:x14	-0.31752	0.039577	-8.023	3.39E-14
x9:x15	-2.9559	0.5444	-5.4296	2.90E-07
x9:x16_1	0.16725	0.086066	1.9433	0.06623
x10:x11	-0.11995	0.014434	-8.3097	4.74E-15
x10:x13	-0.15598	0.022114	-7.0536	2.46E-11
x10:x15	2.2751	0.52492	4.3343	4.52E-05
x10:x16_1	0.3902	0.091601	4.2598	6.20E-05

x11:x13	-0.057548	0.025445	-2.2616	0.03325
x11:x16_1	-0.59073	0.059839	-9.8719	0
x12:x15	0.12277	0.04466	2.749	0.0106
x12:x16_1	0.008566	0.0047225	1.8139	0.08561

Hallucinogens

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.22681	0.094423	-2.4021	0.02479
x1 (ECS)	0.07726	0.032183	2.4007	0.02484
x2 (Age)	0.0022272	0.028569	0.077959	0.94646
x3 (Total Drinks 7 Days)	-0.032389	0.045157	-0.71727	0.51643
x4 (Drink Frequency 7 Days)	-0.0046904	0.036355	-0.12902	0.91591
x5 (Drinks/Day 12 Months)	-0.021125	0.024389	-0.86614	0.42678
x6 (Drink Frequency 12 Months)	0.10623	0.029075	3.6538	0.00061
x7 (Total Tobacco 7 Days)	-0.080646	0.022677	-3.5563	0.00086
x8 (Tobacco Frequency 7 Days)	0.59539	0.12458	4.7793	6.54E-06
x9 (Cocaine)	0.14217	0.068832	2.0654	0.05131
x10 (Opiates)	-0.66893	0.13481	-4.962	2.76E-06
x11 (Sedatives)	0.70246	0.14078	4.9896	2.45E-06
x12 (Stimulants)	0.030857	0.0083792	3.6825	0.00056
x13 (Marijuana)	0.0065073	0.021284	0.30574	0.79094
x14 (Father)	0.06189	0.023545	2.6286	0.01459
x15 (Mother)	0.13255	0.5071	0.26139	0.82224
x16_1 (Gender)	-0.02261	0.058349	-0.38749	0.73988
x1:x4	0.036806	0.020836	1.7665	0.09425
x1:x8	0.050288	0.021219	2.37	0.02642
x1:x13	-0.039922	0.020628	-1.9353	0.06717
x1:x16_1	-0.096729	0.043237	-2.2372	0.03525
x2:x9	0.13239	0.033821	3.9145	0.00024
x2:x11	-0.088736	0.033883	-2.6189	0.01494
x2:x12	0.006427	0.0024571	2.6157	0.01504
x3:x9	0.11837	0.02814	4.2065	7.70E-05
x3:x10	-0.056919	0.029891	-1.9042	0.07149
x4:x5	0.053899	0.025363	2.1251	0.04498
x4:x7	-0.037749	0.021035	-1.7946	0.08891
x4:x9	-0.12396	0.034019	-3.6437	0.00063
x4:x11	0.064681	0.026296	2.4598	0.02183
x4:x13	-0.053216	0.022399	-2.3758	0.02624
x5:x9	0.093314	0.047296	1.973	0.062
x5:x10	-0.11902	0.038521	-3.0898	0.00402
x5:x11	-0.059315	0.028216	-2.1022	0.04736
x6:x8	0.14191	0.026411	5.373	3.86E-07
x6:x10	0.07462	0.03775	1.9767	0.06165
x6:x12	-0.011127	0.0022261	-4.9985	2.37E-06
x6:x16_1	-0.082258	0.036965	-2.2253	0.0361
x7:x8	-0.069254	0.022405	-3.091	0.00401
x7:x11	-0.070195	0.018432	-3.8083	0.00035
x8:x9	0.034384	0.013883	2.4767	0.02099
x8:x13	0.064465	0.027173	2.3724	0.02634
x8:x15	-2.3105	0.67801	-3.4078	0.00145
x9:x11	-0.12545	0.016288	-7.7016	2.99E-13
x9:x12	0.0083335	0.0035499	2.3475	0.02762
x9:x13	0.11842	0.031964	3.7047	0.00052
x9:x14	0.22351	0.04463	5.008	2.26E-06
x10:x12	-0.020697	0.0039431	-5.249	6.99E-07
x10:x14	0.066316	0.024384	2.7197	0.01148
x10:x15	5.9728	0.74249	8.0443	3.02E-14
x10:x16_1	0.58778	0.065169	9.0193	0
x11:x12	0.025602	0.0041578	6.1575	5.78E-09

x11:x14	-0.078412	0.020079	-3.9051	0.00025
x11:x15	-3.9675	0.69157	-5.737	5.78E-08
x11:x16_1	-0.4959	0.072623	-6.8284	1.01E-10
x12:x15	-0.073917	0.043568	-1.6966	0.10813
x12:x16_1	0.019456	0.0050381	3.8618	0.00029

Opiates

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.11094	0.072458	-1.5311	0.14896
x1 (ECS)	0.047756	0.020553	2.3236	0.02913
x2 (Age)	-0.021631	0.020821	-1.0389	0.33681
x3 (Total Drinks 7 Days)	-0.047945	0.050102	-0.95694	0.37649
x4 (Drink Frequency 7 Days)	0.0048795	0.034118	0.14302	0.90631
x5 (Drinks/Day 12 Months)	-0.055308	0.022567	-2.4508	0.02217
x6 (Drink Frequency 12 Months)	-0.011338	0.018814	-0.60266	0.59001
x7 (Total Tobacco 7 Days)	0.035401	0.021771	1.626	0.12424
x8 (Tobacco Frequency 7 Days)	-0.061976	0.13551	-0.45736	0.68937
x9 (Cocaine)	0.48194	0.10954	4.3996	3.41E-05
x10 (Hallucinogens)	1.1487	0.13976	8.2192	8.35E-15
x11 (Sedatives)	-0.86756	0.12755	-6.8017	1.17E-10
x12 (Stimulants)	0.0088007	0.0025997	3.3853	0.00156
x13 (Marijuana)	-0.015363	0.026664	-0.57616	0.60653
x15 (Mother)	-0.029746	0.35588	-0.083582	0.94466
x16_1 (Gender)	0.013537	0.04393	0.30815	0.79001
x1:x7	0.075083	0.020975	3.5796	0.0008
x1:x10	0.18812	0.032419	5.8028	4.02E-08
x1:x11	-0.071123	0.023194	-3.0665	0.00427
x2:x10	0.21161	0.037412	5.6563	8.79E-08
x2:x11	-0.2254	0.032938	-6.8432	9.19E-11
x3:x4	0.037573	0.020335	1.8477	0.08012
x3:x7	0.03691	0.018607	1.9837	0.06093
x3:x8	-0.033757	0.016892	-1.9984	0.05903
x3:x9	0.19832	0.025051	7.9165	6.80E-14
x3:x13	0.042771	0.022177	1.9286	0.0681
x4:x9	-0.1885	0.033153	-5.6856	7.69E-08
x5:x7	-0.047404	0.019996	-2.3707	0.02641
x5:x9	-0.069289	0.0295	-2.3488	0.02759
x5:x10	0.13164	0.031051	4.2394	6.73E-05
x5:x11	-0.14536	0.026273	-5.5329	1.70E-07
x6:x9	-0.077845	0.024543	-3.1719	0.00311
x6:x10	-0.18868	0.028069	-6.7218	1.90E-10
x6:x11	0.21566	0.028283	7.625	5.10E-13
x7:x9	0.1148	0.019752	5.8117	3.84E-08
x8:x9	0.034124	0.01252	2.7255	0.0113
x8:x10	-0.086218	0.01277	-6.7515	1.59E-10
x8:x11	0.04198	0.010919	3.8449	0.00031
x8:x12	-0.016679	0.0040037	-4.1659	9.01E-05
x8:x15	1.6512	0.70148	2.3538	0.0274
x8:x16_1	0.26391	0.072741	3.6281	0.00067
x9:x15	-1.9114	0.52588	-3.6347	0.00065
x9:x16_1	-0.18298	0.071709	-2.5516	0.01764
x10:x12	0.017671	0.0036467	4.8457	4.80E-06
x10:x13	-0.13732	0.020882	-6.5758	4.74E-10
x10:x15	-4.713	0.77258	-6.1002	7.89E-09
x10:x16_1	-0.45154	0.074567	-6.0555	9.98E-09
x11:x15	4.9326	0.61908	7.9676	4.77E-14
x11:x16_1	0.50246	0.080127	6.2707	2.96E-09
x12:x13	-0.0042135	0.0020636	-2.0418	0.05381

Sedatives

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.063295	0.06235	1.0152	0.34706
x1 (ECS)	0.022653	0.017454	1.2979	0.22614
x2 (Age)	-0.0063168	0.017389	-0.36326	0.75515
x3 (Total Drinks 7 Days)	0.16392	0.07281	2.2513	0.03408
x4 (Drink Frequency 7 Days)	-0.060771	0.029101	-2.0882	0.04885
x5 (Drinks/Day 12 Months)	0.0063587	0.019491	0.32624	0.77862
x6 (Drink Frequency 12 Months)	0.026354	0.017924	1.4703	0.1666
x7 (Total Tobacco 7 Days)	0.012226	0.018608	0.65701	0.55494
x8 (Tobacco Frequency 7 Days)	-0.67415	0.11159	-6.0413	1.09E-08
x9 (Cocaine)	0.1641	0.05368	3.057	0.00437
x10 (Hallucinogens)	0.27159	0.072366	3.753	0.00043
x11 (Opiates)	1.1102	0.10969	10.121	0
x12 (Stimulants)	-0.00077495	0.0022157	-0.34976	0.76323
x13 (Marijuana)	0.17497	0.044005	3.9761	0.00019
x14 (Father)	0.0008118	0.017085	0.047516	0.96787
x15 (Mother)	0.18437	0.30614	0.60224	0.59001
x16_1 (Gender)	-0.0027746	0.037746	-0.073507	0.94819
x1:x5	0.037386	0.017247	2.1677	0.04103
x1:x8	0.052648	0.02148	2.451	0.02217
x1:x9	-0.07494	0.02616	-2.8647	0.00763
x1:x11	0.18558	0.023318	7.9586	5.06E-14
x2:x10	-0.18269	0.023965	-7.6233	5.15E-13
x2:x11	0.20098	0.027039	7.4329	1.88E-12
x3:x4	-0.041226	0.017809	-2.3148	0.02972
x3:x8	-0.056293	0.02099	-2.6818	0.01275
x3:x9	-0.20718	0.027782	-7.4572	1.60E-12
x3:x10	-0.11854	0.024863	-4.7676	6.87E-06
x3:x11	0.41456	0.030515	13.585	0
x3:x15	-0.56774	0.30361	-1.87	0.07679
x4:x10	0.15499	0.029285	5.2924	5.79E-07
x4:x11	-0.44184	0.039512	-11.182	0
x5:x6	0.040378	0.014729	2.7415	0.01082
x5:x7	0.032768	0.015676	2.0903	0.04869
x5:x8	-0.1174	0.030016	-3.9111	0.00024
x5:x9	0.40788	0.031176	13.083	0
x5:x10	-0.064984	0.026973	-2.4092	0.0244
x5:x11	-0.23925	0.026356	-9.0774	0
x6:x8	-0.16679	0.027628	-6.0369	1.10E-08
x6:x9	0.079715	0.030611	2.6042	0.01544
x7:x8	0.090469	0.01705	5.3061	5.39E-07
x7:x11	-0.08084	0.017622	-4.5875	1.52E-05
x8:x9	-0.12458	0.013819	-9.0147	0
x8:x10	0.065791	0.012247	5.372	3.86E-07
x8:x11	0.065053	0.012399	5.2466	7.07E-07
x8:x14	-0.07299	0.018513	-3.9426	0.00021
x8:x15	5.6013	0.53161	10.536	0
x8:x16_1	-0.25341	0.062201	-4.0741	0.00013
x9:x10	0.12583	0.015716	8.0068	3.66E-14
x9:x12	0.013536	0.0025597	5.2882	5.85E-07
x9:x13	0.30761	0.024737	12.435	0
x9:x16_1	0.24058	0.060539	3.974	0.00019
x10:x11	-0.12079	0.016654	-7.2531	6.41E-12
x10:x12	-0.017375	0.0027485	-6.3217	2.20E-09
x10:x14	0.10847	0.022908	4.735	7.90E-06
x10:x16_1	0.25037	0.061208	4.0905	0.00012
x11:x13	-0.16537	0.023859	-6.9314	5.31E-11
x11:x15	-5.0012	0.54716	-9.1402	0
x11:x16_1	-0.16684	0.063851	-2.613	0.01511

x13:x15	-0.56159	0.22925	-2.4497	0.0222

Stimulants

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.11131	0.078811	1.4124	0.18556
x1 (ECS)	-0.038318	0.023064	-1.6614	0.11569
x2 (Age)	-0.023192	0.02256	-1.028	0.34208
x3 (Total Drinks 7 Days)	0.12014	0.089307	1.3453	0.20838
x4 (Drink Frequency 7 Days)	-0.085533	0.037819	-2.2616	0.03325
x5 (Drinks/Day 12 Months)	0.12023	0.025063	4.7973	6.02E-06
x6 (Drink Frequency 12 Months)	0.064702	0.02136	3.0291	0.00473
x7 (Total Tobacco 7 Days)	-0.01706	0.024613	-0.69314	0.53072
x8 (Tobacco Frequency 7 Days)	1.3016	0.14539	8.952	0
x9 (Cocaine)	0.32042	0.089353	3.586	0.00078
x10 (Hallucinogens)	-0.12336	0.10733	-1.1493	0.28653
x11 (Opiates)	-0.29382	0.1409	-2.0853	0.04913
x12 (Sedatives)	0.0051447	0.0029203	1.7617	0.09494
x13 (Marijuana)	-0.014103	0.022828	-0.61777	0.58116
x14 (Father)	0.021006	0.060127	0.34936	0.76323
x15 (Mother)	-0.64329	0.4018	-1.601	0.13053
x16_1 (Gender)	0.050649	0.049595	1.0213	0.3451
x1:x8	0.078438	0.026103	3.0049	0.00509
x1:x10	-0.098128	0.031071	-3.1582	0.00325
x1:x11	-0.25233	0.03497	-7.2156	8.18E-12
x2:x5	-0.039931	0.020773	-1.9223	0.06889
x3:x8	-0.24174	0.030043	-8.0463	2.93E-14
x3:x11	0.26766	0.028124	9.5172	0
x3:x12	0.0078777	0.0025577	3.0799	0.00412
x3:x15	-0.70687	0.3854	-1.8341	0.08222
x4:x8	0.1713	0.037185	4.6068	1.41E-05
x4:x11	-0.26031	0.035715	-7.2886	5.00E-12
x5:x8	0.38852	0.035747	10.868	0
x5:x10	0.1064	0.046633	2.2815	0.03199
x5:x11	-0.2372	0.042345	-5.6015	1.18E-07
x6:x10	0.27017	0.029604	9.1263	0
x7:x8	-0.091	0.02046	-4.4478	2.77E-05
x7:x13	-0.03657	0.01851	-1.9757	0.0617
x8:x10	0.035504	0.018747	1.8938	0.07297
x8:x11	-0.056698	0.01254	-4.5213	2.01E-05
x8:x13	-0.099466	0.028304	-3.5141	0.001
x8:x15	-4.6699	0.72641	-6.4287	1.16E-09
x8:x16_1	-0.49516	0.07509	-6.5942	4.24E-10
x9:x10	-0.028789	0.016335	-1.7624	0.09494
x9:x15	-1.4531	0.54717	-2.6556	0.01359
x10:x11	-0.042229	0.014902	-2.8339	0.00831
x10:x15	2.8089	0.61931	4.5356	1.89E-05
x11:x15	2.1681	0.7847	2.7629	0.01024
x11:x16_1	0.5833	0.065393	8.9199	0
x12:x14	0.0088012	0.0023364	3.7669	0.00041
x14:x15	-0.65884	0.25482	-2.5856	0.01626
x14:x16_1	0.10843	0.047345	2.2902	0.03144

Marijuana

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	8.1708	0.2524	32.372	0
x1 (ECS)	-0.0025895	0.23706	-0.010923	0.99247
x2 (Age)	-0.17051	0.23635	-0.72146	0.51427
x3 (Total Drinks 7 Days)	0.84217	0.50579	1.6651	0.11514
x4 (Drink Frequency 7 Days)	0.3732	0.40998	0.91028	0.40216
x5 (Drinks/Day 12 Months)	0.6603	0.27926	2.3645	0.02676
x6 (Drink Frequency 12 Months)	-0.77861	0.25241	-3.0847	0.00406
x7 (Total Tobacco 7 Days)	1.9331	0.24641	7.8453	1.09E-13
x9 (Cocaine)	3.7293	0.29585	12.605	0
x10 (Hallucinogens)	0.98857	0.28357	3.4862	0.00109
x12 (Sedatives)	0.97543	0.27819	3.5064	0.00102
x13 (Stimulants)	0.55419	0.23598	2.3485	0.02759
x14 (Father)	0.35927	0.23685	1.5168	0.15272
x1:x6	0.7277	0.2061	3.5308	0.00094
x1:x13	0.45798	0.23166	1.9769	0.06165
x2:x7	-0.44289	0.23867	-1.8557	0.07881
x3:x9	-0.54216	0.13978	-3.8788	0.00027
x4:x5	-1.0871	0.27811	-3.9088	0.00024
x5:x6	-0.86162	0.20531	-4.1967	7.97E-05
x7:x12	-0.40667	0.16553	-2.4568	0.02193
x10:x14	-0.62233	0.25907	-2.4022	0.02479

<u>Valuation-Control Complex</u>

Cocaine

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.21801	0.092523	2.3562	0.02727
x1 (VCC)	0.0084324	0.028136	0.2997	0.79475
x2 (Age)	0.10037	0.020128	4.9865	2.47E-06
x3 (Total Drinks 7 Days)	-0.10526	0.060086	-1.7519	0.09684
x4 (Drink Frequency 7 Days)	0.064819	0.038244	1.6949	0.10836
x5 (Drinks/Day 12 Months)	0.074469	0.022767	3.2709	0.00224
x6 (Drink Frequency 12 Months)	-0.013122	0.018856	-0.69591	0.53021
x7 (Total Tobacco 7 Days)	0.0068708	0.021783	0.31542	0.7862
x8 (Tobacco Frequency 7 Days)	0.73355	0.092031	7.9707	4.72E-14
x9 (Hallucinogens)	0.46022	0.10924	4.213	7.50E-05
x10 (Opiates)	-0.10223	0.11273	-0.90693	0.4036
x11 (Sedatives)	0.42215	0.046208	9.1359	0
x12 (Stimulants)	-0.021302	0.0084282	-2.5275	0.01868
x13 (Marijuana)	-0.024113	0.019626	-1.2286	0.25296
x14 (Father)	-0.021175	0.021218	-0.99797	0.35535
x15 (Mother)	-1.6435	0.4966	-3.3095	0.00199
x16_1 (Gender)	-0.074189	0.055707	-1.3318	0.21321
x1:x7	0.070669	0.023378	3.0228	0.00481
x1:x9	0.18494	0.0346	5.345	4.43E-07
x1:x10	-0.19881	0.035036	-5.6745	8.10E-08
x1:x11	-0.048278	0.021719	-2.2228	0.03615
x1:x12	-0.0050319	0.0025167	-1.9994	0.05903
x1:x14	-0.056556	0.019263	-2.936	0.00622
x2:x3	0.057609	0.023547	2.4466	0.02235
x2:x8	0.14546	0.029644	4.9069	3.59E-06
x2:x9	0.1923	0.028601	6.7236	1.90E-10
x3:x5	0.074442	0.024544	3.0331	0.00469
x3:x8	0.1322	0.039361	3.3586	0.0017
x3:x9	0.091828	0.041076	2.2355	0.03533
x3:x11	-0.31607	0.036344	-8.6967	0
x3:x12	0.0084922	0.0025254	3.3627	0.00168
x3:x13	0.09164	0.022421	4.0873	0.00012
x4:x8	-0.19438	0.047011	-4.1348	0.0001
x4:x9	-0.15694	0.044449	-3.5307	0.00094
x4:x11	0.38012	0.044505	8.541	0
x5:x7	-0.054669	0.018642	-2.9325	0.00628
x5:x9	0.1659	0.035076	4.7297	8.09E-06
x5:x11	0.12783	0.028815	4.4362	2.89E-05
x6:x8	0.14028	0.025184	5.5701	1.39E-07
x6:x11	-0.19896	0.029957	-6.6416	3.16E-10
x7:x8	0.090095	0.022058	4.0846	0.00012
x7:x9	-0.084401	0.020769	-4.0638	0.00013
x7:x11	0.088186	0.018604	4.7401	7.74E-06
x8:x11	0.066431	0.012139	5.4727	2.31E-07
x8:x14	0.20564	0.024269	8.4733	1.70E-15
x8:x15	-3.4857	0.53358	-6.5327	6.14E-10
x9:x11	0.061953	0.016777	3.6926	0.00054
x9:x14	-0.31718	0.039307	-8.0695	2.46E-14
x9:x15	-3.3182	0.54241	-6.1176	7.21E-09
x9:x16_1	0.21742	0.081011	2.6838	0.0127
x10:x11	-0.11906	0.014158	-8.4095	1.70E-15
x10:x13	-0.1508	0.021665	-6.9606	4.47E-11
x10:x15	2.1964	0.52507	4.1831	8.43E-05
x10:x16_1	0.27352	0.080828	3.384	0.00157
x11:x13	-0.067314	0.025331	-2.6573	0.01359
x11:x16_1	-0.611	0.060297	-10.133	0

x12:x15	0.11112	0.043979	2.5266	0.01869
x12:x16_1	0.012633	0.0049171	2.5692	0.01694

Hallucinogens

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.2302	0.045132	-5.1006	1.43E-06
x1 (VCC)	0.039554	0.021982	1.7994	0.08812
x2 (Age)	0.079544	0.021581	3.6858	0.00055
x3 (Total Drinks 7 Days)	-0.040112	0.052844	-0.75905	0.49093
x4 (Drink Frequency 7 Days)	-0.032562	0.036065	-0.90285	0.40544
x5 (Drinks/Day 12 Months)	-0.014837	0.024371	-0.60879	0.58682
x6 (Drink Frequency 12 Months)	0.049534	0.023831	2.0785	0.04978
x7 (Total Tobacco 7 Days)	-0.081627	0.023138	-3.5278	0.00095
x8 (Tobacco Frequency 7 Days)	0.24265	0.031451	7.7154	2.68E-13
x9 (Cocaine)	0.24036	0.045751	5.2536	6.87E-07
x10 (Opiates)	0.14015	0.038309	3.6584	0.0006
x11 (Sedatives)	-0.014704	0.039698	-0.37039	0.75142
x12 (Stimulants)	0.01958	0.003775	5.1865	9.49E-07
x13 (Marijuana)	-0.011558	0.021676	-0.53324	0.63399
x14 (Father)	0.04771	0.022403	2.1296	0.04461
x16_1 (Gender)	-0.0094269	0.05861	-0.16084	0.89549
x1:x7	-0.048987	0.023284	-2.1039	0.04723
x1:x10	0.18501	0.023589	7.8429	1.13E-13
x2:x5	0.054542	0.019993	2.7281	0.01124
x2:x9	0.18734	0.02561	7.3152	4.15E-12
x3:x4	0.047491	0.022103	2.1486	0.04283
x3:x9	0.13593	0.019939	6.8177	1.07E-10
x3:x10	-0.045186	0.020885	-2.1636	0.04139
x3:x13	-0.058856	0.025261	-2.3299	0.0287
x4:x9	-0.1669	0.031788	-5.2502	6.95E-07
x4:x11	0.060532	0.022372	2.7057	0.01194
x5:x11	-0.12282	0.021667	-5.6684	8.27E-08
x6:x8	0.1384	0.022607	6.1218	7.00E-09
x6:x12	-0.010106	0.002213	-4.5669	1.66E-05
x7:x10	-0.056535	0.018009	-3.1394	0.00345
x7:x11	-0.073854	0.018195	-4.0591	0.00014
x9:x11	-0.10552	-7.7374	2.5757e-14	0.02123
x9:x13	0.11938	0.031575	3.7808	0.00039
x9:x14	0.26035	0.032551	7.9981	3.84E-14
x9:x16_1	0.15462	0.051702	2.9906	0.0053
x11:x12	0.017296	0.0023483	7.3653	2.94E-12
x11:x14	-0.044036	0.017689	-2.4896	0.02039
x12:x16_1	0.016868	0.0048777	3.4582	0.00121

Opiates

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	-0.13254	0.072169	-1.8365	0.08202
x1 (VCC)	0.040253	0.020747	1.9402	0.0666
x2 (Age)	-0.047837	0.021001	-2.2778	0.03225
x3 (Total Drinks 7 Days)	-0.036696	0.050317	-0.72928	0.5097
x4 (Drink Frequency 7 Days)	0.0079581	0.034373	0.23153	0.84409
x5 (Drinks/Day 12 Months)	-0.05452	0.022952	-2.3754	0.02624
x6 (Drink Frequency 12 Months)	-0.0063606	0.019117	-0.33272	0.77445
x7 (Total Tobacco 7 Days)	0.03682	0.022141	1.663	0.11547
x8 (Tobacco Frequency 7 Days)	-0.11278	0.054631	-2.0644	0.05135
x9 (Cocaine)	0.525	0.093189	5.6337	9.84E-08
x10 (Hallucinogens)	0.64991	0.044538	14.592	0
x11 (Sedatives)	-0.60287	0.095112	-6.3385	1.98E-09
x12 (Stimulants)	0.0079989	0.0026187	3.0546	0.00439
x13 (Marijuana)	-0.046051	0.019696	-2.338	0.02823
x15 (Mother)	0.060994	0.35664	0.17102	0.88857
x16_1 (Gender)	0.010387	0.044234	0.23482	0.84248
x1:x5	-0.045838	0.020606	-2.2245	0.03611
x1:x7	0.096573	0.022107	4.3685	3.91E-05
x1:x10	0.08931	0.02511	3.5567	0.00086
x2:x9	0.063258	0.028105	2.2508	0.03408
x2:x11	-0.21705	0.029353	-7.3945	2.42E-12
x3:x4	0.037971	0.020457	1.8561	0.07881
x3:x7	0.044184	0.018199	2.4278	0.0234
x3:x9	0.18642	0.022677	8.2207	8.35E-15
x4:x9	-0.17706	0.034732	-5.098	1.45E-06
x5:x7	-0.0482	0.01983	-2.4307	0.02326
x5:x11	-0.11275	0.021446	-5.2575	6.75E-07
x6:x9	-0.056763	0.023394	-2.4264	0.02345
x6:x10	-0.18749	0.027124	-6.9124	5.91E-11
x6:x11	0.20665	0.027502	7.514	1.08E-12
x7:x9	0.10117	0.019503	5.1874	9.49E-07
x8:x10	-0.056695	0.0082332	-6.8861	6.93E-11
x8:x11	0.047575	0.0094879	5.0142	2.20E-06
x8:x16_1	0.1865	0.062501	2.9839	0.00538
x9:x15	-2.2217	0.44545	-4.9876	2.46E-06
x9:x16_1	-0.20784	0.062155	-3.3439	0.00178
x10:x13	-0.094492	0.017929	-5.2702	6.37E-07
x10:x16_1	-0.21276	0.064134	-3.3174	0.00194
x11:x15	3.3288	0.40243	8.2716	6.22E-15
x11:x16_1	0.38016	0.074412	5.1088	1.38E-06

Sedatives

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.024389	0.062107	0.39269	0.73685
x1 (VCC)	0.013669	0.017229	0.79342	0.47029
x2 (Age)	-0.0066322	0.017301	-0.38333	0.7422
x3 (Total Drinks 7 Days)	0.1868	0.072644	2.5715	0.01687
x4 (Drink Frequency 7 Days)	-0.063683	0.029032	-2.1936	0.03869
x5 (Drinks/Day 12 Months)	0.0020966	0.019542	0.10728	0.93122
x6 (Drink Frequency 12 Months)	0.02655	0.017838	1.4884	0.16114
x7 (Total Tobacco 7 Days)	0.010509	0.018515	0.56762	0.61196
x8 (Tobacco Frequency 7 Days)	-0.64186	0.11007	-5.8316	3.47E-08
x9 (Cocaine)	0.18519	0.080406	2.3032	0.03054
x10 (Hallucinogens)	-0.12577	0.11793	-1.0665	0.32389
x11 (Opiates)	1.284	0.13956	9.2003	0
x12 (Stimulants)	-0.00034913	0.0022039	-0.15841	0.89549
x13 (Marijuana)	0.17437	0.043753	3.9853	0.00018
x14 (Father)	0.0016467	0.017034	0.096672	0.93814
x15 (Mother)	0.30026	0.30469	0.98546	0.36168
x16_1 (Gender)	0.02026	0.037392	0.54183	0.62847
x1:x5	0.039047	0.017251	2.2635	0.03322
x1:x8	0.082159	0.022537	3.6454	0.00063
x1:x10	-0.13514	0.029037	-4.6541	1.15E-05
x1:x11	0.18292	0.022003	8.3134	4.74E-15
x2:x9	-0.16846	0.02444	-6.893	6.79E-11
x2:x11	0.18293	0.02869	6.376	1.60E-09
x3:x4	-0.042658	0.017883	-2.3854	0.02571
x3:x8	-0.050011	0.020661	-2.4205	0.02375
x3:x9	-0.1456	0.027275	-5.3382	7.88E-08
x3:x10	-0.20886	0.027533	-7.586	6.57E-13
x3:x11	0.43029	0.031252	13.768	0
x3:x15	-0.68629	0.30292	-2.2655	0.03319
x4:x9	0.18042	0.030779	5.8618	2.96E-08
x4:x11	-0.46683	0.040112	-11.638	0
x5:x6	0.041472	0.014618	2.8371	0.00824
x5:x7	0.031309	0.015692	1.9952	0.05939
x5:x8	-0.10872	0.030201	-3.6	0.00074
x5:x9	-0.062916	0.027422	-2.2943	0.03115
x5:x10	0.38104	0.031854	11.962	0
x5:x11	-0.24011	0.026315	-9.1248	0
x6:x8	-0.16926	0.027264	-6.2079	4.32E-09
x6:x10	0.07902	0.030093	2.6259	0.01467
x7:x8	0.085114	0.017089	4.9806	2.54E-06
x7:x11	-0.087971	0.0177	-4.9701	2.66E-06
x8:x9	0.06194	0.012688	4.8817	4.07E-06
x8:x10	-0.12195	0.013914	-8.7646	0
x8:x11	0.069098	0.012761	5.4146	3.13E-07
x8:x14	-0.069737	0.018435	-3.783	0.00039
x8:x15	5.4962	0.52859	10.398	0
x8:x16_1	-0.2812	0.06243	-4.5043	2.18E-05
x9:x10	0.11883	0.015933	7.4584	1.60E-12
x9:x11	-0.11238	0.01855	-6.0583	9.89E-09
x9:x12	-0.014053	0.0030793	-4.5635	1.68E-05
x9:x14	0.09909	0.022882	4.3305	4.57E-05
x9:x16_1	0.24428	0.061687	3.96	0.0002
x10:x12	0.015054	0.0026221	5.7414	5.70E-08
x10:x13	0.30791	0.024809	12.411	0
x10:x15	1.3966	0.58817	2.3746	0.02624
x10:x16_1	0.35179	0.068092	5.1663	1.05E-06
x11:x12	-0.0058883	0.0032675	-1.8021	0.08774
x11:x13	-0.16606	0.023821	-6.9713	4.20E-11

	Estimate	Standard Error	t-stat	FDR-corrected p-values
x11:x15	-5.6098	0.6812	-8.2352	7.52E-15
x11:x16_1	-0.16617	0.064809	-2.564	0.0171
x13:x15	-0.56871	0.22808	-2.4935	0.02024

Stimulants

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.12215	0.079536	1.5358	0.14799
x1 (VCC)	-0.026834	0.023023	-1.1655	0.27973
x3 (Total Drinks 7 Days)	0.18837	0.088276	2.1338	0.04429
x4 (Drink Frequency 7 Days)	-0.090726	0.038093	-2.3817	0.02591
x5 (Drinks/Day 12 Months)	0.11509	0.025024	4.5993	1.45E-05
x6 (Drink Frequency 12 Months)	0.057816	0.021674	2.6675	0.01327
x7 (Total Tobacco 7 Days)	-0.018836	0.02483	-0.75862	0.49093
X8 (Tobacco Frequency 7 Days)	0.98933	0.12317	8.0322	3.11E-14
x9 (Cocaine)	0.097101	0.032108	3.0242	0.0048
x10 (Hallucinogens)	-0.20911	0.084441	-2.4764	0.02099
x11 (Opiates)	0.32695	0.076938	4.2496	6.45E-05
x12 (Sedatives)	0.0052459	0.0029582	1.7733	0.09299
x13 (Marijuana)	-0.018284	0.023034	-0.79379	0.47029
x14 (Father)	0.0024552	0.060876	0.040331	0.97223
x15 (Mother)	-0.63112	0.40142	-1.5722	0.1381
x16_1 (Gender)	0.029065	0.049517	0.58698	0.60023
x1:x8	0.062193	0.028374	2.1919	0.03872
x1:x10	-0.089502	0.028517	-3.1386	0.00345
x1:x11	-0.16196	0.034077	-4.7527	7.33E-06
x3:x8	-0.22405	0.028785	-7.7838	1.69E-13
x3:x11	0.25054	0.027678	9.0521	0
x3:x12	0.0077219	0.0026002	2.9697	0.00562
x3:x15	-1.0664	0.36644	-2.9102	0.00668
x4:x8	0.15457	0.035867	4.3096	5.00E-05
x4:x11	-0.25852	0.037241	-6.9417	4.94E-11
x5:x8	0.33811	0.035023	9.6541	0
x5:x11	-0.13356	0.030837	-4.3312	4.56E-05
x6:x10	0.31323	0.038607	8.1132	1.89E-14
x6:x11	-0.074914	0.034177	-2.1919	0.03872
x7:x8	-0.072229	0.021917	-3.2955	0.00207
x7:x9	-0.044806	0.023309	-1.9222	0.06889
x7:x13	-0.03702	0.018824	-1.9666	0.06283
x8:x11	-0.048082	0.012599	-3.8163	0.00034
x8:x12	0.013621	0.0040864	3.3332	0.00184
x8:x13	-0.12984	0.028524	-4.552	1.76E-05
x8:x15	-3.9026	0.49276	-7.9198	6.66E-14
x8:x16_1	-0.55796	0.08209	-6.7969	1.19E-10
x10:x15	3.2924	0.46957	7.0116	3.19E-11
x11:x12	-0.016408	0.0036726	-4.4678	2.54E-05
x11:x16_1	0.45508	0.067202	6.7718	1.40E-10
x12:x14	0.0076174	0.0023789	3.2021	0.00282
x14:x15	-0.57193	0.25856	-2.212	0.0371
x14:x16_1	0.11931	0.0479	2.4908	0.02036

Marijuana

	Estimate	Standard Error	t-stat	FDR-corrected p-values
(Intercept)	0.12215	0.079536	1.5358	0.14799
x1 (VCC)	-0.033762	0.2373	-0.14228	0.90631
x2 (Age)	-0.22125	0.23661	-0.93511	0.38833
x3 (Total Drinks 7 Days)	0.70924	0.507	1.3989	0.18971
x4 (Drink Frequency 7 Days)	0.4761	0.40985	1.1616	0.28115
x5 (Drinks/Day 12 Months)	0.65732	0.27873	2.3583	0.02716
x6 (Drink Frequency 12 Months)	-0.70412	0.2522	-2.7919	0.00939
x7 (Total Tobacco 7 Days)	1.9197	0.24608	7.8011	1.48E-13
x9 (Cocaine)	3.9492	0.31015	12.733	0
x10 (Hallucinogens)	1.0592	0.33828	3.1312	0.00353
x11 (Opiates)	0.90149	0.28567	3.1556	0.00327
x12 (Sedatives)	0.12251	0.33223	0.36876	0.75176
x13 (Stimulants)	0.56188	0.23658	2.375	0.02624
x14 (Father)	0.33238	0.23653	1.4053	0.18776
x1:x6	0.69869	0.20937	3.3371	0.00181
x1:x12	-0.78946	0.26325	-2.9989	0.00518
x1:x13	0.62535	0.24641	2.5378	0.01823
x2:x12	-0.62702	0.25472	-2.4616	0.02175
x3:x9	-0.55758	0.14206	-3.9249	0.00023
x4:x5	-1.0602	0.27735	-3.8226	0.00033
x5:x6	-0.84237	0.20497	-4.1097	0.00011
x7:x11	-0.37699	0.16595	-2.2717	0.03271
x10:x14	-0.82425	0.26834	-3.0716	0.00421

Supplementary Analysis: Generalized Linear Models without Drink Frequency 7 Days

Given that the variables Total Drinks 7 Days and Drink Frequency 7 Days were highly correlated, we attempted to repeat the GLM analysis but after removing Drink Frequency 7 Days as a covariate. Results are shown in Supplementary Table 12. The most telling difference in results after the removal of Drink Frequency 7 Days was the complete absence of ECS connectivity correlated with Hallucinogen use. Seeing as how Drink Frequency 7 Days was originally a significant covariate in Hallucinogen use (via a two-way interaction with ECS connectivity), its removal may have something to do with the reduced relevance of ECS connectivity with regards to Hallucinogen use. Most other models also revealed reduced r^2 values after the removal of Drink Frequency 7 Days as a covariate (except for VS and VCC models correlated with Hallucinogen use and VCC correlated with Marijuana use which all reported increased r^2 values). The removal of a covariate that formed significant interactions with network connectivity from the modelling process may have resulted in the reduced effect sizes reported. We cannot ascertain why the r^2 values increased in Hallucinogen use and VCC – Marijuana use. Despite these changes, many other interactions between network connectivity and substance use remained the same.

Supplementary Table 12: VS, ECS and VCC connectivity correlations with drug use after removing Drink Frequency 7 Days as a covariate. Effect size denoted by r^2 values. Estimates displayed are standardized partial coefficients for each component of the GLM. *p*-values shown here are FDR-corrected. Exact p-values are displayed where possible. If the p-value is too small (less than 0.00001), then it is displayed as p<0.00001. Father, Fathers' Use History; Mother, Mothers' Use History; E, Estimate

Substance	VS	ECS	VCC
Cocaine	r^2 = 0.69672 VS – Total Tobacco (E=0.084613, t=3.5261, p=0.001) VS – Hallucinogens (E=0.11757, t=4.307, p<0.0001) VS – Opiates (E=-0.20471, t=-6.8544, p<0.0001) VS – Father (E=-0.082774, t=-3.5397, p=0.000956)	r^2 = 0.69673 ECS – Total Tobacco (E=0.076192, t=3.4516, p=0.001268) ECS – Hallucinogens (E=0.16808, t=5.6423, p<0.0001) ECS – Opiates (E=-0.28933, t=-8.0022, p<0.0001) ECS – Father (E=-0.058761, t=-3.1112, p = 0.003766)	r^2 = 0.70341 VCC – Total Tobacco (E=0.07735, t=3.3546, p=0.001727) VCC – Hallucinogens (E=0.14465, t=5.065, p<0.0001) VCC – Opiates (E=-0.25404, t=-7.5381, p<0.0001) VCC – Father (E=-0.71339, t=-3.5465, p=0.000935)
Hallucinogens	r^2 = 0.69347 VS – Total Tobacco (E=-0.065359, t=-2.741, p=0.01042) VS – Tobacco Frequency (E=0.1455, t=5.1389, p<0.0001) VS – Cocaine (E=-0.18597, t=-5.3259, p<0.0001) VS – Opiates (E=0.14446. t=4.2153, p<0.0001) VS – Stimulants (E=0.0076568, t=2.945, p=0.006103)		r^2 = 0.67785 VCC (E=0.045612, t=2.0926, p=0.047819) VCC – Opiates (E=0.11685, t=4.0443, p=0.000147)
Opiates	r^2 = 0.69803 VS – Drinks/Day (E=-0.048524, t=-2.3866, p=0.024784) VS – Total Tobacco (E=0.073661, t=-3.2318, p=0.002594) VS – Tobacco Frequency (E=0.1032, t=4.9648, p<0.0001) VS – Sedatives (E=-0.046284, t=-2.3497, p=0.027118)	r^2 = 0.70546 ECS – Total Tobacco (E=0.080977, t=3.7964, p=0.000377) ECS – Cocaine (E=-0.057769, t=-2.327. p=0.028543) ECS – Hallucinogens (E=0.20452, t=6.2304, p<0.0001) ECS – Sedatives (E=-0.064148, t=-2.6901 p=0.0119)	r^2 = 0.68456 VCC – Drinks/Day (E=-0.051342, t=-2.3967, p=0.024255) VCC – Total Tobacco (E=0.098612, t=4.3799, p<0.0001) VCC – Opiates (E=0.070859, t=2.9319, p=0.006302)
Sedatives	r^2 = 0.77945 VS – Total Drinks (E=-0.053132, t=-2.4184, p=0.023129) VS – Drinks/Day (E=0.066042, t=3.4846, p=0.00146) VS – Cocaine (E=0.098801, t=3.7027, p=0.000536) VS – Hallucinogens (E=-0.083114, t=-3.3086, p=0.002012) VS – Opiates (E=0.057487, t=2.9993, p=0.005223)	r^2 = 0.77961 ECS – Drinks/Day (E=0.054261, t=2.9788, p=0.005541 ECS – Opiates (E=0.13513, t=5.5431, p<0.0001)	r^2 = 0.78814 VCC – Drinks/Day (E=0.062482, t=3.3551, p=0.001729) VCC – Opiates (E=0.10598, t=5.3441, p<0.0001)
Stimulants	r^2 = 0.61496 VS – Drink Frequency (E=-0.050943, t=-2.4309, p=0.022471) VS – Opiates (E=-0.12122, t=-4.7914, p<0.0001) VS – Sedatives (E=0.0062615, t=2.3271, p=0.028588) VS – Father (E=-0.12028, t=-4.3835, p<0.0001)	r^2 = 0.62565 ECS – Opiates (E=-0.25182, t=-8.6554, p<0.0001) ECS – Father (E=-0.05043, t=-2.3425, p=0.027599)	r^2 = 0.61818 VCC – Opiates (E=-0.21521. t=-6.5822, p<0.0001) VCC – Father (E=-0.067227, t=-2.9352, p=0.006266)
Marijuana	r^2 = 0.47088 VS – Tobacco Frequency (E=-0.64873, t=-3.1075, p=0.0038) VS – Stimulants (E=0.69977, t=0.26035, p=0.014926)	r^2 = 0.46724 ECS – Drink Frequency (E=0.79532, t=3.8604, p=0.000296)	r^2 = 0.48047 VCC – Drink Frequency (E=0.71204, t=3.4224, p=0.001389) VCC – Hallucinogens (E=-1.0476, t=-3.9359, p=0.000224) VCC – Stimulants (E=0.66555, t=2.7145, p=0.011201)