1 **Title**
2 The impact of early public health interventions on SARS-CoV-2 transmission and evolution
3

4 **Author list**
5 Sebastian Duchene[1‡], Leo Featherstone[1‡], Birgitte Freiesleben de Blasio[2,3], Edward C.
6 Holmes[4], Jon Bohlin[2] and John H.-O. Pettersson[4,5,*]
7 [‡] These authors contributed equally to this work.
8
Author affiliations
[1]Department of Microbiology and Immunology, The University of Melbourne at The Peter
Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia.
[2]Department of Methods Development and Analytics, Division of Infectious Disease Control
and Environmental Health, Norwegian Institute of Public Health, Oslo, Norway.
[3]Department of Biostatistics, Oslo Centre for Biostatistics and Epidemiology, Institute of
Basic Medical Sciences, University of Oslo, Norway
[4]Marie Bashir Institute for Infectious Diseases and Biosecurity, School of Life and
Environmental Sciences and School of Medical Sciences, the University of Sydney, Sydney,
New South Wales 2006, Australia.
[5]Zoonosis Science Center, Department of Medical Biochemistry and Microbiology, University
of Uppsala, SE-751 23 Uppsala, Sweden.
21
22 **Corresponding author:** John H.-O. Pettersson
23 **Corresponding author email:** john.pettersson@imbim.uu.se

24
25 **Author contributions**
26 SD, ECH, JB and JHOP designed the research; SD, LF, BFDB, ECH, JB and JHOP
27 compiled, computed and analysed the data; SD, LF, ECH and JHOP wrote the manuscript.
28
29 **Competing interest statement**
30 All authors have read the manuscript and have no conflict of interest to declare.
31

Funding statement
JHOP is funded by the Swedish research council FORMAS (grant no: 2015-710) and VR
(grant no: 2020-02593). ECH is funded by an ARC Australian Laureate Fellowship
(FL170100022). SD and LF are funded by an ARC Discovery Early Career Award awarded
to SD (DE190100805). The funders had no role in study design, data collection and
interpretation, or the decision to submit the work for publication.

Abstract
Background
Many countries have attempted to mitigate and control COVID-19 through the
implementation of non-pharmaceutical interventions, particularly with the aim of reducing
population movement and contact. However, it remains unclear how the different control
strategies impacted the local phylodynamics of the causative SARS-CoV-2 virus.
Aim
To assess the duration of chains of virus transmission within individual countries and the
extent to which countries export viruses to their geographic neighbours.
Methods
To address core questions in genomic epidemiology and public health we analysed complete
SARS-CoV-2 genomes to infer the relative frequencies of virus importation and exportation,

as well as virus transmission dynamics, within countries of northern Europe. To this end, we examined virus evolution and phylodynamics in Denmark, Finland, Iceland, Norway and Sweden during the first year of the pandemic.

Results

The Nordic countries differed markedly in the invasiveness of control strategies implemented. In particular, Sweden did not initially employ any strict population movement limitations and experienced markedly different transmission chain dynamics, which were more numerous and tended to have more cases, a set of features that increased with time during the first eight months of 2020.

Conclusion

Together with Denmark, Sweden was also characterised as a net exporter of SARS-CoV-2. Hence, Sweden effectively constituted an epidemiological and evolutionary 'refugia' that enabled the virus to maintain active transmission and spread to other geographic localities. In sum, our analysis reveals the utility of genomic surveillance where active transmission chain monitoring is a key metric.

Key words

SARS-CoV-2, Nordic, transmission chain modelling, virus exportation

Introduction

Since its initial description in December 2019, SARS-CoV-2, the causative agent of COVID19 [1, 2], has rapidly led to a major global health crisis. The pandemic has caused more than 150 million infections and over 3.3 million deaths worldwide and continues to accelerate, imposing a significant impact on health care systems, societies and the global economy. Countries are continuously struggling with how to effectively counteract the pandemic, balancing the protection of health with social and economic considerations. During the first year of the pandemic, in the absence of therapeutics and available therapies and vaccines, efforts centred on so-called 'non-pharmaceutical strategies', particularly initial short-term large-scale restrictions to population movement (e.g. 'lock-downs'), increased testing and various levels of social distancing. Analysis of the local epidemiological consequences of different control strategies provides key information on the most effective approaches to reduce the rate of virus transmission within communities.

The Nordic countries, defined here as Denmark, Finland, Norway, Iceland and Sweden, provide a powerful example of geographically, politically and socially related countries that differ markedly in how control strategies for COVID-19 were implemented during the first six months of the pandemic. In particular, Sweden initially took a less restrictive approach in which no general population movement restrictions were enforced, schools for children below the age of 16 years old remained open, no mandatory quarantine was imposed for infected households, and businesses continued almost as usual [3]. In contrast, Norway and Denmark enforced a more invasive population movement restriction that included enforced home office for workers in the public sector, home-bound schooling, targeted private sector close downs, as well as closed international borders for non-residents. Iceland, a relatively small homogenous island population (of ~350,000) never initiated a population movement restriction as Norway and Denmark, but rather focused on large-scale testing and contact tracing to limit virus spread within the community. In relation to population size, Sweden has had a higher number of COVID19-related cases and deaths than all other Nordic countries [4, 5], with a total of around 9,758/137 cases/deaths per 100,000 people, compared to around 4,465/43 in Denmark, 2,169/14 in Norway, 1,608/17 in Finland, and 1,790/8 in

100   Iceland, as of 12[th] of May 2021 [6], with most cases between April and July 2020 and
101   November 2020 to March 2021.
102
Although the relative 'success' of COVID-19 control measures are normally gauged in the
number of cases and deaths at the country level, it is also the case that intervention and
mitigation strategies may lead to marked differences in transmission dynamics among
populations, which may in turn impact the evolution of the virus. Using a comparative
analysis of genome sequence data we addressed whether the different approaches to
COVID-19 control employed by the Nordic countries resulted in differences in virus
transmission dynamics and the relative frequencies of virus importation/exportation (i.e. virus
"phylodynamics") during the first year of the pandemic following introduction of SARS-CoV-2
in the Nordic region. Accordingly, we compiled an extensive data set of SARS-CoV-2 virus
genomes and performed a phylo-epidemiological study to identify any differences in
transmission chain dynamics between these countries.
114
Materials and Methods
Data set construction
We downloaded all SARS-CoV-2 genomes that were complete and had high sequencing
coverage available the 22[nd] March 2021 at the GISAID platform (www.gisaid.org), for which
no ethical approval was needed. We selected all those from the five Nordic countries as
follows: 50,126 from Denmark, 2,374 from Finland, 4,167 from Iceland, 3,388 from Norway,
and 7,863 from Swede). To obtain a representative subset of the global diversity we also
selected 3,437 genomes from the latest NextStrain global build in 22[nd] March [7].
123
Sequencing intensity is markedly different between countries, with Iceland sequencing
around 68% of positive reported cases whereas in Sweden the corresponding number is
about 1.1%. We attempted to control for such sampling bias by subsampling genomes from
the Nordic countries relative to that with the lowest sampling intensity, meaning that all
countries were sampled a rate of 1.1 genomes per hundred cases. For this purpose, we
divided the number of genomes from each country by the cumulative number of cases
reported in each country by mid-March 2021, as recorded in ourworldindata.org [8]. We
repeated this procedure 10 times, and in each obtained a sequence alignment with all the
data from NextStrain and with the number of genomes from the Nordic countries proportional
to the prevalence of the virus (Table 1). The complete data set consisted of 71,355 genomes,
with 67,918 from the Nordic countries. Our 10 alignments with equal sequencing intensity for
Nordic countries had between 15,616 and 15,297 sequences (i.e. between 12,179 and
11,860 Nordic genomes, plus the 3,467 from the NextStrain build). The GISAID accession
numbers of all sequences included in the final sequence alignment are available as
Supplementary Table S1.
139
Estimating a time-scaled phylogenetic tree
We estimated maximum likelihood phylogenetic trees using IQ-TREE v2.0.6 [9] employing
the GTR+ Γ model of nucleotide substitution [10]. We fit a strict molecular clock to the data
using LSDv0.3 [11]. We calibrated the molecular clock by specifying the sequence sampling
times and fixing the evolutionary rate to 1×10^{-3} subs/site/year as estimated previously [12].
We chose this approach because testing for temporal signal and fitting a sophisticated
molecular clock model is unfeasible for a data set of this size. For a small number of
sequences (113, ~0.14% of the data set) only the month and year were available, such that
we assigned them the 15[th] of the corresponding month.
149

Detection of virus transmission chains
We defined transmission chains as monophyletic groups of genomes (at least 2) all collected
from one of the Nordic countries, which is analogous to transmission lineages as defined
previously [13]. We computed key statistics from each transmission chain, specifically: the
duration (i.e. the length of time in days from the first to the last collected genome), the size
(i.e. the number of genomes), and the time of origin. For each country we also calculated an
H_t-index of transmission chains, where H_t is the number of transmission chains with at least
H_t cases (analogous to the H-index citation metric). Larger values reflect increase in the
number and size of lineages. For example, of Sweden's lineages, 38 had at least 38 cases
(Table 1). Because we subsampled the data 10 times, for these statistics we compute the
range of values, as a crude measure of uncertainty due to sampling.
161
We fit linear regressions for transmission chain size against as a function of the detection
date (the date of collection of the first genome within a transmission chain), such that positive
trends would suggest increased community spread over time (i.e. larger transmission chains
over time). We also considered the duration of transmission chains as a function of detection
date, where a positive trend means that transmission chains detected later in the pandemic
tended to last longer. Because the number of cases had substantially decreased around
August 2020 (Supplementary Figure S1) and because around this time most stringent public
health interventions were in place (Supplementary Figure S2–S5), we fit our regressions
across two periods, before and after 31st August 2020. Importantly, this time also preceded
the emergence of lineage B.1.1.7 and its introduction to the Nordic countries by a few
months (Supplementary Figure S6). This so-called 'variant of concern' has been shown to
have increased transmissibility relative to circulating diversity [14]. We only report trends as
residual distributions were strongly skewed..
175
Inferring virus migration dynamics between Nordic countries
To infer the frequency of importation or exportation events of SARS-CoV-2 between the
Nordic countries and from the rest of the world, we employed a Bayesian stochastic mapping
approach, also known as discrete phylogeography, as implemented in BEAST v1.10.4 [15–
17] using guidelines from Dellicour et al. for very large genomic data sets [18]. We fixed the
time-tree described above and assigned a geographic location for each tip, which could be
either of the five Nordic countries or "other" (for those collected in other countries). This
method is broadly similar to that used to infer geographic movement of the virus in Belgium
[12]. We repeated this procedure for the 10 subsampled replicate trees and report median
posterior values and ranges across replicates. Stochastic mapping generates a posterior
distribution of 'type-changes' between locations along the branches.
187
188 We ran a Markov chain Monte Carlo of length 5×10^7 steps, recording every 5000th step.
189 Sufficient sampling from the posterior was assessed by verifying that the effective sample
190 size of all parameters were above 200 as estimated in Tracer v.1.7 [19]. We inferred the
191 posterior number of migration events between the six possible states (Markov jumps) and the
192 amount of time spent at each state, known as Markov rewards.
193
Assessing public health measures and governmental stringency
To provide an overview of public health measures taken and how they were enforced
between February 2020 through March 2021 per Nordic country, we plotted data from the
Oxford COVID-19 Government Response Tracker [20] which included: testing policy, contact
tracing, public information campaigns, international travel control, workplace closure, school
closure, cancelation of public transportation, stay at home requirements, facemask usage,

gathering restrictions, cancelation of public events, restrictions on movement, and
government stringency response index (Supplementary figures S2–S5).
202
Results
The estimated time of emergence of sampled transmission chains provides information about
the onset of community transmission, although such an association is sensitive to sampling
bias, particularly the time-scale of country-specific sequencing efforts. Our phylogenetic
analyses revealed that SARS-CoV-2 was imported to all Nordic countries between January
to late February 2020, with detectable community transmission from early February in
Sweden and from early March in the other Nordic countries as inferred by the presence of
transmission chains (Figure 1). Sustained community transmission continued for all Nordic
countries beyond April, with the exception of Iceland which was characterised by a relatively
short one-month period of community transmission, and Norway, which had a low number of
cases combined with a low sequencing intensity of 0.039 (the second lowest in our study
after Sweden; Table 1).

Table 1. Summary statistics computed for SARS-CoV-2 data from each Nordic country. The median and range (over 10 replicates) is shown for relevant statistics, with the exception of the time to the most recent common ancestor (TMRCA; the putative time of emergence) of the first transmission chain, for which only the range is reported because the average disproportionally affects countries with low prevalence. For regression statistics, the symbol ~ denotes "function of", and in these cases we report the number of replicates with positive regression slopes before and after 31[st] August 2020.

	Denmark	Finland	Iceland	Norway	Sweden
TMRCA range of first detected transmission chain	2020-01-14– 2020-03-05	2020-01-24– 2020-03-04	2020-02-20– 2020-07-13	2020-01-29– 2020-05-20	2020-01-01– 2020-01-29
Median TMRCA of transmission chains	2020-10-22 (2020-10-20– 2020-10-25)	2020-07-26 (2020-07-17– 2020-08-14)	2020-06-29 (2020-06-06– 2020-09-25)	2020-11-13 (2020-10-28– 2020-11-18)	2020-10-13 (2020-10-10– 2020-10-15)
Median number of genomes included after correcting for prevalence	2380 (2341–2457)	773 (746–799)	70 (56–72)	940 (854–988)	7863 (7863–7863)
Median number transmission chains	227 (197–231)	86 (79–103)	4 (3–5)	105 (94–119)	677 (638–690)
Median duration of transmission chains in days	42 (40–44)	26 (21–28)	45 (19–56)	25 (23–27)	31 (30–32)
Median size of largest transmission chain	222 (99–243)	80 (65–135)	44 (34–56)	134 (129–147)	366 (175–368)
H_t-index	21 (20–22)	11 (9–12)	2 (2–3)	11 (10–12)	38 (36–39)
Median number of exportation events	125 (98–203)	36 (20–60)	0.002 (0.00–0.80)	33 (22–42)	552 (479–704)
Median number of importation events	385 (361–412)	248 (230–271)	18 (14–24)	264 (246–273)	579 (491–628)
Exportations-to-importations ratio	0.34 (0.24–0.56)	0.14 (0.0813–0.26)	0.00 (0.00–0.057)	0.13 (0.083–0.17)	0.95 (0.76–1.45)
Number of genomes collected	50126	2374	4167	3388	7863
Genome collection date range	2020-03-02– 2021-02-21	2020-01-29– 2021-02-06	2020-03-02– 2021-01-05	2020-01-29– 2021-03-08	2020-01-31– 2021-03-09
Sequencing intensity	0.22	0.033	0.68	0.039	0.011
Number of cases reported at sampling collection date	226777	72073	6119	87519	744272
Transmission chain ~ detection date (# replicates out of 10 with positive slope; before and after early November 2020)	4, 0	8, 1	NA, NA	6, 1	2, 0
Transmission chain duration ~ detection date (# replicates out of 10 with positive slope; before and after early November 2020)	3, 0	8, 0	NA, NA	3, 1	8, 1

The number of transmission chains observed was around 227 (range: 197–231) for Denmark, 86 (range: 79–103) for Finland, in Iceland (45, range: 19–56), and 105 (range: 94–119) for Norway. Notably, the highest number of transmission chains - 677 (range: 638–690) - was observed in Sweden. The median duration of such transmission chains (number of days from detection to last sampled case) was higher in Iceland (45, range: 19–56) and Denmark (42, range: 40–44) than other localities: 31 (range: 30–32) for Sweden, 26 (range: 21–28) for Finland, and 25 (range: 23–27) for Norway (Table 1).

The largest transmission chain was found in Sweden, with 366 cases (range: 175–368), followed by Denmark with 228 (range: 99–243), Norway with 134 (range: 129–147), Finland with 80 (range: 65–135), and Iceland with 44 (range: 34–56). In addition to having the largest transmission chain, Sweden also had the highest H_t-index (Table 1). The H_t-index for Sweden was 38 (range: 36–39), 21 for Denmark (range: 20–22), 11 for Finland (range: 9–12), 2 for Iceland (range: 2–3), and 11 for Norway (range: 10–12), suggesting that Sweden had the least-interrupted community transmission.

There was inconclusive evidence of an increase in transmission chain size in the first period (prior to the 31st August 2020), with between 2 and 8 of 10 subsampling replicates displaying a positive regression slope. Due to low sample size and number of data points in the second period we did not perform a trend analysis for Iceland. After 31st August 2020 most replicates (9 or more) exhibited a negative trend for all countries, indicating a decrease in transmission chain size over time. Similarly, we did not find consistent trends for the duration of transmission chains in the first period, whereas in the second period they appeared to decline over time, with at most one replicate displaying a positive slope in Denmark, Finland, Norway and Sweden (Table 1; Figure 1). Overall, these results are consistent with stringent public health interventions in the second period examined here that may have limited the spread of existing transmission chains. Thus, the increased number of cases observed in the second half of 2020 was most likely driven by importations that led to limited or ceased transmission, rather than an increasing size and duration of transmission chains. This is consistent with the majority of importation events leading to transmission chains or singletons coming in the second half of 2020 for most countries (Supplementary Figure S7).

Figure 1. Transmission chain summaries. Open circles mark transmission chain size (left) and chain duration (right) against the time of the first sample from each chain for each country. Trend lines are fitted here to one of ten replicates with dotted lines corresponding to trends that were not repeated in at least 9 of 10 replicates. Trends are omitted for Iceland due to low sample size. Trends are split on either side of 31st August 2020.

The total number of exportation events was larger for Sweden (552, range: 479–704) than for Denmark (125, range: 98–203). Most exportation events from Sweden were into Finland, whereas those from Denmark were into non-Nordic countries, followed by Sweden and Iceland (Figure 2, Panel B). An inspection of the amount of time in the tree occupied by each country, known as the Markov rewards, revealed that Sweden occupied the largest portion, consistent with it being the major exporter. In contrast, Norway, had very low Markov rewards, despite having more genomes than Iceland and Finland, both with higher Markov rewards (Supplementary Figure S8).

Figure 2. Circular migration diagram of migration events between the five Nordic countries and other locations. Panel A shows migration events with 'Other' countries included and Panel B shows migration between Nordic countries only. The size of the coloured links denotes the posterior median number of inferred migration events from the Bayesian phylogeographic analysis across 10 replicates. The black arrows represent the migration direction.

Discussion

Until March 2021, Sweden experienced a greater number of COVID-19-related cases and deaths in relation to population size than all other Nordic countries [6]. Our genomic analysis shows that whilst SARS-CoV-2 was imported to all Nordic countries during the same time-period in early 2020, Sweden experienced more transmission chains which also tended to be larger. Importantly, these trends emerged despite Sweden having the lowest virus genome sampling proportion, suggesting that community transmission was well-established during both waves relative to the other Nordic countries. Reassuringly, all Nordic countries displayed a decrease in the size of transmission chains and their duration in the second time period examined here.

Denmark and Sweden were characterised by having acted as exporters of SARS-CoV-2 to their Nordic neighbours. Conversely, Iceland, Norway and in particular Finland showed lower levels of virus exportation. At the same time, our analyses indicate that Finland was the main receiver of the exportations from Sweden, suggesting a near unidirectional mode of exportation between the two countries.

During the first six months of the pandemic Sweden never enforced strict community mobility restrictions. Instead, social distancing, limiting sizes of social events and moving to distance education for students above 16 years were employed as the main mitigation efforts. Conversely, Denmark, Finland and Norway enforced a relatively strict population movement restrictions of their communities, closure of borders and government run facilities [4]. These mitigation efforts were taken during both the first and second wave (Supplementary table S2, Supplementary figures S2–S5). During the second wave, which started in early September 2020, Sweden recommended limiting the number of people that meet within social circles. In

early 2021, Sweden also recommended the use of protective face-masks in public spaces. In
Norway, locally-based contact tracing, testing, isolation, and quarantining, supported by a
national contact tracing team, has been the primary strategy since the summer of 2020. In
contrast, the danish strategy focuses on extensive testing, free-of-charge with readily
accessible testing units (Supplementary figures S2). Both countries enforce local contact-
reducing measures during periods of outbreaks. Iceland, in a similar manner to South Korea,
utilised large-scale testing and contact tracing combined with social distancing and voluntary
home-based quarantine without the need to regulate population movement [21] (see also
Supplementary table S2). Thus, while regulating population movement appears to provide an
efficient mode to reduce community transmission [22, 23], other mitigation efforts are also
viable options under certain circumstances. Norway experienced relatively low numbers of
transmission chains during the first six months of the pandemic, with the bulk of the
transmission chains sampled here emerging in mid-April and towards the end of July.
Although this might in part reflect a delayed start to genome sequencing, which have
increased significantly in all countries since then, and such results should be interpreted with
caution (Figure 1). Indeed, we note that the sequencing intensity of Norwegian cases is lower
than some of the other Nordic countries with similar case burdens (Table 1). Thus,
sequencing proportion bias per country needs to be taken into account when considering our
results. To reduce effect of sequencing bias, we attempted to subsample our data to obtain
even sequencing proportions although this does not account for other sources of bias, such
as if close contacts of a positive case are sequenced or not. When sample size per country
increases with time, the effect of such potential sampling biases are expected to decrease.
365
To what extent then did Sweden's mitigation strategy affect the epidemiological situation
internally and across Nordic region as a whole? Sweden received substantial attention due to
the number of deaths reported. Sweden has been shown to have had one of the longest
durations of excess deaths and highest case numbers between February and May 2020 [5]
and it has been suggested that a greater level of self-isolation likely would have reduced the
number of deaths in Sweden [24].
372
The effect of closing/opening schools are also likely to have contributed to the transmission
dynamics [25, 26]. In the case of the Nordic countries, it can be argued that by closing pre-,
primary, secondary schools and higher education for several weeks during the spring of 2020
(Supplementary table S2, Supplementary figure S3), Denmark and Norway also reduced
contact and virus transmission in the adult population to a higher extent than Sweden where
schools remained open during the same period. However, although Sweden exhibited a
relatively high number of cases compared to the other Nordic countries, and our results are
consistent with a relatively high community transmission for Sweden during the first twelve
months, such an increase in community transmission does not necessarily imply a causal
relationship with excess deaths. Whilst we employ a robust genomic-based modelling
approach to study transmission chain changes, equating or directly relating such changes
with the number of deaths may lead to incorrect conclusions as we have not taken into
account any patient-related epidemiological parameters such as gender or age. Furthermore,
it has been argued that the higher number of deaths in Sweden may also have been
influenced by how intensive care unit admission criteria were applied, particularly for the
elderly population in care homes, during the early phase of the pandemic [24]. Thus, official
mitigation strategies and efforts alone are unlikely to explain epidemiological differences
between countries, which were also impacted by the compliance and practical enforcement
from the general public, businesses, and health care providers.
392

A critical issue, although one that has received little attention, is that if transmission chains
are allowed to remain active, they also provide increased opportunity for the virus to evolve
and adapt to local populations, potentially acquiring mutations that in some way enhance
virus fitness. Indeed, several so-called 'variants of concern' have independently emerged
globally with mutations that are associated with epidemiologically important properties. For
example, the B.1.1.7 variant has acquired mutations that has significantly increased the
transmission potential of the virus [14] and the B.1.351 variant has shown increased
resistance against vaccines [27]. Clearly, mutations routinely appear in the SARS-CoV-2
genome, including viruses sampled from Sweden [28], and it is important to continuously
evaluate their functional relevance. In addition to providing increased evolutionary potential,
sustained transmission chains also provide epidemiological 'refugia' for the virus to be
transmitted to other localities. This is in line with our results showing that Sweden had a
greater frequency of exportation events among neighbouring Nordic countries. As a
consequence, interrupting and ultimately stopping transmission chains is not only important
to minimise virus spread within populations, but also to reduce the chances for the virus to
accumulate beneficial mutations. Fortunately, COVID-19 vaccination has started in all Nordic
countries around the New Year of 2020/2021, which will reduce transmission chains and
protect vulnerable patient groups. However, genome sequencing will continue to play an
important role to monitor for potential vaccine escape mutants.
412
413 Overall, our study highlights the utility of continuous genomic surveillance and retrospective
414 studies to compare and understand differences in pandemic responses with respect to
415 transmission dynamics. In particular, the data presented suggests that transmission chain
416 monitoring may prove to be a useful metric in comparing outbreak mitigation outcome.
417

Acknowledgements
We thank all people who have contributed SARS-CoV-2 genome data to the GISAID
platform. Details on specific contributions are available via www.gisaid.org/. JHOP is funded
by the Swedish research council FORMAS (grant no: 2015-710) and VR (grant no: 2020-
02593). ECH is funded by an ARC Australian Laureate Fellowship (FL170100022). SD and
LF are funded by an ARC Discovery Early Career Award awarded to SD (DE190100805).
424

425 **References**
426
1. Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus associated
with human respiratory disease in China. Nature. 2020;579:265–9.
2. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and
epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding.
Lancet. 2020;395:565–74.
3. Ludvigsson JF. The first eight months of Sweden's COVID-19 strategy and the key actions
and actors that were involved. Acta Paediatr. 2020.
4. Conyon MJ, He L, Thomsen S. Lockdowns and COVID-19 Deaths in Scandinavia. SSRN
Electronic Journal. 2020. doi:10.2139/ssrn.3616969.
5. Kontis V, Bennett JE, Rashid T, Parks RM, Pearson-Stuttard J, Guillot M, et al. Magnitude,
demographics and dynamics of the effect of the first wave of the COVID-19 pandemic on all-
cause mortality in 21 industrialized countries. Nat Med. 2020.
6. WHO. WHO Coronavirus Disease (COVID-19) Dashboard. https://covid19.who.int.
Accessed 12 May 2021.
7. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-
time tracking of pathogen evolution. Bioinformatics. 2018;34:4121–3.
8. Ritchie H, Ortiz-Ospina E, Beltekian D, Mathieu E, Hasell J, Macdonald B, et al.
Coronavirus Pandemic (COVID-19). Our World in Data. 2020.

'https://ourworldindata.org/coronavirus. Accessed 21 Mar 2020.

9. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al.
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic
Era. Mol Biol Evol. 2020;37:1530–4.

10. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol.
2016;2:vew007.

11. To T-H, Jung M, Lycett S, Gascuel O. Fast Dating Using Least-Squares Criteria and
Algorithms. Syst Biol. 2016;65:82–97.

12. Duchene S, Featherstone L, Haritopoulou-Sinanidou M, Rambaut A, Lemey P, Baele G.
Temporal signal and the phylodynamic threshold of SARS-CoV-2. Virus Evol. 2020.
doi:10.1093/ve/veaa061.

13. du Plessis L, McCrone JT, Zarebski AE, Hill V, Ruis C, Gutierrez B, et al. Establishment
& lineage dynamics of the SARS-CoV-2 epidemic in the UK. preprint. Epidemiology; 2020.
doi:10.1101/2020.10.23.20218446.

14. Volz E, Mishra S, Chand M, Barrett JC, Johnson R, Geidelberg L, et al. Assessing
transmissibility of SARS-CoV-2 lineage B.1.1.7 in England. Nature. 2021;593:266–9.

15. Minin VN, Suchard MA. Counting labeled transitions in continuous-time Markov models
of evolution. J Math Biol. 2008;56:391–412.

16. Minin VN, Suchard MA. Fast, accurate and simulation-free stochastic mapping. Philos
Trans R Soc Lond B Biol Sci. 2008;363:3985–95.

17. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian
phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol.
2018;4:vey016.

18. Dellicour S, Durkin K, Hong SL, Vanmechelen B, Martí-Carreras J, Gill MS, et al. A
phylodynamic workflow to rapidly gain insights into the dispersal history and dynamics of
SARS-CoV-2 lineages. preprint. Evolutionary Biology; 2020. doi:10.1101/2020.05.05.078758.

19. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in
Bayesian Phylogenetics Using Tracer 1.7. Syst Biol. 2018;67:901–4.

20. Hale T, Angrist N, Goldszmidt R, Kira B, Petherick A, Phillips T, et al. A global panel
database of pandemic policies (Oxford COVID-19 Government Response Tracker). Nat Hum
Behav. 2021;5:529–38.

21. Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et
al. Spread of SARS-CoV-2 in the Icelandic Population. N Engl J Med. 2020;382:2302–15.

22. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, et al. Estimating the
effects of non-pharmaceutical interventions on COVID-19 in Europe. Nature. 2020;584:257–
61.

23. Geoghegan JL, Ren X, Storey M, Hadfield J, Jelley L, Jefferies S, et al. Genomic
epidemiology reveals transmission patterns and dynamics of SARS-CoV-2 in Aotearoa New
Zealand. Nat Commun. 2020;11:6351.

24. Kamerlin SCL, Kasson PM. Managing COVID-19 spread with voluntary public-health
measures: Sweden as a case study for pandemic control. Clin Infect Dis. 2020.

25. Buja A, Paganini M, Cristofori V, Baldovin T, Fusinato R, Boccuzzo G, et al. Opening
schools and trends in SARS-CoV-2 transmission in European countries. preprint. Public and
Global Health; 2021. doi:10.1101/2021.02.26.21252504.

26. Stage HB, Shingleton J, Ghosh S, Scarabel F, Pellis L, Finnie T. Shut and re-open: the
role of schools in the spread of COVID-19 in Europe. preprint. Epidemiology; 2020.
doi:10.1101/2020.06.24.20139634.

27. Madhi SA, Baillie V, Cutland CL, Voysey M, Koen AL, Fairlie L, et al. Efficacy of the
ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. N Engl J Med. 2021.

28. Ling J, Hickman RA, Li J, Lu X, Lindahl JF, Lundkvist Å, et al. Spatio-Temporal
Mutational Profile Appearances of Swedish SARS-CoV-2 during the Early Pandemic.
Viruses. 2020;12.

Supplementary materials for this manuscript including the following:

Supplementary Figure S1. Number of SARS-CoV-2 genomes sampled with time for all
Nordic countries and global samples (i.e. 'other') included in the final alignment.

504
Supplementary Figure S2. Information on indicators of government response between
February 2020 through February 2021 by country regarding (i) testing policy, (ii) contact
tracing, (iii) public information campaigns, and (iv) international travel control. All indicators
are represented using ordinal scales, showing the highest value in the countries at each time
point. If local measures are used, these values may apply to a limited geographic area.
Testing policy: 0: no; 1: symptomatic and certain criteria; 2: anyone with symptoms; 3: open
public testing; Contact tracing: 0: no; 1: limited; 2: comprehensive; Public information
campaigns: 0: no; 1: public officials cautions; 2: coordinated campaigns; International travel
control (foreign travellers exclusively): 0: no; 1: screening arrival; 2: quarantine arrival; 3 ban
arrivals some regions; 4: total border closure.

Supplementary Figure S3. Information on indicators of government response between February 2020 through February 2021 by country regarding (i) workplace closure, (ii) school closure, (iii) cancelation of public transportation, and (iv) stay at home requirements. All indicators are represented using ordinal scales, showing the highest value in the countries at each time point. If local measures are used, these values may apply to a limited geographic area. Workplace closure: 0: no; 1: recommend work from home; 2: require closing for some sectors; 3: require closing for all-but-essential sectors; School closure: 0: no; 1: recommend closing with alteration; 2: require closing some levels; 3: require closing all levels; Cancel public transport: 0: no; 1: recommend closing; 2: require closing; Stay at home requirements: 0: no; 1: recommended not leaving house.

Supplementary Figure S4. Information on indicators of government response between
February 2020 through February 2021 by country regarding (i) facemasks, (ii) gathering
restrictions, (iii) cancelation of public events, and (iv) restrictions on movement. All indicators
are represented using ordinal scales, showing the highest value in the countries at each time
point. If local measures are used, these values may apply to a limited geographic area. Face
masks: 0: no; 1: recommend; 2: required in specified public spaces; 3: required in all public
spaces when social distancing not possible; Restrictions on gatherings: 0: no; 1: >1000; 2:
>100; 3: >10; 4: 10 or less; Cancel public events: 0: no; 1: recommended; 2: required
closing; Restrictions on movements: 0: no; 1: recommended no travel between regions; 2:
internal movement restrictions.

Supplementary Figure S5. Stringency index of government responses between February

2020 through March 2021 by country. The index is an average of 9 indicators and adjusted

for geographic scope. The following indicators are included: school closure, workplace

closure, cancel public events; restrictions on gatherings; close public transport, stay at home

requirements, restrictions on internal movements, international travel controls, public

information campaigns.

Supplementary Figure S6. Log number of genomes in each of the five most common pango

lineages and B1.1.7 over time in the complete dataset.

Supplementary Figure S7. Number of putative introductions identified by detecting
transmission chains (monophyletic groups of at least 2 genomes from a Nordic country) or
singletons (genomes from a Nordic country sitting outside a transmission chains). Grey bars
correspond to the total number of introductions per fortnight. Orange bars are for singletons
and those in blue are for transmission chains. Here we included all importation events
detected across 10 phylogenetic trees, where each tree contains Nordic genomes sampled
relative to prevalence, such that sequencing intensity is equivalent.

Supplementary Figure S8. Jittered markov rewards for each Nordic country and for the

global diversity in the data (labelled as "other"). The Markov rewards correspond to the

portion of time that all lineages spend at either of the six states in the stochastic mapping.

The y-axis is the Markov rewards as a proportion of total tree length. Note that countries with

most exportation events, Sweden and Denmark are also occupy the largest portion of the

tree.

Supplementary Table S1. Zip file containing a table including all GIASID accession
numbers and replicate information for all sequences SARS-CoV-2 used in the analyses.
565
566 **Supplementary Table S2.** Summary table of mitigation responses taken by the Nordic
567 countries until 2021-03-25 according to the European Centre for Disease Control.