COVID-19 patients and cancer in northern Italy

Lucia Mangone[1], Francesco Gioia[1], Pamela Mancuso[1], Isabella Bisceglia[1], Marta Ottone[1], Massimo Vicentini[1], Carmine Pinto[2], Paolo Giorgi Rossi[1]

[1] Epidemiology Unit, Azienda USL-IRCCS di Reggio Emilia, Via Amendola 2, 42122 Reggio Emilia, Italy.

[2] Medical Oncology, Clinical Cancer Centre, Azienda USL-IRCCS di Reggio Emilia, Viale Risorgimento 80, 42123 Reggio Emilia, Italy

Corresponding author:

Isabella Bisceglia

Epidemiology Unit, Azienda USL-IRCCS di Reggio Emilia, Via Amendola 2, 42122 Reggio Emilia, Italy.

e-mail:isabella.bisceglia@ausl.re.it

Keywords

Cancer prevalence; COVID-19; risk factors; prognostic factors

List of abbreviations

CI= Confidence Interval
COVID-19= COronaVIrus Disease 19
CS= Cancer Survivors
ICU= Intensive Care Unit
IRR= Incidence Rate Ratios
OR= Odds Ratio
RE-CR=Reggio Emilia Cancer Registry
RT-PCR= Real Time Polymerase Chain Reaction
SARS-CoV-2= Severe Acute Respiratory Syndrome COronaVirus 2
UK= United Kingdom
WHO= World Health Organization

Article category: Research Article

Abstract
The aim of this population-based work was to evaluate the impact of having had cancer on COVID-19 risk and prognosis during the first wave of the pandemic (27 February – 13 May 2020) in Reggio Emilia Province. Prevalent cancer cases diagnosed between 1996 and 2020 were linked with the provincial COVID-19 surveillance system. We compared cancer survivors' cumulative incidence of being tested, testing positive for SARS-CoV-2, and dying of COVID-19 with that of the general population; among COVID-19 patients, we compared cancer survivors' risk of dying with that of other patients.

During the study period 15,391 people (1,525 cancer survivors - CS) underwent RT-PCR for SARS-CoV-2, of whom 4,541 (449 CS) tested positive; 549 (114 CS) died of COVID-19. The probability of undergoing testing was 29.5% in the general population as well in CS, while the cumulative incidence of being tested, testing positive, and COVID-19 death were lower in CS: age- and sex-adjusted Incidence Rate Ratios were 0.69 [95%CI 0.65-0.73], 0.55 [95%CI 0.50-0.61], and 0.52 [95%CI 0.42-0.64], respectively. Cancer survivors had worse prognosis when diagnosed with COVID-19, particularly those below the age of 70 (odds ratio (OR) of death 4.91 [95%CI 2.38-10.09]), while the OR decreased after age 80 (1.20 [95%CI 0.87-1.65]). The OR was higher for patients with a recent diagnosis (<2 years OR=2.99 [95%CI 1.52-5.89]) or metastases (OR=2.14 [95%CI 0.89-5.16]).

Cancer patients may have adopted behaviors that protected them from infection, but they were still at higher risk of death once infected.

Novelty and impact:

Cancer survivors during the first wave of the pandemic showed lower COVID-19 cumulative incidence and mortality. When infected, they had worse prognosis, particularly in people younger than age 70, with a recent diagnosis, or with metastases.

Introduction

The novel coronavirus (SARS-Cov-2) has become a worldwide threat, and the related disease, COVID-19, is the major healthcare concern of the year 2020. After China, Italy was one of the first countries that experienced a tragic increase in the incidence of and mortality for COVID-19, with more than 232,000 cases and over 33,000 deaths by the end of May 2020.[1]

The first Chinese study, on a very small group of 18 patients, showed that cancer patients presented a higher risk of SARS-CoV-2 infection and a higher risk of requiring mechanical ventilation or ICU (Intensive Care Unit) admission compared to the general population.[2]

Further studies have found that COVID-19 patients with cancer tend to have much more severe symptoms and a nearly 3-fold higher death rate than COVID-19 patients without cancer.[3] Especially patients with hematologic, lung, and metastatic cancers demonstrated higher rates of severe events compared with patients without cancer.[3] In addition, patients who underwent cancer surgery and contracted SARS-CoV-2 showed higher death rates and higher chances of having critical symptoms.[4,5] Thus, several authors have suggested that immunosuppressed status (whether caused by the disease itself or the treatment) of some cancer patients increases their risk of SARS-CoV-2 infection and of worse outcomes compared with the general population.[6,7]

With more than 18 million new cases per year globally, cancer affects a significant portion of the population. In Italy, over 377,000 new cases of cancer are estimated to occur in 2020, with an estimate of about 3.6 million cancer survivors; this prevalence is estimated to increase dramatically in the next few years.[8]

The aim of this work was to evaluate the impact of having had cancer on COVID-19 risk and prognosis in an Italian province with a high incidence rate of tumors and which also saw a high cumulative incidence of COVID-19 between late February and early May 2020. To reach this aim we compared the risk of undergoing a SARS-CoV-19 test, testing positive, being hospitalized, and dying of COVID-19 of people diagnosed with cancer in the previous 25 years with the same risks of the general population. We also compared cancer survivors' risk of dying of COVID-19 with that of other COVID-19 patients.

Methods

Study design

This is a population-based cohort study using registry data of the Reggio Emilia Province. The study was approved by the Area Vasta Emilia Nord Ethics Committee on 7 April 2020, protocol n.2020/0045199.

Setting

Reggio Emilia is a province in northern Italy with about 532,000 inhabitants. The Local Health Authority, the local public entity of the Italian National Health Service, provides hospital, outpatient, primary, and preventive care to the entire population residing in the province.

With the spread of the SARS-CoV-2 virus in northern Italy, the following measures were adopted in the Province of Reggio Emilia: on February 22, schools were closed and social restrictions were imposed; on 8 March, mobility restrictions were imposed, and on 11 March, only essential services were permitted to remain open. During this phase, all cases with suspicious symptoms (fever, cough, dyspnea) were tested. The first case of SARS-CoV-2 disease (COVID-19) in the province was diagnosed on February 27, 2020.

The period of recruitment of patients involved in the study was from 22 February to 13 May 2020. Up to May 13, there were more than 4500 confirmed cases in the province; the epidemic was still spreading, but at a slower rate, and cumulative incidence reached about 9 per 1000.

Participants

This study analyzed the variables of the residents in the Reggio Emilia Province who underwent the molecular test for COVID-19 between 27 February and 13 May 2020.

Then, information on COVID-19-positive and cancer patients was linked. A COVID-19 patient was diagnosed based on real-time reverse transcription polymerase chain reaction (RT-PCR) of nasopharyngeal specimens according to WHO indications; a cancer patient, excluding non-melanoma skin cancers, was defined if that patient was in the Reggio Emilia Cancer Registry (RE-CR) archive.

Data sources

The Reggio Emilia Cancer Registry registers all new cancer diagnoses occurring in people residing in the Reggio Emilia Province. The main information sources of the RE-CR are the anatomic pathology reports, the hospital discharge records, and mortality data. The RE-CR, which covers all the resident population in the Province of Reggio Emilia, has been active since 2000 and has registered all incident cases from 1996 to 2019, with active follow up for deaths and residence of all prevalent cases updated to 31/12/2019. It collects information on site, morphology, partial staging (presence of metastases), mode of diagnosis, and survival.[9]

All RT-PCR SARS-CoV-2 tests performed in Italy must be recorded in the national case-based integrated COVID-19 surveillance system.[10] This surveillance system contains data on all COVID-19 patients, collected by the Public Health Department of the Local Health Authority during epidemiologic investigations, through: 1) daily reports from COVID-19 labs for all positive RT-PCR tests; 2) an initial epidemiologic investigation conducted through phone interviews with all cases, followed by daily phone calls to patients cared for in outpatient settings, both carried out by

the Public Health Department of each Local Health Authority; 3) daily reports extracted from electronic medical records for hospitalized patients; 4) check of death records to assess mortality, particularly in outpatient settings.

The whole list of cases tested for COVID-19 (positive and negative) was linked with the entire archive of prevalent cases of malignant cancer on 31 December 2019 present in the RE-CR database.

Outcome measures

From the COVID-19 surveillance system data, we report the cumulative incidence of: a) being tested for SARS-CoV-2; b) testing positive for SARS-CoV-2 (for the COVID-19 definition adopted by the Italian Ministry of Health, these are all COVID-19 cases); c) people hospitalized while SARS-CoV-2 positive; d) people dying while SARS-CoV-2 positive. Follow up was closed on 13 May 2020.

Based on the data collected by the Reggio Emilia Cancer Registry, all the tested subjects were grouped according to their oncological history: patients who had never been diagnosed with a malignancy *versus* patients with a clinical history of malignancy at any time since 1996. Cancer patients were then categorized according to the time since cancer diagnosis (< 2 years, 2-5 years, and > 5 years), cancer site, and presence of metastases at diagnosis.

Statistical analysis

Pearson's chi-square test was used to examine differences in the proportions of subjects with and without cancer and tested/not tested for SARS-CoV-2. We report age- and sex-adjusted incidence rate ratios, with relative 95% confidence intervals (95% CI) using Poisson regression, for cumulative incidences. The outcomes of interest for this analysis were: cancer survivors' being tested for SARS-CoV-2, having a positive test, being hospitalized, and dying of COVID-19, compared with the same outcomes of those who had never had a cancer diagnosis. Multivariate analysis was performed using a logistic regression model to measure the odd ratios, with relative 95% CI, of hospitalization and death for COVID-19 patients with cancer, adjusting for age and sex. STATA v.14.1 (StataCorp LP 4905 Lakeway Drive, Texas 77845 USA) was used for all analyses.

Results

On 31 December 2019, there were 43,716 residents of the Province of Reggio Emilia still alive after a cancer diagnosis (excluding non-melanoma skin cancers), for a prevalence of 8.9%. Between 27 February and 13 May 2020, 15,391 residents in the Province of Reggio Emilia underwent molecular testing for SARS-CoV-2 (2.9% of the resident population); testing was more frequent among the elderly and among females; 1,525 cancer survivors underwent testing (3.5% of total

cancer survivors) (Table 1). The age- and sex-adjusted cumulative incidence ratio was 0.69 (95% CI = 0.65, 0.73) (Table 2).

Of the 15,391 tests performed, 4,541 (29.5%) were positive for COVID-19, with no differences between those with a cancer diagnosis (29.4%) and those without (29.5%) (Table 3). The cumulative incidence of SARS-CoV-2-positive tests was 0.8% in the general population and 1.0% in people with cancer (Table 1). The age- and sex-adjusted incidence ratio was 0.55 (95% CI = 0.50, 0.61) (Table 2).

The overall cumulative incidence of hospitalizations was 0.28%, higher in males and in older people; in cancer patients it was 0.5% (236 hospitalized cases) (Table 1), corresponding to an age- and sex-adjusted hospitalization rate ratio of 0.59 (95% CI = 0.51, 0.68) (Table 2).

The overall cumulative mortality was 0.1% in the general population, reaching 1.1% in people over age 80. Crude cumulative mortality was higher in people with cancer (114 deaths, 0.26%), while the age- and sex-standardized mortality ratio was 0.52 (95% CI = 0.42, 0.64) (Table 2).

Table 3 and Figure 1 show the differences between COVID-19 patients with and without cancer. There were slight differences by sex (Fig 1A), while in terms of age, cancer patients were more frequently both between 70-79 years (21.6%) and in the age group > 79 years (46.1%) (Fig.1B). Patients with cancer underwent less self-isolation (47.4% vs 69.8%) but were more often hospitalized (52.6% vs 30.2%) and died (25.4% vs 10.6%) (Fig. 1C). Among hospitalized patients and deaths for COVID-19, there were no differences by sex, but patients with cancer were much older (Fig. 1D and 1E).

The multivariate analysis (Table 4) showed that among COVID-19 patients, females had a lower risk of being hospitalized and of dying (OR 0.61, 95% CI = 0.56, 0.66 and 0.62, 95% CI = 0.54, 0.71, respectively). People ages 70-79 had the highest probability of being hospitalized, while those < age 70 had the lowest. The probability of death increased with age, with ORs for those below age 70 compared to those over 79 of 0.06 (95% CI = 0.04, 0.06) and for age 70-79 vs over 79 of 0.49 (95% CI = 0.41, 0.59). We observed a strong interaction between age and cancer in the effect on fatality rate and, to a lesser degree, also for hospitalization. The OR of being hospitalized for COVID-19 among those below age 70 with a previous diagnosis of cancer was 2.52 (95% CI = 1.76, 3.62) and among those over 79, 1.51 (95% CI = 1.10, 2.07). The OR of dying was 4.91 (95% CI = 2.38, 10.09) for patients below age 70 and 1.20 (95% CI = 0.87, 1.65) for patients over age 79.

Excess mortality in cancer patients was higher in the presence of metastases, but differences could be due to chance (OR 2.14, 95% CI = 0.89, 5.16) (Table 5). Concerning timing, the probability of death was higher for all patients with cancer, but the strongest excess was for cancers diagnosed less than two years before the onset of COVID-19 symptoms (OR 2.99; 95% CI = 1.52, 5.89). The

patterns were similar for hospitalization: among the 449 SARS-CoV-2-positive patients with cancer, the sites most frequently affected were breast (101 cases), digestive organs (86 cases, of which 48 colon, 20 rectum, 13 stomach, 4 pancreas, 1 biliary tract), urinary tract (57 of which: 41 bladder, 14 kidney, 2 upper urinary tract), and male genital organs (56 cases, of which 52 prostate, 3 testis, and 1 penis). COVID-19 patients with lymphomas, other hematological organs, respiratory organs, female genital organs, or urinary tract showed the greatest excess of hospitalizations; other hematological organs, melanoma, and female genital organs showed the strongest excess mortality.

Discussion

SARS-CoV-2 infections and COVID-19-related hospitalizations and deaths were more frequent among cancer patients than in the general population. However, when considering the different age structure of cancer survivors, the adjusted cumulative incidence was about 45% lower in people with cancer, as was hospitalization and mortality. Differences in cumulative incidence were not likely due to differences in testing, since the cumulative probability of being tested was only 30% lower for people with cancer and the probability of testing positive was almost identical in the two populations. This suggests that the criteria for referring to testing identified targets with similar probabilities of disease, regardless of the presence of a previous diagnosis of cancer.

Once having COVID-19, the probability of being hospitalized and of dying was much higher for people with a previous diagnosis of cancer that were below the age of 70 at the time of infection; excess mortality was appreciable only up to the age of 80, while after 80 the excess was largely compatible with random fluctuation. For hospitalization, the excess decreased, though it remained appreciable in all age groups.

Higher fatality and hospitalization rates for COVID-19 patients with a previous diagnosis of cancer have been observed in several studies[11-17], including a recent large registry-based Italian study.[11] Instead, we did not find any studies comparing the risk of SARS-CoV-2 infection and COVID-19 death in cancer survivors with that in the general population. Only a few case series tried to infer the risk of infection, comparing the prevalence of SARS-CoV-2 infection in cancer patients with that in health care providers[12] or with that in the general population,[13-15] or comparing the prevalence of cancer among COVID-19 patients with that in the general population.[16] These studies gave inconsistent results.[12,17]

The observed lower probability of being tested and of testing positive could be related to protective behaviours adopted early by people with previous cancer, i.e. social distancing and wearing masks. It must be taken into consideration that the Italian Government implemented special protection measures for people with chronic diseases, including cancer, such as exemption from any work not allowing social distancing.

While age is a powerful risk factor for death, we observed fewer hospitalizations for the age group ≥ 80 than in the age group 70-80. The observed excess in mortality among the elderly is consistent with all previous studies both in the general population[7,18,19] and in cancer patients.[20] Hospitalization rates may have been influenced by the availability of beds during the epidemic peak and also by the context; as many patients over age 80 were already assisted in nursing homes, there was no need for hospitalization in some cases.

The effect of sex is also consistent with all previous studies, showing more severe disease in males.[20]

Not surprisingly, the proportion of people with a previous cancer diagnosis among COVID-19 patients (about 9%, in our province) was much higher than that reported in other studies from China, ranging between 2.7%[21] and 1%.[2]

In our cohort of COVID-19 patients, a previous diagnosis of cancer increased the risk of death up to the age of 80, after which the impact of having had a cancer was modest or null. A systematic review including about 50,000 COVID-19 patients found exactly the same phenomenon: the impact of previous cancer on survival was important before the age of 65 but was almost null in the elderly.[19] This has also been observed for the impact of all comorbidities on COVID-19 survival, which decreases as age increases.[10]

It is worth noting that the excess risk of death and hospitalization was stronger in patients with a recent diagnosis of cancer and in patients with metastases at diagnosis. This is consistent with the findings of Liang and colleagues,[2] who found an excess only in patients who recently received treatment, and with the results of Dai and colleagues, who reported an excess for patients who had a diagnosis of stage IV cancer.[3] Data from a large cohort of patients with active cancer and COVID-19 in the UK did not confirm an excess risk due to recently undergoing chemotherapy or radiotherapy compared to patients that did not. The excess risk of those with metastases, however, was confirmed.[20]

Thus, it remains unclear whether the increased risk of death was due to the direct effects of cancer or to treatment (surgery and chemotherapy). However, our data plus insights from other studies suggest that the magnitude of the effect on COVID-19 prognosis is greater during the active phase of cancer, after which the impact decreases.

In our study the most represented cancer sites among COVID-19 patients were breast, digestive tract, urinary tract, and male genital organs; less frequent were thyroid, hematologic, female genital organs, melanoma, respiratory organs, lymphoma, head neck, and other sites. In studies from China,[3, 21] the most frequent cancer sites among COVID-19 patients were lung and digestive tract,

while breast was not as frequent as in Italy. These differences reflect the prevalence of these neoplasms in our province, compared with that in the Chinese population.

The strongest excess of death was observed in patients with cancer of the female genital organs, urinary tract, other hematologic organs, and thyroid, while for cancer of the respiratory organs and lymphomas, we observed a considerable excess of hospitalizations but a just a small excess of deaths. It is not easy to compare our results with those of other studies because most included only hospitalized patients; if the risk of hospitalization does not reflect that of death, relative risk among hospitalized patients is affected by a collider bias. Nevertheless, our data are not consistent with those of the large UK cohort, where small excess risk was observed for lymphoma and respiratory cancers.[20]

These results make it difficult to put together a profile of patients at higher risk of dying of COVID-19 based on cancer site and phase of care, for whom particular measures, including delaying treatments, could be taken to reduce the risk of infection, as suggested by some authors.[22,23] The only feature that clearly emerged was recent diagnosis and the presence of metastases.

Our data suggest that public awareness and measures allowing cancer survivors to reduce their risk of infection did reduce the cumulative incidence of COVID-19 and related deaths. This is an important message for the policy makers, physicians, and patients that are trying to define how to better manage patients during this crisis.[24]

The main strength of this study is its population-based design, which eliminates any selection bias occurring in case series. Furthermore, the assessment of exposure, i.e. previous diagnosis of cancer, was conducted through the linkage with a cancer registry with 25 years of prevalence data and timely registration of incident cases (to 31 December 2019). It is worth noting that this information was acquired before the onset of the pandemic and is thus completely independent of outcome occurrence. The main limitation of our study is that we do not have any information on treatment or on comorbidities, which could have influenced outcomes. Furthermore, because we could not include cancer patients with a diagnosis occurring in 2020, we could not observe the phase of diagnosis and disease assessment, which for many cancer sites is very intensive in terms of access to healthcare facilities.

Conclusion

Our population-based study showed that during the peak of the COVID-19 epidemic in northern Italy, cancer survivors had lower cumulative incidence of COVID-19 and related hospitalizations and death. Among COVID-19 patients, the cancer survivors had a higher risk of hospitalization and of death especially in the age group < 70, those with metastatic cancers, or with a recent diagnosis.

Conflicts of interest: The authors have no conflicts of interest to disclose.

References

1. www.salute.gov [consulted on 1[th] July 2020]

2. Liang W, Guan W, Chen R, Wang W, Li J, Xu K, Li C, Ai O, Lu W, Liang H, Li S, He J. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China.Lancet Oncol. 2020 Mar;21(3):335-337. doi: 10.1016/S1470-2045(20)30096-6. Epub 2020 Feb 14.PMID:32066541

3. Dai M, Liu D, Liu M, Zhou F, Li G, Chen Z, Zhang Z, You H, Wu M, Zheng Q, Xiong Y, Xiong H, Wang C, Chen C, Xiong F, Zhang Y, Peng Y, Ge S, Zhen B, Yu T, Wang L, Wang H, Liu Y, Chen Y, Mei J, Gao X, Li Z, Gan L, He C, Li Z, Shi Y, Qi Y, Yang J, Tenen D, Chai L, Mucci LA, Santillana M, Cai H. Patients with Cancer Appear More Vulnerable to SARS-COV-2: A Multicenter Study during the COVID-19 Outbreak. Cancer Discov. 2020 Jun;10(6):783-791. doi: 10.1158/2159-8290.CD-20-0422. Epub 2020 Apr 28

4. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. JAMA 2020;323:1239–42

5. Gosain R, Abdou Y, Singh A, Rana N, Puzanov I, Ernstoff MS. COVID-19 and Cancer: a Comprehensive Review. Curr Oncol Rep 2020 May 8;22(5):53. doi: 10.1007/s11912-020-00934-7

6. International Pharmaceutical Federation. Coronavirus SARS-CoV- 2 outbreak: information and guidelines for pharmacists and the pharmacy workforce. Febuary 12, 2020

7. Zhang L, Zhu F, Xie L, Wang C, Wang J, Chen R, Jia P, Guan HQ, Peng L, Chen Y, Peng P, Zhang P, Chu Q, Shen Q, Wang Y, Xu SY, Zhao JP, Zhou M. Clinical characteristics of COVID-19-infected cancer patients: a retrospective case study in three hospitals within Wuhan, China. Ann Oncol. 2020 Jul;31(7):894-901.doi: 10.1016/j.annonc.2020.03.296. Epub 2020 Mar 26

8. I numeri del cancro in Italia 2020. AIOM-AIRTUM-SIAPEC-IAP. Intermedia Editore, October 2020

9. Cancer Incidence in Five Continents, Vol. XI (electronic version). Lyon: International Agency for Research on Cancer. Available from: https://ci5.iarc.fr, accessed [23th October 2020].

10. Giorgi Rossi P, Ferroni E, Spila Alegiani S, Pitter G, Leoni O, Cereda D, Marino M, Pellizzari M, Sultana J, Trifirò G, Massari M and the ITA-COVID19 working group. Survival of hospitalized COVID-19 patients in Northern Italy: a population-based cohort study by the ITACOVID19 Network. https://doi.org/10.1101/2020.05.15.20103119

11. Rugge M, Zorzi M, Guzzinati S. SARS-CoV-2 infection in the Italian Veneto region: adverse outcomes in patients with cancer Nature Cancer volume 1, pages784–788(2020)2020 Jul;31(7):894-901.Brief Communication Published: 31 July 2020

12. He W, Chen L, Chen L, Yuan G, Fang Y, Chen W, Wu D, Liang B, Lu X, Ma Y, Li L, Wang H, Chen Z, Li Q, Gale RP. COVID-19 in persons with haematological cancers. Leukemia 2020;34(6):1637–45. https://doi.org/10.1038/s41375-020-0836-7

13. Van Dam PA, Huizing M, Mestach G, Dierckxsens S, Tjalma W, Bich Trinh X, Papadimitriou K, Altintas S, Vermorken J, Vulsteke C, Janssens A, Berneman Z, Prenen H, Meuris L, Berghe V, Smits E, Peeters M. SARS-CoV-2 and cancer: Are they really partners in crime?12. 2020 Sep;89:102068.doi: 10.1016/j.ctrv.2020.102068. Epub 2020 Jul 11

14. Yu J, Ouyang W, Chua MLK, Xie C. SARS-CoV-2 Transmission in Patients With Cancer at a Tertiary Care Hospital in Wuhan, China. JAMA Oncol 2020; 25: e200980https: //doi.org/10.1001/jamaoncol. 2020.0980

15. Barlesi F, Foulon S, Bayle A, Gachot B, Pommeret F, Willekens C, Stoclin A, Merad M, GriscelliI F, Micol JB, Sun R, Nihouarn T, Balleygier C, André F, Scotte F, Besse B, Soria JC, Albiges L, Roussy G. Outcome of cancer patients infected with COVID-19, including toxicity of cancer research. Presented at: 2020 virtual annual meeting of the American Association for Cancer Research; April 27–28; 2020

16. ElGohary GM, Hashmi S, Styczynski S, Kharfan-Dabaja MA, Alblooshi RM, de la Cámara R, Mohmed S, Alshaibani A, Cesaro S, Abd El-Aziz N, Almaghrabi R, Gergis U, Yasser M. The risk and prognosis of COVID-19 infection in cancer patients: A systematic review and meta-analysis. Hematol Oncol Stem Cell Ther. 2020 Jul 30:S1658-3876(20)30122-9. doi: 10.1016/j.hemonc.2020.07.005. Online ahead of print

17. Robinson AG, Gyawali B, Evans G. COVID-19 and cancer: do we really know what we think we know? Nat Rev Clin Oncol 2020;17(7):386–8

18. Parohan M, Yaghoubi S, Seraji A, Javanbakht MH, Sarraf P, Djalali M Risk factors for mortality in patients with Coronavirus disease 2019 (COVID-19) infection: a systematic review and meta-analysis of observational studies. Aging Male. 2020 Jun 8:1-9. doi: 10.1080/13685538.2020.1774748. Online ahead of print

19. Giannakoulis VG, Papoutsi E, Siempos II. Effect of Cancer on Clinical Outcomes of Patients With COVID-19: A Meta-Analysis of Patient Data JCO Glob Oncol. 2020 Jun;6:799-808. doi: 10.1200/GO.20.00225

20. Lee L, Cazier JB, Starkey T, Turnbull CD; UK Coronavirus Cancer Monitoring Project Team, Kerr R, Middleton G. COVID-19 mortality in patients with cancer on chemotherapy or other anticancer treatments: a prospective cohort study. Lancet. 2020 Jun 20;395(10241):1919-1926. doi: 10.1016/S0140-6736(20)31173-9. Epub 2020 May 28. PMID: 32473682

21. Ma J, Yin J, Qian Y, Wu Y. Clinical characteristics and prognosis in cancer patients with COVID-19: A single center's retrospective study. J Infect 2020 Aug;81(2):318-356. doi: 10.1016/j.jinf.2020.04.006. Epub 2020 Apr 14

22. Al-Quteimat OM, Amer AM. The Impact of the COVID-19 Pandemic on Cancer Patients. Am J Clin Oncol. 2020 Jun;43(6):452-455. doi: 10.1097/COC.0000000000000712. 15

23. Finley C, Prashad A, Camuso N, Daly C, Aprikian A, Ball CG, Bentley J, Charest D, Fata P, Helyer L, O'Connell D, Moloo H, Seely A, Werier J, Zhong T, Earle CC. Guidance for management of cancer surgery during the COVID-19 pandemic. Can J Surg. 2020 May 1;63(22):S2-S4. doi: 10.1503/cjs.005620

24. Al-Shamsi H, Alhazzani W, Alhuraiji A, Coomes EA, Chemaly RF, Almuhanna M, Wolff RA, Ibrahim NK, Chua MLK, Hotte SJ, Meyers BM, Elfiki T, Curigliano G, Eng C, Grothey A, Xie C. A Practical Approach to the Management of Cancer Patients During the Novel Coronavirus Disease 2019 (COVID-19) Pandemic: An International Collaborative Group. Oncologist 2020 Apr 3. doi: 10.1634/theoncologist.2020-0213. Online ahead of print

Table 1. Distribution of 15,391 patients who tested for Covid 19, by sex, age and cancer history.

	Population	Test Cov-2		Positive for SARS-CoV-2		Hospitalization		Death	
		(n.)	%	(n.)	%	(n.)	%	(n.)	%
Sex									
Males	261,563	6,741	2.6	1,989	0.8	830	41.7	282	14.2
Females	270,328	8,650	3.2	2,552	0.9	641	25.1	267	10.5
Age									
<50	304,808	5,918	1.9	1,334	0.4	167	12.5	3	0.2
50-59	80,406	2,677	3.3	818	1.0	164	20.0	11	1.3
60-69	60,419	1,915	3.2	651	1.1	259	39.8	41	6.3
70-79	48,973	1,733	3.5	527	1.1	313	59.4	89	16.9
>79	37,285	3,148	8.4	1,211	3.2	568	46.9	405	33.4
Cancer									
Yes	43,716	1,525	3.5	449	1.0	236	52.6	114	25.4
No	488,175	13,866	2.8	4,092	0.8	1,235	30.2	435	10.6
All cases	*531,891*	*15,391*	*2.9*	*4,541*	*0.9*	*1,471*	*32.4*	*549*	*12.1*

Table 2. Incidence rate ratio (IRR) of being tested, having a positive test and dying during Covid-19 disease for people with previous diagnosis of cancer compared with the general population without cancer.

	IRR	95% CI	
Test Cov-2	0.69	0.65	0.73
Test Positive	0.55	0.50	0.61
Hospitalization	0.59	0.51	0.68
Death	0.52	0.42	0.64

Table 3. Characteristics of patients who tested for Covid 19 with and without cancer.

Test Cov-2	All cases	Patients with cancer	Patients without cancer	Males with cancer	Males without cancer	Females with cancer	Females without cancer
	n. (%)	n. (%)	n. (%)	n. (%)	n. (%)	n. (%)	n. (%)
Positive	4,541 (29.5)	449 (29.4)	4,092 (29.5)	206 (26.9)	1,783 (29.9)	243 (32.0)	2,309 (29.2)
Negative	10,850 (70.5)	1,076 (70.6)	9,774 (70.5)	560 (73.1)	4,192 (70.1)	516 (68.0)	5,582 (70.8)
All cases	15,391	1,525	13,866	766	5,975	759	7,891
P		0,955		0,09		0,11	

Table 4. Multivariate logistic regression of positive cases by age and sex, and age-stratified multivariate logistic regression for cancer history.

	Hospitalization			Death		
	OR*	95%CI		OR*	95%CI	
Sex						
Male	1			1		
Female	0.61	0.56	0.66	0.62	0.54	0.71
Age						
< 70	0.43	0.39	0.47	0.06	0.04	0.07
70 - 80	1.11	1.01	1.21	0.49	0.41	0.59
> 80	1			1		
Cancer and age						
Age<70						
Without cancer	1			1		
With cancer	2.52	1.76	3.62	4.91	2.38	10.09
Age 70-80						
Without cancer	1			1		
With cancer	1.55	0.99	2.44	1.72	1.05	2.83
Age>80						
Without cancer	1			1		
With cancer	1.51	1.10	2.07	1.20	0.87	1.65

Table 5. Multivariate logistic regression of positive cases related to clinical outcomes

		Hospitalization			Death		
	n	OR*	95%CI		OR*	95%CI	
Cancer							
None	4,092	1			1		
Yes	449	1.66	1.34	2.05	1.45	1.12	1.89
Metastasis							
None	417	1			1		
Yes	27	1.66	0.70	3.93	2.14	0.89	5.16
Cancer diagnosis							
None	4,092	1			1		
<2 years	50	3.28	1.72	6.25	2.99	1.52	5.89
2-5 years	124	1.79	1.21	2.65	1.21	0.72	2.02
>5 years	275	1.61	1.23	2.10	1.40	1.02	1.91
Cancer site							
Breast	101	1.16	0.75	1.79	0.95	0.52	1.73
Digestive organs	86	1.34	0.86	2.09	1.17	0.69	1.97
Urinary tract	57	1.98	1.11	3.51	1.83	0.98	3.42
Male genital organs	56	1.08	0.61	1.91	0.64	0.32	1.30
Thyroid	25	1.52	0.61	3.79	3.08	0.77	12.26
Female genital organs	24	3.04	1.32	7.01	3.40	1.28	9.02
Other haematological	24	3.58	1.46	8.79	4.74	1.84	12.21
Respiratory organs	20	3.63	1.26	10.44	1.64	0.58	4.64
Melanoma	20	1.35	0.52	3.49	3.77	1.15	12.35
Lymphoma	20	5.69	1.93	16.80	1.49	0.43	5.15
Head-neck	8	0.73	0.17	3.17	2.86	0.56	14.69
Other site	6	2.81	0.51	15.49	3.42	0.46	25.55

Fig 1. Characteristics and clinical outcomes of 4,541 COVID-19 positive patients, with and without cancer.