____

The effectiveness of early diagnostic tools for neural lesions in leprosy

An observational cohort study

Luciano Henrique Vieira Cegana[1,¶,*,#a], Susilene Maria Tonelli Nardi[2,&,*,#b], Luciana Pascoeto[3,&], Vania Del Arco Paschoal[4,&]

1 Department of Physical Therapy and Public Health, Municipal Health Office, Public Service, Votuporanga, São Paulo, Brazil

2 Department of Biomedical Sciences, Adolfo Lutz Institute, Health Center, São José do Rio Preto, São Paulo, Brazil.

3 Department of Physical Therapy and Public Health, Specialized Physical Rehabilitation Center – CER II, Public Service, São José do Rio Preto, São Paulo, Brazil

4 Department of Nursing and Public Health, Medical School of São José do Rio Preto – FAMERP, Medical School, São José do Rio Preto, São Paulo, Brazil

* Corresponding author
E-mail: lucegana@gmail.com (LHVC)
E-mail: susilenenardi@gmail.com (SMTN)

____

ABSTRACT

Introduction: Leprosy can cause different lesions in peripheral nerves and inervatory structures. **Objectives:** To analyse the effectiveness of evaluation protocols used to identify neural lesions in leprosy such as Degree of Physical Disability (DPD), Simplified Neurological Assessment (SNA) and propose to use Neurodynamic Assessment (NDA). **Method:** Descriptive analytical study, associative, with 27 individuals treated in two outpatient leprosy clinics in São Paulo State, between 2017 and 2019, and 27 individuals from the paired control group. The Mann-Whitney, Multivariate Linear Regression and association between variables and $P<0.05$ values were used. **Results:** The test that most captured the neurological alterations was the SNA, with 22 (81.5%) in the upper limbs (ULs) and 25 (92.6%) in the lower limbs (LLs), followed by the NDA, with 20 (74.1%) in the ULs and 11 (40.7%) in the LLs. The DPD showed handicap in the hands of 16 (59.2%) individuals and in the feet of 17 (62.9%) individuals, and they have expressed sensitivity. DPD showed agreement with SNA in 21 (77.8%) of the cases in ULs ($p=0.010$) and 19 (70,4%) of the cases in LLs ($p=0.060$). DPD and NDA showed that in 19 (70.4%) of the patients evaluated there was concordance of results in ULs ($p=0.143$); 9 (33.3%) in LLs ($p=0.125$). SNA and the NDA in the ULs found agreement in 21 (77.8%); 11 (40.7%) ($p=0.786$) in the LLs. **Conclusion:** The three assessment instruments are specific and will hardly produce false positive tests. DPD can produce more false negatives than SNA. If there is an instrument to be chosen, it should be the SNA, since it is more sensitive, more accurate and has a less negative likelihood ratio. Neurodynamic tests were positive in 7.4% of individuals while there were still no changes in the SNA; afterwards, these changes appeared.

____

Descriptors: Neural mobilization; Leprosy; Peripheral nerves; Disability; Pain; Physical

therapy.

INTRODUCTION

Among infectious diseases, leprosy is considered one of the main causes of physical

disabilities, due to its potential to cause neural injuries[1].

Worldwide, 208,619 new cases of the disease were reported to the World Health

Organization (WHO) in 2018, 30,957 occurred in the Americas region and 28,660 (92.6% of

the total from the Americas) were reported in Brazil. New cases, which were assessed for its

Degree of Physical Disability (DPD), totaled 86.5% (n=24,780); 2,109 (8.5%) had visible

physical deformities (DPD 2)[2].

The lack of new diagnostic tools and new drugs, limited knowledge about strategic

areas of transmission and unsatisfactory tools for managing the complications caused by

leprosy have been an obstacle to its control. Therefore, more coordinated research efforts are

still needed[3].

The diagnosis of leprosy is based on the presence of at least one of three cardinal

signs: a) lesion(s) and/or area(s) of the skin with changes in thermal and/or pain and/or tactile

sensitivity; or b) thickening of the nerve trunks of the peripheral nerves in the upper and

lower limbs, associated with sensitivity and/or motor changes in extremities, with decreased

muscle strength in the myotomes supplied by these and/or autonomic nerves; or c) presence

of *Mycobacterium leprae* bacilli, confirmed by intradermal smear microscopy or skin

biopsy[4].

Sometimes neurological involvement is preponderant when compared to a

dermatological condition not very significant[5]. The most affected nerves are the ulnar in the

elbow and the fibular in the head of the fibula, followed by the ulnar sensory, superficial and

sural fibular branches[6].

In daily practice, a greater occurrence of sensory changes over motor changes is

found, as well as a slight presence of deformities[7]. The sensory and motor actions of

____

M.leprae in the upper and lower limbs cause secondary injuries such as fingers [8] and toes

in flexion and "fallen hands and foot"[9].

Although the prevalence of leprosy is decreasing, cases that no longer have the active

disease, but which present neurological sequelae, continue to be monitored in health

services[10]. Leprosy reactions are one of the main causes of the onset of disabilities[11]. It is

essential to assess the integrity of neural function at the time of diagnosis, in the occurrence

of reactive states, at discharge due to cure (end of multidrug therapy) and during 5 years after

discharge[7].

Peripheral neuropathies in leprosy are a triggering factor for physical disabilities, and

some evaluation protocols can be used for their prevention, such as the Degree of Physical

Disability of the World Health Organization (DPD) and the Simplified Neurological

Assessment (SNA) of the Brazilian Ministry of Health[12]. The routine protocols

recommended by the Brazilian Ministry of Health are subjective and, in view of that, other

tests could complement the diagnostic examination of peripheral neuropathy such as the

Neurodynamic Assessment (NDA)[7].

The DPD is an epidemiological indicator, not very sensitive, but it can define the

severity of neurological changes. The SNA assesses muscular strength and sensitivity of

eyes, hands and feet, in addition to the palpation of the possibly affected nerves and the

morphological evaluation of the limbs[7].

The Neurodynamic Assessment (NDA) is used for injury diagnosis and treatment of

the peripheral nervous system and the structures innervated by it[13]; the interface between

the musculoskeletal system and the peripheral nervous system is used, so the movements

applied to the musculoskeletal system mobilize the structures of the peripheral nervous

system[14].

____

105   The use of neurodynamic tests has helped in the diagnosis of neural lesions to identify

106   the origin of the lesions, but little is known about the results of the neurodynamic tests applied

107   to patients with leprosy. Thus, we suggest using the mobilization of the nervous system in

108   clinical practice as a tool to assess peripheral nerves that may be affected by leprosy.

____

OBJECTIVES

Analyzing the effectiveness of the evaluation protocols used to identify neural lesions

in leprosy, such as the Degree of Physical Disability (DPD) recommended by the World

Health Organization, the Simplified Neurological Assessment (SNA) and propose to use the

Neurodynamic Assessment (NDA)

Describing the sociodemographic and clinical variables of the groups studied and

check whether there is a statistical association between the results of the evaluation tools: the

Degree of Physical Disability recommended by the World Health Organization (DPD), the

Simplified Neurological Assessment (SNA) and the positive Neurodynamic Assessment

(NDA) in individuals who have or had leprosy, in addition to determining the specificity and

sensitivity of each of the assessment protocols.

____

METHODS

Associative, analytical, descriptive research, following case-control design, approved

by the FAMERP Ethics and Research Committee in accordance with the requirements of

National Resolution No. 196/96, Opinion No. 2,469,355. Participants were asked to sign the

Informed Consent Form (ICF) in accordance with Ordinance No. 466/2012.

The study included all leprosy patients diagnosed between 2017 and 2019, that agreed

to sign ICF, N=27 of both sexes and different ages, in one medium (50 to 100 thousand

inhabitants) and one large (100,001 to 900,000 inhabitants) Brazilian municipality

(Votuporanga and São José do Rio Preto), under treatment or undergoing chemotherapy. A

control group of 27 people, chosen from the population, without a diagnosis of leprosy, was

evaluated pairing age and sex, thus totaling 54 people. Data collection went from December

2017 to November 2019, until three evaluations were completed, with an average interval of

3 months between evaluations.

We selected as exclusion criterion for both groups those with compressive syndromes

of the central nervous system (CNS) and the peripheral nervous system, other diseases of the

CNS such as stroke or degenerative syndromes, diabetics and alcoholics with altered

sensitivity and people/patients who refused to sign the informed consent form.

Initially, a patient profile data sheet, extracted from their medical records, was used,

with the name, address, age, sex, date of the beginning of treatment, date of discharge, clinical

classification of leprosy and type of treatment. In the control group, name, address, age, sex

and existing morbidities were used. Evaluation instruments were applied:

• **Simplified Neurological Assessment (SNA)** is a protocol recommended by the

Brazilian Ministry of Health, and it contemplates dermatoneurological exploration,

the evaluation of eyes, hands and feet sensitivity, and the evaluation of the motor

function[15]. In this clinical examination, the integrity of the skin and its nutrition are

____

assessed. Testing quick perception of a light touch and/or deep pressure[15]. It is

recommended to use the Semmes-Weinstein monofilament set (6 monofilaments:

0.05g, 0.2g, 2g, 4g, 10g and 300g) in the sensitivity assessment points in hands

(palmar surface and back of the hand) and feet (plantar surface and instep near the

hallux and the second toe)[15]. To assess motor strength, it is recommended to perform

manual testing of muscle strength in the muscle tendon unit during movement and

the ability to oppose gravity and manual resistance in each muscle group from a

specific nerve myotome. Ulnar nerve (Abductor digiti minimi), median nerve

(abductor pollicis brevis), radial nerve (wrist extensors), fibular nerve (tibialis

anterior, extensor digitorum longus, extensor hallucis longus, fibularis tertios).

The grading criteria for muscle strength can be expressed from 0 to 5 following the

Kendall scale for the degree of muscle strength[15]. (Appendix 1)

• **WHO Degree of Physical Disability (DPD**) indicates the loss of protective

sensitivity, muscle strength and / or visible deformities in the face, upper and lower

limbs, varies between Degree 0 (no leprosy-related physical disability), I (decreased

strength and/or loss of sensation) or II (presence of visible disabilities and

deformities) according to the severity of sensory and/or motor and/or morphological

changes caused by neural lesions of leprosy[15].

● **Neurodynamic tests**: straight leg raises (SLR), Slump Test and upper limb tension

tests (ULTT) 1, 2b and 3[16].

1. The straight-leg raise test (SLR) is performed with the patient in the supine position, with

the trunk and hips in neutral positions. The examiner places one hand under the Achilles

tendon and the other above the knee. The hip is flexed with the knee held in extension

until it reveals a predetermined symptomatic response, or until it reaches its hip range of

motion (ROM) limit. The ROM value should be compared to the SLR of the contralateral

____

limb, and to what is considered as normal. If pain, especially lower back pain, is reported

during the test, the most likely cause is a herniated disc or a central pathology causing

compression. The normal value of SLR in normal individuals varies from 50° to 120°[16].

But if pain is reported in the limb following the nerve path, we consider the test positive

in individuals with leprosy, as neural tension can be caused by compression or

incarceration of the peripheral nerve. Painful expression or reproduction of symptoms

usually in the course of innervation. Do not confuse with the discomfort of hamstring

stretching, because when maximal knee extension increases posterior thigh pain, it did

not increase muscle tension, measured indirectly using EMG. The test is considered

positive when the patient reports any discomfort/pain (pins and needles and sensation

changes, pulling, tension, pain, numbness or tingling, that radiates inferiorly through the

posterior/lateral sides of the leg, often associated with paresthesia) or when the physical

therapist encounters resistance to movement [17].

2. The Slump Test should be performed with the patient seated, with the thighs fully

supported, the knees together and the hands together on the back. The patient is asked to

flex the thoracic and lumbar spine, keeping the sacrum in a vertical position and, soon

after, also perform cervical flexion. The physiotherapist puts pressure on the cervical

region in order to accentuate flexion. Then, the patient performs an active knee extension,

associated with dorsal flexion of the ankle. Cervical flexion is slowly released, and the

painful response must be carefully evaluated. The symptoms must be noted at each stage

and must also be performed for the other member. ROM and painful responses should be

assessed[16].

3. The upper limb tension test (ULTT1) (median neural tension) is described as a brachial

plexus tension test or a median nerve test. It is performed with the patient in the supine

position. The examiner exerts force to depress the scapular waist, which has an external

____

rotation and 110° abduction of the glenohumeral articulation, elbow extension, radioulnar

supination and wrist and fingers extension, the inclination of the cervical to the opposite

side was suppressed in our assessment to isolate core involvements. ULTT1 tests are

considered positive if present: complaint at deep elongation or pain in the cubital fossa

that extends down the anterior and radial part of the forearm and to the radial side of the

hand; tingling sensation in the first four fingers; stretching in the anterior shoulder

area[16].

4. The upper limb tension tests (ULTT 2b) assess the radial nerve with the patient in the

supine position. The examiner holds the elbow and wrist of the patient. Using the thigh,

the examiner depresses the scapular waist and internally rotates the shoulder, extends the

elbow, flexes the wrist, fingers and thumb[16].

5. The upper limb tension test (ULTT 3) (neural tension of the ulnar) is performed with the

patient in the supine position, keeping the wrist of the patient extended and the forearm

supine, performing an elbow flexion. After positioning the patient, the therapist performs

a shoulder depression associated with an external rotation. Shoulder abduction is added

(so that the hand is close to the ear of the patient). In our assessment, we suppressed

lateral flexion of the cervical to isolate central pathologies. The test is considered positive

when the patient reports any discomfort/pain or when the physical therapist encounters

resistance to movement. Pain is intermittent, deep and burning in quality. Also, a cold,

tingling feeling extended distally from the medial elbow to the little finger [18]. When the

patient has any of this reported symptoms, the test is completed and the physiotherapist

keeps the limb of the patient positioned in the range in which pain or resistance to

movement is reported to perform a goniometer measurement[14].

The leprosy group underwent 3 evaluations, quarterly and the control group was

evaluated in a single moment. None of the subjects were in reaction episode at the time of

____

the evaluation. The application of the data collection instrument took 40 to 50 minutes and

was performed by a single examiner.

Data analysis

● The Simplified Neurological Assessment was assigned with a value "without

changes" or with the value "with changes" for the upper and lower limbs. We considered that

the change in sensitivity in hands decreased when the individual didn't feel the 0,02g green

monofilament. The decrease in plantar sensitivity was considered when didn't feel the 0,5g

blue monofilament. The muscle strength domain was also considered to perform SNA

grouping and was coded as "altered" when muscle strength was less than 5, according to the

Kendall scale, identifying a muscular paresis and therefore a neurological alteration [15]. The

palpation was marked as altered when there was pain, thickening, or both.

● We synthesize the results of the Neurodynamic Assessment in positive and

negative tests, considering the set of tests for the upper and lower limbs one apart to the other.

● The degree of physical disability of the WHO was classified as "*without disability*"

when it was Degree 0, and was grouped as "*disabled*" when its results were Degrees 1 or 2.

After tabulation of the data collected in this work, 2 analytical and statistical functions

were performed: descriptive and inferential. In a descriptive way, the profile of the studied

sample was drawn, considering the analyzed variables and their consequences, and the data

was replicated in an absolute and relative way.

In the inferential scope, the analysis of independence and prediction between

variables was drawn as a statistical objective. In addition, the Mann-Whitney U test and

Multivariate Linear Regression were used with the results of independence between the

proposed variables, taking into account the P-values (significance $P \leq 0.05$). All analyses were

____

obtained using the SPSS Statistics Software (Version 23) and were linked to the features of

the Excel tool (version 2016).

Statistical analysis and synthesis of the results were performed on a Venn-Euler

diagram. To perform the descriptive analysis and verify the association between SNA, DPD

and NDA results, EPI INFO 7.1 statistical software was used.

To validate the diagnostic tests, calculations of sensitivity, specificity, positive and

negative likelihood ratios were performed using the Virtual Health Library (*Biblioteca*

Virtual de Saúde, BVS) virtual calculator[19]. Sensitivity, specificity and accuracy vary from

0-1. A sensitive and specific test is one which is positive in the presence of pathology and

negative in the absence of pathology[18,20].

Likelihood ratios show the performance of a diagnostic test. The Positive Likelihood

Ratio (PLR) varies from 1 to infinity, the higher your result, the better the test. The Negative

Likelihood Ratio (NLR) varies from 1 to 0, the closer to 0, the better the test[21].

____

RESULTS

In this study, all 27 individuals diagnosed with leprosy who were notified in the years

2017 to 2019 participated (8 in the municipality of Votuporanga, 19 in the municipality of

São José do Rio Preto), in addition to 27 volunteers matched for age and sex in the control

group, totaling 54 people. Among the intervention group, 17 (63%) were male. The ages

varied between 23 and 88 years in the control group and between 23 and 87 years in the

intervention group, with a mean age of 53.1 (SD 17.6). The age median in the control group

was 57 while in the intervention group it was 55.

The age distribution was identical in the control and intervention groups due to the

pairing (n=54) (p-value 0.945). There was no difference in sex (p-value 1.00). In the

intervention group there were 19 (70.4%) active workers, in the control group it was 22

(81.5%) (p-value 0.344).

Regarding leprosy, 20 (74.1%) of the participants in the intervention group had the

dimorphous and Virchow's leprosy clinical forms, with multibacillary multidrug therapy

(74.1%). Seven (25.9%) patients had a reaction phenomenon to leprosy during the treatment,

but not at the time of the evaluations. 25 (92,6%) still remained under drug treatment, 11

(40,7%) were evaluated in less than 30 days after diagnosis, 4 (14,8%) were evaluated in less

than 60 days after diagnosis, 1 (3,7%) were evaluated in less than 90 days after diagnosis, 3

(11,1%) before 180 days after diagnosis, 3 (11,1%) before 270 days after diagnosis and 3

(11,1%) before completing 365 days between diagnosis and first evaluation. 2 (7,4%) patients

were evaluated after de discharge.

The participants in the intervention group showed neurological changes, at least in

one site of the upper limbs, assessed by the three types of exams. The test that most captured

data was the SNA with 22 (81.5%) participants, followed by the Neurodynamic Assessment,

____

where 20 (74.1%) were altered, and the DPD showed deficiency in the hands of 16 (59.2%)

individuals.

As assessing the lower limbs, it appears that in the intervention group the participants

showed neurological changes in the three assessments studied. Neurological changes were

more perceived by SNA, attested in 25 (92.6%) participants. The Degree of Physical

Disability showed a disability in 17 (62.9%) individuals. In the Neurodynamic Assessment,

11 (40.7%) showed alterations. In the control group, no neurological changes were identified

by the 3 instruments, confirming the reliability of the sample (Table 1).

In Figure 1, using the Venn-Euler diagram, it can be seen that 51.8% (14 of 27) of

patients were diagnosed with neural problems in the upper limbs using the three analysis

instruments. When we use only two associated assessments, the number of cases with

changes decreases. We also observed that the instruments individually captured different

people with problems, 2 (7.4%) with SNA and 2 with NDA; that is, 2 individuals that only

had their neural lesions identified because they passed through SNA and another 2 different

individuals only had neural injury identified because they were subjected to NDA.

In Figure 2, using the Venn-Euler diagram, it can be seen that 18.5% (5 of 27) of

patients were diagnosed with neural problems in the lower limbs using the three analysis

instruments. When using two instruments, SNA and NDA, the number of identified cases

remains with 5, (18.5%); using NDA and DPD it is 0 (0%), as they do not identify any

individual who is also not identified by SNA. Using the DPD instrument and the SNA

together identifies more individuals with neural lesions 12 (44.4%). We also observed that

the instruments individually captured different people with problems, 3 (11.1%) with SNA

and 1 (3.7%) with NDA; that is, three individuals only had their neural lesions identified

because they passed through SNA and the neural lesion of another different individual was

only identified because he was subjected to NDA.

____

Sensitivity, specificity, positive and negative likelihood ratio

To define the characteristics of the tests as a diagnosis, we used the concepts of

sensitivity, specificity and positive and negative likelihood ratios considering the ULs and

the LLs separately, based on the distribution of the test results, in order to provide evidence

on which clinical practices can be based. (Table 2)

Our sample tells us that the instruments used are specific because the 3 assessments

obtained a value equal to 1 in both the upper limbs and lower limbs, and that they hardly

produce false positives, since the positive likelihood ratio tended to infinity in the 3

assessment instruments[21].

Considering the sensitivity, we saw that SNA is the most sensitive test to assess the

upper limbs in individuals with leprosy, as it had a value of 0.81, followed by the NDA with

0.74. The DPD is considered least sensitive, with a value that was equal to 0.59[20]. In LLs,

SNA remains as the most sensible over other assessment instruments, with 0.92. NDA was

not a tool as sensitive for LLs as it was for ULs because we obtained a sensitivity value equal

to 0.41. DPD obtained a better sensitivity, with 0.63[20].

The best accuracy in ULs was obtained by SNA (0.90), followed by NDA (with 0.87)

and DPD (with 0.80), whereas in LLs, the accuracy of SNA was 0.96, followed by DPD,

with 0.81, and NDA, with 0.70[20].

According to the Negative Likelihood Ratio (NLR) in ULs, SNA obtained the best

value among the evaluations, with 0.18, while the value in the NDA and DPD was 0.26 and

0.40, respectively, showing that false negatives can occur more frequently in the NDA, and

could be still larger in the DPD. In LLs, the NLR in the SNA was 0.07, meaning that it is

difficult for the SNA to produce a false negative; however, the NDA can produce false

____

negatives because the NLR was equal to 0.60; the DPD also could report no deficiency while

in fact there is already a neural alteration because the NLR was equal to 0.37[21].

Degree of hand and foot disabilities compared to the Simplified Neurological

Assessment and Neurodynamic Assessment of upper and lower limbs

The WHO Disability Degree indicator of the hands shows agreement in 21 (77.8%)

of the cases that were assessed by SNA (P-value = 0.010). It should be noted that the SNA

made the diagnosis of neural injury in 6 (22.3%) cases that, according to the DPD, did not

have physical disability (degree zero of physical disability).

Regarding the association of the degree of hand impairment with the Neurodynamic

Assessment in 51.8% (n=14) of the patients evaluated there were agreed positive results; in

18.5% (n=5) there were agreed negative results. The results differed in eight participants,

with the Neurodynamic Assessment more frequently capturing neural changes (n=6/22.2%)

than the Degree of Physical Disability (n=2/7.4%). The association was not significant (p-

value=0.143).

The comparison between the degree of disability of the feet and the SNA of the lower

limbs was acceptable in 62.9% (n=17) of the cases but diverged in 29.6% (n=8) cases,

exposing a greater sensitivity of the SNA in showing the neural damage, despite the

association not showing statistical significance (p-value=0.060).

Verifying the degree of disability of the feet and the Neurodynamic Assessment of

the lower limbs showed agreement of positive results in 18.5% (n=5) of the patients evaluated

and in 14.8% (n=4) of the negative results. In eighteen participants, the results differed; the

Degree of Physical Disability of the feet more frequently captured neural changes

(n=12/44.4%) than the Neurodynamic Assessment (n=6/22.2%). The association was not

significant (p-value=0.125). (Table 3)

____

Simplified Neurological Assessment compared to Neurodynamic Assessment in the

Upper and Lower Limbs

By checking the Simplified Neurological Assessment and the Neurodynamic

Assessment in the ULs, we found agreement on the positive results in 66.7% (n=18) of the

cases with neurological changes and in 11.1% (n=3) of the cases without changes with

negative tests. There was disagreement in six individuals. The SNA was more effective in

verifying the neurological changes of the upper limbs in 14.8% (n=4) cases while the

Neurodynamic Assessment identified it in 7.4% (n=2) cases.

The association between Simplified Neurological Assessment and the Neurodynamic

Assessment of lower limbs showed agreement on positive results in 37% (n=10) of the

patients evaluated and in 3.7% (n=1) of the negative results. In sixteen participants, the results

differed, the SNA of LLs more frequently captured neural changes (n=15/55.5%) than the

Neurodynamic Assessment of the lower limbs (n=1/3.7%). The association was not

significant (p-value=0.786). (Table 4)

____

DISCUSSION

With this study we hoped to identify an evaluation protocol that used only human

resources for its execution, facilitating the identification of neural lesions in leprosy

combined with the daily practice of physical therapists who are not involved in the

management of leprosy, but routinely use Neurodynamic Assessment. In this way, more

suspicions and diagnoses of the disease could be made and would help in cases that present

themselves as purely neural. Concerning leprosy, it is important that not only the physician

suspects the disease, but any health professional.

There was no statistical difference in relation to sex, age and occupation in the two

groups studied, since the pairing was performed to avoid the effect of age and sex, two

recognized confounding factors for the results[22,23], so a comparison was made between

groups, showing similarity and equal care between them[24,25].

In the intervention group, men, active, with an average age of 53.09 years and

multibacillary disease were more frequent, as happened in Santarém, where they had ages

above 51 years old[26], being one of the sociodemographic factors that were associated with

the installation of disabilities[27].

Regarding neurological tests, the one that most captured neural alterations was the

SNA in both upper limbs and lower limbs, highlighting the importance of the process of

identifying the degree of disabilities through SNA, at the beginning of treatment and after

discharge[28]. Studies show worsening of physical disabilities during treatment, mainly in the

multibacillary form, with the lower limbs being the segments that show the most significant

degree of disability evolution, justifying the imperative need for a more careful monitoring

of these cases through routine assessments and interventions[29].

In the univariate and multivariate analysis, there is agreement of the result found in

the DPD with the SNA in the upper limbs, but the SNA detects neural lesions more

____

sensitively. Since disability is only considered when sensitivity is decreased beyond purple

monofilament and/or muscle strength is less than 5, DPD does not detect the subtle changes

that suggest neural distress, and we emphasize that the SNA made the diagnosis of neural

injury in 6 (22.3%) of the cases while, according to the DPD, those cases did not yet present

disabilities, showing the importance of using and applying SNA.

The degree of physical disability was an innovative measure to gradually quantify

physical disability in an index and were not designed to be sensitive to early changes, but

with the increasingly early diagnosis of neural damage, it was necessary to improve the

assessment instruments. Due to the change in the profile of patients today, these instruments

need to be more sensitive to subtle changes that neurological changes show us over the course

of the disease, so SNA is more sensitive in detecting neurological changes and their

influences on socioeconomic and emotional aspects[30].

The DPD is recommended at the beginning of treatment and at discharge, but when

the patient shows a slight improvement, there is no variation in the degree of disability and

the services are unable to assess whether the actions developed are actually being

effective[31]. It must be considered that when the case is carefully evaluated and the results

of the exams recorded in the SNA protocol, the diagnosis of deficiencies in the eyes, hands

and feet is more accurate than when the deficiency is identified using the framework "WHO

degree of physical disability"[28].

As an indicator, there is a fragile and subjective comparison of the DPD with the

limitation of activities, and the social participation of the patients is not included in its results.

An adaptation of the indicator is necessary to develop a more current classification, based on

a more comprehensive concept of disability, as is done by the International Classification of

Functioning (ICF)[32].

____

Nerve damage is associated with physical disabilities. Thus, regular monitoring of

nerve function, through SNA, combined with adequate clinical management of neuritis,

neuropathies and leprosy reactions, are effective strategies to prevent it[33].

In the univariate analysis, as well as in the multivariate, of ULs and LLs, DPD and

NDA, despite their results being relative to neural injury concordance in some cases, were

not statistically significant. The NDA seems to identify 75% more injuries in the upper limbs

than the DPD, and DPD seems to identify 50% more injuries in the lower limbs than the

NDA.

Considering leprosy and NDA, a study of ULs found a positive NDA, mainly

affecting those with a degree of physical disability level 2. Of these, they presented a decrease

in the ROM of elbow flexion in the ulnar neural tension test (ULTT3) on both sides when

compared to control group[14].

Leprosy patients test positive when submitted to Neurodynamic Assessment of lower

limbs[25]. But, even when the tests do not reproduce the symptoms in the affected nerves,

characterizing negative tests, studies suggest that a neuropathy cannot be ruled out yet, this

can mean a more severe lesion with demyelination of the fibres[35] (as occurred in degree 2

patients that tested negative in the lower limbs in our analysis), producing a possible false

negative result[20], thus explaining the low sensitivity of the NDA in lower limbs, because

low sensitivity produces more false negative results[20].

Both in the univariate and multivariate analysis of ULs and LLs, when comparing the

SNA with the NDA, we note that the SNA identifies the neural lesion earlier than the NDA.

According to the evidence available in 2019, it is suggested that neurodynamic tests

should not be used in isolation, as a single test, to diagnose neural distress of the median

nerve. They should be interpreted in the context of a loss of function tests of small fibres in

a domain[34]. Combining anamnesis and clinical history is an important tool to make the

____

differential diagnosis, where the combination of negative neurodynamic test results could be

used to rule out a disorder in the peripheral nerves; with limited evidence, they are relevant

in cervical radiculopathies[35].

The need for adaptation of the peripheral nervous system (PNS) in the face of

mechanical stress, stretching, slips, changes in its diameter and compressions is notorious. If

this dynamic protection mechanism fails, the PNS is vulnerable to edema, ischemia, fibrosis

and hypoxia, which are the causes of neural changes[36].

In this sense, the scientific community highlights the need for more possibilities for

neural injury investigations in leprosy, in order to diagnose and monitor the neurological

changes caused by it. A study in Nigeria points out that 50% of patients who complete

treatment already had neurological changes before diagnosis, according to the EHF score,

but 90%, when receiving assistance and monitoring of injuries, end treatment with less

disability[37]. Neural damage needs to be identified early and current leprosy control efforts

must be intensified to ensure immediate treatment in order to reduce the burden of the

disease, including deficiencies in individuals and the community. When using EHF, greater

sensitivity is offered in the evaluation compared to DPD[31]. Physical disabilities were

diagnosed in 41% of the individuals. The EHF score showed overlapping of impairments in

the examined segments and proved to be more appropriate than the DPD classification system

to describe the extent of the physical disabilities in the patients[38].

In our study, neurodynamic tests were positive in 2 (7.4%) individuals while there

were still no changes in SNA and later these changes appeared, which makes us think of the

association of assessments as a way to complement the diagnosis and monitoring of neural

changes. This would make professionals, when applying NDA, be able to suspect neural

injury caused by leprosy and minimally perform other evaluations, making a more accurate

investigation or even referring it to a specialist in the area.

____

Therefore, a need to create an assessment (validated or not) is necessary so that we

can, through an indicator, reveal the neural lesion more precociously, given that 95% of

patients have neurological changes, with musculoskeletal symptoms, which interfere in their

functional capacities, causing difficulties in performing their activities of daily living and

work when compared to those who have no symptoms. However, the presence of disability

did not prevent or limit them from performing these activities. Even with pain, paresthesia,

decreased strength and other injuries, they still perform their activities[39].

____

Conclusions

We conclude that the 3 assessment instruments are specific and that these assessments

can and should be used in combination to expand the monitoring of neural lesions in leprosy,

as there are changes that are not perceptible with one instrument but can be confirmed by

another.

The Simplified Neurological Assessment (SNA) is the most sensitive instrument,

with greater accuracy and it is unlikely to produce a false negative both in the assessments

of the upper limbs and lower limbs. The Neurodynamic Assessment (NDA) is the second

most sensitive instrument for assessing ULs, followed by the Degree of Physical Disability

(DPD). In LLs, the second most sensitive instrument is DPD, followed by NDA.

Both NDA and DPD can produce false negative results more frequently than SNA,

that is, reporting that nerves are healthy while in reality they already have existing

neurological changes, implying that some individuals may not receive the proper treatment

as early as possible.

The negative likelihood ratio confirms that the DPD produces more false negatives

than SNA, so when the DPD tells us that there are no neurological changes yet, they may

already be present at a more subtle level, so we suggest that the DPD always be performed

after the results of the SNA and not determining the DPD using only the table that helps in

the summary of the SNA data proposed by the guidelines, which considers visible

deficiencies such as a Degree of Physical Disability Level 2 (DPD 2) or a barely visible

Degree of Physical Disability level 1 (DPD 1), since neurological alterations are thus

underestimated.

The WHO grades has been used as an indicator for early case detection and as an

indicator for change in impairments, but depending on the expertise of the team, the degree

of disability will be the only instrument used to monitor and assess the patient, and The WHO

Grades are not designed to be sensitive to early changes.

Greater coverage is achieved, for the early identification of neural injuries, by

applying the 3 assessments studied in this research, opting for SNA, if you choose only one,

as it was the most sensitive and accurate instrument for identifying neural injuries in isolation,

in addition to having the least negative likelihood ratio.

NDA does not establish a relationship of dependency with DPD instruments, nor with

SNA; in its application to investigate neural injury in people who have leprosy, it does not

prove to be such a sensitive tool in isolation, but when associated with clinical anamnesis

and evaluations already used, DPD and SNA, it facilitates diagnosis, impacting the suspicion

of new cases of the disease.

The limitation of this study is found in the patient's lack of access to more complex

tests such as electroneuromyography or ultrasound. The fact that the treatment is still

centralized, makes it difficult for the patient to adhere, especially those who have subtle

neurological changes that already suggest neural damage, but the patient will hardly seek the

health system if he does not notice the symptoms.

The association of leprosy and neurodynamics issues alerts professionals not involved

with leprosy to suspect leprosy neuropathy when they find a positive neurodynamic test in

their clinical practice. Neurodynamic tests were positive in 7.4% of individuals while there

were still no changes in SNA and later these changes appeared. Future studies could clarify

whether NDA can reveal neural injury earlier.

REFERENCES

1. Boletim. Epidemiológico. Situação epidemiológica e estratégias. Boletim Epidemiológico. Secretaria de Vigilância em Saúde. Ministério da Saúde. 2. ed. Brasília: Ministério da Saúde; 2014. 52 p.

2. Secretaria de Vigilância em Saúde. Ministério da Saúde. Boletim Epidemiológico. Hanseníase 2020; Suppl. Jan. 2020.

3. World Health Organization. Global Leprosy Strategy 2016-2020: Accelerating towards a leprosy-free world. 2016.

4. Guidelines for the diagnosis, treatment and prevention of leprosy. New Delhi: World Health Organization, Regional Office for South-East Asia. 2018

5. Freitas RCSD, Pereira, CAZ, Muri NT, Franzon VAZ, Fatuch CAC. Primary Neural Leprosy mimicking rheumatological disorders. Revista da Associação Médica Brasileira, 2019; 65(6):767-770.

6. Santos DFD, Mendonça MR, Antunes DE, Sabino EFP, Pereira RC, Goulart LR, Goulart IMB. Revisiting primary neural leprosy: Clinical, serological, molecular, and neurophysiological aspects. PLOS Neglected Tropical Diseases 2017;11(11)

7. Ministério da Saúde (Brazilian Ministry of Health). Secretaria de Vigilância em Saúde, Departamento de Vigilância das Doenças Transmissíveis. Diretrizes para vigilância, atenção e eliminação da hanseníase como problema de saúde pública. Brasília: Brazilian Ministry of Health. 2016.

8. Carter GT, Weiss MD. Diagnosis and treatment of work-related proximal median and radial nerve entrapment. Physical Medicine and Rehabilitation Clinics. 2015; 26(3):539-549.

9. Karmakar M, Joshua JM. Changes in plantar load distribution and gait pattern following foot drop correction in leprosy affected patients. LeprRev. 2015; 86: 213–219

10. Rohatgi S, Naveen S, Salunke P, Someshwar S, Rajiv HJ, Joshi R. The story of a deformed leprous foot. LeprRev. 2016; 87: 104–108

11. Júnior LGC, Machado GB, & Faria TA. (2015). Reação hansênica tipo dois em paciente multibacilar, forma Virchowiana, em vigência de tratamento: relato de caso. Revista de Medicina, 94(3), 197-200.

12. Ministério da Saúde (Brazilian Ministry of Health), Office of the Minister. Ordinance No. 149 of February 3, 2016. Approves the Guidelines for Surveillance, Attention and Elimination of Leprosy as a Public Health Problem, with the purpose of guiding managers and professionals of health services [Internet]. Brasília: Brazilian Ministry of Health; 2016 [cited 2020 Sept 01]. Available from: http://bvsms.saude.gov.br/bvs/saudelegis/gm/2016/prt0149 _04022016. html

13. Shacklock M. Neurodynamics. Physiotherapy. 1995; 81(1): 9-16.

14. . Scheibe D, de Oliveira JFP, Gonçalves SB, Gomes ARS, de Macedo ACB.
Mensuração do ângulo articular do cotovelo no teste de tensão neural em indivíduos com
hanseníase. Fisioterapia e Pesquisa. 2012;19(2):165-170

15. Brazil. Ministério da Saúde (Ministry of Health). Secretariat of Health Surveillance.
Department of Surveillance of Transmissible Diseases. Guia prático sobre a hanseníase
[recurso eletrônico] / Ministério da Saúde, Secretaria de Vigilância em Saúde,
Departamento de Vigilância das Doenças Transmissíveis. – Brasília: Ministério da Saúde,
2017. 68 p.: il.

16. Butler DS. Mobilização do sistema nervoso. Manole: Barueri (SP). 2003

17. Herrington, L., Bendix, K., Cornwell, C., Fielden, N., & Hankey, K. (2008). What is
the normal response to structural differentiation within the slump and straight leg raise
tests?. Manual Therapy, 13(4), 289-294.

18. Shacklock, M. O. (1996). Positive upper limb tension test in a case of surgically
proven neuropathy: analysis and validity. Manual Therapy, 1(3), 154-161.

19. Biblioteca Virtual em Saúde – BVS (Virtual Health Library), Atenção Primária em
Saúde. Programa nacional Rede Telessaúde Brasil, Calculadoras, abordagem diagnóstica.
[cited 2020 Sep 1]. Available from: https://aps.bvs.br/apps/calculadoras/?page=1.

20. Ferreira JC, Patino CM. Understanding diagnostic tests. Part 1. J Bras Pneumol. 2017
Sep-Oct;43(5):330. doi: 10.1590/S1806-37562017000000330. PMID: 29160379;
PMCID: PMC5790651.

21. Ferreira Juliana Carvalho, Patino Cecilia Maria. Entendendo os testes diagnósticos.
Parte 3. J. bras. pneumol. [Internet]. 2018 Feb [cited 2020 Aug 18]; 44(1): 4-4. Available
from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1806-
37132018000100004 &lng=en. https://doi.org/10.1590/s1806-37562018000000017.

22. Caminha MDFC, Cruz RDSBL, Acioly VMCD, Nascimento RRD, Azevedo
PTÁCC, Lira PICD, Batista Filho M. Fatores de risco para a não amamentação: um estudo
caso-controle. Revista Brasileira de Saúde Materno Infantil. 2015; 2:193-199.

23. Boas práticas clínicas. Documento das Américas. Organização Pan-americana de
Saúde. Organização Mundial de Saúde, Rede Pan-americana para Harmonização da
Regulamentação Farmacêutica Clínicas. 2005 [cited 2020 May 25].

24. Guyatt GH, Sackett DL, Cook DJ, Guyatt G, Bass E, Brill-Edwards P, Haynes B.
Users' guides to the medical literature: II. How to use an article about therapy or
prevention A. Are the results of the study valid? Jama. 1993;21:270.

25. Esse estudo é válido? PEDro Physiotherapy evidence database. 2020; [cited 2020
may 25]. Avaible from: https://www.pedro.org.au/portuguese/tutorial/is-the-trial-valid/

26. de Macêdo Basso ME, da Silva RLF. Perfil clínico-epidemiológico de pacientes
acometidos pela hanseníase atendidos em uma unidade de referência. Revista da
Sociedade Brasileira de Clínica Médica. 2017;1:27-32.

27. da Silva JSR, Palmeira IP, Sá AMM, Nogueira LMV, Ferreira AMR. Fatores sociodemográficos associados ao grau de incapacidade física na hanseníase. Revista Cuidarte. 2018;9(3): 2338-2348.

28. Finez MA, Salotti SRA. Identificação do grau de incapacidades em pacientes portadores de hanseníase através da avaliação neurológica simplificada. J Health Sci Inst. 2011; 29(3):171-5.

29. Vieira CSCA, Mariano MLL, Silva EMA. Prevalência das deficiências físicas da hanseníase após alta medicamentosa em municípios que alcançaram a meta de eliminação. Hansen Int. 2016. 41 (1-2): p. 46-54.

30. Santos AR, Ignotti E. Prevenção de incapacidade física por hanseníase no Brasil: Análise histórica. Cien Saude Colet [journal on the Internet] (2019/Jan). [cited 2020 may 23]. Avaible from: http://www.cienciaesaudecoletiva.com.br/artigos/prevencao-de-incapacidade-fisica-por-hanseniase-no-brasil-analise-historica/17077?id=17077

31. Nardi SMT, Cruz LPD, Pedro HDSP, Marciano LHSC, Paschoal VDA. Avaliação das deficiências físicas em pessoas com hanseníase empregando dois indicadores: grau de incapacidades e Eyes-Hands-Feet. Hansenologia Internationalis (Online). 2011;36(2): 09-15.

32. De Souza VTC, Junior WMS, De Jesus AMR, De Oliveira DT, Raptis HA, De Freitas PHL, Secheinberg S. Is the WHO disability grading system for leprosy related to the level of functional activity and social participation. Lepr Rev. 2016;87(2): 191-201.

33. Silva JSRD, Palmeira IP, Sá AMM, Nogueira LMV, Ferreira AMR. Variáveis clínicas associadas ao grau de incapacidade física na hanseníase. Revista Cuidarte. 2019; 10(1): Jan-Apr.

34. Thoomes EJ, Van Geest S, Van Der Windt DA, Falla D, Verhagen AP, Koes BW, Vleggeert-Lankamp CL. Value of physical tests in diagnosing cervical radiculopathy: a systematic review. The SpineJournal. 2018; 1:179-189.

35. Koulidis K, Veremis Y, Anderson C, Heneghan NR. Diagnostic accuracy of upper limb neurodynamic tests for the assessment of peripheral neuropathic pain: a systematic review. Musculoskelet Sci Pract. 2019 Apr;40:21-33.

36. Ellis RF, Hing WA. Neural mobilization: a systematic review of randomized controlled trials with an analysis of therapeutic efficacy. Journal of manual & manipulative therapy, 2008. 16(1): 8-22.

37. Onyeonoro UU, Aguocha GU, Madukwe SO, Nwokeukwu HI, Nwamoh UN, Aguocha BU. Pattern of disabilities among leprosy patients in Abia State, Nigeria—a retrospective review. Indian J Lepr. 2016; 88(1), 21-28.

38. Raposo MT, Caminha AVDQ, Heukelbach J, Sánchez-González MÁ, Medeiros JLAD, Nemes MIB. Assessment of physical impairments in leprosy patients: a comparison between the world health organization (WHO) disability grade and the eye-hand-foot score. Revista do Instituto de Medicina Tropical de São Paulo. 2011; 53(2): 77-81.

39. Do Prado GD, Prado RB, Marciano LH, Nardi SM, Cordeiro JA., Monteiro HL.
WHO disability grade does not influence physical activity in Brazilian leprosy
patients. Lepr Rev. 2011. 82(3):270-278.

Table 1. Distribution of the results of the Simplified Neurological Assessment, degree of
hand and foot disabilities, neurodynamic test of the upper and lower limbs in the
intervention group. (N=27)

		LIMBS							
TESTS	**RESULTS**	**Intervention Group**				**Control Group**			
		Upper		**Lower**		**Upper**		**Lower**	
		n	%	n	%	n	%	n	%
Simplified Neurological Assessment	No alteration	5	18.52	2	7.41	27	100	27	100
	Alteration	22	81.48	25	92.59	0	0	0	0
Degree of Physical Disability	No disability (Degree zero)	11	40.74	10	37.04	27	100	27	100
	Disability (Degree I and II)	16	59.26	17	62.96	0	0	0	0
Neurodynamic Assessment	Negative	7	25.93	16	59.26	27	100	27	100
	Positive	20	74.07	11	40.74	0	0	0	0

645   **Table 2**. Measures of sensitivity, specificity, accuracy, positive and negative likelihood
646   ratios for Simplified Neurological Assessment, Neurodynamic Assessment (NDA) and
647   Degree of Physical Disability (DPD) in the upper and lower limbs

648
649

Measures	ULs	SNA Test	NDA Test	DPD Test
	Sensitivity	0.81	0.74	0.59
0-1	Specificity	1	1	1
	Accuracy	0.90	0.87	0.80
1 to 0	NLR	0.18	0.26	0.41
1 to infinity	PLR	Infinite	Infinite	Infinite
Measures	**LLs**	**SNA Test**	**NDA Test**	**DPD Test**
	Sensitivity	0.92	0.41	0.63
0-1	Specificity	1	1	1
	Accuracy	0.96	0.70	0.81
1 to 0	NLR	0.07	0.59	0.37
1 to infinity	PLR	Infinite	Infinite	Infinite

Caption: SNA: Simplified neurological assessment; NDA: Neurodynamic Assessment;
DPD: Degree of Physical Disability; ULs: Upper limbs; LLs: Lower limbs NLR: negative
likelihood ratio; PLR: positive likelihood ratio[43]

Table 3. Degree of hand and foot disabilities compared to Simplified Neurological
Assessment and Neurodynamic Assessment in upper limbs and lower limbs when applied
to the intervention group. (n=27)
657

	Simplified Neurological Assessment	**DEGREE OF HANDS DISABILITIES**				**P-value**
		No Disability		**Disability**		
		N	**%**	**n**	**%**	
UIs	No alteration	5	18.51	0	0.00	**0.010**
	Alteration	6	22.33	16	59.25	
	Neurodynamic Assessment	**No Disability**		**Disability**		
		N	**%**	**n**	**%**	
	Negative	5	18.51	2	7.41	**0.143**
	Positive	6	22.22	14	51.85	

	Simplified Neurological Assessment	**DEGREE OF FEET DISABILITY**				
		No Disability		**Disability**		
		N	**%**	**N**	**%**	
LLs	No alteration	2	7.41	0	0.00	**0.060**
	Alteration	8	29.62	17	62.96	
	Neurodynamic Assessment	**No Disability**		**Disability**		
		N	**%**	**n**	**%**	
	Negative	4	14.81	12	44.44	**0.125**
	Positive	6	22.22	5	18.51	

659 **Table 4.** Simplified Neurological Assessment compared to Neurodynamic Assessment,
660 in the upper and lower limbs, when applied to the intervention group.
661

	NEURODYNAMIC ASSESSMENT	**SIMPLIFIED NEUROLOGICAL ASSESSMENT**				**p-value**
		No Alteration		**Alteration**		
		N	**%**	**n**	**%**	
Upper limbs	Negative	3	11.11	4	14.81	**0.059**
	Positive	2	7.41	18	66.67	
		No Alteration		**Alteration**		
Lower limbs	Negative	1	3.70	15	55.56	**0.786**
	Positive	1	3.70	10	37.04	

663   **Figure 1**. Intersection of three assessments used to check upper limbs (ULs) neurological
664   injuries of individuals with leprosy using the Venn-Euler Diagram.
665

666
667
Caption. SNA: Simplified Neurological Assessment; DPD: Degree of Physical
Disability; NDA: Neurodynamic Assessment. SNA identified neural injury in 22
individuals, NDA in 20 individuals and DPD in 16. The intersection zones correspond to
individuals captured by 2 or 3 assessment instruments. The numbers outside the
intersections mean the individuals who are captured only by that specific assessment. A
= individuals captured only by SNA, B = individuals captured only by NDA, C =
individuals captured only by DPD.

688 **Figure 2**. Intersection of the results obtained from the three assessments used to verify
689 neurological injuries in the lower limbs (LLs) of individuals with leprosy

690
691
Caption. SNA: Simplified Neurological Assessment; DPD: Degree of Physical
Disability; NDA: Neurodynamic Assessment. SNA identified neural injury in 25
individuals, NDA in 11 individuals and DPD in 17. The intersection zones correspond to
individuals captured by 2 or 3 assessment instruments. The numbers outside the
intersections mean the individuals who are captured only by that specific assessment. A=
individuals captured only through SNA, B = individuals captured only through NDA, C
= individuals captured only through DPD
699
700

707 **APPENDIX 1**

709 **MEDICAL SCHOOL OF SÃO JOSÉ DO RIO PRETO**
710 **DATA COLLECTION INSTRUMENT**

	PATIENT'S (PERSON) PROFILE DATA
1	Record Number _________________________ 1( )SJRP 2( )Votuporanga
2	Name ____
3	BirthDate ___ /____/____ Age:_____ Years
4	Sexo: 1( ) male 2( ) female
	OFFICIAL ADDRESS
5	Street____ nº____ CEP:___________________ Phone:__________ _____________
6	Ocupation:_________________________________
	EXCLUSION CRITERIA
7	Diabetic? 1( ) Yes 2 ( )No
8	Compressive syndromes of the central and periferal nervous system? 1( ) Yes 2 ( )No. Which? ___________________________________
9	**Alcoholic** 1( ) Yes 2 ( )No
	LEPROSY DATA
10	Clinical Classification: 1( ) LII 2( ) LTT 3( ) LDT 4( ) LDD 5( ) LDV 6( ) LVV 7( ) Pure Neural
11	MDT (Multidrugtherapy): 1( ) PB 2( ) MB
12	Dates: Start of treatment: ___/___/___ dischargue date of PCT: ___/___/__
13	Pacient's Situation: 1( ) Dsicharge 2( ) in treatment
14	Reacional state: 1( ) yes 2( ) no. If yes, which: ___________________________ 1( ) ENL 2( ) RR 3( ) ENL + RR 4( ) Neuritis 6( ) *Lucius Phenomenon* 7( ) Vasculitis 8( ) Orbaneja

712
713

15	Observe the following scale and report the degree of pain at this point: 0( ) 1( ) 2( ) 3( ) 4( ) 5( ) 6( ) 7( ) 8( ) 9( ) 10 ( ).		
	1st Assess	2nd Assess	3rd Assess

715
716
717
718

719
720   **Simplified Neurological Assesment**
721

		1st Assess		2nd Assess		3rd Assess	
16	**Palpation of Nerves Upper Limbs** - 1: normal; 2: thickening; 3: pain; 4: pain + thickening	R	L	R	L	R	L
	Ulnar						
	Median						
	Radial						
17	**Strength Assessment – Upper Limbs** – 0: Grade Zero; 1: Grade I; 2: Grade II; 3: Grade III; 4: Grade IV; 5: Grade V	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
	5th Finger Abduction						
	Thumb abduction						
	Wrist extension						
18	**Inspection UL:** 1 claw; 2 reabsortion of extremities; 3 contracture; 4 ulcers; 5 Normal	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
19	**Palpation of Nerves Lower Limbs** - 1: normal; 2: thickening; 3: pain; 4: pain + thickening	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
	Fibular						
	Posterior Tibial						
20	**Strength Assessment – Lower Limbs** – 0: Grade Zero; 1: Grade I; 2: Grade II; 3: Grade III; 4: Grade IV; 5: Grade V	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
	Hallux Extension						
	Dorsiflexion						
21	**Inspection LL:** 1 claws; 2 reabsortion of extremities; 3 contracture; 4 ulcers; 5 Normal	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L

Sensitivity assessment of Upper Limbs: (1) Normal sensitivity (2) Loss of sensitivity

22

23 **Sensitivity Assessment of Lower Limbs**

1ª / /		2ª / /		3ª / /	
D	E	D	E	D	E

	DPD – Degree of Physical Disability 1( ) Degree zero – without disability 2( )Degree 1- Loss of sensation and/or loss of strength of hands and feet 3( )Degree 2- Visible deformity (hands and feet drop, claw hands, plantar ulcers, reabsorption of the extremities)					
24		1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
	Hand						
	Foot						

	Passive Neck Flexion (PNF)					
25	Goniometry ( ) 65° 1( ) Positive 2 ( ) Negative	1st Assess		2nd Assess		3rd Assess
	Interviewer's observation					

	Straight Leg Raise (SLR)	1st Assess		2nd Assess		3rd Assess	
		R	L	R	L	R	L
26	Expected Goniometry 50-120°						
	Pain: VAS						
	1 Positive 2 Negative						
	Interviewer's observation						

	Test	**Variable**	1st Assess		2nd Assess		3rd Assess		Expected Goniometry
			R	L	R	L	R	L	
27	ULTT1 MEDIAN	Elbow							145° flexion
		Wrist							70° extension
	Pain (VAS)	Elbow							
		Wrist							
	Positive 1 Negative 2								
	Interviewer's observation								

28	ULTT2b RADIAL	Elbow							145º flexion
		Wrist							90º flexion
	Pain (VAS)	Elbow							
		Wrist							
	Positivo 1 Negativo 2								
Interviewer's observation									
29	ULTT3 ULNAR	Elbow							145º flexion
		Wrist							70º extension
	Pain (VAS)	Elbow							
		Wrist							
	Positive 1 Negative 2								
Interviewer's observation									
30	PKB	Knee + Hip Ext							10º H E+ 140 º Knee Flexion
		Knee							140º Knee Flexion
	Pain (VAS)	Knee + Hip Ext							
		Knee							
	Positive 1 Negative 2								
Interviewer's observation									
31	ULTT3 + ABD H + RI ULNAR	Elbow							145º flexion
		Wrist							70º extension
	Pain (VAS)	Elbow							
		Wrist							
	Positive 1 Negative 2								
Interviewer's observation									

	Slump test	1st assess		2nd assess		3rd assess	
		R	L	R	L	R	L
3 2	Knee Extension Goniometry 0º						
	Pain: VAS						
	1 Positive 2 Negative						
	Interviewer's observation						

723