1   **Isolation thresholds for curbing SARS-CoV-2 resurgence**

2   Laith Yakob

3   Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London,

4   England, WC1E 7HT (laith.yakob@lshtm.ac.uk; +44(0)207 927 2684)

Summary

Self-instigated isolation is heavily relied on to curb SARS-CoV-2 transmission. Accounting for

uncertainty in the latent and prepatent periods, as well as the proportion of infections that remain

asymptomatic, the limits of this intervention at different phases of infection resurgence are

estimated. We show that by October, SARS-CoV-2 transmission rates in England had already begun

exceeding levels that could be interrupted using this intervention alone, lending support to the

second national lockdown on November 5[th].

Main text

A general population lockdown occurred in England on 23[rd] March 2020 to reduce SARS-CoV-2

transmission. This drastic intervention successfully inhibited disease spread by rapidly depleting the

opportunities for transmission events between infected and susceptible people remaining in general

circulation [1].

Subsequent to easing out of lockdown from July 4[th] 2020, infections resurged and England entered

its second national lockdown on November 5[th] 2020. The return of millions of (largely susceptible)

people to general circulation underlies the epidemic re-entering an exponential growth phase.

However, also culpable in the current public health emergency is the failure of interventions during

the period following lockdown's release.

Contact tracing endeavours to reduce SARS-CoV-2 transmission have thus far proven ineffective in

England and so isolation is primarily instigated by those responding to symptoms' development in

themselves or their close associations [2]. The mechanism by which this reactive isolation operates is

importantly distinct from pre-emptive mass quarantine (lockdown). Symptoms-prompted, reactive

isolation only applies to individuals who are infected (c.f. the total population), and, more

specifically, to those who register symptoms. Hence infectious individuals who have not yet

experienced symptoms, or who will never experience them, are missed.

The mathematical epidemiology of reactive isolation is fairly nascent yet critical in the context of the

current epidemic. Here, we generate estimates for reactive isolation thresholds that account for

uncertainties in the latent and pre-patent period of infection as well as in the proportion of infected

individuals that register and respond appropriately to symptoms.

Mathematical derivation of reactive self-isolation

Beginning with the simplest derivation for physical isolation: the pre-emptive quarantine threshold

proportion (Q) is $Q > (1 - (1/R))$ where 'R' is the reproduction number [3]. For reactive isolation (Q^*),

this threshold is inflated to account for the leaked infections occurring because of the delay between

becoming infectious and first exhibiting symptoms: $Q^* > (1 - (1/R)) \times [((g-1)/g)^{-(p-l)}]$. Respectively,

p and l are the prepatent and latent period of infection (in days), and g is the mean duration of

infectiousness (12 days on average [4]). If symptoms typically develop at the same time as an

individual becomes infectious, the square-bracket component equals one and the original threshold

(Q) is regained. A further modification can be made to account for the proportion of infections that

never give rise to symptoms (denoted 'a'): $Q^{**} > (1/(1 - a)) \times (1 - (1/R)) \times [((g-1)/g)^{-(p-l)}]$. For

example, if half of infections remained asymptomatic, the proportion of symptomatic infections that

need to be isolated to achieve an equivalent impact must be doubled. As with those who never

develop symptoms, individuals who fail to respond appropriately to developing symptoms – early

indication is that this is not a negligible proportion [5] – will continue to contribute to transmission,

so 'a' could be considered a composite of these two proportions.

Accounting for uncertainty in parametrization

The latent and prepatent periods are quite variable for COVID-19 patients. Instead of single point

estimates for these parameters, collated data form a distribution of reported times. The latent

period is drawn at random from a Weibull distribution and then subtracted from the random draw

from a second Weibull distribution depicting the range of reported prepatent periods. Fig 1A

illustrates these distributions as informed by the clinical and epidemiological literature [6-8]. Also

shown is the distribution of times between development of infectiousness and symptoms onset as

fitted to 10,000 random draws. The distributions of prepatent and latent periods overlap so to avoid

the possibility of symptoms developing prior to infectiousness, random draws whereby

infectiousness trailed the day of symptoms onset were removed and resampled. 10,000 random

draws were then made from this newly derived distribution of the delay between infectiousness and

symptoms, and the isolation threshold (Q**) was estimated for a range of R values and a range of

asymptomatic proportions (Python code: https://github.com/lwyakob/COVIDquarantine).

Isolation thresholds accounting for uncertainty

Fig 1B shows the mean isolation threshold required to control SARS-CoV-2 accounting for the range

of estimates for the prepatent and latent periods. The value for R is dynamic, varying according to

current intervention effectiveness and population-level susceptibility, so the isolation threshold is

shown for a range of plausible R values. The form of the relationship between Q^* and R shows an

isolation threshold that increases asymptotically with reproduction number. However, allowing for

uncertainty in prepatent and latent periods results in a wide 95% prediction interval. The

interpretation is that when accounting for both the uncertainty in estimating the population mean,

plus the random variation of the individual values, reactive isolation cannot interrupt transmission

(at least 95 times out of 100) if R already exceeds a value of ~2.3.

Figure 1. A) Dashed lines indicate distributions for the latent (blue, Weibull(α=4, β=2)) and prepatent

period (red, Weibull(α=6, β=3)) as derived from the COVID-19 literature [6-8]. The solid line is the

resulting distribution for the time difference between the two from which 10,000 random draws

were made (inset). B) The isolation threshold (Q^*) as calculated for the 10,000 random draws along

with the mean (white line) and 95% predictive interval (dashed lines). The blue cross indicates the

theoretical maximum R number for which reactive isolation may interrupt transmission. C) The

maximum asymptomatic proportion of COVID-19 infections that permits transmission interruption

by reactive isolation for a range of R values (using the expression for Q^{**}). The red boxes illustrate

estimates for England as of October 2020 [9, 10].

Reactive isolation is further limited when asymptomatic infections comprise a non-negligible

proportion (alternatively, when those exhibiting symptoms fail to isolate themselves to some

degree). Fig 1C shows the theoretical limits of the proportion of infections that can be asymptomatic

and yet SARS-CoV-2 transmission interrupted through isolating symptomatic individuals (using the

Q** expression). Superimposed on this trade-off between the reproduction number and the

isolation threshold are estimates for R in England as of October 2020 [10], and the 95% confidence

and predictive intervals for the proportion of infections that remain asymptomatic as generated by a

living systematic review [9]. Respectively, by October 30% and 60% of these parameter spaces were

already beyond the level at which reactive isolation can be sufficient to interrupt transmission (i.e.,

these regions fall to the right of the hatched arc in Fig 1C).

Limitations and future work

One limitation of the current analysis is the consideration of transmission and control at the

population level rather than stratified by various risk factors. To address this, results were generated

for a full range of R values. It is important to note that stratification would impact the derivation of R

but not the population-level isolation thresholds calculated for a given R value [11]. Another

limitation is the implicit assumption that, in the absence of intervention, asymptomatically infected

individuals contribute to onwards transmission as much as symptomatically infected individuals. It is

unclear how questionable this assumption is but clinical studies indicate that asymptomatic and

symptomatic individuals have similar viral loads [12]. Should evidence arise of their differential

contributions to transmission, the model and code associated with this study can be modified easily

to account for this feature.

Even during pre-emptive quarantine (i.e., lockdown) the formulae described here continue to apply

to those who remain in general circulation (e.g., essential personnel). Future work should look at

how isolation thresholds can be estimated to inform this intervention combination, among others.

References

(1)   **Nightingale E, Brady OJ, Yakob L.** The importance of saturating density dependence for
predicting SARS-CoV-2 resurgence. *medRxiv* 2020: 2020.2008.2028.20183921.
(2)   **Moon J, et al.** Optimising 'Test and Trace' Systems: Early lessons from a comparative analysis
of six countries. *SSRN* 2020: http://dx.doi.org/10.2139/ssrn.3694441.
(3)   **Anderson RM, May RM**. *Infectious Diseases of Humans: Dynamics and Control*. Oxford:
Oxford University Press, 1991.
(4)   **He X, et al.** Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature*
Medicine 2020; **26**(5): 672-675.
(5)   **Smith LE, et al.** Factors associated with adherence to self-isolation and lockdown measures
in the UK: a cross-sectional survey. *Public Health* 2020; **187**: 41-52.
(6)   **Nishiura H, Linton NM, Akhmetzhanov AR.** Serial interval of novel coronavirus (COVID-19)
infections. *International Journal of Infectious Diseases* 2020; **93**: 284-286.
(7)   **Backer JA, Klinkenberg D, Wallinga J.** Incubation period of 2019 novel coronavirus (2019-
nCoV) infections among travellers from Wuhan, China, 20–28 January 2020. *Eurosurveillance* 2020;
25(5): 2000062.
(8)   **Kretzschmar ME, et al.** Impact of delays on effectiveness of contact tracing strategies for
COVID-19: a modelling study. *The Lancet Public Health* 2020; **5**(8): e452-e459.
(9)   **Buitrago-Garcia D, et al.** Occurrence and transmission potential of asymptomatic and
presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLOS Medicine*
2020; **17**(9): e1003346.
(10)   **Riley S, et al.** High prevalence of SARS-CoV-2 swab positivity and increasing R number in
England during October 2020: REACT-1 round 6 interim report. *medRxiv* 2020:
2020.2010.2030.20223123.
(11)   **Fine P, Eames K, Heymann DL.** "Herd Immunity": A Rough Guide. *Clinical Infectious Diseases*
2011; **52**(7): 911-916.
(12)   **Lee S, et al.** Clinical Course and Molecular Viral Shedding Among Asymptomatic and
Symptomatic Patients With SARS-CoV-2 Infection in a Community Treatment Center in the Republic
of Korea. *JAMA Internal Medicine* 2020; **180**(11): 1447-1452.

Financial support: The author received no financial support for this work.

Conflicts of interests: None.