HIV infection alters SARS-CoV-2 responsive immune parameters but not clinical outcomes in COVID-19 disease

Farina Karim[1,2 †], Inbal Gazy[2,3 †], Sandile Cele[1,2 †], Yenzekile Zungu[1 †], Robert Krause[1,2 †], Mallory Bernstein[1], Yashica Ganga[1], Hylton Rodel[1,4], Ntombifuthi Mthabela[1], Matilda Mazibuko[1], Khadija Khan[1], Daniel Muema[1,2], Dirhona Ramjit[1], Gila Lustig[5], Thumbi Ndung'u[1,4,6,7], Willem Hanekom[1,4], Bernadett I. Gosnell[9], COMMIT-KZN Team[§], Emily Wong[1,8], Tulio de Oliveira[3], Mahomed-Yunus S. Moosa[9], Alasdair Leslie[1,4*], Henrik Kløverpris[1,4,10*], Alex Sigal[1,2,7*]

[1] *Africa Health Research Institute, Durban 4001, South Africa.* [2] *School of Laboratory Medicine and Medical Sciences, University of KwaZulu-Natal, Durban 4001, South Africa.* [3] *KwaZulu-Natal Research Innovation and Sequencing Platform, Durban 4001, South Africa.* [4] *Division of Infection and Immunity, University College London, London WC1E 6BT, UK.* [5] *Centre for the AIDS Programme of Research in South Africa, Durban 4001, South Africa.* [6] *HIV Pathogenesis Programme, The Doris Duke Medical Research Institute, University of KwaZulu-Natal, Durban 4001, South Africa.* [7] *Max Planck Institute for Infection Biology, Berlin 10117, Germany.* [8] *Division of Infectious Diseases, University of Alabama at Birmingham, Birmingham, AL 35294, USA.* [9] *Department of Infectious Diseases, Nelson R. Mandela School of Clinical Medicine, University of KwaZulu-Natal, Durban 4001, South Africa.* [10] *Department of Immunology and Microbiology, University of Copenhagen, Copenhagen 2200N, Denmark.*

† Equal contribution
§ The names/affiliations of COMMIT-KZN Team members not listed separately appear at end of paper
* Corresponding authors: al.leslie@ahri.org, henrik.kloverpris@ahri.org, alex.sigal@ahri.org

1 Abstract

HIV infection alters the immune response and can compromise protective immunity to multiple pathogens
following vaccination. We investigated the impact of HIV on the immune response to SARS-CoV-2 using
longitudinal samples from 124 participants from KwaZulu-Natal, South Africa, an area of extremely
high HIV prevalence. 44% of participants were people living with HIV (PLWH) and commonly had
other co-morbidities, including obesity, hypertension, and diabetes. The majority of PLWH but not HIV
negative participants showed CD8 T cell expansion above the normal range post-SARS-CoV-2. Yet,
in participants with HIV suppressed by antiretroviral therapy (ART), CD8 expansion was associated
with milder COVID-19 disease. There were multiple differences in T cell, B cell, and natural killer cell
correlations in PLWH compared to HIV negative participants, including lower tissue homing CXCR3+
CD8 T cells in the presence of SARS-CoV-2 RNA in PLWH but not HIV negative and a pronounced
early antibody secreting cell (ASC) expansion in HIV negative but not PLWH. These changes were
COVID-19 associated: low CXCR3 correlated with increased COVID-19 disease severity across groups,
and high ASC correlated with increased disease severity in HIV negative participants and waned when
SARS-CoV-2 was cleared. Despite the altered response of immune cell subsets, COVID-19 disease in
PLWH was mostly mild and similar to HIV negative participants. This likely reflects the heterogeneity
of an effective COVID-19 immune response. Whether the differences in immune cell dynamics in PLWH
will lead to different long-term consequences or compromise vaccination is yet to be determined.

Introduction

More severe COVID-19 disease correlates with lymphopenia and low T cell concentrations [1–3] and mild disease correlates with a robust T cell response to the infection [2, 4–7]. Neutralizing antibodies and associated expansion of antibody secreting B cells are also elicited in most SARS-CoV-2 infected individuals [8–10], although the effectiveness of this response may be partially compromised by the loss of CD4 T follicular helper cells [11] and increased extrafollicular B cell expansion in severe COVID-19 disease [8]. The immune response also includes innate immune cytokine release, including IL6, IL8, and CXCL10/IP10 [5, 12, 13]. IP-10 strongly correlates with severe outcomes [5, 12, 13], and CXCR3, the receptor for IP-10, is known to be involved in antiviral T cell trafficking to tissue compartments [14–16].

Table 1: **Participant Characteristics by HIV status**

	All Participants (n=124)	HIV- (n=69, 55.6%)	HIV+ (n=55, 44.4%)	p-value*
Demographic Characteristics				
Age years, median (IQR)	45.0 (35.0-57.4)	48.8 (34.3-61.8)	41.6 (35.3-50.5)	0.07[#]
Female, n (%)	94 (75.8)	55 (79.7)	39 (70.9)	0.29
Comorbidity, n (%)				
Hypertension	34 (27.4)	23 (33.3)	11 (20.0)	0.11
Diabetes	28 (22.6)	20 (29.0)	8 (14.5)	0.08
Obesity	51 (41.1)	33 (47.8)	18 (32.7)	0.10
History TB	18 (14.5)	3 (4.3)	15 (27.3)	**0.0005**
HIV Associated Parameters				
HIV viremic, n (%)	-	-	13 (23.6)	
Years ART, median (IQR)	-	-	8.3 (3.1-13.1)	
CD4 at enrollment, cells/ml (IQR)	618 (338-974)	880 (543-1140)	497 (279-706)	**<0.0001[#]**
CD4/CD8 ratio at enrollment, (IQR)	1.2 (0.8-1.9)	1.6 (1.2-2.2)	0.8 (0.4-1.2)	**<0.0001[#]**
Disease Severity, n (%)				
Asymptomatic	8 (6.5)	5 (7.2)	3 (5.5)	>0.99
Mild disease	82 (66.1)	46 (66.7)	36 (65.4)	>0.99
Require Supplemental Oxygen	34 (27.4)	18 (26.1)	16 (29.1)	0.84
Symptom, n (%)				
Fever	7 (5.6)	3 (4.3)	4 (7.3)	0.70
Shortness of breath	86 (69.4)	49 (71.0)	37 (67.3)	0.70
Loss of taste and/or smell	56 (45.2)	34 (49.3)	22 (40.0)	0.36
Cough	84 (67.7)	49 (71.0)	35 (63.6)	0.44
Fatigue	68 (54.8)	41 (59.4)	27 (49.1)	0.23
Muscle aches	48 (38.7)	28 (40.6)	20 (36.4)	0.71
Joint ache	49 (39.5)	33 (47.8)	16 (29.1)	**0.04**
Diarrhoea	32 (25.8)	19 (27.5)	13 (23.6)	0.68
Heart palpitations	26 (21.0)	15 (21.7)	11 (20.0)	>0.99
Chest pain or pressure	20 (6.1)	11 (15.9)	9 (16.4)	>0.99
Headache	58 (46.8)	35 (50.7)	23 (41.8)	0.37
Chills	46 (37.1)	28 (40.6)	18 (32.7)	0.45
Sore throat	45 (36.3)	27 (39.1)	18 (32.7)	0.57
Nausea	30 (24.2)	19 (27.5)	11 (20.0)	0.40
Dizziness	29 (23.4)	20 (29.0)	9 (16.4)	0.13

*p-value calculated by Fisher's exact test except for [#] calculated by Mann-Whitney U test.

Less is understood about how HIV modulates immunity to COVID-19. HIV leads to a dysregulation
of T cell responses, including of the highly HIV infectable CD4 T follicular helper cells required for
germinal center formation and antibody affinity maturation [17–19], and causes B cell dysregulation and
dysfunction [20]. Specific T cell parameters such as trafficking, activation, and exhaustion are modulated
by HIV infection [21–23]. HIV co-infection is therefore likely to change the COVID-19 immune response.
Whether and how this occurs is important to understand since HIV is known to interfere with protective
vaccination to multiple pathogens [24–27]. These effects tend to persist despite ART and involve a
sub-optimal antibody response to the vaccine.
Results from epidemiological studies on the interaction of HIV and SARS-CoV-2 are mixed. Several
large studies observed that mortality risk is increased by approximately 1.5 to 3-fold with HIV infection
[28–31] while the majority of studies have found no statistically significant differences in clinical
presentation, adverse outcomes, or mortality in PLWH [31–39].
Here we aimed to determine the effects of HIV on the immune response in the face of COVID-19
in KwaZulu-Natal, South Africa, which may be important in understanding long-term consequences of
COVID-19 infection and the response to vaccination in this population.

Results

We performed a longitudinal observational cohort study to enroll and track patients with a positive
COVID-19 qPCR test presenting at two hospitals. Patients presented because of either COVID-19
symptoms or because they were known contacts of confirmed COVID-19 cases. Enrollment was during the
phase of the KwaZulu-Natal pandemic which peaked in July 2020 (Fig S1). Out of 124 study participants,
55 (44%) were PLWH (Table 1). 94% of study participant were of African descent. Hypertension,
diabetes and obesity, known risk factors for more severe COVID-19 disease [39, 40], were common:
Hypertension and obesity were present in 27%, and 41% of study participants respectively, a similar
prevalence to that reported in KwaZulu-Natal [41, 42]. Diabetes prevalence in our study was 23%,
compared to 13% reported for South Africa [43]. In PLWH, 13 or 24% of PLWH were HIV viremic at
any point in the study. For individuals on ART, median ART duration was 8 years. As expected, both
the CD4 T cell concentration and the CD4 to CD8 T cell ratio was significantly lower in PLWH relative
to HIV negative participants (Table 1).
A minority of study participants were asymptomatic (Table 1), and about a third these were PLWH.
Because some of the study participants were asymptomatic, we used days from diagnostic swab as our
timescale. The date of diagnostic swab was tightly distributed for symptomatic participants at a median
of 3 days post-symptom onset (Figure S1).
The majority of symptomatic participants in the study (67%) did not progress beyond mild disease.
This did not differ between the HIV negative and PLWH (Table 1). 27% of participants required
supplemental oxygen at some point in the COVID-19 disease course. In HIV viremic PLWH, the fraction
of participants requiring oxygen was 38%, but this difference relative to HIV negative was not significant.
Common symptoms in the study population included shortness of breath, loss of taste/smell, cough,
fatigue, muscle aches, and joint aches. There was no significant difference in the frequency of symptoms
between HIV negative and PLWH except for joint aches, which was lower in PLWH.
We next assessed immune cell phenotypes which were responsive to SARS-CoV-2 infection. We
used three approaches to ascertain that we measured a response to SARS-CoV-2 infection, and not
COVID-19 independent differences between PLWH and HIV negative: 1) Detection of temporal changes
post-SARS-CoV-2 infection; 2) detection of changes before versus after SARS-CoV-2 RNA was cleared
from the upper respiratory tract (URT); 3) detection of differences in immune cell phenotypes at different
levels of disease severity. To preserve statistical power, we amalgamated outcomes 4 to 8 in the WHO
ordinal scale and scored disease severity per participant at each time-point as 1: asymptomatic, 2: mild,
and 3: requiring supplemental oxygen.

Figure 1: **Longitudinal SARS-CoV-2 viremia and immune cell response in PLWH and HIV negative participants.** (A) SARS-CoV-2 was detected using qPCR. Plotted is the mean Ct value of SARS-CoV-2 ORF1ab, N gene and S gene. A Ct value of 40, indicated by the dotted red line, was the detection limit. Inconclusive results, with only one SARS-CoV-2 gene detected, were not included. Blue points are HIV negative, red are PLWH with HIV viremia suppressed by ART, purple are HIV viremic participants, where HIV viremia was above the limit of detection at > 40 HIV RNA copies/ml. (B) Concentrations of total lymphocytes (first column), CD8 T cells (second column), CD4 T cells (third column), and the CD4/CD8 T cell ratio (forth column) as a function of time post-diagnostic swab. (C) Percentage of participant samples with concentration of lymphocytes, concentration of CD4 T cells, or the CD4/CD8 T cell ratio below the normal range, or concentration of CD8 T cells above the normal range at the indicated time-point. (D) Concentrations of total lymphocytes, CD8 T cells, CD4 T cells, or the CD4/CD8 T cell ratio in the presence or absence of SARS-CoV-2 RNA. (E) Concentrations of total lymphocytes, CD8 T cells, CD4 T cells, or the CD4/CD8 T cell ratio as a function of disease severity. (F) Percentage of participant samples with concentration of lymphocytes, concentration of CD4 T cells, or the CD4/CD8 T cell ratio below the normal range, or concentration of CD8 T cells above the normal range as a function of disease severity. Disease severity is scored as 1: asymptomatic, 2: mild, and 3: requiring supplemental oxygen. Dashed lines in (B), (D), and (E) denote upper and lower limits of the clinically accepted normal range for the South African population, and red dashed lines denote the lower (for lymphocytes, CD4 T cells, and CD4/CD8 ratio) and upper (CD8 T cells) limits used for panels (C) and (F). p-values are $* < 0.05$; $** < 0.01$; $*** < 0.001$ as determined by Kruskal-Wallis test with Dunn's multiple comparison correction, Mann-Whitney U-test for paired comparisons, or Fisher's exact test for pairwise comparison of frequencies.

To assess whether or not SARS-CoV-2 was cleared from the URT, we measured SARS-CoV-2 RNA
levels using RT-qPCR on combined oropharyngeal and nasophryngeal swabs. We measured frequencies
of immune cell subsets from freshly isolated peripheral blood mononuclear cells (PBMCs). We performed
a total of 387 measurements with up to 5 weekly longitudinal time-points per participant. Time-points
were binned as 0 to 6 days, 7 to 13 days, 14 to 20 days, and 21 days and over post-diagnostic swab,
with a median of 3 longitudinal measurements per study participant (Fig S1). About 60% of study
participants (52% HIV negative, 67% PLWH) who were sampled up to 6 days post-diagnostic swab
had detectable SARS-CoV-2 RNA and this fraction decreased with time with no significant differences
between HIV negative and PLWH (Fig 1A).

Lymphopenia was present early in the course of COVID-19 disease in about 10% of HIV negative and
30% of PLWH (Fig 1B). The total lymphocyte count increased and lymphopenia decreased over time
in both groups (Fig 1C). When the total lymphocyte count was examined in relation to the presence
of SARS-CoV-2 RNA, lymphocyte numbers tended to recover as SARS-CoV-2 RNA was cleared from
URT for both HIV negative and PLWH. This increase was not significant in PLWH who were ART
suppressed (Fig 1D). About 40% of HIV negative and ART suppressed PLWH showed total lymphocyte
numbers below the normal range during more severe COVID-19 disease relative to >5% with mild or
asymptomatic infection (Fig 1E,F). The exception to this trend was PLWH with detectable viremia. In
this group, there was no clear decline in lymphocyte numbers with more severe disease (Fig 1E,F).

There was an increase in the absolute concentration of CD8 T cells post-SARS-CoV-2 infection that
was striking in PLWH (Fig 1B). About 55% of PLWH had abnormally high CD8 T cell concentrations
at the later time-points post-SARS-CoV-2 infection relative to about 15% of HIV negative participants
(Fig 1C). There was a trend towards increasing CD8 T cell numbers once SARS-CoV-2 was cleared, but
this was only significant for the HIV negative group (Fig 1D). CD8 T cell levels were also associated
with disease severity. Participants with time-points showing more severe COVID-19 disease had strongly
decreased CD8 counts relative to mild and asymptomatic participants. This occurred both in the HIV
negative and ART suppressed PLWH groups (Fig 1E,F). These trends were absent in the viremic PLWH,
where CD8 T cells counts were elevated regardless of SARS-CoV-2 clearance or COVID-19 disease
severity.

The CD4 T cell subset showed a more attenuated increase post-SARS-CoV-2 infection relative to
CD8 T cells in PLWH. In HIV negative participants, the concentration of CD4 T cells was relatively
constant as a function of time from SARS-CoV-2 infection (Fig 1B,C). However, CD4 T cell concentration
did increase after SARS-CoV-2 RNA clearance in the HIV negative and HIV suppressed PLWH (Fig
1D). Similarly to CD8 T cells, the number of CD4 cells was dramatically lowered with more severe

COVID-19 disease relative to when disease was asymptomatic or mild in HIV negative participants or ART suppressed PLWH. In contrast, CD4 levels were low regardless of COVID-19 disease state in HIV viremic (Fig 1E,F). The ratio of CD4 to CD8 T cells is often used as a prognostic indicator in PLWH [44]. The CD4/CD8 ratio was lowered throughout the measured SARS-CoV-2 infection course in PLWH (Fig 1B), with the majority of PLWH showing an abnormally low CD/CD8 (Fig 1C). There was also no change in the CD4/CD8 ratio as a function of time post-SARS-CoV-2 infection and with disease severity (Fig 1D-F).

Figure 2: CD8+ T-cell phenotype changes in HIV negative and PLWH during SARS-CoV-2 infection. (A) Gating strategy for CD8+ T-cell subsets. (B) CXCR3 expression on CD8 T-cells as a function of HIV status, time in days from diagnostic swab, presence or absence of SARS-CoV-2 RNA, and disease severity. Blue points are HIV negative, red are PLWH with HIV viremia suppressed by ART, purple are HIV viremic where HIV viremia was above the limit of detection at > 40 HIV RNA copies/ml, and green are all participants. Disease severity is scored as 1: asymptomatic, 2: mild, and 3: requiring supplemental oxygen. (C) HLA-DR or (D) PD-1 expression on CD8 T-cells as a function of HIV status, time in days from diagnostic swab, and presence or absence of SARS-CoV-2 RNA. p-values are * <0.05; ** <0.01; *** < 0.001 as determined by Kruskal-Wallis test with Dunn's multiple comparison correction or Mann-Whitney U-test for paired comparisons.

115 We also examined T cell canonical phenotypes, including CXCR3 involved in T cell trafficking,
HLA-DR for immune activation, and PD-1 for immune exhaustion (Fig 2A, Fig S2, Fig S3A). We found
similar CD8 T cell CXCR3 expression in HIV negative compared to ART suppressed PLWH, but lower
expression in HIV viremic PLWH (Fig 2B). When we investigated expression as a function of time, there
was a significant increase in CXCR3 expressing CD8 T cells with time post-SARS-CoV-2 infection in
HIV negative individuals that was delayed in PLWH. CXCR3 expression was lowered in PLWH in the
presence of SARS-CoV-2 RNA and increased to the level of expression similar to that in HIV negative
upon SARS-CoV-2 clearance. In contrast, in HIV negative, CXCR3 expression on CD8 T cells was not
strongly affected by the presence of SARS-CoV-2 (Fig 2B). When we correlated CXCR3 expression on
CD8 T cells with disease severity, we found reduced levels in more severe disease in PLWH and when the
HIV negative and PLWH groups were amalgamated (Fig 2B). CD4 T cells showed similar patterns (Fig
S3B).

Figure 3: **Changes in B cell phenotypes in HIV negative and PLWH during SARS-CoV-2 infection.**
(A) CD19+ B cells were gated on CD27 and CD38 and included naïve (CD27-), memory (CD27+CD38lo/-) and
antibody secreting cells (ASC, CD27+CD38+). (B) Total CD19 count and frequencies of naïve, memory, and
ASC in the CD19+ population. (C) The ASC response as a function of time post-diagnostic swab, presence of
SARS-CoV-2 RNA, and disease severity. Disease severity is scored as 1: asymptomatic, 2: mild, and 3: requiring
supplemental oxygen. Blue points are HIV negative, red are PLWH with HIV viremia suppressed by ART, purple
are HIV viremic, and green are all participants. p-values are * <0.05; ** <0.01; *** <0.001 as determined by
Kruskal-Wallis test with Dunn's multiple comparison correction or Mann-Whitney U-test for paired comparisons.

To analyse T cell activation and exhaustion, we examined HLA-DR and PD-1. We observed elevated
HLA-DR expression in CD8 T cells, indicating increased activation, in HIV viremic PLWH relative to
HIV negative (Fig 2C). In CD4 T cells, the increased activation in ART suppressed PLWH relative
to HIV negative became significant (Fig S3C). CD8 T cell activation increased with time in the HIV
negative group but not in PLWH (Fig 2C). We found no significant changes in the expression of the
exhaustion marker PD-1 on T cells during longitudinal SARS-CoV-2 sampling irrespective of HIV status
in CD8 (Fig 2D) or CD4 (Fig S3D) T cells. We also examined CCR7 expression, predominantly expressed
on naïve T cells, and this showed no change in CD8 and CD4 T cells (Fig S4, Fig S3E).
We examined the longitudinal B cell response in PLWH versus HIV negative participants. CD19+
B cells were analysed for the B cell maturation markers CD27 and CD38, to identify naïve (CD27-),
memory (CD27+CD38-) and antibody secreting cell (ASC, CD27+CD38++) populations (Fig 3A).
The concentration of CD19+ B cells was lower in viremic PLWH, but not those on suppressive ART,

compared to HIV negative (Fig 3B). ART suppressed PLWH displayed more naïve and fewer memory B cells. However, this likely represents a pre-existing state in PLWH since there was no significant change as a function of time post-SARS-CoV-2 infection (Fig S5).

There was no difference overall in the fraction of ASCs between the HIV negative and PLWH groups (Fig 3B). However, ASCs were significantly expanded at the earliest time-point post-SARS-CoV-2 infection for both HIV negative and PLWH. The expansion was more pronounced in HIV negative individuals and decreased with time (Fig 3C). Consistent with this, ASC were significantly elevated in individuals with detectable SARS-CoV-2 compared to those who have cleared the virus, and this association was stronger in HIV negative relative to PLWH (Fig 3C). We also observed an association between increased disease severity and a higher fraction of ASCs. However, and consistent with a reduced ASC response in PLWH, this effect was only significant in the HIV negative group (Fig 3C).

Figure 4: **Immune cell and clinical correlates in HIV negative and PLWH.** Spearman rank correlation values (ρ) are shown from red (1.0) to blue (-1.0). p-values per correlation are *< 0.5; **< 0.01; ***< 0.001. The number of matched pairs for HIV negative participants ranged from 80 to 125 and for PLWH from 48 to 81. Rectangles represent regions where a set of correlations is present in one participant group (solid rectangles) and absent in the other (dashed rectangles). Black dashed lines represent the divide between clinical and cellular parameters.

We also determined the ratio of plasma cells to plasmablasts based on CD138 expression on ASCs. Plasmablasts are short-lived ASCs which may either die or terminally differentiate into antibody secreting plasma cells. We observed a non-statistically significant trend for a reduced plasma cell to plasmablast ratio in PLWH at the earliest time-points (Fig S6). This trend, however, was not apparent when participants were grouped according to SARS-CoV2 viremia (Fig S6). Taken together, the CXCR3 and ASC results show that HIV infection can skew the T cell and B cell responses in SARS-CoV2 infection.

To examine the differences in the immune response more broadly, we calculated the correlations between measured phenotypes and clinical parameters in HIV negative and PLWH. Overall, we observed different significant correlations between T cell, B cell, and natural killer (NK) cell phenotypic markers, as well as disease severity in PLWH compared to HIV negative individuals (Fig 4).

For HIV negative participants, there were significant negative and positive correlations between CD4 T cell parameters, and between these and the CD8 T cell count and the CCR7+CD8 T cell fraction (Fig

4, orange box). There were also correlations between CD4 and CD8 T cell phenotypes and CD56+CD16+ NK cells (Fig 4, purple box). These correlations were lost in PLWH.

New correlations arose in PLWH, particularly involving CD8 T cells: CXCR3+CD8 T cells became negatively correlated with disease severity (Fig 4 black box). CD8 T cell activation (HLA-DR+) also gained correlations with CD19 B cell phenotypes (green box), and the plasma cell to plasmablast ratio gained correlations to CD4 and CD8 T cell phenotypes (Fig 4, blue box).

Taken together, these differing correlations may point to an overall alternative regulation of immune cell subsets in response to SARS-CoV-2 in PLWH, despite the majority of PLWH having undetectable HIV viral loads due to effective suppression by ART.

Discussion

We have shown that HIV infection changed the immune cell response but not clinical outcomes or symptoms of COVID-19 infection. We did not pre-select individuals with severe COVID-19 disease, for which HIV has been reported as a risk factor in South Africa [28]. The prevalence of HIV in our study of 44% is high relative to the overall prevalence in KwaZulu-Natal (18%, [45]) but may be roughly consistent with the expected prevalence given the age and gender distribution of the study participants [46].

Immune cell subsets which we observed to be responsive to SARS-CoV-2 infection showed differences between the HIV negative and PLWH groups. This included: 1) A CD8 T cell expansion in PLWH above the clinically accepted normal limit in a majority of PLWH (Fig 1); 2) a sharp decrease in the tissue homing marker CXCR3 in the presence of SARS-CoV-2 in PLWH (Fig 2); 3) an attenuated expansion of ASCs in PLWH relative to HIV negative participants (Fig 3). We also found an association between low CXCR3 expression on CD8 and CD4 T cells and COVID-19 disease severity, which was strongest in PLWH. To our knowledge, This association has not been previously described. It is consistent with the known role of the CXCR3 ligand CXCL10/IP-10 in the COVID-19 immune response [5, 12, 13] and the role of CXCR3 in the antiviral response in other infections [14–16]. Other studies have reported a positive correlation between ASC frequency and COVID-19 disease severity [6, 8]. However, consistent with a reduced ASC response in HIV co-infected participants, this effect was only significant in the HIV negative group (Fig 3). Importantly, the correlations between the T cell and B cell phenotypes and disease severity were different between HIV negative and PLHW (Fig 4).

We used three approaches determine if the changes in immune cell phenotypes were a response to SARS-CoV-2 infection, and not COVID-19 independent differences between HIV negative and PLWH. First, we detected changes as a function of time post-SARS-CoV-2 infection. Second, we detected changes before versus after SARS-CoV-2 RNA was cleared. Third, we detected changes as a function of COVID-19 disease severity. A limitation of the study is we did not examine antigen specific responses. Further studies, using techniques such as analysis of activation induced marker (AIM) assays for T cells, could add insight into the SARS-CoV-2 specific response [2, 4–7, 47].

We also observed that HIV viremia changes the immune response to COVID-19 in PLWH, including consistently elevated CD8 T cells levels regardless of presence or absence of SARS-CoV-2 RNA or the degree of disease severity (Fig 1), and lack of recovery of CD4 T cells after SARS-CoV-2 clearance (Fig 1). Therefore, effective ART suppression would be expected to play a role in attenuating the effects of HIV infection on COVID-19 immune response.

Participants in this study generally showed mild COVID-19 outcomes, despite high frequencies of co-morbidities and HIV infection. Given that COVID-19 infection outcomes were similar in PLWH relative to HIV negative participants, the differences in immune response between the groups may indicate an alternative, as opposed to dysregulated, immunity to SARS-CoV-2 in PLWH. One indication of this is that although CD8 T cells were elevated to abnormally high numbers in PLWH in response to COVID-19, the lack of such expansion correlated with a worse COVID-19 infection outcome in PLWH. The clinical consequences of this remain unclear, but should be considered in the long-term repercussions of COVID-19 infection and the response to a vaccine.

Material and Methods

Ethical statement and study participants

The study protocol was approved by the University of KwaZulu-Natal Institutional Review Board (approval BREC/00001275/2020). Adult patients (>18 years old) presenting either at King Edward VIII or Clairwood Hospitals in Durban, South Africa, between 8 June to 25 September 2020, diagnosed to be SARS-CoV-2 positive as part of their clinical workup and able to provide informed consent were eligible for the study. Written informed consent was obtained for all enrolled participants.

Clinical laboratory testing

An HIV rapid test and viral load quantification was performed from a 4ml EDTA tube of blood at an accredited diagnostic laboratory (Molecular Diagnostic Services, Durban, South Africa) using the RealTime HIV negative1 viral load test on an Abbott machine. CD4 count, CD8 count, and a full blood count panel were performed by an accredited diagnostic laboratory (Ampath, Durban, South Africa).

qPCR detection of SARS-CoV-2

RNA was extracted from combined oropharyngeal and nasophryngeal swabs from 140 μl viral transport medium using the QIAamp Viral RNA Mini kit (cat. no. 52906, QIAGEN, Hilden, Germany) according to manufacturer's instructions, and eluted into 100 μl AVE buffer. To detect SARS-CoV-2 RNA, 5 μl RNA was added to the TaqPath 1-step RT-qPCR mastermix. 3 SARS-CoV-2 genes (ORF1ab, S and N) were amplified using the TaqPath COVID-19 Combo Kit and TaqPath COVID-19 CE-IVD RT-PCR Kit (ThermoFisher Scientific, Massachusetts, United States) in a QuantStudio 7 Flex Real-Time PCR system (ThermoFisher Scientific). Data was analysed using the Design and Analysis software (ThermoFisher Scientific). For positive samples, Ct values are represented as the average of the Ct values of all three genes. A sample was scored positive where at least 2 out of the 3 genes were detected, and inconclusive if only 1 of the genes was detected.

PBMC isolation and immune phenotyping by flow cytometry

PBMCs were isolated by density gradient centrifugation using Histopaque 1077 (Sigma-Aldrich, St. Louis, Missouri, United States) and SepMate separation tubes (STEMCELL Technologies, Vancouver, Canada). For T cell and NK cell phenotyping, 10^6 fresh PBMCs were surface stained in 50l antibody mix with the following antibodies from BD Biosciences, (Franklin Lakes, NJ, USA): anti-CD45 Hv500 (1:100 dilution, clone HI30, cat. 560777); anti-CD8 BV395 (1:50 dilution, clone RPA-T8, cat. 563795); anti-CD4 BV496 (1:25 dilution, clone SK3, cat. 564651); anti-PD1 BV421 (1:50 dilution, clone EH12.1, cat. 562516); anti-CXCR3 PE-CF594 (1:25 dilution, clone 1C6/CXCR3, cat. 562451). The following antibodies were from BioLegend (San Diego, CA, USA): anti-CD19 Bv605 (1:100 dilution, clone HIB19, cat. 302244); anti-CD16 Bv650 (1:50 dilution, clone 3G8, cat. 302042); anti-CD56 Bv711 (1:50 dilution, clone HCD56, cat. 318336); anti-CD3 Bv785 (1:25 dilution, clone OKT3, cat. 317330); anti-CXCR5 FITC (1:25 dilution, clone J252D4, cat. 356914); anti-HLA-DR PE (1:50 dilution, clone L243, cat. 307606); anti-CCR7 PerCP-Cy5.5 (1:25 dilution, clone G043H7, cat. 353220); anti-CD38 PE-Cy7 (1:25 dilution, clone HIT2, cat. 303516); anti-ICOS APC (1:25 dilution, clone C398.4A, cat. 313510) and anti-CD45RA AF700 (1:25 dilution, clone HI100, cat. 304120). PBMCs were incubated with antibodies for 20 minutes at room temperature. For B-cell phenotyping, the following antibodies were used: (all from BioLegend) anti-CD45 APC (1:25 dilution, clone HI30, cat. 304012); anti-CD3 Bv711 (1:50 dilution, clone OKT3, cat. 317328), anti-CD14 Bv711 (1:25 dilution, clone M5E2, cat. 301838); anti-CD19 Bv605 (1:50 dilution, clone HIB19, cat. 302244); anti-CD27 Hv500 (1:50 dilution, clone O323, cat. 302836); anti-CD38 PE-Cy7 (1:25 dilution, clone HIT2, cat. 303516) and anti-CD138 BV785 (1:25 dilution, clone MI15, cat. 356538). Cells were then washed twice in PBS and fixed in 2% paraformaldehyde and stored at 4^{o}C before acquisition on FACSAria Fusion III flow cytometer (BD) and analysed with FlowJo software version 9.9.6 (Tree Star).

Statistical analysis

Data is described with the non-parametric measures of median and interquartile range, and significance determined using the non-parametric Mann-Whitney U test for pairwise comparisons, Fisher exact test for pairwise comparisons of frequencies, and the Kruskal-Wallis test with multiple comparison correction by the Dunn Method for comparisons involved more than two populations. All tests were performed using Graphpad Prism 8 software.

Acknowledgements

This work was supported by the Bill and Melinda Gates Investment INV-018944 to AS.

COMMIT-KZN Team

Moherndran Archary, Department of Paediatrics and Child Health, University of KwaZulu-Natal

Kaylesh J. Dullabh, Department of Cardiothoracic Surgery, University of KwaZulu-Natal

Jennifer Giandhari, KwaZulu-Natal Research Innovation and Sequencing Platform

Philip Goulder, Africa Health Research Institute and Department of Paediatrics, Oxford

Guy Harling, Africa Health Research Institute and the Institute for Global Health, University College London

Rohen Harrichandparsad, Department of Neurosurgery, University of KwaZulu-Natal

Kobus Herbst, Africa Health Research Institute and the South African Population Research Infrastructure Network

Prakash Jeena, Department of Paediatrics and Child Health, University of KwaZulu-Natal

Thandeka Khoza, Africa Health Research Institute

Nigel Klein, Africa Health Research Institute and the Institute of Child Health, University College London

Richard Lessells, KwaZulu-Natal Research Innovation and Sequencing Platform

Rajhmun Madansein, Department of Cardiothoracic Surgery, University of KwaZulu-Natal

Mohlopheni Marakalala, Africa Health Research Institute and Division of Infection and Immunity, University College London

Mosa Moshabela, College of Health Sciences, University of KwaZulu-Natal

Kogie Naidoo, Centre for the AIDS Programme of Research in South Africa

Zaza Ndhlovu, Africa Health Research Institute and the Ragon Institute of MGH, MIT and Harvard

Kennedy Nyamande, Department of Pulmonology and Critical Care, University of KwaZulu-Natal

Nesri Padayatchi, Centre for the AIDS Programme of Research in South Africa

Vinod Patel, Department of Neurology, University of KwaZulu-Natal

Theresa Smit, Africa Health Research Institute

Adrie Steyn, Africa Health Research Institute and Division of Infectious Diseases, University of Alabama at Birmingham

References

1. Lucas, C., Wong, P., Klein, J., Castro, T. B., Silva, J., Sundaram, M., Ellingson, M. K., Mao, T., Oh, J. E., Israelow, B., *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* **584,** 463–469 (2020).

2. Sekine, T., Perez-Potti, A., Rivera-Ballesteros, O., Strålin, K., Gorin, J.-B., Olsson, A., Llewellyn-Lacey, S., Kamal, H., Bogdanovic, G., Muschiol, S., *et al.* Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *Cell* (2020).

3. Chen, G., Wu, D., Guo, W., Cao, Y., Huang, D., Wang, H., Wang, T., Zhang, X., Chen, H., Yu, H., *et al.* Clinical and immunological features of severe and moderate coronavirus disease 2019. *The Journal of clinical investigation* **130** (2020).

4. Grifoni, A., Weiskopf, D., Ramirez, S. I., Mateus, J., Dan, J. M., Moderbacher, C. R., Rawlings, S. A., Sutherland, A., Premkumar, L. & Jadi, R. S. Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell.* ISSN: 0092-8674 (2020).

5. Moderbacher, C. R., Ramirez, S. I., Dan, J. M., Grifoni, A., Hastie, K. M., Weiskopf, D., Belanger, S., Abbott, R. K., Kim, C., Choi, J., *et al.* Antigen-specific adaptive immunity to SARS-CoV-2 in acute COVID-19 and associations with age and disease severity. *Cell* (2020).

6. Mathew, D., Giles, J. R., Baxter, A. E., Oldridge, D. A., Greenplate, A. R., Wu, J. E., Alanio, C., Kuri-Cervantes, L., Pampena, M. B., D'Andrea, K., *et al.* Deep immune profiling of COVID-19 patients reveals distinct immunotypes with therapeutic implications. *Science* **369** (2020).

7. Mateus, J., Grifoni, A., Tarke, A., Sidney, J., Ramirez, S. I., Dan, J. M., Burger, Z. C., Rawlings, S. A., Smith, D. M., Phillips, E., *et al.* Selective and cross-reactive SARS-CoV-2 T cell epitopes in unexposed humans. *Science* **370,** 89–94 (2020).

8. Woodruff, M. C., Ramonell, R. P., Nguyen, D. C., Cashman, K. S., Saini, A. S., Haddad, N. S., Ley, A. M., Kyu, S., Howell, J. C., Ozturk, T., Lee, S., Suryadevara, N., Case, J. B., Bugrovsky, R., Chen, W., Estrada, J., Morrison-Porter, A., Derrico, A., Anam, F. A., Sharma, M., Wu, H. M., Le, S. N., Jenks, S. A., Tipton, C. M., Staitieh, B., Daiss, J. L., Ghosn, E., Diamond, M. S., Carnahan, R. H., Crowe, J. E., Hu, W. T., Lee, F. E.-H. & Sanz, I. Extrafollicular B cell responses correlate with neutralizing antibodies and morbidity in COVID-19. *Nature Immunology.* ISSN: 1529-2916. https://doi.org/10.1038/s41590-020-00814-z (2020).

9. Robbiani, D. F., Gaebler, C., Muecksch, F., Lorenzi, J. C., Wang, Z., Cho, A., Agudelo, M., Barnes, C., Finkin, S., Hagglof, T., *et al.* Convergent Antibody Responses to SARS-CoV-2 Infection in Convalescent Individuals. *bioRxiv* (2020).

10. Quinlan, B. D., Mou, H., Zhang, L., Guo, Y., He, W., Ojha, A., Parcells, M. S., Luo, G., Li, W., Zhong, G., *et al.* The SARS-CoV-2 receptor-binding domain elicits a potent neutralizing response without antibody-dependent enhancement. *IMMUNITY-D-20-00389* (2020).

11. Kaneko, N., Kuo, H.-H., Boucau, J., Farmer, J. R., Allard-Chamard, H., Mahajan, V. S., Piechocka-Trocha, A., Lefteri, K., Osborn, M., Bals, J., *et al.* Loss of Bcl-6-expressing T follicular helper cells and germinal centers in COVID-19. *Cell* (2020).

12. Laing, A. G., Lorenc, A., Del Barrio, I. D. M., Das, A., Fish, M., Monin, L., Muñoz-Ruiz, M., McKenzie, D. R., Hayday, T. S., Francos-Quijorna, I., *et al.* A dynamic COVID-19 immune signature includes associations with poor prognosis. *Nature medicine* **26,** 1623–1635 (2020).

13. Yang, Y., Shen, C., Li, J., Yuan, J., Wei, J., Huang, F., Wang, F., Li, G., Li, Y., Xing, L., *et al.* Plasma IP-10 and MCP-3 levels are highly associated with disease severity and predict the progression of COVID-19. *Journal of Allergy and Clinical Immunology* (2020).

14. Groom, J. R. & Luster, A. D. CXCR3 in T cell function. *Experimental cell research* **317,** 620–631 (2011).

15. Stiles, L. N., Hosking, M. P., Edwards, R. A., Strieter, R. M. & Lane, T. E. Differential roles for CXCR3 in CD4+ and CD8+ T cell trafficking following viral infection of the CNS. *European journal of immunology* **36,** 613–622 (2006).

16. Zhang, B., Chan, Y. K., Lu, B., Diamond, M. S. & Klein, R. S. CXCR3 mediates region-specific antiviral T cell trafficking within the central nervous system during West Nile virus encephalitis. *The Journal of Immunology* **180,** 2641–2649 (2008).

17. Okoye, A. A. & Picker, L. J. CD 4+ T-cell depletion in HIV infection: mechanisms of immunological failure. *Immunological reviews* **254,** 54–64 (2013).

18. Pallikkuth, S., Parmigiani, A., Silva, S. Y., George, V. K., Fischl, M., Pahwa, R. & Pahwa, S. Impaired peripheral blood T-follicular helper cell function in HIV-infected nonresponders to the 2009 H1N1/09 vaccine. *Blood, The Journal of the American Society of Hematology* **120,** 985–993 (2012).

19. Perreau, M., Savoye, A.-L., De Crignis, E., Corpataux, J.-M., Cubas, R., Haddad, E. K., De Leval, L., Graziosi, C. & Pantaleo, G. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1 infection, replication, and production. *Journal of Experimental Medicine* **210,** 143–156 (2013).

20. Moir, S. & Fauci, A. S. Insights into B cells and HIV-specific B-cell responses in HIV-infected individuals. *Immunological reviews* **254,** 207–224 (2013).

21. Day, C. L., Kaufmann, D. E., Kiepiela, P., Brown, J. A., Moodley, E. S., Reddy, S., Mackey, E. W., Miller, J. D., Leslie, A. J., DePierres, C., *et al.* PD-1 expression on HIV-specific T cells is associated with T-cell exhaustion and disease progression. *Nature* **443,** 350–354 (2006).

22. Deeks, S. G., Kitchen, C. M., Liu, L., Guo, H., Gascon, R., Narváez, A. B., Hunt, P., Martin, J. N., Kahn, J. O., Levy, J., *et al.* Immune activation set point during early HIV infection predicts subsequent CD4+ T-cell changes independent of viral load. *Blood* **104,** 942–947 (2004).

23. Mavigner, M., Cazabat, M., Dubois, M., L'Faqihi, F.-E., Requena, M., Pasquier, C., Klopp, P., Amar, J., Alric, L., Barange, K., *et al.* Altered CD4+ T cell homing to the gut impairs mucosal immune reconstitution in treated HIV-infected individuals. *The Journal of clinical investigation* **122,** 62–69 (2012).

24. Avelino-Silva, V. I., Miyaji, K. T., Mathias, A., Costa, D. A., de Carvalho Dias, J. Z., Lima, S. B., Simoes, M., Freire, M. S., Caiaffa-Filho, H. H., Hong, M. A., *et al.* CD4/CD8 ratio predicts yellow fever vaccine-induced antibody titers in virologically suppressed HIV-infected patients. *JAIDS Journal of Acquired Immune Deficiency Syndromes* **71,** 189–195 (2016).

25. Carson, P. J., Schut, R. L., Simpson, M. L., O'Brien, J. & Janoff, E. N. Antibody class and subclass responses to pneumococcal polysaccharides following immunization of human immunodeficiency virus-infected patients. *Journal of infectious diseases* **172,** 340–345. ISSN: 1537-6613 (1995).

26. Cooper, C., Thorne, A., Klein, M., Conway, B., Boivin, G., Haase, D., Shafran, S., Zubyk, W., Singer, J., Halperin, S., *et al.* Immunogenicity is not improved by increased antigen dose or booster dosing of seasonal influenza vaccine in a randomized trial of HIV infected adults. *PloS one* **6,** e17758 (2011).

27. Fuster, F., Vargas, J. I., Jensen, D., Sarmiento, V., Acuña, P., Peirano, F., Fuster, F., Arab, J. P., Martınez, F., Soto, S., *et al.* CD4/CD8 ratio as a predictor of the response to HBV vaccination in HIV-positive patients: A prospective cohort study. *Vaccine* **34,** 1889–1895 (2016).

28. Davies, M.-A. *et al.* HIV and risk of COVID-19 death: a population cohort study from the Western Cape Province, South Africa. *medRxiv* (2020).

29. Geretti, A. M., Stockdale, A. J., Kelly, S. H., Cevik, M., Collins, S., Waters, L., Villa, G., Docherty, A., Harrison, E. M., Turtle, L., *et al.* Outcomes of COVID-19 related hospitalisation among people with HIV in the ISARIC WHO Clinical Characterisation Protocol UK Protocol: prospective observational study (2020).

30. Bhaskaran, K., Rentsch, C. T., MacKenna, B., Schultz, A., Mehrkar, A., Bates, C., Eggo, R. M., Morton, C. E., Bacon, S., Inglesby, P., *et al.* HIV infection and COVID-19 death: population-based cohort analysis of UK primary care data and linked national death registrations within the OpenSAFELY platform. *medRxiv* (2020).

31. Vizcarra, P., Pérez-Elıas, M. J., Quereda, C., Moreno, A., Vivancos, M. J., Dronda, F., Casado, J. L., Moreno, S., Pérez-Elıas, M. J., Fortún, J., *et al.* Description of COVID-19 in HIV-infected individuals: a single-centre, prospective cohort. *The Lancet HIV* (2020).

32. Huang, J., Xie, N., Hu, X., Yan, H., Ding, J., Liu, P., Ma, H., Ruan, L., Li, G., He, N., *et al.* Epidemiological, virological and serological features of COVID-19 cases in people living with HIV in Wuhan City: A population-based cohort study. *Clinical Infectious Diseases* (2020).

33. Sigel, K., Swartz, T., Golden, E., Paranjpe, I., Somani, S., Richter, F., De Freitas, J. K., Miotto, R., Zhao, S., Polak, P., *et al.* COVID-19 and people with HIV infection: outcomes for hospitalized patients in New York City. *Clinical Infectious Diseases* (2020).

34. Shalev, N., Scherer, M., LaSota, E. D., Antoniou, P., Yin, M. T., Zucker, J. & Sobieszczyk, M. E. Clinical Characteristics and Outcomes in People Living With Human Immunodeficiency Virus Hospitalized for Coronavirus Disease 2019. *Clinical Infectious Diseases* (2020).

35. Stoeckle, K., Johnston, C. D., Jannat-Khah, D. P., Williams, S. C., Ellman, T. M., Vogler, M. A., Gulick, R. M., Glesby, M. J. & Choi, J. J. *COVID-19 in Hospitalized Adults With HIV* in *Open Forum Infectious Diseases* **7** (2020), ofaa327.

36. Dandachi, D., Geiger, G., Montgomery, M. W., Karmen-Tuohy, S., Golzy, M., Antar, A. A., Llibre, J. M., Camazine, M., Santiago, D., Carlucci, P. M., *et al.* Characteristics, Comorbidities, and Outcomes in a Multicenter Registry of Patients with HIV and Coronavirus Disease-19. *Clinical Infectious Diseases* (2020).

37. Haerter, G., Spinner, C. D., Roider, J., Bickel, M., Krznaric, I., Grunwald, S., Schabaz, F., Gillor, D., Postel, N., Mueller, M. C., *et al.* COVID-19 in people living with human immunodeficiency virus: a case series of 33 patients. *Infection,* 1 (2020).

38. Karmen-Tuohy, S., Carlucci, P. M., Zervou, F. N., Zacharioudakis, I. M., Rebick, G., Klein, E., Reich, J., Jones, S. & Rahimian, J. Outcomes among HIV-positive patients hospitalized with COVID-19. *Journal of acquired immune deficiency syndromes (1999)* (2020).

39. Richardson, S., Hirsch, J. S., Narasimhan, M., Crawford, J. M., McGinn, T., Davidson, K. W., Barnaby, D. P., Becker, L. B., Chelico, J. D., Cohen, S. L., *et al.* Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *Jama* (2020).

40. Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J., Wang, Y., Song, B. & Gu, X. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The lancet.* ISSN: 0140-6736 (2020).

41. Van Heerden, A., Barnabas, R. V., Norris, S. A., Micklesfield, L. K., van Rooyen, H. & Celum, C. High prevalence of HIV and non-communicable disease (NCD) risk factors in rural KwaZulu-Natal, South Africa. *Journal of the international AIDS society* **20,** e25012 (2017).

42. Malaza, A., Mossong, J., Bärnighausen, T. & Newell, M.-L. Hypertension and obesity in adults living in a high HIV prevalence rural area in South Africa. *PloS one* **7,** e47761 (2012).

43. Federation, I. D. IDF diabetes atlas ninth edition 2019 (2019).

44. Serrano-Villar, S. & Deeks, S. G. CD4/CD8 ratio: an emerging biomarker for HIV. *The lancet HIV* **2,** e76–e77 (2015).

45. Simbayi, L., Zuma, K., Zungu, N., Moyo, S., Marinda, E., Jooste, S., Mabaso, M., Ramlagan, S., North, A., Van Zyl, J., *et al.* South African National HIV Prevalence, Incidence, Behaviour and Communication Survey, 2017: towards achieving the UNAIDS 90-90-90 targets (2019).

46. Kharsany, A. B., Cawood, C., Khanyile, D., Lewis, L., Grobler, A., Puren, A., Govender, K., George, G., Beckett, S. & Samsunder, N. Community-based HIV prevalence in KwaZulu-Natal, South Africa: results of a cross-sectional household survey. *The Lancet HIV* **5,** e427–e437. ISSN: 2352-3018 (2018).

47. Reiss, S., Baxter, A. E., Cirelli, K. M., Dan, J. M., Morou, A., Daigneault, A., Brassard, N., Silvestri, G., Routy, J.-P., Havenar-Daughton, C., *et al.* Comparative analysis of activation induced marker (AIM) assays for sensitive identification of antigen-specific CD4 T cells. *PLoS One* **12,** e0186998 (2017).

Supplementary Figures

Fig S 1: **Study characteristics.** (A) Study timeline. (B) Time from symptom onset or diagnostic swab to study enrollment for symptomatic participants. Left panel shows time elapsed for HIV negative (blue points) and PLWH (red points) between symptom onset and enrollment. Middle panel shows days from the time the diagnostic swab was taken to enrollment for HIV negative and PLWH. Right panel shows time difference between symptom onset and diagnostic swab. (C) Percentage of participants attending follow-up visits for all participants (green), HIV negative (blue), and PLWH (red). (D) Number of visits (from a total of 5 study time-points per participant) for each participants. Median and IQR for all participants (green), HIV negative (blue), and PLWH (red).

Fig S 2: **CD8+ T-cell phenotype marker distribution.** Relative distribution of each phenotype marker measured simultaneously as a proportion of total CD8+ T-cells as arc pie chart (left) or plot with each point representing one sample (right) and y-axis showing frequency of each marker combination.

Fig S 3: CD4+ T-cell phenotype changes in HIV negative and PLWH during SARS-CoV-2 infection. (A) Gating strategy for CD4+ T-cell subsets. (B) CXCR3 expression on CD4 T-cells as a function of HIV status, time in days from diagnostic swab, presence or absence of SARS-CoV-2 RNA, and disease severity. Blue points are HIV negative, red are PLWH with HIV viremia suppressed by ART, purple are HIV viremic, and green are all participants. Disease severity is scored as 1: asymptomatic, 2: mild, and 3: requiring supplemental oxygen. (C) HLA-DR or (D) PD-1, or CCR7 (E) expression on CD4 T-cells as a function of HIV status, time in days from diagnostic swab, and presence or absence of SARS-CoV-2 RNA. p-values are determined by Kruskal-Wallis test with Dunn's multiple comparison correction or Mann-Whitney U-test for paired comparisons.

● All ● HIV- ● HIV+ (all) ● HIV+ suppressed ● HIV+ viremic

Fig S 4: **CCR7 expression on CD8+ T-cells in HIV negative and PLWH during SARS-CoV-2 infection.** (A) Gating strategy. (B) CCR7 expression on CD8 T-cells as a function of HIV status, time in days from diagnostic swab, presence or absence of SARS-CoV-2 RNA, and disease severity. Blue points are HIV negative, red are PLWH with HIV viremia suppressed by ART, purple are HIV viremic, and green are all participants. Disease severity is scored as 1: asymptomatic, 2: mild, and 3: requiring supplemental oxygen. p-values are determined by Kruskal-Wallis test with Dunn's multiple comparison correction or Mann-Whitney U-test for paired comparisons.

Fig S 5: **No significant changes in the frequencies of memory and naive B cells with time in HIV negative and PLWH.** Fraction of CD19 positive cells which are naive (A) or memory (B) B cells. Green points are all participants, blue are HIV negative, and red are PLWH.

Fig S 6: **The plasma cell to plasmablast ratio post-SARS-CoV-2 infection.** (A) Gating strategy for differentiating ASCs into CD138+ plasma cells (PC) and CD138- plasmablasts (PB). (B) the PC to PB ratio as a function of time post-SARS-CoV-2 infection and in the presence or absence of SARS-CoV-2 RNA in URT. Blue points are HIV negative, red are PLWH.

Fig S 7: **NK cell frequency with time post-SARS-CoV-2 infection.** (A) Gating strategy from live/CD45+/CD3-/CD19- lymphocytes for CD56hiCD16lo and CD56midCD16hi NK cells. (B) Frequency of CD56hiCD16lo and (C) CD56midCD16hi NK cells in HIV negative (left, blue points) and HIV positive (right, red points) at different times post-diagnostic swab. Horizontal bars represent median values.