

The economic value of quarantine is higher at lower case prevalence, with quarantine justified at lower risk of infection

James Petrie^{1,2,*} and Joanna Masei^{2,3}

¹Applied Mathematics, University of Waterloo; ²WeHealth Solutions PBC; ³Ecology & Evolutionary Biology, University of Arizona; * corresponding author
jimpetrie@uwaterloo.ca

1. Abstract

Some infectious diseases, such as COVID-19, are so harmful that they justify broad scale social distancing. Targeted quarantine can reduce the amount of indiscriminate social distancing needed to control transmission. Finding the optimal balance between targeted vs. broad scale policies can be operationalized by minimizing the total amount of social isolation needed to achieve a target reproductive number. Optimality is achieved by quarantining on the basis of a risk threshold that depends strongly on current disease prevalence, suggesting that very different disease control policies should be used at different times or places. Aggressive quarantine is warranted given low disease prevalence, while populations with a higher base rate of infection should rely more on social distancing by all. The total value of a quarantine policy rises as case counts fall, is relatively insensitive to vaccination unless the vaccinated are exempt from distancing policies, and is substantially increased by the availability of modestly more information about individual risk of infectiousness.

2. Introduction

Control of SARS-CoV-2 transmission has not been achieved by interventions such as improved ventilation alone, but has also required social distancing. Social distancing can take the form either of population-wide measures, ranging from extreme, mandatory lockdowns to more modest voluntary behavior change, or it can be targeted via effective testing, tracing, quarantine, and isolation. Social distancing to control the spread of SARS-CoV-2 has imposed immense social costs, and is difficult to maintain for long periods. Targeting quarantine to those at higher risk of being infectious has the potential to reduce the total harms of social distancing across the population. Here we calculate which combination of population-wide social distancing plus targeted quarantine will minimize harms while controlling transmission. We operationalize this question by comparing the harms of different policy combinations that all achieve the same target value for the effective reproduction number. Figure 1 demonstrates how quarantining a greater fraction of infectious individuals allows overall social distancing to be reduced while holding the number of expected transmissions constant.




Fig. 1. Two scenarios of quarantine and social distancing that give rise to the same number of expected transmissions. A) One out of three infectious individuals is quarantined, and the general population is at 1/3 of normal social activity. B) Two out of three infectious individuals are quarantined, and the general population is at 2/3 of normal social activity. By finding and quarantining infectious people, social distancing imposed on the general population is significantly reduced.

Our approach, at the interface of infectious disease epidemiology and economics (1), solves a number of problems previously encountered at that interface. First, it is difficult to decide on an exchange rate between social restrictions / livelihoods versus lives (2); compounding this is the fact that selfish agents may count only the cost to their own life, and not that of others they infect (3). Difficulties determining an exchange rate lead some to present only idealized curves without a concrete recommendation (4, 5). Idealized curves can lead to abstract insights, but are of limited use to practical decision-making.

We avoid the need for an exchange rate between social restrictions and infections by instead comparing one form of (population-wide, partial) social restrictions with another form of (targeted, more stringent) social restrictions. This allows the lowest cost strategy to be selected without having to monetarize the cost of health outcomes, the latter being held

constant in our analysis. The policy optimization problem is thus framed as a minimization of a cost function across different degrees of isolation for different individuals.

Second, much prior work integrates across an entire sweep of a pandemic modelled using an SIR approach (6). These projections have often been inaccurate. We instead focus on a moment in time, an approach that is enabled by a control theory view. At the beginning of the COVID-19 pandemic, populations experienced rapid exponential growth, e.g. doubling in as little as 2.75 days in New York City (7). Through some combination of top-down control measures and individual behavior modifications, many populations subsequently achieved relatively flat case counts, i.e. an effective R value remarkably close to 1 (8, 9). Even locations that subsequently lost control of the pandemic have tended to eventually issue stay at home orders leading to exponential decline. Over the very long term, a geometric mean of R not much greater than 1 is inevitable (see Appendix A). This implies that there is a form of control system (whether intentional or not) using social distancing to regulate transmission rate.

Third, many models consider decision-making that is informed by perfect, instantaneous knowledge (6). This is problematic given that key indicators such as hospital usage or even positive tests have a marked lag relative to infections. Our approach focuses instead on making optimal use of available, probabilistic information. Past economic approaches to modelling the transmission of SARS-CoV-2 treat more stark examples of information, e.g. testing to find out who is infected and should isolate, as opposed to who is exposed to what degree and should quarantine (6, 10). But quantitative information about individual risk can be obtained from a variety of sources. For example, the risk of infection with SARS-CoV-2 can be estimated based on proximity and duration of contact with a known case, and their estimated infectiousness as a function of timing relative to symptom onset date (11). Similarly, setting local rather than global shutdown policies, in the light of differences between regions, can lower costs (12).

Our work ultimately describes the value of information, specifically information about who is at high enough risk of being infected to quarantine strictly rather than merely to conform to population-wide restrictions. Given an individual estimated to be infectious with probability ‘r’, we propose a method for deciding whether to quarantine this individual. We do so by weighing the cost of quarantine against the degree to which indiscriminately applied social distancing would need to be increased to achieve the same reduction in transmission. Our approach informs choice of the lowest cost strategy needed to achieve epidemiological targets. In particular, we formalize the intuition that populations with low prevalence should widen the net of who they quarantine.

3. Model

A. Optimal Risk Threshold. Consider a well-mixed population of size P , of which I people are currently both infected and infectious, and S people are susceptible. According to a standard continuous, deterministic SIR model approach (13), the effective reproduction number $R_t = R_0 * S/P$ depends on the fraction of people still susceptible (S/P). Here we define R_0 and R_t to explicitly exclude interventions such as social distancing or quarantine, because these are the interventions whose magnitudes are being optimized. However, R_t is in-

tended to include fixed cost interventions such as improved ventilation.

Population-wide social distancing is parameterized as a value D that varies between 0 and 1 such that the reproductive number is proportional to $(1 - D)$. I.e., a value of $D = 0$ indicates normal social contact, while $D = 1$ indicates complete isolation. Reductions in D come from a variety of behavior changes including working from home and reducing social contact, and also includes measures to mitigate the danger given contact, such as wearing masks, meeting outside and/or at greater physical distance.

Let Q_i denote the number of infectious people who are quarantined or isolated and Q_n denote the number of non-infectious people who are quarantined. The benefit from a targeted quarantine policy depends on $\frac{I - Q_i}{I}$, the fraction of infectious cases who are not successfully quarantined and then isolated. The effective reproduction number R'_t after both social distancing and quarantine interventions are applied is given by:

$$R'_t = R_t * (1 - D) * \frac{I - Q_i}{I} \quad [1]$$

R'_t gives the expected number of onward transmissions per infected case in the general population. Equation 1 is undefined when the number of locally transmitted cases $I = 0$. Even for a low but non-zero value of I , a different, stochastic treatment warranted - this is discussed in Appendix B.

The per-individual cost of reduced contact is represented by the function $f(x)$, where x is the individual counterpart to population-wide D , and where $f(x)$ is assumed to be strictly increasing (from 0 cost with normal contact, to the maximum cost with complete isolation). The exact functional form of $f(x)$ is unknown, but in Model Section B we show that the results have a bounded dependence on its form. The total cost to a population is given by Equation 2, based on $(Q_i + Q_n)$ people in complete isolation, and $(P - Q_i - Q_n)$ people with contact reduced by D .

$$f(1)(Q_i + Q_n) + f(D)(P - Q_i - Q_n) \quad [2]$$

The amount of social distancing needed to meet a target reproductive number, R_{target} , given fixed values for Q_i and Q_n , is given by:

$$D = \max(0, 1 - \frac{R_{target}}{R_t} \frac{I}{I - Q_i}) \quad [3]$$

This value is bounded at 0; if $R_t \frac{I - Q_i}{I} \leq R_{target}$, the target has already been met and no social distancing is required.

In the real world, it is not possible to adjust D instantaneously or exactly in response to changes in the impact of quarantine policy. However, in Appendix A, we show that Equation 3 is insensitive to time and control uncertainty, so long as caseload and control measures are fairly steady over time, as has been the case in many regions for at least some substantial period of time (8, 9).

Combining Equations 2 and 3, the total cost of a quarantine policy, in combination with a degree of population-wide distancing that keeps caseload steady, is given by:

$$J(Q_i, Q_n) = f(1)(Q_i + Q_n) + f(\max(0, 1 - \frac{R_{target}}{R_t} \frac{I}{I - Q_i}))(P - Q_i - Q_n) \quad [4]$$

170 This equation can be used to compare the overall cost reduction
171 from different quarantine policies.

172 We assume that an estimated risk of infectiousness is avail-
173 able when considering whether to recommend quarantine to
174 an individual. Risk could be estimated using proximity and
175 duration of contact with a known case as described in (11).
176 Or it could be estimated for members of a subpopulation like
177 a workplace by rapid testing of a random sample of that sub-
178 population and projecting the proportion positive onto the
179 remainder. When a person with risk of infectiousness r is
180 quarantined, Q_i is incremented by 1 with probability r , and
181 Q_n is incremented by 1 with probability $1 - r$. Quarantining
182 this person is worthwhile if the expected change in cost given
183 by Equation 5 is less than 0:

$$184 \quad rJ(Q_i + 1, Q_n) + (1 - r)J(Q_i, Q_n + 1) - J(Q_i, Q_n) < 0 \quad [5]$$

185 Solving the corresponding equation for r , the optimal risk
186 threshold is given by:

$$187 \quad r_{\text{thresh}} = \frac{J(Q_i, Q_n) - J(Q_i, Q_n + 1)}{J(Q_i + 1, Q_n) - J(Q_i, Q_n + 1)} \quad [6]$$

188 To apply Equation 6, we first consider the simplest equation
189 for the cost of isolation, namely the linear function $f(x) = x$.
190 Using this equation in a situation where some social distancing
191 is necessary ($\frac{R_{\text{target}}}{R_t} * \frac{I}{I - Q_i - 1} < 1$), Equation 6 simplifies to:

$$192 \quad r_{\text{thresh}} = \frac{I - Q_i - 1}{P - Q_i - Q_n - 1} \quad [7]$$

193 In this simple case, the interpretation of Equation 7 is
194 straightforward and intuitive. The optimal risk threshold for
195 quarantine is equal to the disease probability in the community,
196 excluding from that community both those already isolated or
197 quarantined, and the focal individual. This implies that in re-
198 gions with very low prevalence rates, much stricter quarantine
199 requirements should be used, making the general population
200 less impacted by disease-control measures. In populations
201 with higher disease prevalence in the community, there is less
202 benefit to quarantining a single individual at low risk, because
203 they represent a smaller fraction of the overall disease risk to
204 the population.

205 **B. Sensitivity to Isolation Cost Function.** We do not know the
206 shape of the isolation cost function $f(x)$ as the extent of
207 isolation x varies. However, some non-linearity is expected.
208 E.g., to achieve total rather than partial isolation can require
209 a specialized quarantine facility in addition to a more extreme
210 loss of individual utility. D is defined such that benefits to
211 disease transmission are linear in D . When considering a
212 reduction in the extent of social activity, benefits are likely
213 somewhere between linear and quadratic (5). When people
214 seek out places that have variable levels of crowding, benefits
215 are quadratic; when they seek out people who are still willing
216 to meet with them, benefits are linear. By defining D to
217 make benefits linear in D , we pass this uncertainty to the cost
218 function. Disparate considerations all suggest a concave-up
219 shape for $f(x)$.

220 In Appendix C, we show that the deviation from Equation
221 7 is relatively small for reasonable choices of $f(x)$. In particu-
222 lar, if the threshold r_1 is computed using $f_1(x) = x$ and the

223 threshold r_2 is computed using any $f_2(x)$ that is strictly in-
224 creasing and concave up, then $r_2 \geq r_1$ and $\frac{r_2}{r_1} \leq \frac{\max_{x \in [0,1]} f_2'(x)}{\min_{x \in [0,1]} f_2'(x)}$.
225 In practice, the value of r_2 is often much closer to r_1 than
226 indicated by the upper bound. For example, while the ratio be-
227 tween optimal risk thresholds for the two cost functions shown
228 in Figure 2 is bounded at a 10-fold difference, we numerically
229 find it to be at most 4.1-fold.




Fig. 2. Non-linearity in the cost of isolation (A) leads to a modest four-fold change in optimal risk threshold, or less, depending for example on the proportion of infectious people quarantined (B).

230 Both the cost of isolation and the benefit of isolation can vary
231 among individuals. For example, the cost of isolation is higher for
232 essential workers than for those that can easily work from home,
233 while the benefit of isolation is higher for those whose jobs expose
234 more and/or more vulnerable people. Equation 7 can be modified to
235 accommodate knowledge about an individual. If isolation is m times
236 more expensive for this person than for the general population, and
237 the potential danger to others if they are actually infectious is d times
238 that of an average case, then the risk threshold for this person
239 should be modified to $r_{\text{thresh}} * m/d$. Note that this approach differs
240 substantially from suggestions to target isolation directly to the
241 elderly and other high medical risk individuals (5). We instead
242 recommend increasing the stringency of quarantine among their
243 contacts.
244

For a given individual, the cost of quarantine is likely to depend more than linearly on quarantine length, due both to logistical and to psychological factors. To accommodate this, the cost of quarantine could be modelled as time-dependent by scaling $f(x) * m(t)$. The benefit of quarantine is also not equal for each possible day of quarantine, given a probability distribution for incubation period and a quantitative timecourse of infectiousness (11). Consideration of these two factors will lead shorter recommended quarantine durations, targeted to the most infectious days. Testing individuals in quarantine could further shorten its duration (14).

The remarkable thing about Equation 7 is its insensitivity to R_t and R_{target} . To determine the degree to which this might generalize, in Appendix B, we derive Equation 8:

$$\frac{1}{r_{thresh}} \frac{f(1) - f(D_0)}{1 - D_0} \approx \frac{1}{r_{pop}} f'(D_0) \quad [8]$$

Equation 8 shows that r_{thresh} is chosen so that the risk-weighted cost of quarantine is equal to the marginal cost of distancing for the general population. Here, r_{thresh} depends indirectly on R_t and R_{target} because they influence the set point, D_0 . However, for a linear cost function, the slope ($f'(D_0)$) and average slope ($\frac{f(1)-f(D_0)}{1-D_0}$) are constant, and therefore r_{thresh} only depends on the disease prevalence in the general population. In other words, when the cost per individual is a linear function of distancing, then the cost is the same whether that distancing is all directed at making one individual isolate completely instead of following current distancing norms, versus distributing the additional distancing among all individuals. For a non-linear function, these two costs are no longer equal. D_0 is proportional to R_{target}/R_t , and so Equation 8 describes the degree to which non-linearity makes r_{thresh} depend on these terms. This dependence will be most pronounced when the local slope of $f(x)$ around current distancing levels and the mean slope from current distancing to $D = 1$ are substantially different.

Because the optimal risk threshold isn't very sensitive to the cost function, we focus our analysis on the more convenient linear cost function (with solution given by Equation 7).

4. Results

A. Applications in time and place. In a large population with many cases, the risk threshold for quarantine r_{thresh} is closely related to the base rate of infection in the population. Here we give several examples in which we apply this finding, to illustrate how dramatically this causes optimal policy to vary across different places and times.

In British Columbia in October 2020, rapid testing was widespread, but only when symptoms were present, making official case counts of around 100/day a good estimate of symptomatic cases, such that total cases can be estimated in proportion. (When testing is less adequate, estimating the true number of cases is more complex, involving projections from death rates and/or testing rates, but has been attempted e.g. at (8) or (15).) Assuming another 20 asymptomatic or other undiscovered cases per day, each with a 10 day infectious window, yields 200 non-isolated infectious individuals on any given day. In addition, we estimate 4 infectious days per discovered case prior to isolation, yielding another 400 non-isolated infectious individuals on any given day. The base rate of infectiousness, conditional on not having been isolated,

is thus of order $600/5,000,000 = O(10^{-4})$. From Equation 7, a quarantine risk threshold of 10^{-4} is appropriate, or a little higher if a non-linear cost function is assumed in Equation 6. This implies that quarantine recommendations for entire schools or workplaces are worthwhile if they are expected to isolate one more infected person than would be achieved via traditional contact tracing.

In contrast, North Dakota in October 2020 had around 400 detected cases per day and perhaps another ~ 400 undetected in a population of $\sim 800,000$. This yields ~ 5000 undetected infectious individuals on a given day, or a 0.6% base rate and corresponding risk threshold. Several studies have found the secondary attack rate to be on the order of $1/10 - 1/100$ (16–18), so this risk threshold aligns well with standard guidelines for the minimum contact considered to be close for contact tracing purposes. We note however that a 14 day quarantine is more than is needed to put the conditional probability of infectiousness, given lack of symptoms, of a non-household close contact below the base rate (11).

Imported and exported cases introduce complications. Policy is best set for regions that are relatively self-contained, and do not e.g. cut off commuter communities from one another. When imported and exported cases make up only a small fraction of total cases, then subtleties in using case counts to estimate R'_t (discussed in Appendix B) have negligible effects. When local transmissions are abundant enough such that elimination is not a near-term goal, then incoming individuals can therefore be quarantined by comparing their infectiousness risk to r_{thresh} on the basis of base rates of infection in the location they arrived from, plus the risk of travel. These considerations apply e.g. to the need to quarantine following travel from one part of the U.S. to another.

Some locations, e.g. British Columbia, achieved very low levels of local transmission, but with a degree of economic connectedness with harder hit locations that made infeasible sufficiently strict quarantine to put the risk from incomers below r_{thresh} as described above (19). This is because the marginal cost of quarantine applies not just to the quarantined individuals, but to trucking routes and other aspects of the economic system. At this point, some significant expected number of imported cases acts as a forcing function outside the exponential dynamics of R'_t . In this case, it might be better to set a target value for the expected number of cases per day, and then to set R_{target} to achieve this in combination with imported cases.

If local elimination is achieved, including successfully quarantining all imported cases, then R_{target} can be relaxed up to R_t , removing all social distancing, with huge social gain. The disproportionate benefits warrant careful attention to avoid even a single imported non-quarantined case. How best to achieve this is best quantified by a stochastic model, outside the scope of this paper. Qualitatively, we note that the cost of letting a single case slip through, and hence having to return to social distancing, is higher for larger populations. The harm is also magnified by delays in realizing that an outbreak is underway, and hence the extent of harm depends strongly on local surveillance and contact tracing capabilities.

When different regions come under shared political control, importation risk can be controlled at the source rather than the destination. An example is Vietnam in March 2020. Viewed as an entire country instead of as smaller regions, it is optimal

364 to quarantine at a threshold of $1/10^6$. This would motivate
 365 quarantining entire cities so that the rest of the country doesn't
 366 have to substantially modify their behaviour. This is similar
 367 to the approach Vietnam actually took, with up to 80,000
 368 people in quarantine at a time from regional lockdowns and
 369 aggressive contact tracing (20).

370 **B. Quantifying the Value of Quarantine.** The value of a test,
 371 trace, and quarantine policy can be quantified in terms of
 372 the reduction it enables in full or partial person-days of isola-
 373 tion across a population. By definition, the marginal value of
 374 quarantining an individual of marginal risk r_{thresh} is 0. This
 375 implies, for a cost function that is uniform across individuals
 376 and neglects complications due to compliance, that quarantining
 377 one definitely infectious person for one day yields a benefit
 378 of C/r_{thresh} where e.g. C is the assessed cost of quarantine.
 379 Taking into account the quarantined individual, the net benefit
 380 is $C/(1/r_{thresh} - 1)$.

381 In many countries, compliance not just with social dis-
 382 tancing, but also with quarantine has been low. This is to
 383 be expected, given the uncompensated cost of quarantine to
 384 individuals, who are currently asked (or in cases coerced) to
 385 sacrifice for the public good. We advocate that governments
 386 align incentives, by guaranteeing e.g. 150% the individual's
 387 normal daily income. This approach also serves to convert
 388 the costs and benefits calculated here, whose units are of
 389 person-days of isolation, to a dollar value for quarantine. On
 390 the basis of median income, and taking into account that
 391 some individuals are able to work from home and so will not
 392 draw the full 150%, this would in an affluent nation amount
 393 to something on the order of USD\$150 per day of complete
 394 quarantine. This is the implied utility sacrificed by either total
 395 quarantine or equivalent social distancing D . I.e., a dollar
 396 value on the of order USD\$150 per day is what one would need
 397 to pay people in order to incentivize their compliance, i.e. to
 398 achieve neutrality among preferences. Because this neutrality
 399 between quarantine plus payment vs. no payment and no
 400 quarantine does not include the dangers posed by COVID-19
 401 itself, the existence of such payments would not incentivize
 402 deliberate exposure. Redistributive payments move costs from
 403 the quarantined individuals to the taxpayer.

404 With this cost applying to both indiscriminate distancing
 405 and targeted quarantining, we can assess the value of a quar-
 406 antine policy from its ability to reduce the total amount of
 407 distancing required to achieve R_{target} . Given the changing na-
 408 ture of a pandemic, we calculate how this total value depends
 409 on case prevalence and on the extent of population immunity.

410 When case prevalence is low, quarantining a single indi-
 411 vidual makes a larger proportional difference to the risk to
 412 others, enabling a greater relaxation of social distancing. We
 413 see in Figure 3B that in the pertinent range of 0.1%-5% case
 414 prevalence, the value of effective quarantine policies increases
 415 as case counts drop. Results are similar for the differently
 416 shaped distributions of assessable risk shown in Figure 3A.
 417 For lower case prevalence, the value flattens out because the
 418 benefit from quarantining someone infectious rises at the same
 419 rate that the number of total infectious people drops, while
 420 the number of infectious contacts quarantined per index case
 421 remains almost constant.

422 Vaccination has enormous value in reducing the amount
 423 of distancing required to achieve R_{target} . But unless the
 424 immune are exempted from distancing norms, the marginal




Fig. 3. The benefits from quarantine policies are highest when case prevalence is low. Each case one serial interval prior has on average ten contacts, of whom one is infected. We assume that case investigation and contact tracing jointly succeed in reaching 10% of all contacts. A) Assessable risk among contacts is modeled as a gamma distribution, in all cases with mean 0.1, and with different shape parameters to explore the importance of resolution among low-risk individuals. B) $Q_i = 0.4I$, $Q_n = 4Q_i$, $C = \$150$, $R_t = 4$, $R_{target} = 1$, $f(x) = x$.

425 value of quarantine, in helping sustain R_{target} with even less
 426 distancing, is unchanged. The effective reproductive number
 427 before including distancing, R_t , is defined to be proportional
 428 to the susceptible fraction, S/P . Under a linear cost function,
 429 we see from Equation 7 that the threshold for quarantine is
 430 insensitive to R_t . This means that the value of a quarantine
 431 policy is also insensitive to R_t and hence to this change in
 432 the vaccination status of the population. However, if people
 433 who have immunity are exempt from social distancing, not
 434 only is the amount of social distancing per person reduced,
 435 but also the number of people who have to distance. This
 436 has an additional impact on r_{thresh} : the effective size of the
 437 population is reduced from P to S . Reducing the effective size
 438 of the population increases r_{thresh} by replacing P with S
 439 in the denominator of Equation 7.

440 In Figure 4, the value of an example exposure notification
 441 system is shown as a function of the susceptible fraction of
 442 the population. If the immune are required to social distance,
 443 the benefit of the program is independent of the proportion




Fig. 4. With vaccinated people exempt from distancing, the benefit of a contact tracing system is approximately proportional to the susceptible fraction (dashed line). When distancing is required, with the immune following the same distancing norms as the susceptible, the benefit does not depend on the susceptible fraction (solid line). $I/P = 0.0055$, $Q_i = 0.4 * I$, $Q_n = 4 * Q_i$, $C = \$150$, $R_0 = 4$, $R_{target} = 1$. Assumptions about contact tracing are as for Figure 3.

444 immune. If the immune are exempt, the benefit of the program
 445 is nearly proportional to the susceptible fraction. Intuitively,
 446 this is because quarantining infectious cases only allows an
 447 increase in social activity for the susceptible.

448 **C. Marginal Value of Information.** With a method available to
 449 assess the value of a quarantine policy, we can now assess the
 450 value of information in enabling better quarantine/isolation
 451 policies. We first consider random surveillance using a test with
 452 50% sensitivity and near-100% specificity (the latter achieved
 453 via a follow-up PCR test). The value per test is given by the
 454 probability of testing positive * expected number of additional
 455 infectious days in isolation * daily benefit of each additional
 456 isolation day. If a positive test leads to 3 extra isolation days,
 457 and assuming a large enough population to neglect the -1
 458 term in Equation 7, the direct benefit (not including onward
 459 contact tracing) per test is $\frac{I-Q_i}{P-Q_i+Q_n} * 0.5 * 3 * \$150/r_{thresh}$.
 460 With $r_{thresh} \approx \frac{I-Q_i}{P-Q_i+Q_n}$, the per-test benefit is insensitive
 461 to the risk threshold and equal to $\approx \$225$. With a non-linear
 462 cost function, the benefit might be up to ten times less.

463 This makes random surveillance clearly worthwhile for rapid
 464 testing technologies that can be made available for as little as
 465 \$5 per test (21). The benefit of surveillance testing further
 466 increases with effective contact tracing the ability to quaran-
 467 tine close contacts provides additional benefit. The benefit
 468 is reduced if a fraction of people who test positive do not
 469 effectively isolate.

470 When testing is too expensive to be used for random surveil-
 471 lance, or merely too slow to roll out at sufficient scale, it can
 472 achieve higher marginal value for populations at higher risk.
 473 We note that it can be hard to justify strict quarantine on indi-
 474 viduals of modestly above-average risk when the risk threshold
 475 is low in absolute terms, especially when elimination does not
 476 yet seem within reach. An alternative is daily testing instead

of quarantine. The economic benefit from testing increases 477
 with risk of the focal individual relative to the average, and 478
 can thus justify more expensive and more sensitive daily tests. 479

D. Application to Exposure Notification. Exposure notifica- 480
 tion use smartphone apps to rapidly recommend quarantine, 481
 while using sophisticated algorithms to assess risk. Here we 482
 consider the economic value of using a better rather than 483
 a worse risk assessment algorithm, comparing choices that 484
 were implemented in different jurisdictions for version 1 of the 485
 Google/Apple Exposure Notification system. 486

We simulate a range of encounters using the COVID Tuner 487
 risk score configuration tool (22), following a 3D grid with 488
 distance distributed uniformly between 0 and 5 meters, dura- 489
 tions uniformly between 5 and 30 minutes, and date relative to 490
 symptom onset date uniformly between -10 days and +10 days. 491
 This tool, on the basis of the current public health definition of 492
 close contact (2 metres for 15 minutes from day -2 to day +9) 493
 estimates an implicit risk threshold of 0.002. We simulate the 494
 Swiss, Irish, and Arizonan configurations included in this tool, 495
 plus a “baseline” configuration that is equivalent to the Arizo- 496
 nan configuration but with only two rather than 6 non-zero 497
 levels of infectiousness. The Swiss and Irish configurations use 498
 only one non-zero level of infectiousness. 499

We assume that the cost of quarantine is \$150 per day, 500
 and that the average quarantine is 10 days long. This gives 501
 an expected cost of quarantine of \$1500 per individual that 502
 a particular configuration succeeds in getting to quarantine. 503
 At the margin, this indicates the value that society implicitly 504
 places, i.e. that removing a 0.002 risk is worth \$1500. The 505
 benefit of quarantine is calculated in terms of “excess risk” 506
 above 0.002 having a value where each 0.002 of excess is worth 507
 \$1500. 508

Net benefit was calculated for the specific scenario of a 509
 population of 10 million, 2% of whom test positive over the 510
 time period of interest. We assume that 20% use the app, of 511
 whom 50% enter their diagnosis into the app, thus notifying 512
 20% of contacts that the app scores as above threshold risk, 513
 who go on to infect half as many people as they would were they 514
 not contacted. We assume the cost of quarantine is similarly 515
 only half what it would be given complete compliance. We 516
 assume that each has 30 contacts following the distribution 517
 described above. Net benefit comes out to \$6.1 million for the 518
 Irish app, \$6.2 million for the Swiss app, \$11.4 million for the 519
 baseline app and \$14.7 million for the Arizonan app. The 520
 differences between these figures illustrates the importance of 521
 assigning different infectiousness values to different days. It is 522
 unfortunate in this regard that version 2 of the Google/Apple 523
 exposure notification system restricts the number of non-zero 524
 levels of infectiousness down to two. 525

We note that the value of an exposure notification app 526
 would be dramatically higher if, instead of a blanket 14 day 527
 quarantine duration, a shorter quarantine, targeted to the 528
 days of greatest risk, were used (11). Further gains would 529
 come from using negative test results to shorten quarantine. 530

5. Discussion 531

Here we showed that the socially optimal policy is to quaran- 532
 tine an individual if their risk of infectiousness is even mildly 533
 above that of the average person in the population who is 534
 not under quarantine. How much above depends slightly on 535

536 the non-linearity of the cost of isolation with the strictness
537 of isolation. While only order of magnitude calculations are
538 sometimes possible, given that case prevalences vary over sev-
539 eral orders of magnitudes across different populations, our
540 rough calculations are nevertheless instructive. Some tools,
541 like exposure notification apps, have quantitative sensitivities
542 that can be tuned in real time in ways informed by such calcu-
543 lations. With more information about risk, quarantine policy
544 can make a greater contribution to returning to a more normal
545 life. The value of quarantine policy in doing so is higher for
546 low case prevalence. It depends on vaccination prevalence
547 only if the vaccinated follow different distancing norms than
548 the unvaccinated, or if population immunity on its own is
549 extensive enough to achieve $R_t < 1$.

550 To simplify the optimal trade-off between targeted quar-
551 antine and broad social distancing, we have neglected two
552 complications. First, we have based recommendations as to
553 who should quarantine on the assumption that they will do
554 so completely. In reality, quarantined individuals often re-
555 duce rather than eliminate non-household contact, let alone
556 all contact. More nuanced messaging regarding the degree of
557 quarantine may be difficult to manage.

558 Second, how to choose R_{target} is not specified here. In the
559 Introduction and Appendix A, we discuss a control theory
560 perspective on the fact that social distancing adjusts to keep
561 the long-term geometric mean of R_t near 1. A well-functioning
562 control system is one with a negative feedback loop such that
563 the error in achieving a desired outcome feeds in as a correcting
564 input (23). Feedback can come either from top-down policy or
565 from changes in individual decisions in response to conditions.

566 A control system can be explained as a combination of
567 proportional, integral, and derivative control (23). Derivative
568 control would set social distancing policy based on whether
569 exponential growth has resumed, and if so with what doubling
570 time. This is unlikely to occur spontaneously, but rather
571 requires government action in response to epidemiological
572 reports not yet posing significant personal risk to the average
573 individual. Spontaneous control by self-interested individuals
574 responding to local conditions would under ideal circumstances
575 follow proportional control, i.e. reflect current case counts.
576 More likely, individuals will respond to the integral of case
577 counts over time, e.g. as reflected in hospital occupancy
578 rates. Targeting hospital occupancy rates (a combination of
579 proportional and integral control) is also the explicit policy of
580 some governments, e.g. the State of Arizona (24). If the goal
581 is a managed approach to naturally acquired herd immunity,
582 hospital usage is a reasonable target. But since the amount
583 of social distancing (whether mandated or voluntary) needed
584 to keep R_t near 1 is the same against a background of low
585 case counts as with high case counts, there are obvious health
586 benefits to doing so with low case counts. Integral control will
587 lead to significant fluctuations with a high total case burden.

588 We recommend mostly derivative control, with sufficient
589 proportional control to ensure return to low case counts follow-
590 ing fluctuations. The optimal extent of proportional control
591 depends on whether shorter, sharper action vs. moderate
592 decline with more social freedom achieves the least harm in
593 transitioning between different case prevalences. The choice
594 of R_{target} to control the speed with which case numbers are
595 brought to low levels must also place a value on illness and
596 death in comparison with social restrictions, a difficult prob-

lem.

597
598 Regardless of the form of feedback, compliance with trans-
599 mission control measures such as size limits to gathering,
600 compulsory masking etc. can be hard to predict, and hence
601 the degree of top-down control is limited. This underlies our
602 decision in this manuscript to focus on marginal costs given
603 prevailing policy and behaviors. Our approach exploits the
604 fact that some unspecified form of negative feedback keeps R_t ,
605 or at least its geometric mean, near 1.

606 If testing and tracing improves, then less population-wide
607 distancing will be required to achieve the same value of R_{target} ,
608 and if the same level of distancing is maintaining, this will
609 manifest as a drop in R'_t . This might on occasion lead to a
610 reconsideration of R_{target} to a lower level once that seems
611 achievable at lower cost, but otherwise, reductions in social
612 distancing should again ideally be triggered via derivative
613 control, reducing the burden on the population.

614 While some form of negative feedback can be inferred from
615 the empirical tendency of R_t to stay near 1, positive feedback
616 can also occur: as case counts rise, contact tracing becomes
617 more difficult (25), potentially accelerating the outbreak. Con-
618 sciously adjusting the risk threshold that triggers quarantine,
619 whether regarding incoming travellers, or risk thresholds in
620 an exposure notification app, contributes to this positive feed-
621 back loop. However, conscious management of this positive
622 feedback loop can manage the process to focus contact tracers'
623 attention where it can do the most good, in a manner that is
624 under straightforward real-time control.

625 Because of the positive feedback associated with contact
626 tracing workload and optimal quarantine policies, control of R'_t
627 via negative feedback, e.g. to bring it back down to its target
628 given a rise due to seasonal conditions or simply pandemic
629 fatigue, is better exerted through population-wide measures
630 to affect D . Our work shows that this is the socially optimal
631 approach, and indeed if performed adequately, should be ac-
632 companied by a perhaps unintuitive relaxing of quarantine.
633 Our formal development assumes a perfect control system, and
634 focuses on one-individual perturbations to an equilibrium. It
635 is likely that D will not reliably adjust in a completely timely
636 manner, but a proactive approach can improve the chances of
637 this occurring.

638 The value of quarantine policies is highest when case counts
639 are low. However if low cases counts are a consequence of
640 immunity (whether from natural infection or from vaccination),
641 then the value of a quarantine policy will go down when those
642 with immunity follow different distancing norms from those
643 without. There is a direct benefit from relaxing distancing for
644 the immune, whether this behavior is self-regulated or whether
645 it comes in the form of a more official "immunity passport".
646 Immunity passports were once problematic because they would
647 have set up an adverse incentive in favor of contracting SARS-
648 CoV-2. However, with increasing vaccine availability, the
649 benefits from immunity passports have risen at the same time
650 as an adverse incentive to get infected has been converted
651 into a pro-socially aligned incentive to get vaccinated. Our
652 approach can be used to merge immunity history, quarantine
653 policies, and even recent negative test results into an integrate
654 metric of "risk to others". The socially optimal policy is to use
655 all available information to target restrictions to where they
656 will do the most good, and restore as much social activity as
657 possible while still controlling transmission.

6. Acknowledgements

James Petrie acknowledges support from the Ontario Graduate Scholarship and thanks his advisor, Prof. Stephen Vavasis, for giving him the freedom to work on this research project. We thank Amanda Wilson for helpful comments on an earlier draft, Kevin Murphy at Google for access to his COVID Tuner risk score configuration tool, and the entire WeHealth team for their ethical commitment to maximizing benefits while minimizing harm, including via the recommendations of this paper. The opinions expressed in this publication are those of the authors and do not necessarily reflect those of WeHealth Solutions PBC.

7. Competing Interests

Joanna Masel consults for WeHealth, who distribute exposure notification solutions to Arizona and Bermuda. James Petrie is likely to soon sign a contract giving him an equity interest in WeHealth.

1. A Brodeur, D Gray, A Islam, SJ Bhuiyan, A literature review of the economics of covid-19, (Institute of Labor Economics (IZA)), IZA Discussion Papers 13411 (2020).
2. O Yakusheva, E van den Broek-Altenburg, G Brekke, A Atherly, The cure is not worse than the disease—a humanitarian perspective. *Available at SSRN 3638575* (2020).
3. ZA Bethune, A Korinek, Covid-19 infection externalities: Trading off lives vs. livelihoods, (National Bureau of Economic Research), Technical report (2020).
4. H d'Albis, E Augeraud-Véron, Optimal prevention and eradication of epidemics. (2020).
5. D Acemoglu, V Chernozhukov, I Werning, MD Whinston, Optimal targeted lockdowns in a multi-group sir model. *NBER Work. Pap.* **27102** (2020).
6. Y Bar-On, T Baron, B Ofer Cornfeld, R Milo, Modelling covid19: Epidemiological evidence and model misspecifications. (2020).
7. MN Lurie, J Silva, RR Yorlets, J Tao, PA Chan, Coronavirus Disease 2019 Epidemic Doubling Time in the United States Before and During Stay-at-Home Restrictions. *The J. Infect. Dis.* **222**, 1601–1606 (2020).
8. (<https://covid19-projections.com/>) (2020).
9. JS Weitz, SW Park, C Eksin, J Dushoff, Awareness-driven behavior changes can shift the shape of epidemics away from peaks and toward plateaus, shoulders, and oscillations. *Proc. Natl. Acad. Sci.* **117**, 32764–32771 (2020).
10. MS Eichenbaum, S Rebelo, M Trabandt, The macroeconomics of testing and quarantining, (National Bureau of Economic Research), Technical report (2020).
11. AM Wilson, et al., Quantifying sars-cov-2 infection risk within the google/apple exposure notification framework to inform quarantine recommendations. (2020).
12. VA Karatayev, M Anand, CT Bauch, Local lockdowns outperform global lockdown on the far side of the covid-19 epidemic curve. *Proc. Natl. Acad. Sci.* **117**, 24575–24580 (2020).
13. RM Anderson, RM May, *Infectious diseases of humans: dynamics and control*. (Oxford university press), (1992).
14. CR Wells, et al., Optimal covid-19 quarantine and testing strategies. *medRxiv* (2020).
15. (<https://pandemicnavigator.oliverwyman.com/forecast?mode=country®ion=United%20States&panel=mortality>) (2020).
16. YJ Park, et al., Contact tracing during coronavirus disease outbreak, south korea, 2020. *Emerg. infectious diseases* **26**, 2465–2468 (2020).
17. R Laxminarayan, et al., Epidemiology and transmission dynamics of covid-19 in two indian states. *Science* **370**, 691–697 (2020).
18. W Chen, N Zhang, J Wei, HL Yen, Y Li, Short-range airborne route dominates exposure of respiratory infection during close contact. *Build. Environ.* **176**, 106859 (2020).
19. (<https://globalnews.ca/news/7416775/canada-coronavirus-border-quarantine/>) (2020).
20. Emerging covid-19 success story: Vietnam's commitment to containment (<https://ourworldindata.org/covid-exemplar-vietnam>) (2020) Accessed: 2021-03-30.
21. Abbott's fast, \$5, 15-minute, easy-to-use covid-19 antigen test (<https://abbott.mediaroom.com/2020-08-26-Abbotts-Fast-5-15-Minute-Easy-to-Use-COVID-19-Antigen-Test-Receives-FDA-Emergency-Use-Authorization-Mobile-App-Displays-Test-Results-to-Help-Our-Return-to-Daily-Life-Ramping-Production-to-50-Million-Tests-a-Month>) (2020) Accessed: 2021-03-30.
22. (<https://risk-score-tuner.appspot.com/>) (2020).
23. RC Dorf, RH Bishop, Modern control systems. (1998).
24. Arizona ended covid restrictions because hospitals have bed space, health chief says (https://tucson.com/news/local/arizona-ended-covid-restrictions-because-hospitals-have-bed-space-health-chief-says/article_c35f294e-4cd8-5ca3-b6c9-abc61b15ec4b.html) (2020) Accessed: 2021-03-30.
25. (<https://covidmodelling.pwias.ubc.ca/blog/>) (2020).
26. A Endo, S Abbott, AJ Kucharski, S Funk, et al., Estimating the overdispersion in covid-19 transmission using outbreak sizes outside china. *Wellcome Open Res.* **5**, 67 (2020).

Appendices

A. Fluctuating R'_t

In Model Section A, we assumed that D instantaneously adapts to achieve R_{target} given prevailing conditions and current quarantine policies. This is motivated by the observation of relatively stable R'_t averaged over longer timescales. In this section, we show that Equation 1, and hence subsequent findings, are still a good approximation even when including stochastic effects, time delays, and uncertainty about system state.

To make this argument, we assume that for a certain region and time frame, the virus is neither eliminated nor creates herd immunity, and that control measures are relatively stable. Over N serial intervals between times t_0 and t_N , the initial and final case prevalence are related by $I(t_N) = I(t_0) \prod_{n=0}^{N-1} R'_t(t_n)$. Equivalently, the geometric mean of R'_t over this time period can be expressed using the initial and final case prevalence, $(\prod_n (R'_t(t_n)))^{1/N} = (\frac{I(t_N)}{I(t_0)})^{1/N}$. For most regions, case prevalence doesn't vary by more than a couple orders of magnitude on the scale of months, which yields a (geometric) mean R'_t value quite close to 1. As an example, with $I(t_N)/I(t_1) = 100$, and $N = 36$ (6 months with a serial interval of 5 days), $100^{1/36} = 1.14$. Approximating the geometric mean of the effective reproductive number as 1, and expanding to include distancing and quarantine terms:

$$1 \approx \left(\prod_{n=0}^{N-1} (R'_t(t_n)) \right)^{1/N} = \left(\prod_{n=0}^{N-1} (R_t(t_n)(1-D(t_n)) \frac{I(t_n) - Q_i(t_n)}{I(t_n)}) \right)^{1/N} \quad [9]$$

$$\log(1) \approx \frac{1}{N} \sum_{n=0}^{N-1} [\log(R_t(t_n)) + \log(1-D(t_n)) + \log\left(\frac{I(t_n) - Q_i(t_n)}{I(t_n)}\right)] \quad [10]$$

$$0 \approx E[\log(R_t(t))] + E[\log(1-D(t))] + E[\log\left(\frac{I(t) - Q_i(t)}{I(t)}\right)] \quad [11]$$

Assuming control measures don't change drastically over time, we replace the mean of the log with the log of the mean:

$$0 \approx \log(E[R_t(t)]) + \log(E[1-D(t)]) + \log\left(E\left[\frac{I(t) - Q_i(t)}{I(t)}\right]\right) \quad [12]$$

$$1 \approx E[R_t(t)] * E[1-D(t)] * E\left[\frac{I(t) - Q_i(t)}{I(t)}\right] \quad [13]$$

This is equivalent to Equation 1 with R'_t set to 1, showing that by averaging over time, the requirements for precise knowledge and control of system state can be relaxed.

B. Imported Cases and Stochasticity

For regions where imported cases make up a significant fraction of total cases, policy-makers should consider the modified dynamics when choosing R_{target} . Imported cases act as a forcing term, with the number of cases at a future timepoint being the sum of locally obtained infections and imported infections. The mechanics of choosing a risk threshold do not change, but it is important to highlight that R_{target} only determines a portion of future cases. A complete solution must also consider the effect of border policies.

When the total number of cases is small, e.g. in a region with excellent local control for which rare imported cases are the primary concern, discrete and stochastic effects become important.

Quantitative models of tiny case count scenarios are challenging, because of stochasticity in the length of time needed to return to elimination, combined with curvature in the harms of shutdown as a function of its duration. This is further exacerbated by superspreader dynamics (26). Modelling the number of secondary cases with a negative binomial distribution with an overdispersion parameter 0.1 and mean 2.5 yields a variance of 65. In locations not aiming for local elimination, stochasticity averages out over time, and expected values can be used. When local elimination is being attempted, it would be useful to make use of discrete, stochastic models when choosing R_{target} , e.g. to calculate the probability distribution of time to elimination and the expected damage from shutdowns of different severity. With the choice of R_{target} set, the previous arguments mostly still hold (although some consideration may be given to the difference in variance in transmissions based on broad vs. targeted distancing).

If local elimination has already been achieved, then a still more complicated case is to consider the risk of an outbreak from an imported case, the time delay and growth in the outbreak before it is noticed, and the subsequent shutdowns needed to re-achieve local elimination and thus relax general social distancing. It is likely that the considerable costs of such an outbreak warrant strict quarantine for incoming individuals. A robust testing and contact tracing system capable of quickly containing any new outbreak is also key.

C. Cost Function Dependence

Consider the situation where r_1 is calculated using $f_1(x) = x$, and r_2 is calculated using $f_2(x)$, where $f_2(x)$ is strictly increasing, concave up, and differentiable. For both of these calculations, define the distancing set point $D_0 = \max(0, 1 - \frac{R_{target}}{R_t} \frac{I}{I - Q_i})$, the required distancing if one more infectious person were isolated $D_- = \max(0, 1 - \frac{R_{target}}{R_t} \frac{I}{I - Q_i - 1})$, and $Q = Q_i + Q_n$. By expanding Equation 6, the two risk thresholds are given by Equations 14 and 15:

$$r_1 = \frac{f_1(1) - f_1(D_0)}{f_1(D_0) - f_1(D_-)} \frac{1}{P - Q - 1} \quad [14]$$

$$r_2 = \frac{f_2(1) - f_2(D_0)}{f_2(D_0) - f_2(D_-)} * \frac{1}{P - Q - 1} \quad [15]$$

Substituting $f_1(x) = x$, and taking the ratio of the two thresholds gives:

$$\frac{r_2}{r_1} = \frac{\frac{f_2(1) - f_2(D_0)}{1 - D_0}}{\frac{f_2(D_0) - f_2(D_-)}{D_0 - D_-}} \quad [16]$$

Since $\frac{f_2(1) - f_2(D_0)}{1 - D_0} \leq \max_{x \in [0,1]} f'_2(x)$, and $\frac{f_2(D_0) - f_2(D_-)}{D_0 - D_-} \geq \min_{x \in [0,1]} f'_2(x)$, then Equation 17 gives a bound on the ratio of the two thresholds.

$$\frac{r_2}{r_1} \leq \frac{\max_{x \in [0,1]} f'_2(x)}{\min_{x \in [0,1]} f'_2(x)} \quad [17]$$

If $f_2(x)$ is concave up (indicating that additional isolation is more costly), then $\frac{f_2(1) - f_2(D_0)}{1 - D_0} \geq \frac{f_2(D_0) - f_2(D_-)}{D_0 - D_-}$, and therefore $r_2 \geq r_1$.

Substituting Equation 7 into Equation 16, we obtain:

$$r_{thresh} = \frac{\frac{f(1) - f(D_0)}{1 - D_0}}{\frac{f(D_0) - f(D_-)}{D_0 - D_-}} \frac{I - Q_i - 1}{P - Q_i - Q_n - 1} \quad [18]$$

For differentiable $f(x)$ and relatively large $I - Q_i$, $D_0 - D_-$ is small, and $\frac{f(D_0) - f(D_-)}{D_0 - D_-}$ can be approximated as $f'(D_0)$. Using the risk of the unquarantined population $r_{pop} = \frac{I - Q_i}{P - Q_i - Q_n} \approx \frac{I - Q_i - 1}{P - Q_i - Q_n - 1}$, we obtain Equation 8.