Airway antibodies emerge according to COVID-19 severity and wane rapidly but

reappear after SARS-CoV-2 vaccination

Alberto Cagigi[1], Meng Yu[1], Sara Falck-Jones[1], Sindhu Vangeti[1], Björn Österberg[1], Eric

Åhlberg[1], Lida Azizmohammadi[1], Ryan Falck-Jones[2,3], Pia C Gubisch[1], Mert Ödemis[1],

Farangies Ghafoor[1], Julia Svensson[1], Mona Eisele[1], Klara Lenart[1], Max Bell[2,3], Niclas

Johansson[4,5], Jan Albert[6,7], Jörgen Sälde[8], Deleah Pettie[9,10], Michael Murphy[9,10], Lauren

Carter[9,10], Neil P King[9,10], Sebastian Ols[1], Anna Färnert[4,5], Karin Loré[1*] and

Anna Smed-Sörensen[1*]

[1]Division of Immunology and Allergy, Department of Medicine Solna, Karolinska Institutet, Karolinska

University Hospital, Stockholm, Sweden. [2]Department of Physiology and Pharmacology, Karolinska

Institutet, Stockholm, Sweden. [3]Department of Perioperative Medicine and Intensive Care, Karolinska

University Hospital, Stockholm, Sweden. [4]Division of Infectious Diseases, Department of Medicine

Solna, Center for Molecular Medicine, Karolinska Institutet, Sweden. [5]Department of Infectious

Diseases, Karolinska University Hospital Solna, Stockholm, Sweden. [6]Department of Microbiology,

Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden. [7]Clinical Microbiology, Karolinska

University Hospital Solna, Stockholm, Sweden. [8]Närakut SLSO, Karolinska University Hospital Solna,

Stockholm, Sweden. [9]Department of Biochemistry, University of Washington, Seattle, WA, United

States. [10]Institute for Protein Design, University of Washington, Seattle, WA, United States.

[*]Equal contribution.

Correspondence to: Karin Loré and Anna Smed-Sörensen, Division of Immunology and

Allergy, Department of Medicine Solna, Karolinska Institutet, Visionsgatan 4, BioClinicum

J7:30, Karolinska University Hospital, 171 64 Stockholm, Sweden.

E-mail addresses: karin.lore@ki.se; anna.smed.sorensen@ki.se

Conflict of interest statement

The authors have declared that no conflict of interest exists.

Abstract

Understanding the presence and durability of antibodies against SARS-CoV-2 in the airways

is required to provide insights on the ability of individuals to neutralize the virus locally and

prevent viral spread. Here, we longitudinally assessed both systemic and airway immune

responses upon SARS-CoV-2 infection in a clinically well-characterized cohort of 147 infected

individuals representing the full spectrum of COVID-19 severity; from asymptomatic

infection to fatal disease. In addition, we evaluated how SARS-CoV-2 vaccination influenced

the antibody responses in a subset of these individuals during convalescence. Not only

systemic but also airway antibody responses correlated with the degree of disease severity

with increasing levels in patients with mild, moderate and severe disease, respectively.

However, while systemic IgG levels were durable for up to 8 months, airway IgG and IgA had

declined significantly within 3 months. In contrast, antigen-specific memory B cells were well

maintained and comparable across disease severity. After vaccination, there was an increase

in both systemic and airway antibodies, in particular IgG, often exceeding the levels found

during acute disease. In contrast to plasma, airway antibody levels were significantly

elevated after the boost vaccination, highlighting the importance of prime and boost

vaccination also for previously infected individuals to obtain optimal mucosal protection.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection that causes

coronavirus disease 2019 (COVID-19) presents with a wide range of disease severity from

asymptomatic to fatal (1, 2). Individuals of advanced age and/or those with comorbidities

are overrepresented among patients who develop severe disease (3). However, the majority

of SARS-CoV-2 infected individuals experience asymptomatic infection or only mild disease

(4).

Systemic antibodies against the SARS-CoV-2 nucleocapsid (N) and the viral surface

glycoprotein spike (S) as well as against the receptor binding domain (RBD) (5, 6) of the S

protein have been studied extensively (7-11). Responses against the internal N protein are

often readily detectable but their contribution to protection and control of disease is not

clear (8, 10). In contrast, antibody responses against S and, in particular, against the RBD

result in virus neutralization (12). Responses against the RBD are thus likely necessary for

protection from re-infection or prevention of symptomatic disease. However, the presence

and durability of antibodies during COVID-19 in the airways is still not well understood.

The respiratory tract is the initial site of viral infection and replication. The availability

of antibodies at this site could therefore determine the ability to neutralize the virus locally

in case of (re-) exposure and prevent viral spread. Generally, antibodies present in the

circulation and at local sites are the result of secretion from short-lived plasmablasts and/or

terminally differentiated plasma cells in the bone marrow or mucosal sites (13). However,

the response to a secondary infection once antibody titers have waned below protective

levels mostly relies on the presence of resting antigen-specific memory B cells that are

rapidly activated upon antigen re-exposure (13). Whether vaccination against SARS-CoV-2

also elicits systemic antibody responses in addition to local antibodies in the airways of

individuals who recovered from COVID-19, and via which mechanism, is currently unknown.

In this study we present data from a cohort of patients that we have followed since

mid-March 2020, which was the start of the pandemic in Sweden. We show longitudinal

data on virus-specific systemic and airway antibody and B cell memory responses generated

in this clinically well-characterized cohort of individuals with SARS-CoV-2 infection (n=147)

ranging from asymptomatic SARS-CoV-2 infection to fatal COVID-19 disease. In addition, we

show how subsequent SARS-CoV-2 vaccination during the convalescent phase significantly

boosts not only the systemic but also airway antibody responses.

Results

Patient enrollment, assessment of disease severity and timeline

Individuals were sampled longitudinally in blood and airways during acute

infection/symptomatic disease and during convalescence (median 3 and 8 months from

symptom onset). Donor-matched plasma, peripheral blood mononuclear cells (PBMC),

nostril swabs (NSW) and nasopharyngeal aspirates (NPA) were collected from all patients

across disease severities whereas endotracheal aspirates (ETA) were only collected from

intubated patients receiving intensive care (Figure 1). Disease severity was assessed daily,

using a seven-point scale derived from the respiratory domain of the sequential organ failure

assessment (SOFA) score (14, 15), with additional levels for non-admitted and fatal cases

(Table 1). Patients were grouped based on peak disease severity, which may differ from

disease severity at time of sampling (Table 1 and Figure 1B). In addition, pre-pandemic

healthy controls (PPHC) (n=30) as well as individuals with influenza-like symptoms, and

possible SARS-CoV-2 exposure, but with negative diagnostic PCR results (PCR-) (n=9) were

sampled in the same way and included as controls. Generally, severe patients were sampled

later after symptom onset as compared with individuals with mild disease resulting in a large

time frame of study inclusion with respect to symptom onset (Table 1 and Figure 1B) (16).

For simplicity, the sampling period/study inclusion during ongoing infection and

hospitalization (for those hospitalized) is referred to as the "acute" phase. Samples collected

at the first follow-up visit during convalescence (range 46-168 days from symptom onset;

median 108 days) are referred to as the "3 months" timepoint whereas those collected at

the second follow-up visit (range 187-344 days; median 245 days) are referred to as the "8

months" timepoint. Time of the first convalescent follow-up sampling from acute sampling

ranged 33-159 days; median 90 days (Table 1).

Plasma IgG and IgA responses to N, S and RBD across COVID-19 severity during acute

disease and after recovery

We first assessed systemic IgG and IgA responses against N, S and RBD at the time of study

inclusion that ranged between 0-54 days from onset of symptoms; median 16 days for the

whole cohort (Table 1). Both IgG and IgA levels against all viral proteins followed the degree

of disease severity with increasing levels in patients with mild, moderate and severe disease

respectively (Figure 2A). In line with previous reports, IgG against N were the most elevated

in patients who had severe disease or a fatal outcome (8, 10). The degree of disease severity

also associated with the levels of systemic inflammation as indicated by the levels of C-

reactive protein (CRP) in blood and by the neutrophil-lymphocyte ratio (NLR) (Figure 2B).

Interestingly, the levels of neutrophils also specifically associated with disease severity

(Figure 2D) and with all of the systemic antibody responses during acute disease (Figure 2D

and Supplementary figure 1). The levels of IgG during acute disease, and to a lower extent

IgA, against all tested antigens, exhibited a positive correlation with the days from onset of

symptoms (Supplementary figure 2A). This difference in antibody titers over time might be

slightly accentuated by the fact that in our cohort the patients with moderate/severe

disease, and even fatal outcome, for whom we initially observed low IgG titers against RBD,

had an early study inclusion (on average 13 days from onset of symptoms). In fact, these

patients showed significantly higher titers later during the acute phase (on average 19 days)

(Supplementary figure 2B-C). Nonetheless, patients with mild disease displayed lower levels

of plasma IgG against RBD as compared with more severe patients, also when samples were

taken after similar duration of symptoms (Supplementary figure 2D). After 3 months from

symptom onset, the IgG levels remained high in the plasma of patients recovering from

moderate and severe disease, while the levels had further increased in the individuals who

had a mild disease (Figure 3A). However, despite this increase over time, the antibody levels

in mild patients never reached the levels observed for moderate and severe patients or for

those who had a fatal outcome (Figure 3 and Supplementary figure 3A).

The IgG levels had significantly waned from 3 to 8 months in patients who recovered from

moderate and severe disease, but the decline was smaller in patients who experienced a

mild disease (Figure 3B and Supplementary figure 3). In contrast to IgG, IgA levels from the

acute phase, against all antigens, waned substantially in most patients already after 3

months (Figure 3 and Supplementary figure 3). Antibody titers during acute disease

correlated with peak disease severity as well as with disease severity at time of sampling

(Supplementary figure 4). The correlation between antibody titers and peak disease severity

was maintained also when analyzing the antibodies at the 3- and 8-month follow-up visits

(Supplementary figure 4) as also observed in another study (17).

Airway IgG and IgA responses and assessment of B cell frequencies in the respiratory tract

We next measured the levels of IgG and IgA in the upper and lower airways and compared

with levels in plasma at matched time points. Due to limited respiratory sample volumes, we

focused our analyses on IgG and IgA responses against the RBD since these responses are

most critical for virus neutralization. We found that RBD-specific antibodies could be

detected in nasal swabs (NSW) (Figure 4A) and nasopharyngeal aspirates (NPA) (Figure 4B)

during the acute phase across all disease severities (Figure 4A, B and C). In agreement with

our observations in plasma, antibody levels in the upper respiratory tract were higher in

patients with moderate or severe disease as compared with individuals with mild disease.

Both IgG and IgA levels had declined significantly already after 3 months, with IgG declining

to almost undetectable levels (Figure 4A- C). RBD antibody levels during acute infection were

on average higher in NPA compared to NSW for both IgG and IgA across disease severity

(Figure 4A- C) suggesting that antibody titers may increase not only with disease severity but

also with sampling at different depths of the upper airways. To address this, we compared

donor-matched NSW (peripheral nostril), NPA (upper airway) and ETA (trachea) collected at

the same time point during acute disease from intubated patients from whom we had both

peripheral, upper and lower airway samples. Interestingly, we still found significantly higher

levels of IgA against the RBD in NPA as compared with NSW and ETA (Figure 4D).

Furthermore, nasopharyngeal antibody levels (both IgG and IgA) showed a strong correlation

with plasma antibody responses (Figure 4E). We also assessed the presence of B cells in the

respiratory tract of COVID-19 patients by analyzing the lymphocytes that could be retrieved

from NPA and ETA as compared with NPA from three healthy controls (HC). Despite

generally obtaining a significantly lower cell yield from NPA as compared with ETA,

lymphocyte frequencies did not differ in NPA and ETA from COVID-19 patients but both were

lower as compared with NPA from HC. Instead, the proportion of B cells in NPA was higher

as compared with ETA in COVID-19 patients and similar to NPA from HC (Figure 5A-B).

Expansion of SARS-CoV-2-specific memory B cells

As mentioned above, the virus-specific B cell memory pool will be essential to remount a

rapid antibody response in the case of re-exposure. To assess the establishment of antigen-

specific memory B cells, donor-matched PBMC from acute disease and convalescence were

analyzed side-by-side using fluorescently labelled S and RBD probes (18-20). Patients with

moderate/severe disease showed the presence of Ig-switched memory B cells specific to S in

the acute phase and the memory B cell pool had further expanded after 3 months (ranging

from 0.009 to 1.35%; mean 0.42% during convalescence) (Figure 5C-F). Individuals with mild

disease showed lower frequencies of S-specific memory B cells during acute disease than the

patients with moderate/severe disease. In fact, the frequencies of S-specific memory B cells

in the mild patients during the acute phase were not different from those observed in the

PCR- individuals or in the PPHC (Figure 5C and E). However, the frequencies of S-specific

memory B cells had substantially increased in the mild patients after 3 months (ranging from

0.17% to 0.64%; mean 0.35% during convalescence) and were comparable to frequencies

among severe patients. In addition, the levels were well maintained between 3 and 8

months in all groups (Figure 5E and F). Further phenotyping of the S-specific memory B cells

indicated that the majority of these cells may be specific for epitopes on S outside of the

RBD (Figure 5D). S-specific memory B cells in the circulation were predominantly IgG+, rather

than IgA+ (Figure 5D).

The effect of vaccination on systemic and airway antibody levels

We finally evaluated the influence of SARS-CoV-2 vaccination on the systemic and airway

antibody responses (Figure 6A). A subset of 12 individuals, 2 that recovered from mild, 4

from moderate and 6 from severe COVID-19 one year earlier, were sampled after receiving

their scheduled vaccination (range 270-358 days; median 326 days from symptom onset)

(Table 2). Donor-matched plasma, NSW and NPA were collected at different timepoints

after prime (7-16 days) from 10 patients and after boost (7-28 days) from 7 patients and

analyzed for the presence of IgG and IgA against RBD. Antibodies against N were also

measured in plasma as a negative control as the vaccines used were based on the S protein.

After vaccination, all individuals demonstrated a significant increase of both plasma IgG and

IgA against the RBD (Figure 6B and Supplementary figure 5) but, as expected, not against N

(Figure 6B). The plasma antibody levels, especially IgG, to RBD further increased from prime

to boost vaccination in the majority of individuals analyzed. However, due to the low

numbers of individuals (n=5) who had this sample set available, this did not reach statistical

significance (p=0.06). While the IgG levels to RBD after boost vaccination exceeded the levels

detected during the acute phase, the IgA levels were equally high (Figure 6C). The airway IgG

levels to RBD also showed a noticeable increase after the boost vaccination in particular. In

fact, the IgG levels in the airway samples, both nasal swabs and NPA, were in most

individuals significantly higher after boost vaccination than they were in the acute stage of

the disease (Figure 6C). In contrast, this was not noted for IgA levels to RBD.

Discussion

By now, it is well documented that higher systemic antibody levels are generated in severe

as compared with mild COVID-19 (7-11, 21-23). In contrast, the presence and durability of

antibodies against SARS-CoV-2 in the airways is much less understood. Nor is it known if and

how respiratory antibody levels are influenced by SARS-CoV-2 vaccination in humans. In this

study, we performed longitudinal analyses of systemic and upper and lower airway antibody

responses in a clinically well-characterized and relatively large cohort of individuals with

SARS-CoV-2 infection representing the full spectrum of COVID-19 severity ranging from

asymptomatic infection to fatal disease. Matched analyses in blood and in the airways

enabled us not only to address the magnitude and durability of systemic antibodies to SARS-

CoV-2 but also to gain insights into the prospects of protective capacity locally in the mucosa

at virus re-entry. This is one key aspect still largely unknown yet critical for our

understanding of immunity to and protection from SARS-CoV-2. Furthermore, we studied

how the systemic versus airway antibody levels were affected by vaccination. Collectively,

this data will contribute to a better understanding of long-term protective effects and

whether vaccination is important to boost the capacity of virus neutralization in the airways

and thus reducing re-infection and virus spread.

Airway mucus along the respiratory tract is thought to serve as a barrier that can trap

respiratory viruses via virus glycoprotein-mucin interactions (24). However, it has been

shown that local immobilization of respiratory viruses such as influenza viruses in the

airways mostly occurs by binding with virus-specific antibodies present in the mucus (25). As

the respiratory tract is the initial site of viral infection and replication, the levels of IgG and

IgA against the RBD in the upper and lower airways are likely critical for SARS-CoV-2

neutralization and could therefore help predict the ability of individuals to neutralize the

virus locally in case of re-exposure. Low but detectable levels of antibodies to SARS-CoV-2

have previously been reported in saliva during convalescence (26). In our study we found

that IgG and IgA against the RBD can be readily detected in the upper and lower airway

during acute disease and that such levels correlated with the systemic response at the same

time point and also followed disease severity. However, for all the patients across disease

severities, airway antibodies waned to low levels much faster than those in plasma during

convalescence. Whether these low antibody levels observed at respiratory sites will be

sufficient for preventing virus re-entry or for protection is not known. The correlation

between systemic and airway antibody levels during acute disease raises questions on

whether the low levels of antibodies in the airways during convalescence are due to

decreased antibody generation locally at mucosal sites or are rather caused by decreased

dissemination from the periphery once systemic antibody levels start to wane. Antibodies in

the upper respiratory tract have been shown to be dominated by secretory IgA which are

mostly produced by plasma cells in the lamina propria of mucosa-associated lymphoid tissue

(MALT) (27, 28). We detected high levels of IgA in the upper airways early during acute

COVID-19 that rapidly declined during convalescence, following the pattern observed for

systemic IgA levels here and in other reports (29-31). This suggests that at least some IgA

disseminated into the airways from the circulation. In contrast, the dynamics of IgG were

different in the respiratory samples compared to plasma with airway IgG following the same

kinetics as IgA, while systemic IgG were well maintained up to 8 months.

When we assessed the presence of lymphocytes in the different airway compartments

during acute disease, we observed higher proportion of B cells along with high antibody

levels, especially IgA, in the nasopharynx, as compared with the nostril or the endotracheal

space. It has previously been shown that the majority of antibody secreting cells generated

after intranasal immunization with live-attenuated vaccines in rodents may reside in the

respiratory tract rather than in the spleen and bone marrow (32) and that these cells secrete

IgA early after a later challenge with the vaccination pathogen (33-35). Therefore, it is

possible that B cells generated during SARS-CoV-2 infection also reside locally in the airways

and contribute to antibody levels in the nasopharynx. While the antibody content in NPA

and ETA could be partially influenced by differences in sampling methods and sample

volumes, these data suggest that antibody abundance and possibly virus neutralization via

IgA differ along the respiratory tract and may be more pronounced in the nasopharynx

compared to the lower airways. Altogether, our observations demonstrate that moderate

and severe COVID-19 result in high levels of circulating antibodies and despite that IgG levels

are well-maintained, antibody levels in the airways decline significantly after the acute

phase.

Once antibody titers have waned below protective levels, the response to a secondary

infection will mostly rely on the presence of resting antigen-specific memory B cells that can

rapidly activate upon antigen re-exposure (13). Therefore, similar to other studies (18-20),

we investigated the induction and maintenance of S-specific memory B cells. Importantly,

because of the comprehensive distribution of disease severity represented in our cohort, we

were able to compare the opposite ends of the COVID-19 disease spectrum by focusing on

individuals with mild disease as compared with patients with moderate/severe disease who

had the highest circulating IgG and IgA levels. Strikingly, despite the fact that these patients

were at the opposite ends of the disease severity spectrum, they had comparable levels of S-

specific memory B cells during convalescence. These appeared to be specific for epitopes on

S outside of the RBD and were predominantly IgG+, rather than IgA+, which may affect the

proportions of different isotypes subsequently produced in the event of antigen re-

exposure.

Immunization at mucosal sites such as for example intranasal administration of live-

attenuated influenza vaccines generally elicits mucosal immune responses (36). However,

several studies, primarily performed with DNA and virus-like particles (VLP) vaccines, have

shown that intradermal, subcutaneous and intramuscular immunization also can result in

local mucosal responses that protect from mucosal challenge (37). It has been speculated

that this could be due to free antigen or B cells migrating from the vaccine draining lymph

nodes to the MALT (37-39). A two-dose regimen of Moderna's mRNA-1273 vaccine

administered intramuscularly and followed by intranasal and intratracheal challenge with

SARS-CoV-2 in rhesus macaques has indeed shown to result in local virus neutralization in

the airways (40). Antibodies in bronchoalveolar lavage and nasal swabs were elicited in a

vaccine dose-dependent manner assessed after the boost vaccination (41).

Whether the systemic and/or mucosal immunity generated during natural infection is

boosted by vaccination and results in a similar or enhanced magnitude of responses would

be important knowledge to acquire for planning the best vaccination strategies for SARS-

CoV-2 as well as for other respiratory viruses. Our results on individuals recovering from

COVID-19 and subsequently receiving vaccination indicated a marked increase of both IgG

and IgA levels systemically but also strikingly in the airways, which in the majority of cases

exceeded the levels observed during acute disease. Notably, the antibody increase observed

between prime and boost vaccination was more prominent in the airways than systemically.

Recent studies on systemic antibody responses after SARS-CoV-2 vaccination in individuals

who recovered from COVID-19 have shown a significant increase in antibody levels after one

vaccine dose with no or only a small increase after the second dose (42-46). This suggests

that one vaccine dose may be sufficient to protect these individuals from disease in case of

re-infection which is important for vaccine dose management at the population level.

However, our data indicate that only assessing the systemic antibody levels after vaccination

is to some extent misleading as respiratory antibody levels, and likely virus neutralization,

may be substantially better with a prime-boost vaccination strategy rather than with one

single dose. The higher levels of airway antibodies after two vaccine doses may be explained

by even a small increase in circulating antibodies after a second vaccine dose causing a

substantial extravasation from the bloodstream into mucosal sites. Interestingly, two recent

studies showed the presence of antibodies in the upper respiratory tract of individuals

vaccinated against SARS-CoV-2 with higher levels and even neutralizing activity in small

subsets of individuals who had recovered from COVID-19 before vaccination (47, 48).

In summary, here we show that COVID-19 disease severity not only determines the

magnitude of systemic but also airway antibody levels with efficient generation of virus-

specific memory B cells against SARS-CoV-2 also occurring upon mild disease. While plasma

IgG levels were generally well detectable after acute disease in all groups, there was a

significant decline in airway antibodies during convalescence. This suggests that antibodies

in the airways may not be maintained at levels that prevent local virus entry upon re-

exposure. However, our data indicate that the majority of infected individuals have the

ability to generate anamnestic responses via the memory B cell pool and that vaccination

against SARS-Cov-2 resulted in a substantial rebound of both systemic and airway antibodies

in patient who recovered from COVID-19. These data indicate a positive effect of vaccination

for increased virus neutralization in the airways and prospects of reduced virus spread,

which further supports following the full vaccination schedule also in this population.

Methods

Study design, patient enrollment and sample collection

One hundred and forty-seven (147) PCR-confirmed SARS-CoV-2 infected patients were

enrolled at the Karolinska University Hospital and Haga Outpatient Clinic (Haga Närakut),

Stockholm, Sweden during March-May 2020 (acute phase) in a time that ranged from 0 to

54 days from onset of symptoms as self-reported by individual patients; and during April-

September 2020 (3 months) in a time that ranged from 46 to 168 days and during November

2020 to February 2021 (8 months) continuing from the previous counts. Patients were

enrolled at various settings, ranging from primary to intensive care. In order to recruit

asymptomatic and mild cases, household contacts of COVID-19 patients were enrolled and

screened with PCR to identify SARS-CoV-2 positive individuals. A small subset of these

individuals who experienced influenza-like symptoms and were possibly exposed to SARS-

Cov-2 but had a negative diagnostic PCR (PCR-) (n=9 of whom 3 were household contacts of

confirmed patients with 1 experiencing fever, and 6 were included based on suspected

infection with 4 experiencing fever) were sampled in the same way and included as controls

alongside with 30 pre-pandemic healthy control samples (PPHC) from 2016-2018.

Respiratory failure was categorized daily according to the respiratory domain of the

Sequential Organ Failure Assessment score (SOFA)(14). The modified SOFA score (mSOFA)

was calculated when arterial partial pressure of oxygen (PaO_2) was not available. In this case

peripheral transcutaneous hemoglobin saturation (SpO_2) was used instead (15). Estimation

of the fraction of inspired oxygen (FiO_2) based on O_2 flow was calculated as per the Swedish

Intensive Care register definition (49). Patients were categorized based on the peak

respiratory SOFA or mSOFA value with the 4-point respiratory SOFA score being extended

with additional levels to distinguish between admitted and non-admitted mild cases (both

respiratory SOFA score 0) and to include fatal outcome. Ten (10) patients with fatal outcome

had peak disease severity score 6 prior to death and 2 patients had scores of 4 and 5. For

convenience, the resulting 7-point composite peak disease severity (PDS) was condensed

into a broader classification consisting of mild (1-2), moderate (3-4), severe (5-6), and fatal

(7). Demographics and additional data were collected from medical records, including

clinical history and risk factors such as BMI and co-morbidities. Total burden of comorbidities

was assessed using the Charlson co-morbidity index (CCI) (50) (Table 1). Additional clinical

information on this patient cohort including the modulation of disease from time to study

inclusion to peak severity can be found in Falck-Jones et al (16).

Blood was collected in EDTA-containing tubes from all patients except those admitted to the

intensive care unit (ICU) for whom blood was pooled from heparin-coated blood gas syringes

discarded in the last 12 hours. For some ICU patients, additional venous blood was also

collected in EDTA tubes. Nostril swabs (NSW) and nasopharyngeal aspirates (NPA) were

collected from the majority of the patients whereas endotracheal aspirates (ETA) were only

collected from patients with mechanical ventilation intubated in the ICU. Admitted patients

were sampled during acute disease at up to four timepoints and ICU patient material was

collected up at to ten timepoints. For this study, unless otherwise stated, the measurements

referring to acute disease were performed with samples collected at the time of study

inclusion and when patients returned for their follow-up visits at 3 and 8 months from

symptom onset. At follow-up sampling, all study individuals had been discharged (if

hospitalized) from the infectious diseases ward but some individuals (<10) who recovered

from severe COVID-19 were still in a hospital aftercare ward at the first follow-up sampling.

All study participants were confirmed SARS-CoV-2 negative by PCR at the time of follow-up

sampling, with the exception of 5 individuals who were PCR + but with high Ct values (>34).

The study was approved by the Swedish Ethical Review Authority, and performed according

to the Declaration of Helsinki. Written informed consent was obtained from all patients and

controls. For sedated patients, the denoted primary contact was contacted and asked about

the presumed will of the patient and to give initial oral and subsequently signed written

consent. When applicable, retrospective written consent was obtained from patients with

non-fatal outcomes.

Enzyme-linked immunosorbent assay (ELISA)

The presence of IgG or IgA binding against the SARS-CoV-2 Nucleocapsid (N) and Spike (S)

trimer or the Receptor Binding Domain (RBD) monomer (5, 6) in plasma and airway samples

was assessed by enzyme-linked immunosorbent assay (ELISA). Recombinant proteins were

received through the global health-vaccine accelerator platforms (GH-VAP) funded by the

Bill & Melinda Gates Foundation. Briefly, 96-half well plates were coated with 50ng/well of

the respective protein. Plates were incubated with a selected duplicate dilution that did not

provide background noise against ovalbumin used as a negative control (data not shown)

(i.e. 1:20 for plasma samples, 1:2 for NSW and NPA, and 1:5 for ETA in 5% milk/PBS buffer).

Duplicate 7-point serial dilutions were initially performed for measuring plasma IgG against

RBD during acute disease and after vaccination. The half maximal effective concentration

(EC$_{50}$) or the endpoint titer (dilution at the set OD value of 0.1) were calculated using

GraphPad Prism 9. However, since for several samples with low antibody concentration

(mostly from the asymptomatic/mild category) the EC50 was below the highest dilution used

(of 1:20) and therefore below the limit of detection (Supplementary figure 6A), the maximal

optical density (OD) at 1:20 dilution was used for most of the analyses. The relation between

maximal OD and EC50 was verified in a subset of patients with high IgG and IgA against S

(Supplementary figure 6B). To be able to compare pre- and post-vaccination antibody levels

that would, in some instances, fall below and above the lower and upper limits of detection,

the endpoint titer was used instead. The relation between EC50 and endpoint titer for these

samples, is shown in Supplementary figure 5. Detection was performed with mouse and goat

anti-human IgG or IgA HRP-conjugated secondary antibodies (clone G18-145 from BD

Biosciences and polyclonal from ThermoFisher, respectively) followed by incubation with

TMB substrate (BioLegend) which was stopped with a 1M solution of sulfuric acid. Blocking

with 5% milk/PBS buffer and washing with 0.1% Tween-20/PBS buffer were performed

between each step. Absorbance was read at 450nm and background correction at 550nm

using an ELISA reader. Data were reported as maximal absorbance i.e. OD, as stated above,

and plotted using GraphPad Prism 9. All of the antibody measurements in plasma and

respiratory samples from SARS-CoV-2 patients were run alongside with samples from two

different control groups as described above. Interestingly, low but readily detectable IgA

reactivity against S was detected in the pre-pandemic healthy controls and in the PCR-

individuals (Supplementary figure 6C). After having verified the specificity and sensitivity of

our ELISA assay for IgA detection with limiting sample dilutions (Supplementary figure 6D-E),

we hypothesize that this might be due to cross-reactivity on the shared portions of the S

protein between SARS-CoV-2 and other common cold coronaviruses. Reports have shown

that cross-reactivity between coronaviruses exists (51, 52).

Flow cytometry

Staining of cells from airway samples was performed fresh. Briefly, samples were centrifuged

at 400 g for 5 min at room temperature and cells were washed with sterile PBS. Mucus was

removed using a 70 μm cell strainer and cells were subsequently stained with the

appropriate combination of fluorescently labelled monoclonal antibodies as illustrated in

Figure 5A. Staining of PBMC was performed on previously cryopreserved samples. The

appropriate combination of fluorescently labelled monoclonal antibodies binding to

different cell surface markers and with fluorescently labelled S and RBD proteins used as

probes for antigen-specific B cells is illustrated in Figure 5C. Probes were prepared from

biotinylated proteins using a 4:1 molar ratio (protein:fluorochrome-labelled streptavidin)

considering the molecular weight of protein monomers and of the streptavidin only. The

probes were prepared using streptavidin conjugated to PE and APC for S and with BV421 for

the RBD. The gating strategy for the identification of antigen-specific memory B cells is

shown in Figure 5C. Briefly, after identification of lymphocytes in single suspension, live B

cells, (i.e. cells not expressing CD3/,CD14/CD16/CD56) were gated. From this gate, B cells

were further isolated by expression of CD19 and CD20 and then switched memory B cells

were identified as IgD-IgM-. From these, S-specific switched memory B cells were identified

by binding to both S protein probes. Further characterization was then carried out by

analyzing IgG expression (IgA+ switched memory B cells are assumed to mirror IgD-IgM-IgG-

B cells) and fluorescently labelled RBD. Stained cells from airway samples were acquired

using a BD LSRFortessa while stained PBMC were acquired using a BD FACSAria Fusion both

interfaced with the BD FACSDiva Software. Results were analyzed using BD FlowJo version

10.

Statistical analyses

All statistical analyses were performed using GraphPad Prism 9. Spearman correlation was

used to assess the interdependence of 2 different non-categorical parameters across

individuals whereas Wilcoxon matched-pairs signed rank or Mann–Whitney U tests as

appropriate, were used to assess differences or similarities for one single parameter

between 2 different groups. Kruskal -Wallis with Dunn's multiple comparisons test was used

when assessing comparison between multiple groups.

Author contributions

Experimental study design: A.C., Ka.L. and A.S-S. Clinical concept design: A.S-S., M.B., N.J.,

J.A., J.S. and A.F. Acquisition and sample processing: A.C., M.Y., S. F-J., S.V., B.Ö., E.Å., L.A.,

R.F-J., M.Ö., F.G., J.S. and M. E. Generation of data: A.C., M.Y., S. F-J., S.V., L.A. and P.C.G.

Provision of custom reagents: D.P., M.M., L.C., and N.P.K. Analysis and interpretation of

data: A.C., Ka.L. and A.S-S. Critical revision of the manuscript: all authors. Statistical analysis:

A.C., K.L. and S.O. Ka.L. and A.S-S. contributed equally to the study.

Acknowledgments

We thank the patients and healthy volunteers who have contributed to this study. We would

also like to thank medical students and hospital staff for assistance with patient sampling

and collection of clinical data, the Biomedicum BSL3 core facility, Karolinska Institutet and

Fredrika Hellgren for assistance with English editing. This work was supported by grants from

the Swedish Research Council, the Swedish Heart-Lung Foundation, the Bill & Melinda Gates

Foundation, the Knut and Alice Wallenberg Foundation through SciLifeLab and Karolinska

Institutet.

References

1.   Gandhi RT, Lynch JB, and Del Rio C. Mild or Moderate Covid-19. *N Engl J Med.*

2020;383(18):1757-66.

2.   Berlin DA, Gulick RM, and Martinez FJ. Severe Covid-19. *N Engl J Med.*

2020;383(25):2451-60.

3.   Brodin P. Immune determinants of COVID-19 disease presentation and severity. *Nat*

Med. 2021;27(1):28-33.

4.   Moghadas SM, Fitzpatrick MC, Sah P, Pandey A, Shoukat A, Singer BH, et al. The

implications of silent transmission for the control of COVID-19 outbreaks. *Proc Natl*

Acad Sci U S A. 2020.

5.   Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, and Veesler D. Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell.*

2020;181(2):281-92 e6.

6.   Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al. Cryo-EM

structure of the 2019-nCoV spike in the prefusion conformation. *Science.*

2020;367(6483):1260-3.

7.   Gaebler C, Wang Z, Lorenzi JCC, Muecksch F, Finkin S, Tokuyama M, et al. Evolution of

antibody immunity to SARS-CoV-2. *Nature.* 2021;591(7851):639-44.

8.   Atyeo C, Fischinger S, Zohar T, Slein MD, Burke J, Loos C, et al. Distinct Early

Serological Signatures Track with SARS-CoV-2 Survival. *Immunity.* 2020;53(3):524-32

e4.

9.   Gudbjartsson DF, Norddahl GL, Melsted P, Gunnarsdottir K, Holm H, Eythorsson E, et

al. Humoral Immune Response to SARS-CoV-2 in Iceland. *N Engl J Med.*

2020;383(18):1724-34.

10.   Guthmiller JJ, Stovicek O, Wang J, Changrob S, Li L, Halfmann P, et al. SARS-CoV-2

Infection Severity Is Linked to Superior Humoral Immunity against the Spike. *mBio.*

2021;12(1):e02940-20.

11.   Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody Responses to SARS-CoV-2

in Patients With Novel Coronavirus Disease 2019. *Clin Infect Dis.* 2020;71(16):2027-

34.

12.   Piccoli L, Park YJ, Tortorici MA, Czudnochowski N, Walls AC, Beltramello M, et al.

Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2 Spike Receptor-

Binding Domain by Structure-Guided High-Resolution Serology. *Cell.* 2020.

13.   Zielinski CE, Corti D, Mele F, Pinto D, Lanzavecchia A, and Sallusto F. Dissecting the

human immunologic memory for pathogens. *Immunol Rev.* 2011;240(1):40-51.

14.   Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The

SOFA (Sepsis-related Organ Failure Assessment) score to describe organ

dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of

the European Society of Intensive Care Medicine. *Intensive Care Med.*

1996;22(7):707-10.

15.   Grissom CK, Brown SM, Kuttler KG, Boltax JP, Jones J, Jephson AR, et al. A modified

sequential organ failure assessment score for critical care triage. *Disaster Med Public*

Health Prep. 2010;4(4):277-84.

16.   Falck-Jones S, Vangeti S, Yu M, Falck-Jones R, Cagigi A, Badolati I, et al. Functional

530   monocytic myeloid-derived suppressor cells increase in blood but not airways and

531   predict COVID-19 severity. *J Clin Invest.* 2021;131(6).

532   17.   Hansen CB, Jarlhelt I, Perez-Alos L, Hummelshoj Landsy L, Loftager M, Rosbjerg A, et

533   al. SARS-CoV-2 Antibody Responses Are Correlated to Disease Severity in COVID-19

534   Convalescent Individuals. *J Immunol.* 2020.

535   18.   Dan JM, Mateus J, Kato Y, Hastie KM, Yu ED, Faliti CE, et al. Immunological memory

536   to SARS-CoV-2 assessed for up to 8 months after infection. *Science.* 2021.

537   19.   Rodda LB, Netland J, Shehata L, Pruner KB, Morawski PA, Thouvenel CD, et al.

538   Functional SARS-CoV-2-Specific Immune Memory Persists after Mild COVID-19. *Cell.*

539   2021;184(1):169-83 e17.

540   20.   Juno JA, Tan HX, Lee WS, Reynaldi A, Kelly HG, Wragg K, et al. Humoral and

541   circulating follicular helper T cell responses in recovered patients with COVID-19. *Nat

542   Med.* 2020.

543   21.   Rydyznski Moderbacher C, Ramirez SI, Dan JM, Grifoni A, Hastie KM, Weiskopf D, et

544   al. Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and

545   Associations with Age and Disease Severity. *Cell.* 2020.

546   22.   Lynch KL, Whitman JD, Lacanienta NP, Beckerdite EW, Kastner SA, Shy BR, et al.

547   Magnitude and kinetics of anti-SARS-CoV-2 antibody responses and their relationship

548   to disease severity. *Clin Infect Dis.* 2020.

549   23.   Robbiani DF, Gaebler C, Muecksch F, Lorenzi JCC, Wang Z, Cho A, et al. Convergent

550   antibody responses to SARS-CoV-2 in convalescent individuals. *Nature.* 2020.

551   24.   Fahy JV, and Dickey BF. Airway mucus function and dysfunction. *N Engl J Med.*

552   2010;363(23):2233-47.

553   25.   Wang YY, Harit D, Subramani DB, Arora H, Kumar PA, and Lai SK. Influenza-binding

554   antibodies immobilise influenza viruses in fresh human airway mucus. *Eur Respir J.*

555   2017;49(1).

556   26.   Isho B, Abe KT, Zuo M, Jamal AJ, Rathod B, Wang JH, et al. Persistence of serum and

557   saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci*

558   *Immunol.* 2020;5(52).

559   27.   Reynolds HY. Immunoglobulin G and its function in the human respiratory tract.

560   *Mayo Clin Proc.* 1988;63(2):161-74.

561   28.   Pakkanen SH, Kantele JM, Moldoveanu Z, Hedges S, Hakkinen M, Mestecky J, et al.

562   Expression of homing receptors on IgA1 and IgA2 plasmablasts in blood reflects

563   differential distribution of IgA1 and IgA2 in various body fluids. *Clin Vaccine Immunol.*

564   2010;17(3):393-401.

565   29.   Sterlin D, Mathian A, Miyara M, Mohr A, Anna F, Claer L, et al. IgA dominates the

566   early neutralizing antibody response to SARS-CoV-2. *Sci Transl Med.* 2020.

567   30.   Cervia C, Nilsson J, Zurbuchen Y, Valaperti A, Schreiner J, Wolfensberger A, et al.

568   Systemic and mucosal antibody responses specific to SARS-CoV-2 during mild versus

569   severe COVID-19. *J Allergy Clin Immunol.* 2020.

570   31.   Staines HM, Kirwan DE, Clark DJ, Adams ER, Augustin Y, Byrne RL, et al. Dynamics of

571   IgG seroconversion and pathophysiology of COVID-19 infections. *medRxiv.* 2020.

572   32.   Sealy R, Webby RJ, Crumpton JC, and Hurwitz JL. Differential localization and function

573   of antibody-forming cells responsive to inactivated or live-attenuated influenza virus

574   vaccines. *Int Immunol.* 2013;25(3):183-95.

575   33.   Rudraraju R, Surman S, Jones B, Sealy R, Woodland DL, and Hurwitz JL. Phenotypes

576   and functions of persistent Sendai virus-induced antibody forming cells and CD8+ T

cells in diffuse nasal-associated lymphoid tissue typify lymphocyte responses of the

gut. *Virology.* 2011;410(2):429-36.

34.   Surman SL, Rudraraju R, Sealy R, Jones B, and Hurwitz JL. Vitamin A deficiency

disrupts vaccine-induced antibody-forming cells and the balance of IgA/IgG isotypes

in the upper and lower respiratory tract. *Viral Immunol.* 2012;25(4):341-4.

35.   Surman SL, Jones BG, Sealy RE, Rudraraju R, and Hurwitz JL. Oral retinyl palmitate or

retinoic acid corrects mucosal IgA responses toward an intranasal influenza virus

vaccine in vitamin A deficient mice. *Vaccine.* 2014;32(22):2521-4.

36.   Krammer F. The human antibody response to influenza A virus infection and

vaccination. *Nat Rev Immunol.* 2019;19(6):383-97.

37.   Su F, Patel GB, Hu S, and Chen W. Induction of mucosal immunity through systemic

immunization: Phantom or reality? *Hum Vaccin Immunother.* 2016;12(4):1070-9.

38.   Or-Guil M, Wittenbrink N, Weiser AA, and Schuchhardt J. Recirculation of germinal

center B cells: a multilevel selection strategy for antibody maturation. *Immunol Rev.*

2007;216:130-41.

39.   Lycke NY, and Bemark M. The regulation of gut mucosal IgA B-cell responses: recent

developments. *Mucosal Immunol.* 2017;10(6):1361-74.

40.   Corbett KS, Flynn B, Foulds KE, Francica JR, Boyoglu-Barnum S, Werner AP, et al.

Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in Nonhuman Primates. *N*

Engl J Med. 2020;383(16):1544-55.

41.   Corbett KS, Nason MC, Flach B, Gagne M, O' Connell S, Johnston TS, et al. Immune

Correlates of Protection by mRNA-1273 Immunization against SARS-CoV-2 Infection

in Nonhuman Primates. *bioRxiv.* 2021:2021.04.20.440647.

42. Krammer F, Srivastava K, Alshammary H, Amoako AA, Awawda MH, Beach KF, et al.

Antibody Responses in Seropositive Persons after a Single Dose of SARS-CoV-2 mRNA

Vaccine. *N Engl J Med.* 2021;384(14):1372-4.

43. Saadat S, Rikhtegaran Tehrani Z, Logue J, Newman M, Frieman MB, Harris AD, et al.

Binding and Neutralization Antibody Titers After a Single Vaccine Dose in Health Care

Workers Previously Infected With SARS-CoV-2. *JAMA.* 2021;325(14):1467-9.

44. Goel RR, Apostolidis SA, Painter MM, Mathew D, Pattekar A, Kuthuru O, et al. Distinct

antibody and memory B cell responses in SARS-CoV-2 naive and recovered individuals

following mRNA vaccination. *Sci Immunol.* 2021;6(58).

45. Levi R, Azzolini E, Pozzi C, Ubaldi L, Lagioia M, Mantovani A, et al. A cautionary note

on recall vaccination in ex-COVID-19 subjects. *medRxiv.* 2021:2021.02.01.21250923.

46. Samanovic MI, Cornelius AR, Wilson JP, Karmacharya T, Gray-Gaillard SL, Allen JR, et

al. Poor antigen-specific responses to the second BNT162b2 mRNA vaccine dose in

SARS-CoV-2-experienced individuals. *medRxiv.* 2021:2021.02.07.21251311.

47. Planas D, Bruel T, Grzelak L, Guivel-Benhassine F, Staropoli I, Porrot F, et al.

Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing

antibodies. *Nat Med.* 2021.

48. Mades A, Chellamathu P, Lopez L, Kojima N, MacMullan MA, Denny N, et al.

Detection of persistent SARS-CoV-2 IgG antibodies in oral mucosal fluid and upper

respiratory tract specimens following COVID-19 mRNA vaccination. *medRxiv.*

2021:2021.05.06.21256403.

49. Intensivvårdsregistret S. SIR:s riktlinje för registrering av SOFA. .

https://www.icuregswe.org/globalassets/riktlinjer/sofa.pdf.

50.   Charlson ME, Pompei P, Ales KL, and MacKenzie CR. A new method of classifying

624   prognostic comorbidity in longitudinal studies: development and validation. *J Chronic*

625   *Dis.* 1987;40(5):373-83.

626   51.   Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, et al. Targets

627   of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease

628   and Unexposed Individuals. *Cell.* 2020;181(7):1489-501 e15.

629   52.   Song G, He WT, Callaghan S, Anzanello F, Huang D, Ricketts J, et al. Cross-reactive

630   serum and memory B cell responses to spike protein in SARS-CoV-2 and endemic

631   coronavirus infection. *bioRxiv.* 2020.

Table 1. Clinical characterization of the SARS-CoV-2 infected cohort

Peak disease severity	Mild		Moderate		Severe		Fatal
	1	**2**	**3**	**4**	**5***	**6***	**7**
Resp. SOFA score	0	0	1	2	3	4	
Admitted (Y/N)	N	Y	Y	Y	Y	Y	Y
PFI (kPa)	> 53	> 53	< 53	< 40	< 27	< 13	-
SFI	> 400	> 400	≤ 400	≤ 315	≤ 235	< 150	
No. of individuals (%)	13 (8.8)	6 (4.1)	10 (6.8)	48 (33)	19 (13)	39 (27)	12 (8.2)
Age, mean (Range)	44 (24-72)	60 (41-72)	56 (46-78)	55 (24-76)	57 (42-74)	61 (25-77)	66 (52-78)
Male (%)	5 (38)	2 (33)	6 (60)	38 (79)	15 (79)	34 (87)	9 (75)
Days from symptoms to admission – median (Range)	-	10 (0-14)	8.5 (4-14)	10 (3-21)	7 (2-14)	10 (2-35)	7 (1-28)
Days from symptoms to inclusion "Acute" – median (Range)	9 (3-44)	11 (0-20)	13.5 (6-18)	13 (4-32)	21 (5-40)	22 (7-54)	13 (8-44)
Days from symptoms to 3-Mo follow-up – median (Range)	102 (88-136)	99,5 (82-103)	112 (81-127)	109 (46-155)	109 (48-130)	120 (53-168)	-
Days from symptoms to 8-Mo follow-up – median (Range)	232 (187-264)	238 (212-250)	245 (227-303)	247 (233-314)	241 (220-270)	254 (224-344)	-
VL (Ct value) median (Range)	27.5 (40-14)	25.0 (29-14)	26.7 (36-15)	26.8 (36-12)	25.8 (36-19)	24.0 (37-14)	20.5 (32-13)
CCI, mean (SD)	1 (2)	2 (1)	1 (1)	2 (2)	2 (1)	2 (1)	3 (1)
BMI, mean (SD)	24.1 (4.5)	25.1 (2.2)	26.0 (3.2)	30.3 (4.2)	29.2 (5.3)	28.6 (4.7)	28.6 (2.4)
Hypertension (%)	1 (8.3)	0 (0)	2 (20)	20 (42)	8 (42)	15 (38)	9 (75)
Diabetes (%)	2 (17)	0 (0)	1 (10)	14 (29)	5 (26)	9 (23)	3 (25)
Current smokers (%)	0 (0)	0 (0)	2 (20)	5 (11)	1 (5.3)	2 (5.3)	0 (0)
ACE-I (%)	0 (0)	0 (0)	0 (0)	5 (10)	1 (5.3)	4 (10)	1 (9.1)
IS drugs (%)	1 (7.7)	0 (0)	0 (0)	4 (8.3)	2 (11)	5 (13)	1 (8.3)

Resp. SOFA: Respiratory Sequential Organ Failure Assessment

PFI: PaO_2/FiO_2-index

SFI: SpO_2/FiO_2-index

VL: Viral Load

CCI: Charlson comorbidity index

BMI: body mass index

ACE-I: angiotensin converting enzyme-inhibitors

IS: immunosuppressive

*Requires mechanical ventilation

646 **Table 2.** Peak disease severity, age, gender and longitudinal sampling timeline of patients

647 vaccinated against SARS-CoV-2.

| Peak disease severity | Days from onset of symptoms | | | | Days from | | |
	Acute	3 Mo	8 Mo	Vacc.	Prime	Boost	Vaccine
1	24	129	240	338	7		AstraZeneca*
1	9	113	266	308	7		AstraZeneca*
3	15	90	250	358	7		AstraZeneca*
3	14	127	238	322	11		AstraZeneca*
4	10	100	254	345	16	13	Pfizer-BioNTec**
4	23	84	252	339	9		AstraZeneca*
6	22	98	244	329	8	8	Pfizer-BioNTec**
6	34	139	260	308		7	Pfizer-BioNTec **
6	38	144	301	324	14	28	Pfizer-BioNTec**
6	43	142	254	339	11	13	Pfizer-BioNTec**
6	21	124	226	270		21	Pfizer-BioNTec**
6	30	92		286	8	10	Moderna***

648 M=male, F=female
649 * Vaxzevria (AstraZeneca)
650 ** Comirnaty (Pfizer-BioNTec)
651 *** mRNA-1273 (Moderna)
652

Figure 1. Study and sampling overview. (A) Overview of study cohort and controls, timeline of longitudinal sampling, hospital admission/discharge, level of care and outcome for each patient. Patients are group based on peak disease severity (PDS); mild (PDS 1 and 2), moderate (PDS 3 and 4), severe (PDS 5 and 6) and fatal (PDS 7). Individual inclusion sample for each patient is color-coded based on disease severity at the time of sampling. **(B)** Overview of the anatomical compartments analyzed, and the measurements performed.

Figure 2. Systemic antibody responses, inflammation markers and other clinical parameters in relation to COVID-19 severity during acute disease. (A) Plasma IgG and IgA responses against N, S and RBD are shown together with the levels of **(B)** C-reactive protein and the neutrophil-lymphocyte ratio as a measure of systemic inflammation and with **(C)** the levels of lymphocytes, monocytes and neutrophils. Black lines indicate medians. Differences were assessed using Kruskal-Wallis with Dunn's multiple comparisons test and considered statistically significant at p<0.05. ** p<0.01, *** p<0.001, **** p<0.0001. The dashed lines indicate the normal thresholds or range values. **(D)** Correlation matrix summarizing the interrelationship observed between the clinical parameters, inflammation markers, blood corpuscles and data from systemic antibody levels measured during acute disease as indicated. The P and R values (Spearman) are shown separately in the mirrored halves of the matrix and have been color-coded as indicated.

Figure 3. Longitudinal systemic antibody responses across COVID-19 severity from acute disease up to 8 months from symptom onset. (A) Individual levels of plasma IgG and IgA (from left to right) in SARS-CoV-2 infected individuals with different peak disease severity (PDS). Black lines indicate medians and dotted lines indicate the average background level from pre-pandemic healthy controls. Mann-Whitney U was used to compare the groups and considered statistically significant at p<0.05. ** p<0.01, *** p<0.001, **** p<0.0001. **(B)** Compiled patient-matched longitudinal data from acute, 3-month- and 8-month follow-ups are shown for the levels of plasma IgG and plasma IgA against RBD. For each chart, the black dotted pattern refers to the median value whereas the grey lines connect data points from the same individuals. The number in each graph indicates the slope of the line connecting the median OD values at 3- and 8-months sampling as an estimate of the antibody decay. The asterisks indicate the statistical significance, calculated as per in Figure 3A.

Figure 4. Longitudinal airway antibody responses to RBD across COVID-19 severity from acute disease up to 8 months from symptom onset. Levels of IgG and IgA to RBD in **(A)** nostrils swabs (NSW) and **(B)** nasopharyngeal aspirates (NPA). The black lines indicate median values. Mann-Whitney U was used to compare the group and considered statistically significant at p<0.05. *p<0.05 ** p<0.01, *** p<0.001, **** p<0.0001. **(C)** Heat map generated grouping patients according to PDS showing acute and convalescent IgG and IgA titers against N, S and RBD (plasma) and RBC (NSW, NPA and ETA) for each patient. The heat map also includes data from PPHC and PCR-individuals (indicated with PDS 0). Missing data and not available samples are shown in black. **(D)** Comparison of the levels of RBD IgG/A in patient-matched NSW, NPA, endotracheal aspirates (ETA) and plasma collected at the same time point. **(E)** Spearman correlation for NPA versus plasma immunoglobulins against the RBD during acute disease. The black lines connect data points from the same individuals. Wilcoxon test was used to compare the groups separately and considered statistically significant at p<0.05. **** p<0.0001. In A) the line overlaps with not detected (ND) for IgG levels.

Figure 5. Assessment of frequencies of B cells in the respiratory tract and of circulating S-specific memory B cells. (A) Representative example with gating strategy for the identification of lymphocytes (identified as negative for CD14/16/123/66) and of total B cells (CD3-CD19+) in respiratory NPA and ETA samples. **(B)** Levels of lymphocytes and of total B cells in NPA and ETA in a subset of patients alongside with NPA from healthy controls. Mann-Whitney test was used to compare the groups and considered statistically significant at p<0.05. ** p<0.01. **(C)** Representative examples with gating strategy of SARS-Cov-2 S-specific memory B cells from one pre-pandemic healthy control, 3-month follow-up samples from one SARS-CoV-2 PCR- individual and one mild and one moderate/severe COVID-19 patient. Further characterization of S-positive memory B cells on RBD binding and B cell isotype (IgG+ or IgA+ assumed to correspond to IgD-IgM-IgG- B cells). **(D)** Bar charts show the cumulative proportion (frequency) of Spike (blue) and RBD (yellow) specific memory B cell as well as the proportion of IgG (green) vs. IgA (red) isotypes among the Spike specific memory B cells in longitudinal samples from mild (n=6) and moderate/severe (n=8) COVID-19 patients. **(E)** Frequencies of S-specific memory B cells in matched acute (filled) and 3-month follow-up (filled with black lining) PBMCs in relation to days in the subset of individuals analyzed color-coded according to PDS. Dotted lines on indicate the average background staining from PCR- and PPHC. **(F)** Levels of circulating Spike+ switched memory B cells during acute disease and convalesce in the subset of patients analyzed, as well as PPHC, color-coded according to PDS. Circles with black lining refer to data during the convalescent phase. Black triangles symbolize the PPHC. Differences were assessed using Kruskal -Wallis with Dunn's multiple comparisons test and considered statistically significant at p<0.05. ** p<0.01.

Figure 6. Vaccination and systemic and airway antibody level rebound. (A) Overview of vaccinated patients with respect with peak disease severity during COVID-19 and sampling timeline after prime and boost. The anatomical compartments analyzed and the measurements performed are also shown. Compiled patient-matched longitudinal data from acute, 3-month- and 8-month follow-ups are shown together with data from after prime and after boost for **(B)** the levels of IgG and IgA against N and RNB for plasma and **(C)** against RBD for NSW and NPA. The grey lines connect data points from the same individuals. Data are color-coded according to peak disease severity during COVID-19 and differences were assessed using Kruskal-Wallis with Dunn's multiple comparisons test and considered statistically significant at p<0.05. ** p<0.01, *** p<0.001, **** p<0.0001.