Neutralising antibodies drive Spike mediated SARS-CoV-2 evasion

Kemp SA[1]*, Collier DA[1,2,3]*, Datir R[1,2,3], Gayed S[4], Jahun A[5], Hosmillo M[5], Ferreira IATM[2,3], Rees-Spear C[1], Mlcochova P[2,3], Ines Ushiro Lumb[6], David Roberts[6], Anita Chandra[2,3], Temperton N[7], The CITIID-NIHR BioResource COVID-19 Collaboration[8], The COVID-19 Genomics UK (COG-UK) Consortium[9], Sharrocks K[4], Blane E[3], Briggs JAG[10], van Gils MJ[11], Smith KGC [2,3], Bradley JR[3,12], Smith C[14], Goldstein RA[1], Goodfellow IG[5], Smielewska A[5,13], Skittrall JP [4,14,15], Gouliouris T[4], Gkrania-Klotsas E[4], Illingworth CJR[14,16], McCoy LE[1], Gupta RK[2,3,16,17]

[1]Division of Infection and Immunity, University College London, London, UK.

[2] Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Cambridge, UK.

[3]Department of Medicine, University of Cambridge, Cambridge, UK.

[4]Department of Infectious Diseases, Cambridge University NHS Hospitals Foundation Trust, Cambridge, UK.

[5]Department of Pathology, University of Cambridge, Cambridge

[6]NHS Blood and Transplant, London, UK

[7]Viral Pseudotype Unit, Medway School of Pharmacy, University of Kent, UK

[8]The CITIID-NIHR BioResource COVID-19 Collaboration, see appendix for author list

[9]The COVID-19 Genomics UK (COG-UK) Consortium, https://www.cogconsortium.uk. Full list of consortium names and affiliations are in the appendix

[10]Medical Research Council Laboratory of Molecular Biology, Cambridge, UK.

[11]Department of Medical Microbiology, Academic Medical Center, University of Amsterdam, Amsterdam Institute for Infection and Immunity, Amsterdam, Netherlands

[12] NIHR Cambridge Clinical Research Facility, Cambridge, UK.

[13]Department of Virology, Cambridge University NHS Hospitals Foundation Trust

[14]Department of Applied Mathematics and Theoretical Physics, University of Cambridge, UK

[15]Clinical Microbiology and Public Health Laboratory, Addenbrookes' Hospital, Cambridge, UK

[16] MRC Biostatistics Unit, University of Cambridge, Cambridge, UK

[17]Africa Health Research Institute, Durban, South Africa

*equal contribution

Address for correspondence:

Ravindra K. Gupta

Cambridge Institute for Therapeutic Immunology and Infectious Diseases

Jeffrey Cheah Biomedical Centre

Puddicombe Way

Cambridge CB2 0AW, UK

Tel: +44 1223 331491

rkg20@cam.ac.uk

Key words: SARS-CoV-2; COVID-19; antibody escape, Convalescent plasma; neutralising

antibodies; mutation; evasion; resistance; immune suppression

Abstract

SARS-CoV-2 Spike protein is critical for virus infection via engagement of ACE2, and amino acid

variation in Spike is increasingly appreciated. Given both vaccines and therapeutics are

designed around Wuhan-1 Spike, this raises the theoretical possibility of virus escape,

particularly in immunocompromised individuals where prolonged viral replication occurs. Here

we report fatal SARS-CoV-2 escape from neutralising antibodies in an immune suppressed

individual treated with convalescent plasma, generating whole genome ultradeep sequences by

both short and long read technologies over 23 time points spanning 101 days. Little

evolutionary change was observed in the viral population over the first 65 days despite two

courses of remdesivir. However, following convalescent plasma we observed dynamic virus

population shifts, with the emergence of a dominant viral strain bearing D796H in S2 and

ΔH69/ΔV70 in the S1 NTD of the Spike protein. As serum neutralisation waned, viruses with the

escape genotype diminished in frequency, before returning during a final, unsuccessful course

of convalescent plasma. *In vitro*, the Spike escape variant conferred decreased sensitivity to

multiple units of convalescent plasma/sera from different recovered patients, whilst

maintaining infectivity similar to wild type. These data reveal strong positive selection on SARS-
CoV-2 during convalescent plasma therapy and identify the combination of Spike mutations
D796H and ΔH69/ΔV70 as a broad antibody resistance mechanism against commonly occurring
antibody responses to SARS-CoV-2.

Introduction

SARS-CoV-2 is an RNA betacoronavirus, with closely related viruses identified in pangolins and

bats[1,2]. RNA viruses have inherently higher rates of mutation than DNA viruses such as

Herpesviridae[3]. The capacity for successful adaptation is exemplified by the Spike D614G

mutation, that arose in China and rapidly spread worldwide[4], now accounting for more than

90% of infections. The mutation appears to increase infectivity and transmissibility in animal

models[5]. Although the SARS-CoV-2 Spike protein is critical for virus infection via engagement of

ACE2, substantial Spike amino acid variation is being observed in circulating viruses[6]. Logically,

mutations in the receptor binding domain (RBD) of Spike are of particular concern due to the

RBD being targeted by neutralising antibodies and therapeutic monoclonal antibodies.

Deletions in the N-terminal domain (NTD) of Spike S1 are also being increasingly recognised,

both within hosts[7] and across individuals[8]. The evolutionary basis for the emergence of

deletions is unclear at present but could be related to escape from immunity or to enhanced

fitness/transmission. The most notable deletion in terms of frequency is ΔH69/ΔV70. This

double deletion has been detected in multiple unrelated lineages, including the recent 'Cluster

5' mink related strain in the North Jutland region of Denmark (https://files.ssi.dk/Mink-cluster-

5-short-report_AFO2). There it was associated with the RBD mutation Y453F in almost 200

individuals. Another European cluster in GISAID includes ΔH69/ΔV70 along with the RBD

mutation N439K.

Although ΔH69/ΔV70 has been detected multiple times, within-host emergence remains

undocumented and the reasons for its selection are unknown. Here we document real time

SARS-CoV-2 emergence of ΔH69/ΔV70 in response to convalescent plasma therapy in an

immunocompromised human host, demonstrating broad antibody escape in combination with

the S2 mutation D796H.

Results

Clinical case history of SARS-CoV-2 infection in setting of immune-compromised host

Please contact authors for clinical details. Some indices are given in Supplementary table 1 and

Supplementary Figure 1 and Supplementary Figure 2). Figure 1 summarises the timing of

treatment.

Negative SARS-CoV-2 specific antibodies and requirement for convalescent plasma (CP)

Given the history of B cell depletion therapy and hypogammaglobulinemia we measured serum

SARS-CoV-2 specific antibodies over the course of the admission. Total serum antibodies to

SARS-CoV-2 were tested at days 44 and 50 by S protein immunoassay (Siemens). Results were

negative. Three units (200mL each) of convalescent plasma (CP) from three independent

donors were obtained from the NHS Blood and Transfusion Service and administered on

compassionate use basis. These had been assayed for antibody titres using the validated

Euroimmun assay (Supplementary figure 3). Patient serum was subsequently positive for SARS-

CoV-2 specific antibodies by S protein immunoassay (Siemens) in the hospital diagnostic

laboratory on days 68, 90 and 101.

Virus genomic comparative analysis of 23 sequential respiratory samples over 101 days

The majority of samples were respiratory samples from nose and throat or endotracheal

aspirates during the period of intubation. Ct values ranged from 16-34 and all 23 respiratory

samples were successfully sequenced by standard long read approach as per the ARTIC protocol

implemented by COG-UK; of these 20 additionally underwent short-read deep sequencing using

the Illumina platform. There was generally good agreement between the methods, though as

expected nanopore demonstrated greater error at low variant frequencies (<10%)

(Supplementary Figure 4). We detected no evidence of recombination, based on two

independent methods.

Maximum likelihood analysis of patient-derived whole genome consensus sequences
demonstrated clustering with other local sequences from the same region (Figure 2A). The
infecting strain was assigned to lineage 20B bearing the D614G Spike variant. Environmental
sampling showed evidence of virus on surfaces such as telephone and call bell but not in air on
days 59, 92 and 101. Sequencing of these surface viruses showed clustering with those derived
from the respiratory tract (Figure 2B). All samples were consistent with having arisen from a
single viral population. In our phylogenetic analysis, we included sequential sequences from
three other local patients identified with persistent viral RNA shedding over a period of 4 weeks
or more (Supplementary Table 2). Viruses from these individuals showed very little divergence
in comparison to the case patient (Figure 2B) and none showed amino acid changes in Spike
over time. We additionally inferred a maximum likelihood phylogeny comparing sequences
from these three local individuals and two long term immunosuppressed SARS-CoV-2 'shedders'
recently reported[7,9], (Figure 2B). While the sequences from Avanzato et al showed a pattern of
evolution more similar to two of the three other local patients, the case patient showed
significant diversification with a mutation rate of 30 per year (Supplementary table 2).

Further investigation of the sequence data suggested the existence of an underlying structure
to the viral population in our patient, with samples collected at days 93 and 95 being rooted
within, but significantly divergent from the original population (Figure 2B and 3A). The
relationship of the divergent samples to those at earlier time points rules out the possibility of
superinfection. The increased divergence of sequences does not necessarily indicate selection;
a spatially compartmentalised subset of viruses, smaller in number than the main viral
population, would be expected to evolve more quickly than the main population due to the
increased effect of genetic drift [10,11].

Virus population structure changes following convalescent plasma and remdesivir
All samples tested positive by RT-PCR and there was no sustained change in Ct values
throughout the 101 days following the first two courses of remdesivir (days 41 and 54), or the

first two units of convalescent plasma (days 63 and 65). According to nanopore data, no
polymorphisms occurred over the first 60 days at consensus level (Figure 3A). However, short
read deep sequence Illumina data revealed minority polymorphisms in the viral population
during this period (Figure 3B). For example, T39I in ORF7a reached a majority frequency of 77%
on day 44, arising *de novo* and increased in frequency during the first period of the infection
(Supplementary Figure 6).
152
In contrast to the early period of infection, between days 66 and 82 a dramatic shift in virus
population structure was observed, with the near-fixation of D796H in S2 along with
ΔH69/ΔV70 in the S1 N-terminal domain (NTD) at day 82. This was identified in a nose and
throat swab sample with high viral load as indicated by Ct of 23 (Figure 4). The deletion was
not detected at any point prior to the day 82 sample, even as minority variants by short read
deep sequencing.
159
On Days 86 and 89, viruses collected were characterised by the Spike mutations Y200H and
T240I, with the deletion/mutation pair observed on day 82 having fallen to very low frequency.
Sequences collected on these days formed a distinct branch at the bottom of the phylogeny in
Figure 4 but were clearly associated with the remainder of the samples, suggesting that they
did not result from superinfection (Supplementary Figure 7), and further were not significantly
divergent from the bulk of the viral population (Supplementary Figure 5).
166
Sequencing of a nose and throat swab sample at day 93 again showed D796H along with
ΔH69/ΔV70 at <10% abundance, along with an increase in a virus population characterised by
Spike mutations P330S at the edge of the RBD and W64G in S1 NTD. This new lineage reached
near 100% abundance at day 93. Viruses with the P330S variant were detected in two
independent samples from different sampling sites, ruling out the possibility of contamination.
The divergence of these samples from the remainder of the population, noted above, suggests
the possibility of their resulting from the stochastic emergence, in the upper respiratory tract,
of a previously unobserved subpopulation of viruses (Supplementary Figure 5).

Following the third course of remdesivir (day 93) and third CP (day 95), we observed a re-
emergence of the D796H + ΔH69/ΔV70 viral population. The inferred linkage of D796H and
ΔH69/ΔV70 was maintained as evidenced by the highly similar frequencies of the two variants,
suggesting that the third unit of CP led to the re-emergence of this viral strain under renewed
positive selection. Ct values remained low throughout this period with hyperinflammation,
eventually leading to multi-organ failure and death at day 102. The repeated increase in
frequency of the novel viral strain during CP therapy strongly supports the hypothesis that the
deletion/mutation combination conferred antibody escape properties.

ΔH69/ΔV70 + D796H confers impaired neutralisation by multiple convalescent plasma units
and sera from recovered COVID-19 patients
Using lentiviral pseudotyping we expressed wild type and ΔH69/ΔV70 + D796H mutant Spike
protein in enveloped virions and compared neutralisation activity of CP against these viruses.
This system has been shown to give similar results to replication competent virus[12,13]. We first
tested infection capacity over a single round of infection and found that ΔH69/ΔV70 + D796H
had similar infectivity to wild type (both in a D614G background, Figure 5A, B). The ΔH69/ΔV70
+ D796H mutant was partially resistant to the first two CP units (Figure 5C, Table 1A). In
addition, patient derived serum from days 64 and 66 (one day either side of CP2 infusion)
similarly showed lower potency against the mutant (Figure 5C, Table 1A). The repeated
observation of D796H + ΔH69/ΔV70 emergence and positive selection strengthens the
hypothesis that these variants were the key drivers of antibody escape. Experimentally, the
D796H + ΔH69/ΔV70 mutant also demonstrated reduced susceptibility to the CP3 administered
on day 95, explaining its re-emergence (Figure 5C, Table 1A).

Given reduced susceptibility of the mutants to at least two units of CP, and the expansion of
sequences bearing ΔH69/ΔV70, we hypothesised that this represented a broad escape
mechanism. We therefore screened antiviral neutralisation activity in sera from five recovered
patients against the mutant and wild type viruses (Figure 5D). We observed that the mutant

was indeed significantly less susceptible to four of five randomly selected sera, with the fifth

showing reduced susceptibility that did not reach statistical significance (Table 1B). Fold change

reductions in susceptibility of the mutant were as high as ten-fold compared to wild type (Table

1B).

In order to probe the impact of the D796H and ΔH69/ΔV70 mutations on potency of

monoclonal antibodies (mAbs) targeting Spike, we screened neutralisation activity of a panel of

seven neutralizing mAbs across a range of epitope clusters[13] (Figure 5E). We observed no

differences in neutralisation between single mutants and wild type, suggesting that the

mechanism of escape was likely outside these epitopes.

In order to understand the mechanisms that might confer resistance to antibodies we

examined a published Spike structure and annotated it our residues of interest (Figure 6). This

analysis showed that ΔH69/ΔV70 is in a disordered, glycosylated loop at the very tip of the

NTD, and therefore could alter binding of antibodies. ΔH69/V70 is close to the binding site of

the polyclonal antibodies derived from COV57 plasma, indicating the sera tested here may

contain similar antibodies [14,15]. D796H is in an exposed loop in S2 (Figure 6) and appears to be in

a region frequently targeted by antibodies[16], despite mutations at position 796 being rare

(Supplementary table 4).

Discussion

Here we have documented a repeated evolutionary response by SARS-CoV-2 against antibody

therapy during the course of a persistent and eventually fatal infection in an

immunocompromised host. The observation of potential selection for specific variants

coinciding with the presence of antibodies from convalescent plasma is supported by the

experimental finding of reduced susceptibility of these viruses to plasma. Further, we were

able to document real-time emergence of a variant ΔH69/ΔV70 in the NTD of Spike that has

been increasing in frequency in Europe. In the case we report that it was not clear that the

emergence of the antibody escape variant was the primary reason for treatment failure.

However, given that both vaccines and therapeutics are aimed at Spike, our study raises the
possibility of virus evasion, particularly in immune suppressed individuals where prolonged viral
replication occurs.

Our observations represent a very rare insight, and only possible due to lack of antibody
responses in the individual following administration of the B cell depleting agent rituximab for
lymphoma, and an intensive sampling course undertaken due to concerns about persistent
shedding and risk of nosocomial transmission. Persistent viral replication and the failure of
antiviral therapy allowed us to define the viral response to convalescent plasma, similar to a
recent report on asymptomatic long term shedding with four sequences over 105 days[9],
although the reported shifts in genetic composition of the viral population could not be
explained phenotypically. Another common finding is the very low neutralisation activity in
serum post transfusion of CP with waning as expected. Apart from the difference in the
outcome of infection (severe, fatal disease versus asymptomatic disease and clearance),
critically important differences in our study include: 1. The intensity of sampling and use of both
long and short read sequencing to verify variant calls, thereby providing a unique scientific
resource for longitudinal population genetic analysis. 2. The close alignment between the
genetic composition of the viral population and CP administration, with an experimentally
verified resistant strain emerging, falling to low frequency, and then rising again under CP
selection. 3. Real time detection of emergence of a variant, ΔH69/ΔV70, that is increasing in
frequency in Europe, and shown here to affect neutralization by multiple COVID-19 patient
derived sera.

We have noted in our analysis the potential influence of compartmentalised viral replication
upon the sequences recovered in upper respiratory tract samples. Both population genetic and
small animal studies have shown a lack of reassortment between influenza viruses within a
single host during an infection, suggesting that acute respiratory viral infection may be
characterised by spatially distinct viral populations[17,18]. In the analysis of data, it is important to
distinguish genetic changes which occur in the primary viral population from apparent changes

that arise from the stochastic observation of spatially distinct subpopulations in the host. While

the samples we observe on days 93 and 95 of infection are genetically distinct from the others,

the remaining samples are consistent with arising from a consistent viral population, supporting

the finding of a reversion and subsequent regain of antibody resistance. We note that in a

study of SARS-CoV-2, Choi et al reported the detection in postmortem tissue of viral RNA not

only in lung tissue, but also in the spleen, liver, and heart[7]. Mixing of virus from different

compartments, for example via blood, or movement of secretions from lower to upper

respiratory tract, could lead to fluctuations in viral populations at particular sampling sites.

Experiments in animal models with sampling of different replication sites could allow a better

understanding of SARS-CoV-2 population genetics and enable prediction of escape variants

following antibody based therapies.

This is a single case report and therefore limited conclusions can be drawn about

generalisability.

In addition to documenting the emergence of SARS-CoV-2 Spike ΔH69/ΔV70 + D796H *in vivo*,

conferring broad reduction in susceptibility to serum/plasma polyclonal, but no effect on a set

of predominantly RBD-targeting monoclonal antibodies, these data highlight that infection

control measures need to be specifically tailored to the needs of immunocompromised

patients. The data also highlight caution in interpretation of CDC guidelines that recommend 20

days as the upper limit of infection prevention precautions in immune compromised patients

who are afebrile[19]. Due to the difficulty with culturing clinical isolates, use of surrogates for

infectious virus such as sgRNA are warranted[20]. However, where detection of ongoing viral

evolution is possible, this serves as a clear proxy for the existence of infectious virus. In our case

we detected environmental contamination whilst in a single occupancy room and the patient

was moved to a negative-pressure high air-change infectious disease isolation room.

The clinical efficacy of CP has been called into question recently[21], and our data suggest caution

in use of CP in patients with immune suppression of both T cell and B cell arms. In such cases,

the antibodies administered have little support from cytotoxic T cells, thereby reducing chances

of clearance and raising the potential for escape mutations. Whilst we await further data, CP
administered for clinical need in immune suppression, should ideally be undertaken in single
occupancy rooms with enhanced infection control precautions, including SARS-CoV-2
environmental sampling and real-time sequencing.

Table 1: Neutralisation activity of convalescent plasma and patient sera taken in between CP
doses against SARS-CoV-2 viral variants observed in A. patient case. B. five unselected
patients with recovered COVID-19. Shown are IC50 dilution titres - the CP or serum reciprocal
dilution at which 50% of virus infectivity is lost against wild type (baseline) or D796H + Δ69/70
mutant Spike protein expressed on the surface of a lentiviral particle. Fold change is the
difference between the two viruses, and values below 1 indicate that the D796H/Δ69/70
mutant is less sensitive to the CP or serum. Only the D66 serum sample failed to demonstrate
significantly higher resistance of the D796H/Δ69/70 mutant.

A.	IC50 as reciprocal dilution (95% CI)		Fold Change IC50 (95% CI)
	WT	D796H + Δ69/70	
CP1	2480.0 (2104.0 - 2965.0)	924.2 (634.1 - 1316.0)	0.37 (0.24 - 0.56)
D64 serum	163.7 (106.6 - 256.4)	55.6 (29.5 - 100.6)	0.27 (0.12 - 0.77)
CP2	447.1 (305.2 - 670.1)	196 (129.8 - 294.7)	0.47 (0.25 - 0.86)
D66 serum	175.7 (90.7 - 410.4)	85.5 (48.2 - 149.8)	0.82 (0.29 - 1.96)
CP3	280.2 (172.8 - 480.3)	115 (75.0 - 175.5)	0.47 (0.24 - 0.91)

B.	IC50 as reciprocal dilution (95% CI)		Fold Change IC50 (95% CI)
	WT	D796H + Δ69/70	
Serum 1	994.1 (852.7-1159.0)	755.8 (426.7-1269.0)	0.71 (0.39-1.33)
Serum 2	2107.0 (1680.0-2662.0)	1045.0 (745.9-1387.0)	0.42 (0.27-0.70)

Serum 3	3241.0 (2377.0-4643.0)	347.4 (234.4-505.4)	0.11 (0.07-0.17)
Serum 4	17368.0 (14357.0-22130.0)	2965.0 (2287.0-3767.0)	0.26 (0.20-0.32)
Serum 5	52735.0 (34276.0-93957.0)	3871.0 (3525.0-4252.0)	0.20 (0.17-0.24)

Ethics

The study was approved by the East of England – Cambridge Central Research Ethics Committee (17/EE/0025). Written informed consent was obtained from both the patient and family. Additional controls with COVID-19 were enrolled to the NIHR BioResource Centre Cambridge under ethics review board (17/EE/0025).

Acknowledgements

We are immensely grateful to the patient and his family. We would also like to thank the staff at CUH and the NIHR Cambridge Clinical Research Facility. We would like to thank Dr Ruthiran Kugathasan and Professor Wendy Barclay for helpful discussions and Dr Martin Curran, Dr William Hamilton, and Dr. Dominic Sparkes. We would like to thank Prof Andres Floto and Prof Ferdia Gallagher. We thank Dr James Voss for the kind gift of HeLa cells stably expressing ACE2. COG-UK is supported by funding from the Medical Research Council (MRC) part of UK Research & Innovation (UKRI), the National Institute of Health Research (NIHR) and Genome Research Limited, operating as the Wellcome Sanger Institute. RKG is supported by a Wellcome Trust Senior Fellowship in Clinical Science (WT108082AIA). LEM is supported by a Medical Research Council Career Development Award (MR/R008698/1). SAK is supported by the Bill and Melinda Gates Foundation via PANGEA grant: OPP1175094. DAC is supported by a Wellcome Trust Clinical PhD Research Fellowship. CJRI acknowledges MRC funding (ref: MC_UU_00002/11). This research was supported by the National Institute for Health Research (NIHR) Cambridge Biomedical Research Centre, the Cambridge Clinical Trials Unit (CCTU) and by the UCL Coronavirus Response Fund and made possible through generous donations from UCL's supporters, alumni, and friends (LEM). JAGB is supported by the Medical Research Council

(MC_UP_1201/16). IG is a Wellcome Senior Fellow and supported by the Wellcome Trust

(207498/Z/17/Z).

Author contributions

Conceived study: RKG, SAK, DAC, AS, TG, EGK

Designed experiments: RKG, SAK, DAC, LEM, JAGB, EGK, AC, NT, AC, CS

Performed experiments: SAK, DAC, LEM, RD, CRS, AJ, IATMF, KS, TG, CJRI, BB, JS, MJvG

Interpreted data: RKG, SAK, DAC, PM, LEM, JAGB, PM, SG, KS, TG, JB, KGCS, IG, CJRI, JAGB, IUL,

DR, JS, BB, RAG

Conflicts of interest

The authors declare no conflicts of interest

Methods

Clinical Sample Collection and Next generation sequencing

Serial samples were collected from the patient periodically from the lower respiratory tract

(sputum or endotracheal aspirate), upper respiratory tract (throat and nasal swab), and from

stool. Nucleic acid extraction was done from 500μl of sample with a dilution of MS2

bacteriophage to act as an internal control, using the easyMAG platform (Biomerieux, Marcy-

l'Étoile) according to the manufacturers' instructions. All samples were tested for presence of

SARS-CoV-2 with a validated one-step RT q-PCR assay developed in conjunction with the Public

Health England Clinical Microbiology [22]. Amplification reaction were all performed on a

Rotorgene™ PCR instrument. Samples which generated a CT of ≤36 were considered to be

positive.

Sera from recovered patients in the COVIDx study[23] were used for testing of neutralisation

activity by SARS-CoV-2 mutants.

For viral genomic sequencing, total RNA was extracted from samples as described. Samples

were sequenced using MinION flow cells version 9.4.1 (Oxford Nanopore Technologies)

following the ARTICnetwork V3 protocol (https://dx.doi.org/10.17504/protocols.io.bbmuik6w)

and BAM files assembled using the ARTICnetwork assembly pipeline

(https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html). A representative set of

10 sequences were selected and also sequenced using the Illumina MiSeq platform. Amplicons

were diluted to 2 ng/µl and 25 µl (50 ng) were used as input for each library preparation

reaction. The library preparation used KAPA Hyper Prep kit (Roche) according to manufacturer's

instructions. Briefly, amplicons were end-repaired and had A-overhang added; these were then

ligated with 15mM of NEXTflex DNA Barcodes (Bio Scientific, Texas, USA). Post-ligation products

were cleaned using AMPure beads and eluted in 25 µl. Then, 20 µl were used for library

amplification by 5 cycles of PCR. For the negative controls, 1ng was used for ligation-based

library preparation. All libraries were assayed using TapeStation (Agilent Technologies,

California, USA) to assess fragment size and quantified by QPCR. All libraries were then pooled

in equimolar accordingly. Libraries were loaded at 15nM and spiked in 5% PhiX (Illumina,

California, USA) and sequenced on one MiSeq 500 cycle using a Miseq Nano v2 with 2x 250

paired-end sequencing. A minimum of ten reads were required for a variant call.

Bioinformatics Processes

For long-read sequencing, genomes were assembled with reference-based assembly and a

curated bioinformatics pipeline with 20x minimum coverage across the whole-genome [24]. For

short-read sequencing, FASTQs were downloaded, poor-quality reads were identified and

removed, and both Illumina and PHiX adapters were removed using TrimGalore v0.6.6 [25].

Trimmed paired-end reads were mapped to the National Center for Biotechnology Information

SARS-CoV-2 reference sequence MN908947.3 using MiniMap2-2.17 with arguments -ax and sr

[26]. BAM files were then sorted and indexed with samtools v1.11 and PCR optical duplicates

removed using Picard (http://broadinstitute.github.io/picard). Single nucleotide polymorphisms

(SNPs) were called using Freebayes v1.3.2 [27] with a ploidy setting of 1, minimum allele

frequency of 0.20 and a minimum depth of five reads. Finally, a consensus sequences of nucleic

acids with a minimum whole-genome coverage of at least 20× were generated with BCFtools

using a 0% majority threshold.

Phylogenetic Analysis

All available full-genome SARS-CoV-2 sequences were downloaded from the GISAID database

(http://gisaid.org/) [28] on 17th September. Duplicate and low-quality sequences (>5% N regions)

were removed, leaving a dataset of 138,472 sequences with a length of >29,000bp. All

sequences were sorted by name and only sequences sequenced with United Kingdom / England

identifiers were retained. From this dataset, a subset of 250 sequences were randomly

subsampled using seqtk (https://github.com/lh3/seqtk). These 250 sequences were aligned to

the 23 patient sequences, as well as three other control patients (persistent long-term shedders

from the same hospital) (Supplementary Table 2) and the SARS-CoV-2 reference strain

MN908947.3, using MAFFT v7.473 with automatic flavour selection [29]. Major SARS-CoV-2 clade

memberships were assigned to all sequences using the Nextclade server v0.8

(https://clades.nextstrain.org/).

Maximum likelihood phylogenetic trees were produced using the above curated dataset using

IQ-TREE v2.1.2 [30]. Evolutionary model selection for trees were inferred using ModelFinder [31]

and trees were estimated using the GTR+F+I model with 1000 ultrafast bootstrap replicates [32].

All trees were visualised with Figtree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/), rooted

on the SARS-CoV-2 reference sequence and nodes arranged in descending order. Nodes with

bootstraps values of <50 were collapsed using an in-house script.

Molecular substitution (clock) rates for the index patient, as well as three long-term shedders

and two recently described immunocompromised patients from literature, were estimated

using BEAST v2.6.3 [34] using a HKY substitution model with 4 rate categories drawn from a

gamma distribution, a strict clock and a coalescent exponential population tree prior. MCMC

was run for 100 million iterations excluding a 15% burn-in. Tracer v1.7.1 was used to analyse

the BEAST trace in order extract the clock rate and ensure convergence had occurred.

In-depth allele frequency variant calling

The SAMFIRE package[35] was used to call allele frequency trajectories from BAM file data. Reads
were included in this analysis if they had a median PHRED score of at least 30, trimming the
ends of reads to achieve this if necessary. Nucleotides were then filtered to have a PHRED
score of at least 30; reads with fewer than 30 such reads were discarded. Distances between
sequences, accounting for low-frequency variant information, was also conducted using
SAMFIRE. The sequence distance metric, described in an earlier paper [11], combines allele
frequencies across the whole genome. Where L is the length of the genome, we define q(t) as a
4 x L element vector describing the frequencies of each of the nucleotides A, C, G, and T at each
locus in the viral genome sampled at time t. For any given locus i in the genome we calculate
the change in allele frequencies between the times t_1 and t_2 via a generalisation of the
Hamming distance

$$d\big(q_i(t_1), q_i(t_2)\big) = \frac{1}{2} \sum_{a \in \{A,C,G,T\}} |q_i^a(t_1) - q_i^a(t_2)|$$

435 where the vertical lines indicate the absolute value of the difference. These statistics were then
436 combined across the genome to generate the pairwise sequence distance metric

$$D\big(\boldsymbol{q}(t_1), \boldsymbol{q}(t_2)\big) = \sum_i d\big(q_i(t_1), q_i(t_2)\big)$$

The Mathematica software package was to conduct a regression analysis of pairwise sequence
distances against time, leading to an estimate of a mean rate of within-host sequence
evolution. In contrast to the phylogenetic analysis, this approach assumed the samples
collected on days 93 and 95 to arise via stochastic emission from a spatially separated
subpopulation within the host, leading to a lower inferred rate of viral evolution for the bulk of
the viral population.

Inference of selection

Under the assumption of a large effective population size, a deterministic one-locus model of

selection was fitted to genome sequence data describing changes in the frequency of the

variant T39I. Where q(t) represents the frequency of a single variant at time t, we used a

maximum likelihood method to infer the initial variant frequency at time t=1, and the selection

coefficient s, for times from the initial time point to the disappearance of the variant from the

population. Specifically, where $n(t_i)$ and $N(t_i)$ were the number of observations of the variant

and the total read depth at that locus at time t_i, we fitted the model

$$q(t) = \frac{q(1)e^{st}}{1 - q(1) + q(1)e^{st}}$$

so as to maximise the binomial likelihood

$$L = \sum_i \log \binom{N_i}{n_i} q(t_i)^{n_i} \big(1 - q(t_i)\big)^{N - n_i}$$

459 A similar calculation was performed to estimate the mean effective selection (incorporating

460 intrinsic selection for the variant plus linkage with other selected alleles) acting upon the

461 variant D796H during the final period of CP therapy. In this case selection was modelled as

462 being time-dependent, kicking in with the administration of therapy. We fitted the model:

$$q(t) = q \ if \ t < \tau$$
$$q(t) = \frac{qe^{s(t-\tau)}}{1 - q + qe^{s(t-\tau)}} \, if \ t \geq \tau$$

to the data, setting $\tau = 95$.

Recombination Detection

All sequences were tested for potential recombination, as this would impact on evolutionary

estimates. Potential recombination events were explored with nine algorithms (RDP, MaxChi,

SisScan, GeneConv, Bootscan, PhylPro, Chimera, LARD and 3SEQ), implemented in RDP5 with

default settings [36]. To corroborate any findings, ClonalFrameML v1.12 [37] was also used to infer

recombination breakpoints. Neither programs indicated evidence of recombination in our data.

Structural Viewing

The Pymol Molecular Graphics System v2.4.0 (https://github.com/schrodinger/pymol-open-

source/releases) was used to map the location of the four spike mutations of interested onto a

SARS-CoV-2 spike structure visualised by Wrobel et al (PDB: 6ZGE) [38].

Generation of Spike mutants

Amino acid substitutions were introduced into the D614G pCDNA_SARS-CoV-2_Spike plasmid

as previously described[39] using the QuikChange Lightening Site-Directed Mutagenesis kit,

following the manufacturer's instructions (Agilent Technologies, Inc., Santa Clara, CA).

Pseudotype virus preparation

Viral vectors were prepared by transfection of 293T cells by using Fugene HD transfection

reagent (Promega). 293T cells were transfected with a mixture of 11ul of Fugene HD, 1μg of

pCDNAΔ19Spike-HA, 1ug of p8.91 HIV-1 gag-pol expression vector[40,41], and 1.5μg of pCSFLW

(expressing the firefly luciferase reporter gene with the HIV-1 packaging signal). Viral

supernatant was collected at 48 and 72h after transfection, filtered through 0.45um filter and

stored at -80˚C. The 50% tissue culture infectious dose ($TCID_{50}$) of SARS-CoV-2 pseudovirus

was determined using Steady-Glo Luciferase assay system (Promega).

Serum/plasma pseudotype neutralization assay

Spike pseudotype assays have been shown to have similar characteristics as neutralisation

testing using fully infectious wild type SARS-CoV-2[12].Virus neutralisation assays were performed

on 293T cell transiently transfected with ACE2 and TMPRSS2 using SARS-CoV-2 Spike

pseudotyped virus expressing luciferase[42]. Pseudotyped virus was incubated with serial dilution

of heat inactivated human serum samples or convalescent plasma in duplicate for 1h at 37˚C.

Virus and cell only controls were also included. Then, freshly trypsinized 293T ACE2/TMPRSS2

expressing cells were added to each well. Following 48h incubation in a 5% CO_2 environment at

37°C, the luminescence was measured using Steady-Glo Luciferase assay system (Promega).

mAb pseudotype neutralisation assay

Virus neutralisation assays were performed on HeLa cells stably expressing ACE2 and using

SARS-CoV-2 Spike pseudotyped virus expressing luciferase as previously described[43].

Pseudotyped virus was incubated with serial dilution of purified mAbs[13] in duplicate for 1h at

37°C. Then, freshly trypsinized HeLa ACE2- expressing cells were added to each well. Following

48h incubation in a 5% CO_2 environment at 37°C, the luminescence was measured using Bright-

Glo Luciferase assay system (Promega) and neutralization calculated relative to virus only

controls. IC50 values were calculated in GraphPad Prism.

Data Availability

Long-read sequencing data that support the findings of this study have been deposited in the

NCBI SRA database with the accession codes SAMN16976824 - SAMN16976846 under

BioProject PRJNA682013 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA682013). Short reads

and data used to construct figures were deposited at https://github.com/Steven-

Kemp/sequence_files.

References

1 Zhang, T., Wu, Q. & Zhang, Z. Probable pangolin origin of SARS-CoV-2 associated with the

COVID-19 outbreak. *Current Biology* (2020).

2 Xiao, K. *et al.* Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins. *Nature* **583**,

286-289, doi:10.1038/s41586-020-2313-x (2020).

3 Sanjuán, R. & Domingo-Calap, P. Mechanisms of viral mutation. *Cell Mol Life Sci* **73**, 4433-4448,

doi:10.1007/s00018-016-2299-6 (2016).

4 Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases

Infectivity of the COVID-19 Virus. *Cell*, doi:10.1016/j.cell.2020.06.043 (2020).

5 Yurkovetskiy, L. *et al.* Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein

Variant. *Cell* **183**, 739-751 e738, doi:10.1016/j.cell.2020.09.032 (2020).

6   Korber, B. *et al.* Tracking changes in SARS-CoV-2 Spike: evidence that D614G increases infectivity
of the COVID-19 virus. *Cell* **182**, 812-827. e819 (2020).
7   Choi, B. *et al.* Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *N Engl*
J Med, doi:10.1056/NEJMc2031364 (2020).
8   McCarthy, K. R. *et al.* Natural deletions in the SARS-CoV-2 spike glycoprotein drive antibody
escape. *bioRxiv*, 2020.2011.2019.389916, doi:10.1101/2020.11.19.389916 (2020).
9   Avanzato, V. A. *et al.* Case Study: Prolonged infectious SARS-CoV-2 shedding from an
asymptomatic immunocompromised cancer patient. *Cell* (2020).
10   Wright, S. Size of population and breeding structure in relation to evolution. *Science*, 430-431
(1938).
11   Lumby, C. K., Zhao, L., Breuer, J. & Illingworth, C. J. A large effective population size for
established within-host influenza virus infection. *Elife* **9**, doi:10.7554/eLife.56915 (2020).
12   Schmidt, F. *et al.* Measuring SARS-CoV-2 neutralizing antibody activity using pseudotyped and
chimeric viruses. 2020.2006.2008.140871, doi:10.1101/2020.06.08.140871 %J bioRxiv (2020).
13   Brouwer, P. J. M. *et al.* Potent neutralizing antibodies from COVID-19 patients define multiple
targets of vulnerability. *Science* **369**, 643-650, doi:10.1126/science.abc5902 (2020).
14   Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent individuals.
Nature **584**, 437-442, doi:10.1038/s41586-020-2456-9 (2020).
15   Barnes, C. O. *et al.* Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal Common
Epitopes and Recurrent Features of Antibodies. *Cell* **182**, 828-842 e816,
doi:10.1016/j.cell.2020.06.025 (2020).
16   Shrock, E. *et al.* Viral epitope profiling of COVID-19 patients reveals cross-reactivity and
correlates of severity. *Science*, doi:10.1126/science.abd4250 (2020).
17   Sobel Leonard, A. *et al.* The effective rate of influenza reassortment is limited during human
infection. *PLoS pathogens* **13**, e1006203, doi:10.1371/journal.ppat.1006203 (2017).
18   Richard, M., Herfst, S., Tao, H., Jacobs, N. T. & Lowen, A. C. Influenza A Virus Reassortment Is
Limited by Anatomical Compartmentalization following Coinfection via Distinct Routes. *Journal*
of virology **92**, doi:10.1128/JVI.02063-17 (2018).
19   CDC. *Discontinuation of Transmission-Based Precautions and Disposition of Patients with COVID-*
19 in Healthcare Settings (Interim Guidance), <https://www.cdc.gov/coronavirus/2019-
ncov/hcp/disposition-hospitalized-patients.html> (2020).

20   Boshier, F. A. T. *et al.* Remdesivir induced viral RNA and subgenomic RNA suppression, and
evolution of viral variants in SARS-CoV-2 infected patients. *medRxiv*, 2020.2011.2018.20230599,
doi:10.1101/2020.11.18.20230599 (2020).

21   Simonovich, V. A. *et al.* A Randomized Trial of Convalescent Plasma in Covid-19 Severe
Pneumonia. *N Engl J Med*, doi:10.1056/NEJMoa2031304 (2020).

22   Meredith, L. W. *et al.* Rapid implementation of SARS-CoV-2 sequencing to investigate cases of
health-care associated COVID-19: a prospective genomic surveillance study. *The Lancet*
Infectious Diseases **20**, 1263-1272, doi:10.1016/S1473-3099(20)30562-4 (2020).

23   Collier, D. A. *et al.* Point of Care Nucleic Acid Testing for SARS-CoV-2 in Hospitalized Patients: A
Clinical Validation Trial and Implementation Study. *Cell Rep Med*, 100062,
doi:10.1016/j.xcrm.2020.100062 (2020).

24   Loman, N., Rowe, W. & Rambaut, A.   (v1, 2020).

25   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal **17**, 10-12 (2011).

26   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics (Oxford, England)*
34, 3094-3100, doi:10.1093/bioinformatics/bty191 (2018).

27   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing. *arXiv*
preprint arXiv:1207.3907 (2012).

28   Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from vision to
reality. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European*
communicable disease bulletin **22**, 30494, doi:10.2807/1560-7917.ES.2017.22.13.30494 (2017).

29   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).

30   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Molecular Biology and Evolution* **37**, 1530-1534, doi:10.1093/molbev/msaa015
(2020).

31   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587-589,
doi:10.1038/nmeth.4285 (2017).

32   Minh, B. Q., Nguyen, M. A. T. & von Haeseler, A. Ultrafast Approximation for Phylogenetic
Bootstrap. *Molecular Biology and Evolution* **30**, 1188-1195, doi:10.1093/molbev/mst024 (2013).

33 Sagulenko, P., Puller, V. & Neher, R. A. TreeTime: Maximum-likelihood phylodynamic analysis. *Virus evolution* **4**, vex042-vex042, doi:10.1093/ve/vex042 (2018).

34 Bouckaert, R. *et al.* BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Comput Biol* **10**, e1003537 (2014).

35 Illingworth, C. J. SAMFIRE: multi-locus variant calling for time-resolved sequence data. *Bioinformatics* **32**, 2208-2209, doi:10.1093/bioinformatics/btw205 (2016).

36 Martin, D. P., Murrell, B., Golden, M., Khoosal, A. & Muhire, B. RDP4: Detection and analysis of recombination patterns in virus genomes. *Virus evolution* **1** (2015).

37 Didelot, X. & Wilson, D. J. ClonalFrameML: efficient inference of recombination in whole bacterial genomes. *PLoS Comput Biol* **11**, e1004041 (2015).

38 Wrobel, A. G. *et al.* SARS-CoV-2 and bat RaTG13 spike glycoprotein structures inform on virus evolution and furin-cleavage effects. *Nature Structural & Molecular Biology* **27**, 763-767, doi:10.1038/s41594-020-0468-7 (2020).

39 Gregson, J. *et al.* HIV-1 viral load is elevated in individuals with reverse transcriptase mutation M184V/I during virological failure of first line antiretroviral therapy and is associated with compensatory mutation L74I. *The Journal of infectious diseases*, doi:10.1093/infdis/jiz631 (2019).

40 Naldini, L., Blomer, U., Gage, F. H., Trono, D. & Verma, I. M. Efficient transfer, integration, and sustained long-term expression of the transgene in adult rat brains injected with a lentiviral vector. *Proceedings of the National Academy of Sciences of the United States of America* **93**, 11382-11388 (1996).

41 Gupta, R. K. *et al.* Full-length HIV-1 Gag determines protease inhibitor susceptibility within in vitro assays. *Aids* **24**, 1651-1655, doi:10.1097/QAD.0b013e3283398216 (2010).

42 Mlcochova, P. *et al.* Combined point of care nucleic acid and antibody testing for SARS-CoV-2 following emergence of D614G Spike Variant. *Cell Rep Med*, 100099, doi:10.1016/j.xcrm.2020.100099 (2020).

43 Seow, J. *et al.* Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nat Microbiol* **5**, 1598-1607, doi:10.1038/s41564-020-00813-8 (2020).

Figure 1

Figure 1. Analysis of 23 Patient derived whole SARS-CoV-2 genome sequences in context of national sequences and other cases of chronic SARS-CoV-2 shedding. A. Circularised maximum-likelihood phylogenetic tree rooted on the Wuhan-Hu-1 reference sequence, showing a subset of 250 local SARS-CoV-2 genomes from GISAID. This diagram highlights significant diversity of the case patient (green) compared to three other local patients with prolonged shedding (blue, red and purple sequences). All SARS-CoV-2 genomes were downloaded from the GISAID database and a random subset of 250 local sequences selected. **B.** Close-view maximum-likelihood phylogenetic tree indicating the diversity of the case patient and three other long-term shedders from the local area (red, blue and purple), compared to recently published sequences from Choi et al (orange) and Avanzato et al (gold). Control patients generally showed limited diversity temporally, though the Choi et al sequences were found to be even more divergent than the case patient. Environmental samples are indicated. 1000 ultrafast bootstraps were performed and support at nodes is indicated.

Figure 2. Virus genetics and population structure in chronic SARS-CoV-2 infection. A. Highlighter plot indicating nucleotide changes at consensus level in sequential respiratory samples compared to the consensus sequence at first diagnosis of COVID-19. Each row indicates the timepoint the sample was collected (number of days from first positive SARS-CoV-2 RT-PCR). Black dashed lines indicate the RNA-dependent RNA polymerase (RdRp) and Spike regions of the genome. Of particular interest, at consensus level, there were no nucleotide substitutions on days 1-57, despite the patient receiving two courses of remdesivir. The first major changes in the spike genome occurred on day 82, following convalescent plasma given on days 63 and 65. The double amino acid deletion in S1, ΔH69/V70 is indicated by black lines. All samples are nose and throat swabs unless indicated with ETA (endotracheal aspirate). **B.** Whole genome amino acid variant trajectories based on Illumina short read ultra deep sequencing at 1000x coverage. All variants which reached a frequency of at least 10% in at least two samples were plotted. Black dashed line represents Δ69/70. CP, convalescent plasma; RDV, Remdesivir.

Figure 3

A

Figure 3. Longitudinal variant frequencies and phylogenetic relationships for virus populations bearing six Spike (S) mutations A. At baseline, all four S variants (Illumina sequencing) were absent (<1% and <20 reads). Approximately two weeks after receiving two units of convalescent plasma (CP), viral populations carrying ΔH69/V70 and D796H mutants rose to frequencies >90% but decreased significantly four days later. This population was replaced by a population bearing Y200H and T240I, detected in two samples over a period of six days. These viral populations were then replaced by virus carrying W64G and P330S mutations in Spike, which both reached near fixation at day 93. Following a 3[rd] course of remdesivir and an additional unit of convalescent plasma, the ΔH69/V70 and D796H virus population re-emerged to become the dominant viral strain reaching variant frequencies of >90%. Pairs of mutations arose and disappeared simultaneously indicating linkage on the same viral haplotype. **B.** Maximum likelihood phylogenetic tree of the case patient with day of sampling indicated. Spike mutations defining each of the clades are shown ancestrally on the branches on which they arose. Number at node denotes support by ultrafast bootstrapping consisting of 1000 replicates.

Figure 4

Figure 4: Spike mutant D796H + ΔH69/V70 has infectivity comparable to wild type but is less sensitive to multiple units of convalescent plasma (CP) and sera from recovered individuals. A. Reverse transcriptase activity of virus supernatants containing lentivirus pseudotyped with SARS-CoV-2 Spike protein (WT versus mutant) **B.** Single round Infectivity of luciferase expressing lentivirus pseudotyped with SARS-CoV-2 Spike protein (WT versus mutant) on 293T cells co-transfected with ACE2 and TMPRSS2 plasmids. **C.** Patient serum (taken at indicated Day (D)) and convalescent plasma (CP units 1-3) neutralization potency against Spike mutant D796H + ΔH69/V70 measured using lentivirus pseudotyped with SARS-CoV-2 Spike protein (WT versus mutant). Indicated is serum dilution required to inhibit 50% of virus infection. **D.** Neutralization potency of sera from 5 unselected convalescent patients (with previous confirmed SARS-CoV-2 infection) against WT versus mutant virus as in panel C. **E** Neutralization potency of a panel of monoclonal antibodies against lentivirus pseudotyped with SARS-CoV-2 Spike protein (WT Wuhan versus single mutants D614G, D796H, ΔH69/V70). Data are representative of at least two independent experiments * $p < 0.05$

Figure 5

Figure 5. Location of Spike mutations ΔH69/Y70 in S1 and D796H in S2. Amino acid residues H69 and Y70 deleted in the N-terminal domain (red) and D796H in subunit 2 (orange) are highlighted on a SARS-CoV-2 spike trimer (PDB: 6ZGE Wrobek et al., 2020). Each of the three protomers making up the Spike homotrimer are coloured separately in shades of grey (centre). Close-ups of ΔH69/Y70 (above) and D796H (below) are shown in cartoon, stick representation. Both mutations are in exposed loops.