1   Double Mammary In Situ: Predicting Feasibility of Right Mammary Artery In Situ for

2   Circumflex Coronary Artery System

3   Rafik Margaryan, MD, PhD[1,2], Daniele Della Latta[2], Giacomo Bianchi, MD, PhD[1], Nicola

4   Martini[2], Andrea Gori[2], Andrea Ripoli[2], & Marco Solinas[1]

5   [1] Adult Cardiac Surgery, Ospedale Del Cuore, Fondazione G Monasterio

6   [2] Deep Learning Unit, Ospedale Del Cuore, Fondazione G Monasterio

7   Author Note

8   Correspondence concerning this article should be addressed to Rafik Margaryan, MD,

9   PhD, Via Aurelia Sud 303. E-mail: margaryan@ftgm.it

Abstract

Arterial revascularization of the myocardium is well established and it is the gold

standard.Double mammary artery in situ revascularisation seems less attractive to surgeons

because of limited mammary length, scare, or no means of its length estimation. Here we

hypothesized that the right mammary artery length is relative to chest anatomy, mainly to

its type. We sought to estimate the feasibility, hence the mammary artery length using

machine learning (neural networks). The predictive model was able to predict a positive

outcome with 95% percent accuracy. (p < 0.01). Model's sensitivity and specificity were 96%

and 95% respectively.Neural networks can be used to predict double mammary feasibility

using chest x-rays. Model is capable of predicting positive outcomes with 95% accuracy. Last

update date "2020-12-04 15:43:45"

Keywords: bypass grafting, arterial revascularisation, double mammary, deep learning

Word count: 1099

²³ Double Mammary In Situ: Predicting Feasibility of Right Mammary Artery In Situ for

²⁴ Circumflex Coronary Artery System

Introduction

²⁶ The use of double (bilateral) internal mammary arteries (BIMAs) is scarcely diffused in

²⁷ the real world (ElBardissi et al., 2012; LaPar et al., 2015), despite being supported by robust

²⁸ evidence over the last two decades. Lack of standardization for the appropriate use of the

²⁹ right internal mammary artery (RIMA) has limited its routine use in myocardial

³⁰ revascularization. RIMA can be anastomosed in situ to the left anterior descending artery

³¹ (LAD) or to the proximal branches of the circumflex artery, but its length may pose

³² technical challenges. Although skeletonization is a technique that provides better internal

³³ mammary artery (IMA) length, it is not widely accepted. As a result, it is not always

³⁴ possible to choose the best anastomotic site, and we are forced to graft the RIMA according

³⁵ to its length rather than the most suitable coronary artery segment. Alternatively, the

³⁶ RIMA can be used as a free graft, which is anastomosed to the left internal mammary artery

³⁷ (LIMA) as a composite conduit. This option is more technically demanding and can expose

³⁸ to higher rates of competitive flow with increased graft failure (Pevni et al., 2007) though no

³⁹ definitive data are available to confirm this finding (Glineur et al., 2008). Mid-term clinical

⁴⁰ outcome seems to be similar regardless of BIMA configuration, either in situ or as a

⁴¹ composite conduit (Glineur et al., 2008). In situ double mammary grafting is safe and

⁴² durable bypass strategy when circumference artery system has suitable targets, although

⁴³ personal judgment of right mammary length, hence, RIMA in situ feasibility is subjective.

⁴⁴ We sought do develop prediction model using chest X ray (antero-posterior projection) and

⁴⁵ deep neural networks in orders to estimate feasability of right mammary artery to circumflex

⁴⁶ artery targets (early obtuse marginal branches and intermediate/early diagonal arteries)

Material and Methods

Patients

We have selected patients from our electronic health record system, undergoing double mammary artery grafting from 2007 until agust 2019 (n = 292) were selected fror this study. All patients underwent successful on-pump bypass surgery. For this analysis we've ignored other types of bypasses. We devided the population into two groups:

- In situ bypass grafting
- Grafting using y- or t-grafts 127 were double mammary grafts in-situ and 165 were double mammary grafts Y grafts. Mean age of the patients was 61.4 ± 7.7. All other demographic data is presentein in Table 1. All patients had preoperative chest x-ray (antero-posterior and lateral).

Data Acquisition and Processing

292 chest digital X-ray examinations (DX) were performed on patients enrolled for valvular surgery. X-ray images of lateral latero (LL) projections were used to train a deep convolutional neural network (DNN). All DX were taken with standing patient placed 180 cm from the X-ray tube using a digital flat detector panel with a spatial resolution of 0.139×0.139 mm^2. The size of the image matrix is variable (2500x2500-4000x4000 px) depending on the size of the patient's chest as well as the X-ray exposure. The potential difference (kV) and exposure current (mAs) parameters are set automatically of the X-ray scanner in order to obtain an adequate contrast independently of the variation of the attenuation of the body of each patient. The dataset was splitted into 183 images to train the DNN and 22 to test the learning. The size of all the images was readjusted to 2000x2000 with a centered crop operation so as to maintain the DNN input layer with fixed size and

partially mask the background that does not provide an useful information for the

classification task. Finally, to minimize the computational load for network training, the

images were scaled to 512x512 px. Whole data was divided randomly into two parts: train

dataset and test dataset. Train dataset was uesed to create the model and validate it. Test

dataset was used for only external validation and never has been used for tuning or

improving the algorithm.

Network Architecture

A GoogLeNet network (Szegedy et al., 2014) was used to perform the classification task.

The outline of the proposed architecture was reported in Table 1. The network comprises 22

trainable layers and is made by nine Inception module stacked upon each other. A single

Inception module represents a combination of 1x1, 3x3 and 5x5 convolutional kernels, used

in parallel with the aim to progressively cover a bigger area and extract details (1x1) as more

general features (5x5). The results are then concatenated with the outcome of a max-pooling

layer and passed to the next layer. The presence of 1x1 convolutions before the 3x3, 5x5

convolutions and after the pooling layer help to reduce the number of parameters in the

Inception module and add more non-linearities to the network. To apply the GoogLeNet

network architecture to the surgical task some implementation changes had to be made: 1)

the input layer dimensions, set at 512 x 512, with a number of channels equals to one instead

of three, being the inputs grayscale images; 2) in the output layer only two units were used,

as we had to classify the images according the two possible locations of surgical incision; The

network was designed and trained with Tensorflow (Workshop on Wireless Traffic

Measurements and Modeling, USENIX Association, ACM SIGMOBILE, ACM Special

Interest Group in Operating Systems, & ACM Digital Library, 2016), running for 500 epochs

on a single NVIDIA Titan Xp. Stochastic gradient descent and Adam optimizer were used to

minimize a categorical cross-entropy for the network parameters update.

Schematic representation of the GoogLeNet network used to classify the input images.

For a detailed description of the network architecture see (Szegedy et al., 2014).

The size of all the images was readjusted to 2000x2000 with a centered crop operation

so as to maintain the DNN input layer with fixed size and partially mask the background

that does not provide an useful information for the classification task. Finally, to minimize

the computational load for network training, the images were scaled to 512x512 px. To

evaluate the progress of learning the accuracy was calculated between the predicted and the

real "point of incision".

<div align="center">

Results

</div>

104   There were no hospital or 30 days mortality. Model has predicted with 0.96 accuracy p

105 < 0.01). There were no significant differences in comorbidited divieded by type of graft and

106 sex group. Model's sensitivity and specificity were 0.96 and 0.95.Area under curve (AUC)

107 was 0.9561.

Discussion

In this study we were able to build a predictive model using chest x-rays with very high performes. To our knowlage it is first of its kind. Deep Learning models in general have very high good perfomance, and in this clinical scenario it has very high accuracy and AUC values for predicting possible double mammary in-situ bypass grafting. It proves to have hit sensitivity and specificity. It is particularly valuable for total arterial re-vascularisation.

Conclusion

Neural networks can be used to predict double mammay fiesibility using chest x-rays. Model is capable of predicting positive outcome with 95% accuracy.

Tables

References

ElBardissi, A. W., Aranki, S. F., Sheng, S., O'Brien, S. M., Greenberg, C. C., & Gammie, J.
    S. (2012). Trends in isolated coronary artery bypass grafting: An analysis of the
    Society of Thoracic Surgeons adult cardiac surgery database. *The Journal of*
    Thoracic and Cardiovascular Surgery, 143(2), 273–281.
    https://doi.org/10.1016/j.jtcvs.2011.10.029

Glineur, D., Hanet, C., Poncelet, A., D'hoore, W., Funken, J.-C., Rubay, J., . . . El Khoury,
    G. (2008). Comparison of bilateral internal thoracic artery revascularization using in
    situ or Y graft configurations: A prospective randomized clinical, functional, and
    angiographic midterm evaluation. *Circulation, 118*(14 Suppl), S216–221.
    https://doi.org/10.1161/CIRCULATIONAHA.107.751933

LaPar, D. J., Crosby, I. K., Rich, J. B., Quader, M. A., Speir, A. M., Kern, J. A., . . .
    Investigators for the Virginia Cardiac Surgery Quality Initiative. (2015). Bilateral
    Internal Mammary Artery Use for Coronary Artery Bypass Grafting Remains
    Underutilized: A Propensity-Matched Multi-Institution Analysis. *The Annals of*
    Thoracic Surgery, 100(1), 8–14; discussion 14–15.
    https://doi.org/10.1016/j.athoracsur.2015.02.088

Pevni, D., Hertz, I., Medalion, B., Kramer, A., Paz, Y., Uretzky, G., & Mohr, R. (2007).
    Angiographic evidence for reduced graft patency due to competitive flow in composite
    arterial T-grafts. *The Journal of Thoracic and Cardiovascular Surgery, 133*(5),
    1220–1225. https://doi.org/10.1016/j.jtcvs.2006.07.060

Szegedy, C., Liu, W., Jia, Y., Sermanet, P., Reed, S., Anguelov, D., . . . Rabinovich, A.
    (2014). Going Deeper with Convolutions. *arXiv:1409.4842 [Cs]*. Retrieved from
    http://arxiv.org/abs/1409.4842

143 Workshop on Wireless Traffic Measurements and Modeling, USENIX Association, ACM

144   SIGMOBILE, ACM Special Interest Group in Operating Systems, & ACM Digital

145   Library. (2016). *Papers presented at the Workshop on Wireless Traffic Measurements*

146   *and Modeling: June 5, 2016,Seattle, WA, USA*. Berkeley, CA: USENIX Association.

147   Retrieved from http://portal.acm.org/toc.cfm?id=1072430

Table 1

Summary of demographic data by sex.

	Female			Male		
	In Situ	Y-graft	p.overall	In Situ	Y-graft	p.overall
	N=8	N=12		N=119	N=153	
Age	67.3 (5.77)	59.6 (11.4)	0.062	60.8 (7.79)	61.7 (7.35)	0.363
BMI	25.2 (2.67)	27.9 (5.40)	0.181	26.7 (2.96)	26.6 (3.56)	0.785
Hypertension:			0.619			0.743
No	1 (12.5%)	3 (25.0%)		45 (37.8%)	62 (40.5%)	
Yes	7 (87.5%)	9 (75.0%)		74 (62.2%)	91 (59.5%)	
Hypercholesterolemia:			0.167			0.757
No	5 (62.5%)	3 (25.0%)		49 (41.2%)	67 (43.8%)	
Yes	3 (37.5%)	9 (75.0%)		70 (58.8%)	86 (56.2%)	
Diabetes:			0.603			0.493
No	7 (87.5%)	8 (66.7%)		96 (80.7%)	117 (76.5%)	
Yes	1 (12.5%)	4 (33.3%)		23 (19.3%)	36 (23.5%)	
Diabetes on Insulin:			.			0.390
No	8 (100%)	12 (100%)		118 (99.2%)	149 (97.4%)	
Yes	0 (0.00%)	0 (0.00%)		1 (0.84%)	4 (2.61%)	
Preoperative Creatinine	0.81 (0.19)	1.32 (1.83)	0.355	0.94 (0.21)	0.92 (0.20)	0.425
ON Pump:			1.000			0.454
No	1 (12.5%)	1 (8.33%)		14 (11.8%)	24 (15.7%)	
Yes	7 (87.5%)	11 (91.7%)		105 (88.2%)	129 (84.3%)	
CPB Time	102 (22.8)	89.0 (30.7)	0.330	101 (31.1)	102 (37.3)	0.823
XC Time	69.4 (19.0)	63.3 (23.5)	0.551	70.5 (23.8)	71.7 (27.1)	0.705

Table 2

Summary of demographic data by sex.

	Female			Male		
	In Situ	Y-graft	p.overall	In Situ	Y-graft	p.overall
	N=8	N=12		N=119	N=153	
predicted:			<0.001			<0.001
In Situ	8 (100%)	0 (0.00%)		114 (95.8%)	8 (5.23%)	
Y-graft	0 (0.00%)	12 (100%)		5 (4.20%)	145 (94.8%)	