Adam M. Ćmiel[1], Bogdan Ćmiel[2]

[1] Institute of Nature Conservation, Polish Academy of Sciences, al. A. Mickiewicza 33, 31-120

Kraków, Poland

[2] Faculty of Applied Mathematics, AGH University of Science and Technology, al. A. Mickiewicza

30, 30-059 Kraków, Poland

corresponding author: cmiel@iop.krakow.pl

9 A new, simple method of describing the COVID-19 trajectory and dynamics in any country

10 based on Johnson Cumulative Distribution Function fitting.

12 Abstract

A simple method is described to study and compare COVID-19 infection dynamics between countries,

based on curve fitting to publicly shared data of confirmed COVID-19 infections in them. The method

was tested using data from 80 countries from 6 continents. We found that Johnson Cumulative

Distribution Functions (CDF) were extremely well fitted to the data ($R^2>0.99$) and that Johnson CDFs

were much better fitted to the data at their tails than either the commonly used Normal or Lognormal

CDFs. Fitted Johnson CDFs can be used to obtain basic parameters of the infection wave, such as the

percentage of the population infected during an infection wave, the days of the start, peak and end of

the infection wave, as well as the durations of the infection wave of the wave's increase and decrease.

These parameters can be easily interpreted biologically and used both for describing the infection

wave dynamics and in further statistical analysis. The usefulness of the parameters obtained was

analysed with respect to the relation between the Gross Domestic Product (GDP) per capita and the

population density, and the percentage of the population infected during an infection wave, the starting

day and the duration of the infection wave in the 80 countries. We found that all the above parameters

were significantly dependent on the GDP per capita, but only the percentage of the population infected

was significantly dependent on the population density in these countries. If used with caution, this

method has a limited ability to predict the future trajectory and parameters of an ongoing infection

wave.

31 Introduction

A highly contagious disease, COVID-19 is caused by the SARS-CoV-2 coronavirus. This virus was

first detected in Wuhan (central China) in December 2019, but as early as mid-January 2020, the virus

had quickly spread throughout China. On 13 January 2020, the first case outside China was confirmed

and on 24 January, the first case in Europe was reported. In the second half of February 2020,

outbreaks with hundreds of cases erupted in South Korea, Italy and Iran (Skórka et al., 2020), and

COVID-19 was declared a pandemic by the World Health Organization on 11 March 2020

(Ducharme, 2020). To date, over 64 million infections and almost 1.5 million deaths have been

reported globally (WHO, 2020).

Since the very beginning of the pandemic, many models have been proposed to understand the

outbreak dynamics of COVID-19 (e.g. IHME, 2020; UGSDSC, 2020; LANL, 2020; Ferguson et al.,

2020; Kissler et al., 2020; Aleta et al.; Hellewell et al., 2020) and have been used by policymakers

(e.g. the US Government) for allocating resources or planning interventions. Some of them, such as

the early IHME model, have received a fair amount of criticism (Jewell et al., 2020). COVID-19

modelling studies generally follow one of two general approaches – forecasting models and

mechanistic models – although there are also hybrid approaches (Holmdahl and Buckee, 2020).

Forecasting models are often statistical in nature, fitting a line or a curve to data and extrapolating

from there, without incorporating the process that produces the pattern (Holmdahl and Buckee, 2020),

while mechanistic models simulate the outbreak through interacting disease mechanisms by using
local nonlinear population dynamics and the global mixing of populations (Hethcote, 2000). Purely
statistical models are reliable only within a short time window and may be useful for making rapid
short-term recommendations, whereas mechanistic modelling can be useful for exploring how the
course of the pandemic might change under various assumptions and political interventions (Kuhl,
2020).
Since its onset, the COVID-19 pandemic has generated a huge amount of data and is probably
the best documented disease in history. New cases, active cases, deaths and numbers of tests
performed are usually published daily by official sources (e.g. governments), gathered and publicly
shared as freely accessible datasets (e.g. Hasell et al., 2020). This offers researchers an opportunity to
focus on analysing the pandemic and its dynamics in fields other than epidemiology. Although the
above-mentioned models provide many pandemic parameters for predicting different scenarios of
future infections, the probability and duration of future pandemic peaks, which is extremely useful for
policymakers in planning interventions, they may not be very useful in fields other than epidemiology.
An urgent need thus arose to develop methods of describing the trajectories of pandemic waves. Such
methods should be simple to apply and should provide parameters describing the trajectory and
dynamics of the epidemic that are easy to interpret and employ in further statistical analyses by
researchers in other fields, like sociology, biology and ecology, which can deepen our understanding
of the COVID-19 pandemic.
The aim of this paper is to present a new and simple method based on curve fitting to the
reported data on confirmed cases of infection, and to study and compare the infection dynamics
between countries (or regions). This method, based on Johnson Cumulative Distribution Function
(CDF) fitting, was tested using data from 80 countries from 6 continents (Africa, Asia, Europe,
Oceania, as well as North and South America). Also, Johnson CDFs were used to calculate basic
parameters of the infection wave dynamics, such as the percentage of the population infected during
the infection wave, the days of the start, peak and end of the infection wave, the duration of the
infection wave, and the durations of the wave's increase and decrease. These parameters are easy to
interpret and can be used in further statistical analyses of epidemic dynamics. This is exemplified by
the influences of Global Domestic Product (GDP) per capita and of population density on the
percentage of infections and the first day and duration of the first infection wave in the countries
concerned. The method presented here and the techniques employed are both straightforward and
well known; they illustrate how simple techniques can be used to solve otherwise difficult problems,
such as describing an epidemic wave.
82
Materials and methods
The data used in this study were obtained from the Our World in Data COVID-19 dataset (Hasell et
al., 2020) from December 2019 to 19 November 2020. The method was tested on 80 countries from 6
regions: 1) Africa (Democratic Republic of Congo, Egypt, Ethiopia, Kenya, Morocco, Nigeria,
Somalia, South Africa, South Sudan, Sudan and Zimbabwe); 2) Asia (Afghanistan, Bangladesh,
Cambodia, China, India, Indonesia, Iran, Iraq, Israel, Japan, Lebanon, Myanmar, Pakistan, Philippines,
Saudi Arabia, Singapore, South Korea, Sri Lanka, Syria, Taiwan, Thailand, Turkey, Vietnam); 3)
Europe (Austria, Belgium, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus, Czechia, Finland,
France, Germany, Greece, Hungary, Ireland, Italy, Netherlands, North Macedonia, Norway, Poland,
Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine, United
Kingdom); 4) North America (Canada, Jamaica, Mexico, United States of America); 5) Oceania
(Australia, Fiji, New Zealand, Papua New Guinea); and 6) South America (Argentina, Bolivia, Brazil,
Chile, Colombia, Paraguay, Peru, Uruguay, Venezuela).

In order to make the data comparable between countries, the number of infections on each day
of the pandemic for each country were standardized – they are presented here as a percentage of the
population of a given country infected, i.e. the number of confirmed infections in a given
country/country population*100%. Also, a five-day moving average was calculated using the
percentage of infections to smooth the data and to minimize the effect of a smaller number of tests
performed and a smaller number of confirmed infections during some short periods like weekends.
This makes the loss function more regular, i.e. it has less relative extrema, and so it is easier to find the
global extremum. Nevertheless, all the presented R^2 for the Johnson CDFs obtained were calculated
using raw (not smoothed) data.
105
Fitting Johnson CDF by moments
Johnson (1949) described a system of frequency curves that represents transformations of the standard
normal curve (detailed description in Hahn and Shapiro, 1967). Applying these transformations to a
standard normal variable allows a unique distribution to be derived for whatever combination of mean,
standard deviation, skewness and kurtosis occurs for a given set of observed data. The standard
method of fitting Johnson curves is to use four coefficients defining a Johnson distribution: two shape
coefficients (γ, δ), as well as a location (ξ) and a scale (λ) coefficient:

$$F(x) = \Phi\left(\gamma + \delta \, sinh^{-1}\left(\frac{x-\xi}{\lambda}\right)\right) \quad (1),$$

where Φ is the cumulative distribution function of the standard normal distribution. This method is not
intuitive, however, as it is difficult to set up starting points from the data in order to perform numerical
fitting. Thus, we selected an alternative method for fitting Johnson curves, using the first four
moments of an empirical distribution (mean, variance, skewness and kurtosis; detailed descriptions in
Hahn and Shapiro (1967) and Hill et al. (1976)). All the statistical fits in the paper were performed
using the Levenberg-Marquardt algorithm (Moré, 1978) to solve the corresponding non-linear least
square optimization problem. The convergence criterion was set at $1.0E^{-10}$.
121
Fitting Johnson CDFs to epidemic waves
There is no strict definition of what is or is not an epidemic wave or phase. The intuitive definition of
a pandemic wave traces the development of an epidemic over time and/or space. During an epidemic,
the number of new cases of infection increases (often rapidly) to a peak and then falls (usually more
gradually) until the epidemic wave is over.
The epidemic's dynamics may differ greatly between countries. Since the beginning of the
pandemic, there has been only one epidemic wave in some countries (e.g. Afghanistan, Argentina),
while in others two have occurred (e.g. Australia); in yet others, even more have taken place, which
may have overlapped and interfered with each other, as in Croatia, where there were four overlapping
and interfering waves. The authorities in many countries, moreover, have imposed lockdowns of
varying severity in order to slow down or "flatten" the infection curve. Hence, the epidemic waves
may not follow Farr's law (which states that epidemics tend to rise and fall in a roughly symmetrical
pattern or bell-shaped curve) and may be asymmetrical.
The basic assumption is that each epidemic wave W in a given country can be described by a
five-parameter scaled Johnson CDF: scale parameter (s), and the above-mentioned moments–
expected value (mean, E), variance (V), skewness (S) and kurtosis (K)

$$W(t)=s*F_{E,V,S,K}(t) \quad (2),$$

139 where t is the time measured since the day of the beginning of the pandemic, and the function $F_{E,V,S,K}$ is
140 the Johnson CDF with parameters γ, δ, ξ, λ assuming the mean, variance, skewness and kurtosis to be
141 equal to E,V,S,K, respectively (see Hahn and Shapiro,1967; Hill et al.,1976). Parameters S and K were

142 expected to improve the curve fit at the tails of the epidemic wave if it was not symmetrical or heavy
143 tailed.
144
145 **Obtaining basic epidemic wave parameters and their biological interpretation**
146 Once the Johnson CDFs were fitted to each pandemic wave in a given country, the basic parameters
147 for obtaining the wave dynamics, i.e. 2.5% quantile ($Q_{2.5\%}$), 50% quantile (median; $Q_{50\%}$), 97.5%
148 quantile ($Q_{97.5\%}$), could be calculated:

$$Q_{2.5\%} = F_{E,V,S,K}^{-1}(2.5\%) \qquad (3)$$

$$Q_{50\%} = F_{E,V,S,K}^{-1}(50\%) \qquad (4)$$

$$Q_{97.5\%} = F_{E,V,S,K}^{-1}(97.5\%) \qquad (5)$$

152 The disadvantage of fitting the Johnson curve by its moments is that it is not possible to
153 determine its mode analytically. Thus, the mode of each Johnson CDF was determined numerically:

$$M = \arg\max f_{E,V,S,K}(x) \qquad (6),$$

where $f_{E,V,S,K}$ is the Johnson Probability Density Function (PDF).
These parameters have an intuitive biological interpretation (Fig. 1). The scale parameter *s*
indicates the total percentage of infections during a given epidemic wave (P_{inf}), $Q_{2.5\%}$ indicates the day
when the infection wave started, while $Q_{97.5\%}$ indicates its end. The median ($Q_{50\%}$) indicates the day
when half the total percentage of infections during a given wave was reached. Finally, the mode (*M*)
indicates the day when the peak occurred. In addition, one can easily obtain the wave duration (*T*)

$$T = Q_{97.5\%} - Q_{2.5\%} \qquad (7),$$

162 the duration of wave increase (t_i)

$$t_i = M - Q_{2.5\%} \qquad (8),$$

164 and the duration of the wave decrease (t_d)

$$t_d = Q_{97.5\%} - M \qquad (9).$$

166 Also, the parameter measuring the asymmetry of the infection wave (*A*) is easily obtained as the ratio

$$A = t_i/t_d \qquad (10).$$

All of the above-mentioned parameters can be easily used in further statistical analyses, as
exemplified by 1) the relationship between Gross Domestic Product (GDP) per capita and basic
parameters describing the dynamics of the first wave of infections: *M*, *T* and P_{inf}, and 2) the relation
between population density and basic parameters describing the dynamics of the first wave of
infections: *M*, *T* and P_{inf}. Only the first wave of infections in each country has been taken into account
here, because in some countries, second (and subsequent) waves did not take place and would have to
be excluded from the analysis.
175
176 **Comparing curves: Johnson vs Normal and Lognormal CDFs**
177 The differences between the Johnson, Normal and Lognormal CDFs were highlighted on the basis of
178 data from Afghanistan, where only one epidemic wave took place, by comparing parameters R^2, P_{inf},
179 $Q_{2.5\%}$, *M* and $Q_{97.5\%}$. Both the 2.5% and the 97.5% quantiles for normal and lognormal distributions
180 were obtained using inverse Normal and inverse Lognormal PDFs, respectively.
181
Fitting Johnson's CDF to the ongoing wave and forecast possibilities
Fitting Johnson's curve to the ongoing wave yields parameters that can also be interpreted as a
forecast of the future shape and dynamics of the infection wave. In such a case, P_{inf}, *M* and $Q_{97.5\%}$
indicate the predicted percentage of infections, the predicted day of the peak and the predicted day of
the end of the ongoing wave, respectively, which can also be used to calculate the predicted times of
the increase, decrease and duration of the ongoing infection wave. Because this method is intended to
describe the infection's dynamics rather than to predict its ultimate outcome, the accuracy of the

forecast is evaluated only on the basis of data from the first wave of infection recorded in the United
Kingdom (see Supplementary Materials).
191
192 **Examples of application**
193
194 **The relationships between Gross Domestic Product (GDP) per capita, and population density**
195 **and the dynamics of the first wave of COVID-19 infections**
196
The data on the GDP per capita and population density in the 80 countries analysed here were
obtained from the Our World in Data COVID-19 dataset (Hasell et al., 2020).
The relationship between GDP per capita, and population density and the basic parameters
describing the dynamics of the first wave of infections (M, T and P_{inf}), obtained using the presented
method of Johnson CDF fitting, was tested using the quantile dependence function method, described
in detail in Ćmiel and Ledwina (2020).This method was designed for measuring, visualizing the
dependence structure, and testing the independence of two random variables. It exploits a recently
introduced local dependence measure (quantile dependence function q), which gives a detailed picture
of the underlying dependence structure and provides a means by which the local association structure
can be minutely examined at different quantile levels (Ćmiel and Ledwina 2020).
207
Results
Fig. 2 illustrates examples of Johnson curves fitted to the data from countries where one ongoing
infection wave (Argentina), one infection wave (Afghanistan), two infection waves (Australia) and
four overlapping and interfering infection waves (Croatia) occurred. Fig. 3A shows Johnson CDFs
fitted to the four waves of infections observed in Croatia, showing where the waves overlapped and
interfered.
Johnson CDF fitting, tested using data from 80 different countries, showed that all the curves
were extremely well fitted: the lowest R^2 was 0.995 (Fiji), the highest R^2 was 0.99997 (Iraq), and the
mean and median R^2 were 0.9995 and 0.9997, respectively. The functions fitted with R^2 and the
COVID-19 trajectory plots with fitted functions for each country are illustrated in the Supplementary
Materials (Table S1; Figures S1-S6).
Fitting Johnson, Normal and Lognormal distribution curves to the single wave of infection that
took place in Afghanistan showed that the Johnson CDF (R^2=0.9998) was the best fitted, whereas the
Normal (R^2=0.9980) and Lognormal (R^2=0.9989) distributions were not so well fitted, mainly at the
tails of the infection wave (Fig. 3B). The parameters $Q_{2.5\%}$, M and $Q_{97.5\%}$ obtained for the infection
wave in Afghanistan using Johnson CDF fitting were 59, 100 and 209 respectively, whereas the same
parameters obtained using Normal CDF fitting and Lognormal CDF fitting were 57, 105 and 152, and
65, 98 and 167, respectively. The percentages of the population infected during the infection wave
obtained using the scale parameters (s) of the fitted Johnson, Normal and Lognormal distributions
were 0.1028%, 0.0984% and 0.0997% respectively.
Seventeen (21.3%) of the 80 countries analysed were described by fitting one infection wave,
while 35 (43.8%), 24 (30%) and 4 (5%) were described by fitting two, three and four infection waves,
respectively (Table S1).
The basic statistics for the skewness parameters of the Johnson CDFs fitted to the first
pandemic waves in the 80 counties showed that in the majority of them, the first wave of infection was
skewed (median S=1.5; minimum S=0; maximum S=141.5). The first wave of infection was
symmetrical in 16 countries (20%; $A<1.05$). Also, the basic statistics for parameter A showed that the
duration of wave decrease was longer than that of wave increase (mean A= 4.7; median A=2.9;
minimum A=1.0; maximum A=22.4).

Analysis of the associations between GDP per capita and parameters M, T and P_{inf} showed that
the percentage of confirmed infections during the first epidemic wave in the 80 countries was
dependent on the GDP per capita (p=0.0147; Fig 4A), the time of the peak (M; p=0.0002; Fig. 4B) and
the duration of the first epidemic wave (T; p=0.0087; Fig. 4C). The relation between the percentage of
infections and GDP per capita tended towards a global positive dependence (Fig. 4A), which means
that the higher the GDP per capita, the greater the percentage of infections during the first epidemic
wave. The relation between the time of the peak and GDP per capita showed a local negative
dependence for countries where the peak occurred late (above median; Fig. 4B). This means that the
very early occurrence of a peak was not really correlated with GDP per capita; but when the peak did
not occur early, then the higher the GDP per capita, the earlier the peak occurred. A similar situation
prevailed for the relation between the duration of the infection wave and GDP per capita (Fig. 4C), i.e.
a very short duration of the first epidemic wave was not really correlated with GDP per capita, but in
the case when the first epidemic wave was not of short duration, then the higher the GDP per capita,
the shorter the first epidemic wave.
Analysis of the associations between population density and parameters M, T and P_{inf} showed
that the percentage of infections during the first epidemic wave in the 80 countries depended on
population density (p=0.0079; Fig 4D), whereas the day of the peak and the duration of the first
epidemic wave did not (T: p=0.4243; Fig. 4E; M: p=0.5924; Fig. 4F).The relation between the
percentage of infections and population density showed a local negative dependence (Fig. 4D), e.g. in
cases when the population density was not very high but the percentage of infections was quite high.
In such cases, the greater the population density, the smaller the percentage of infections.
258
Discussion
The method presented in this paper gives an indication of the spread of COVID-19 in any
particular country that publicizes daily numbers of infected cases. Both this method and the techniques
employed are all straightforward, well known and easy to use, since Johnson CDF fitting is available
in many statistical/calculus packages, e.g. R, Statistica, MATLAB, MS Excel. By using an alternative
method of fitting using moments instead of shape, location and scale parameters, it is easier to set
starting points for numerical fitting, e.g. by visually analysing the scatter plot of the number of
infected cases in time. The curves are extremely well fitted: this is exemplified by the data from 80
different countries on 6 continents. Also, the parameters are easy to interpret and ready to use in
further analyses, such as finding associations between them and other variables that may be associated
with COVID-19 dynamics, i.e. GDP per capita, population density.
Some researchers have used curve-fitting with a normal distribution to respond to a real time
request, applying it to COVID-19 in Wuhan (Tomie, 2020), since it was known that flu epidemics
follow a normal distribution, whereas other researchers noticed the COVID-19 profile had a
characteristically asymmetric tail and applied Lognormal distribution curve fitting (Nishimoto and
Inoue, 2020). Our results show that the first wave of infections was highly skewed in 79% of the
countries analysed. This suggests that unlike flu epidemics, the COVID-19 epidemic is not following a
normal distribution and should not be modelled in this manner. In such a case, log-normal distribution
fitting appears better; but as the example of Afghanistan shows, the apparent differences in R^2 between
the Johnson, Normal and Lognormal CDFs are small, but are in fact ca one order of magnitude in
favour of the Johnson CDF. Moreover, one can see that both the Normal and Lognormal CDFs are less
well fitted at the tails of the infection wave than the Johnson CDF (Fig. 3B), and both reveal a smaller
number of infections than was actually recorded (raw data) and lower than that obtained using
Johnson's CDF. Moreover, the fitted Lognormal curve starts to increase later than the Normal and
Johnson distribution curves, which would in consequence lead to an incorrect estimate of the
beginning of the wave (11 days later than when obtained using Johnson's distribution), whereas the

Normal distribution is far worse fitted at the right tail than the Johnson and Lognormal distributions,
because the infection wave in Afghanistan was not symmetrical. Apart from the fact that using the
Normal distribution would preclude any estimate of the true duration of the wave decrease (by
definition equal to the time of the wave increase), it also leads to a much lower estimate of the day
when the wave of infections ends (57 days earlier than estimated using the Johnson distribution),
which is caused by the "too fast" flattening of the Normal CDF (Fig. 3B). The extremely high R^2
values obtained for the 80 countries (see Supplementary Materials) suggest that the Johnson curve
class is flexible enough to almost perfectly follow the course of the epidemic in these countries. This is
because both skewness and kurtosis are estimated during the Johnson curve fitting procedure, whereas
the shapes of other commonly used curves (Normal, Lognormal, Weibull) are more or less imposed.
This result also suggests that the Johnson distribution should be the preferred curve-fitting approach
for COVID-19 data.
The curve fitting method presented here was designed primarily to obtain easily interpretable
parameters describing past trajectories of COVID-19 infections, but the parameters describing the
current wave of infection, especially in its early stages (before the peak), can be interpreted as a
forecast of the future course of the pandemic. In such a case, however, extreme caution is advisable
(see Jewell et al., 2020). This method is a purely statistical model that does not incorporate the process
producing the pattern of the number of infections and does not account for parameters governing
transmission, disease and immunity. In addition, curve fitting techniques cannot predict the occurrence
of future peaks. Thus, for long-term forecasts and modelling future scenarios of the pandemic, it is
recommended to use more reliable methods, such as those based on SEIR models. Nevertheless, some
short-term predictions can be made using the Johnson CDF method, which may be useful to
policymakers for planning rapid, short-term interventions. One must bear in mind, however, the
method's limitations (see above), as well as those resulting from the data collection and reporting,
which are discussed later in this section.
The results obtained from this application of parameters describing COVID-19 dynamics have
shown that the higher the GDP per capita, the higher the percentage of the population infected. This is
quite an unexpected result, but consistent with a very recent report by Liu et al. (2020), who found a
positive correlation between the human development index (HDI) and the risk of infection and death
from COVID-19 in Italy. Other results have shown that, excluding countries where the infection wave
peaked very early and its duration was short, the higher the GDP per capita, the earlier the peak and
the shorter the first epidemic wave. This result, in turn, is similar to that reported in another very
recent paper, in which the date of the first COVID-19 cases was shown to co-vary positively with
GDP across countries, most probably because of their closer involvement in global tourism and
transport (Jankowiak et al., 2020). Another example showed that the higher the population density, the
lower the percentage of a population infected during the first wave of infections. This, too, seems
unexpected; but this negative dependence results from the fact that infections are presented as a
percentage, which does not scale proportionally with population density. A further possible
explanation is that in countries with a high population density, like China and Singapore, very strict
(full) lockdowns were immediately applied (China, Kretschmer and Yang, 2020; Singapore, Cheong,
2020), which could have resulted in fewer people being infected than in countries with a lower
population density, where lockdowns were only partial, if imposed at all. Moreover, some researchers
report a positive correlation between population density and the number of infections and related
mortality, e.g. in India (Bhadra et al., 2020),whereas others provide no evidence that population
density is linked with COVID-19 cases and deaths, e.g. the USA (Carozzi et al., 2020). Nevertheless,
these examples demonstrate the usefulness of our method. The very recent papers by Liu et al. (2020)
and Jankowiak et al. (2020) have also shown that the amount of research on COVID-19, other than
purely epidemiological modelling of future pandemic scenarios, is increasing. This indicates that

simple methods of obtaining parameters describing infection waves, such as are presented in this
paper, may be very useful and can help to deepen our understanding of the COVID-19 pandemic.
The last, but not least important issue which has to be addressed is a key limitation in
understanding the COVID-19 pandemic, namely, that the true number of infections is not known and
the only infections known are those confirmed by tests. Moreover, testing strategies differ between
countries: in some countries only symptomatic cases are tested, while in others mass testing is carried
out. Moreover, most COVID-19 cases are asymptomatic and remain unreported (Peirlinck et al.,
2020). Consequently, mortality data are generally considered more reliable than the testing-dependent
confirmed case counts that are used in COVID-19 epidemic modelling (e.g. Chikobvu and Sigauke,
2020). However, some countries only report COVID-19 deaths occurring in hospitals, whereas others
report COVID-19 deaths when a test has confirmed the infection (this makes the number of mortalities
testing-dependent as well). On the other hand, if laboratory diagnosis is not required, as in the U.K.
(UK Guidance), it is possible that deaths due to other diseases with COVID-19-like symptoms are
being reported as COVID-19 deaths. It may also be difficult to specify the cause of death in cases
where patients had other diseases, e.g. an advanced stage of cancer, co-existing with COVID-19.
Taking all of the above into account, it is very likely that the real number of deaths is also higher than
the reported number of deaths, something that has been noticed in some countries, e.g. Italy (Foresti,
2020; Stancati and Sylvers, 2020) and China (Long et al., 2020). It is may well be that the numbers of
both confirmed new cases and confirmed deaths are unreliable, but on the other hand, no other data are
available. Some models, e.g. IHME 2020, are capable of estimating the true number of infections, but
this involves making a number of additional assumptions, and is based partly on the reporting of
testing-dependent data. Also, the relation between the true number of infections and the number of
deaths has not been well studied to date and requires a number of assumptions. Using the number of
infections seems to be the easiest way of obtaining basic data on the COVID-19 infection dynamics in
a given country, so long as one is aware that publicly shared data show the number of confirmed cases
and not the number of real infections and takes this into account when interpreting the results.
In conclusion, the method based on fitting Johnson CDF curves to the cumulative number of
confirmed cases is straightforward, well-known and easy to use. It yields curves that are extremely
well fitted to the data, and the basic parameters of COVID-19 infection dynamics thus obtained are
easy to interpret and use in further statistical analyses by researchers from fields other than
epidemiology, e.g. sociology, biology and ecology. While deepening our understanding of the
COVID-19 pandemic, it may also be useful for making short-term predictions, although caution is
advisable in such cases.
366

367 **Acknowledgements**
368 Both authors made equal contributions to this study and are listed alphabetically. This study was
369 financed by the statutory funds of the Institute of Nature Conservation, Polish Academy of Sciences,
370 and the Faculty of Applied Mathematics, AGH University of Science and Technology. We thank
371 Magdalena Lenda and Piotr Skórka for their useful comments and suggestions.
372

References
Bhadra, A., Mukherjee, A., Sarkar, K. 2020. Impact of population density on Covid-19 infected and
mortality rate in India. Modeling Earth Systems and Environment. https://doi.org/10.1007/s40808-
020-00984-7
Carozzi F., Provenzano S., Roth S. 2020. Urban density and COVID-19, discussion paper series, IZA
Institute of Labor economics, IZA DP No. 13440.
Cheong D. 2020. Coronavirus: Most workplaces to close, schools will move to full home-based
learning from next week, says PM Lee. The Straits Times. Singapore. 3 April 2020.Accessed at

https://www.straitstimes.com/singapore/health/most-workplaces-to-close-schools-will-move-to-
full-home-based-learning-from-next on 29 November 2020.
Chikobvu D., Sigauke C. 2020. Statistical distribution fitting to the number ofCOVID-19 deaths in
South Africa. Preprint, DOI: 10.21203/rs.3.rs-32411/v1
Ćmiel B., Ledwina T. 2020. Validation of association. Insurance: Mathematics and Economics 91: 55-
67. https://doi.org/10.1016/j.insmatheco.2019.12.003
Johnson, N. L. 1949. Systems of Frequency Curves Generated by Methods of translation.
Biometrika36: 149–176.
Ducharme J. 2020. TheWHOJustDeclaredCoronavirusCOVID-19a Pandemic. Time
2020;11March2020. Accessed at https://time.com/5791661/who-coronavirus-pandemic-
declaration/ on 28 November 2020.
Ferguson N.M., Laydon D., Nedjati-Gilani G., Imai N., Ainslie K., Baguelin M., Bhatia S., Boonyasiri
A., Cucunubá Z., Cuomo-Dannenburg G., Dighe A., Dorigatti I., Fu H., Gaythorpe K., Green W.,
Hamlet A., Hinsley W., Okell L.C., van Elsland S., Thompson H., Verity R., Volz E., Wang H.,
Wang Y., Walker P.G.T, Walters C., Winskill P., Whittaker C., Donnelly C.A., Riley S., Ghani
A.C. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality
and healthcare demand. Imperial College COVID-19 Response Team. Accessed at
https://doi.org/10.25561/77482 on 29 November 2020.
Foresti C.C.L. 2020. The real death toll for Covid-19 is at least 4 times the official numbers. Corriere
Della Serra. 26 March 2020. Accessed at www.corriere.it/politica/20_marzo_26/the-real-death-toll-
for-covid-19-is-at-least-4-times-the-official-numbers-b5af0edc-6eeb-11ea-925b-
a0c3cdbe1130.shtml?refresh_ce-cp. on 28 November 2020.
Hahn G.J., Shapiro S.S. 1967. Statistical Models in Engineering. Wiley series on systems engineering
and analysis, Wiley, ISSN 0084-019X, pp. 199-220.
Hasell J., Mathieu E., Beltekian D., Macdonald B., Giattino C., Ortiz-Ospina E., Roser M., Ritchie H.
2020. A cross-country database of COVID-19 testing. Scientific Data 7,
345.https://doi.org/10.1038/s41597-020-00688-8
Hellewell J., Abbott S., Gimma A., Bosse N.I., Jarvis C.I., Russell T.W., Munday J.D., Kucharski
A.J., Edmunds J., Sun F., Flasche S., Quilty B.J., Davies N., Liu Y., Clifford S., Klepac P., Jit M.,
Diamond C., Gibbs H., van Zandvoort K., Funk S., Eggo R.M. 2020. Feasibility of
controllingCOVID-19 outbreaks by isolation of cases and contacts. The Lancet Global Health,
8:e488-e496. https://doi.org/10.1016/S2214-109X(20)30074-7.
Hethcote H.W. The mathematics of infectious diseases. SIAM Review, 42: 599-653
Hill I.D., Hill R., Holder R.L. 1976. Algorithm AS 99: Fitting Johnson Curves by Moments. Applied
Statistics 25: 180-189. https://doi.org/10.2307/2346692
Holmdahl S.M., Buckee C. 2020. Wrong but Useful — What Covid-19 Epidemiologic Models Can
and Cannot Tell Us. The New England Journal of Medicine, 383:303-305.
DOI:10.1056/NEJMp2016822
IHME. 2020. Institute for Health Metrics and Evaluation COVID-19 Projections. Accessed at
https://covid19.healthdata.org on 29 November 2020.
Jankowiak Ł., Rozsa L., Tryjanowski P., Møller A.P. 2020. Strong negative covariation between
toxoplasmosis and CoVID-19 at a global scale: a spurious indirect effect? Scientific Reports 10:
12512 (2020). https://doi.org/10.1038/s41598-020-69351-x
Jewell N.P, Lewnard J.A., Jewell B.L. 2020. Caution Warranted: Using the Institute for Health
Metrics and Evaluation Model for Predicting the Course of the COVID-19. Annals of Internal
Medicine, 173: 226-227. https://doi.org/10.7326/M20-1565

Kissler S.M., Tedijanto C., Goldstein E., Grad Y.H., Lipsitch M. 2020. Projecting the transmission
dynamics of SARS-CoV-2 through the postpandemic period. Science *22* May 2020: Vol. 368, Issue
6493, pp. 860-868 DOI: 10.1126/science.abb5793
Kretschmer F., Yang W. 2020. Wuhan lockdown: China takes extreme measures to stop virus spread.
Deutsche Welle. 23 January 2020. Accessed at https://www.dw.com/en/wuhan-lockdown-china-
takes-extreme-measures-to-stop-virus-spread/a-52120126 on 29 November 2020.
Kuhl E. 2020. Data-driven modeling of COVID-19—Lessons learned. Extreme Mechanics
Letters40,100921. https://doi.org/10.1016/j.eml.2020.100921
LANL. 2020. Los Alamos National Laboratory COVID-19 Cases and Deaths Forecasts. Accessed at
https://covid-19.bsvgateway.org. on 28 November 2020.
Liu K., He M., Zhuang Z., He D., Li H. 2020. Unexpected positive correlation between human
development index and risk of infections and deaths of COVID-19 in Italy. One Health, 10:
100174. https://doi.org/10.1016/j.onehlt.2020.100174.
Long Q., Siu-fung L., Mudie L. Estimates Show Wuhan Death Toll Far Higher Than Official Figure.
Radio Free Asia. 27 March 2020. Accessed at www.rfa.org/english/news/china/wuhan-deaths-
03272020182846.html. on 28 November 2020.
Moré J.J. 1978. The Levenberg-Marquardt algorithm: Implementation and theory. In: Watson G.A.
(ed.) Numerical Analysis. Lecture Notes in Mathematics, 630. Springer, Berlin, Heidelberg.
Nishimoto Y., Inoue K. Curve-fitting approach for COVID-19 data and its physical background.
medRxiv 2020.07.02.20144899. doi: https://doi.org/10.1101/2020.07.02.20144899
Peirlinck M., Linka K., Costabal F.S., Bhattacharya j., Bendavid E., Ioannidis J.P.A., Kuhl E.
2020.Visualizing the invisible: The effect of asymptomatic transmission on the outbreak dynamics
of COVID-19. Computer Methods in Applied Mechanics and Engineering 372: 113410.
Skórka P., Grzywacz B., Moroń D., Lenda M. 2020. The macroecology of the COVID-19 pandemic in
the Anthropocene. PLoS ONE 15:e0236856.https://doi.org/10.1371/journal.pone.0236856
Stancati M., Sylvers E. 2020. Italy's Coronavirus Death Toll Is Far Higher Than Reported. The Wall
Street Journal. 1 April 2020. Accessed at www.wsj.com/articles/italys-coronavirus-death-toll-is-
far-higher-than-reported-11585767179. on 28 November 2020.
Tomie T. Understanding the present status and forecasting of COVID—19 in Wuhan. medRxiv
2020.02.14. https://doi.org/10.1101/2020.02.13.20022251
UGSDSC. 2020. University of Geneva and Swiss Data Science Center COVID-19 Daily Epidemic
Forecasting. https://renkulab.shinyapps.io/COVID-19-Epidemic-Forecasting
UK Guidance. Guidance for doctors completing Medical Certificates of Cause of Death in England
and Wales. HM Passport Office, Office for National Statistics. Accessed at
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/8
77302/guidance-for-doctors-completing-medical-certificates-of-cause-of-death-covid-19.pdf. on 28
November 2020.
WHO 2020. World Health Organization Coronavirus Disease (COVID-19) Dashboard. Accessed at
https://covid19.who.int/ on 4 December 2020.
466

468
Figure 1. Graphical presentation of the interpretation of the parameters obtained from Johnson
Cumulative Distribution Function fitting, describing the dynamics of the two infection waves observed
in Australia. P_{inf} – the total percentage of infections in a given infection wave, $Q_{2.5\%}$ – the day when
the infection wave started, $Q_{97.5\%}$ – the day when the infection wave ended, $Q_{50\%}$ – the day when half of
the total percentage of infections during a given wave was reached, M – the day when the infection
wave peaked, T – the duration of the wave, t_i – the duration of the wave increase, t_d – the duration of
the wave decrease, P_t – the total percentage of the population infected after two infection waves.
476

477
478 Fig. 2. Examples of fitted distributions in four scenarios of COVID-19 infection dynamics. A – one
479 ongoing infection wave (before the peak), B – one full wave, C – two waves and D – four overlapping
480 and interfering waves. Open circles – raw data, red lines – fitted Johnson Cumulative Distribution
481 Functions.
482

483
Fig. 3.A – Trajectory of four Johnson Cumulative Distribution Functions fitted to the four waves of
infections reported in Croatia, showing the areas where the waves overlapped and interfered. B – the
differences between Johnson (red line), Normal (green line) and Lognormal Cumulative Distribution
(blue line) Functions fitted to the raw data from Afghanistan (black dots).

Fig. 4. Heat maps showing the local association structure between variables at different quantile levels
obtained using the quantile dependence function *q*.